

Unrecognized Diabetes Among Hospitalized Patients

CLARESA S. LEVETAN, MD
MAUREEN PASSARO, MD
KATHLEEN JABLONSKI, PHD

MARY KASS, MD
ROBERT E. RATNER, MD

OBJECTIVE — To evaluate the hospital care rendered to hyperglycemic individuals who did not have a diagnosis of diabetes before admission.

RESEARCH DESIGN AND METHODS — A total of 1,034 consecutively hospitalized adult patients at a 750-bed inner-city teaching hospital were evaluated. Patients with one or more plasma glucose values >200 mg/dl were identified by the laboratory data system on a daily basis. Patients without a diagnosis of diabetes at the time of admission were evaluated to determine if and how physicians addressed the hyperglycemia, whether a new diagnosis of diabetes was made during admission, and whether follow-up was planned to address the hyperglycemia.

RESULTS — After excluding patients who were admitted for a primary diagnosis of diabetes, 37.5% of all hyperglycemic medical patients and 33% of hyperglycemic surgical patients were without a diagnosis of diabetes at the time of admission. These patients had a mean peak glucose of 299 mg/dl, and 66% had two or more elevated values during their hospitalization. Fifty-four percent received insulin therapy, and 59% received bedside glucose monitoring, yet 66% of daily patient progress notes failed to comment on the presence of hyperglycemia or diabetes. Diabetes was documented in only three patients (7.3%) as a possible diagnosis in the daily progress notes.

CONCLUSIONS — Despite marked hyperglycemia, most medical records made no reference to the possibility of unrecognized diabetes. Given the average delay of a decade between the onset and diagnosis of type 2 diabetes, further evaluation of hyperglycemic hospitalized patients may present an important opportunity for earlier detection and the initiation of therapy.

Half of the 16 million Americans with diabetes are undiagnosed (1–3). The diagnosis of diabetes is frequently not considered until another medical problem occurs and hyperglycemia is found incidentally. As a result, the diagnosis of type 2 diabetes is estimated to be delayed by an average of 10 years after the actual onset of the disease (4,5).

Despite the fact that more than 4 million Americans with a known diagnosis of diabetes are hospitalized annually in this country, little is known about the prevalence of hyperglycemia within the hospitalized population who do not have a diagnosis of diabetes before admission (6). We set forth to evaluate the hospital care rendered to these hyperglycemic individuals without a prior diagnosis of diabetes

and to determine whether such patients are appropriately evaluated and treated.

RESEARCH DESIGN AND METHODS

We prospectively evaluated all 1,034 consecutively hospitalized adult patients during a single week at an inner-city tertiary-care teaching hospital with a capacity of 750 beds. Patients with one or more plasma glucose values >200 mg/dl were identified by the laboratory data system on a daily basis. In addition to the glucose value, the following data were obtained: name, medical record number, medical service, patient age, and sex.

The medical records of these patients were evaluated after their discharge to determine if diabetes was a diagnosis that was present at the time of admission to the hos-

pital. Those patients in whom diabetes was the primary reason for their hospital admission were excluded from further analysis. Medical records of hyperglycemic patients without a diagnosis of diabetes at the time of admission were further reviewed.

A checklist was developed for the purposes of data collection. Frequency tables were run on demographic and outcomes data, and percentages were calculated based on the following information:

1. Presence of diabetes in the medical problem list or included in the initial history by the admitting physician.
2. Presence of more than one glucose value >200 mg/dl.
3. Value of the peak glucose.
4. Documentation in the daily progress notes of diabetes as a possible diagnosis.
5. Documentation of hyperglycemia in the daily progress notes.
6. Orders for medical therapy for the treatment of hyperglycemia.
7. Orders for bedside glucose monitoring.
8. Documentation of hyperglycemia in the discharge summary.
9. Documentation of follow-up plans for further diabetes work-up or therapy in the final progress note or the discharge summary.

RESULTS — Of the 1,034 hospitalized adult patients, 130 (12.6%) had one or more documented plasma glucose values >200 mg/dl. Fifteen patients (11.5%) had diabetes documented as the principal reason for hospital admission and were excluded from further study; the other 115 with hyperglycemia served as the study population for further analysis. Those with preexisting diabetes plus those recognized as having new-onset diabetes at the time of admission constituted 64% (74/115) of the hyperglycemic population (Fig. 1).

Table 1 describes the distribution of hyperglycemic patients by medical service. Thirty-six percent (41/115) of patients were previously unrecognized as having a disorder of glucose metabolism. Of the 41 patients with hyperglycemia in whom there was no documentation of diabetes existing before hospital admission, 23 (56.1%) were men and 18 (43.9%) were women (Table 2). Of

From the Medlantic Research Institute (C.S.L., M.P., K.J., R.E.R.) and the Washington Hospital Center (M.K.), Washington, DC.

Address correspondence and reprint requests to Dr. Claresa Levetan, Medlantic Research Institute, 650 Pennsylvania Ave., SE, Suite 50, Washington, DC 20003.

Received for publication 21 May 1997 and accepted in revised form 21 October 1997.




Figure 1—Schematic of hospitalized patients evaluated for hyperglycemia.

the 41 patients, 27 (65.9%) had two or more documented glucose values >200 mg/dl with a mean peak glucose of 299 mg/dl with a range of 202–503 mg/dl. The mean age among these patients was 61 years.

There was no physician documentation that hyperglycemia or diabetes was present in 27 (65.9%) of the 41 records (Fig. 2). Eleven records (26.8%) did not use the word diabetes, but described “hyperglycemia” or an “increased finger stick glucose” value. Only three records (7.3%) mentioned diabetes as a diagnostic possibility. Other than the three records in which diabetes was considered, none of the daily progress notes or discharge summaries discussed plans for the further evaluation or specific management of hyperglycemia after hospital discharge.

Although 66% of the medical records did not mention diabetes or hyperglycemia, 22 (54%) of the 41 patients received both bedside glucose monitoring and regular

insulin coverage for their elevated glucose values, of which two patients (5%) were subsequently placed on definitive diabetes therapy during admission (Table 3). Two patients (4.9%) received bedside glucose monitoring but no insulin coverage, and seventeen patients (41.5%) had no medical orders reflecting the recognition or treatment of hyperglycemia.

CONCLUSIONS— We found that one-third of all hyperglycemic patients on both the medical and surgical services had no prior history of diabetes. Sixty-six percent of these hyperglycemic individuals had two or more elevated glucose values. Despite therapeutic intervention with insulin in more than half of our patients, there was almost never documentation regarding the hyperglycemia and rarely was diabetes a diagnostic consideration.

We believe that physicians assumed that the hyperglycemia was a transient finding that resulted from the stress of acute illness rather than considering the diagnosis of unrecognized diabetes. Given the profound delay in the diagnosis of type 2 diabetes nationwide, we believe that it is a common occurrence for physicians to overlook hyperglycemia, and our findings are, therefore, not unique to this institution or geographic region (4,5).

There are no unique diagnostic criteria or recommendations for making a definitive diagnosis of diabetes in the stressed state. All of the patients identified had glucose values >200 mg/dl and met the laboratory criteria for diabetes established by the National Diabetes Data Group, the World Health Organization, and the American Diabetes Association (7–11).

Despite the well-defined pathophysiology of intercurrent illness and surgery on serum glucose, stress hyperglycemia has eluded consistent documentation (12–27). Although impaired carbohydrate metabolism resulting in hyperglycemia is seen in patients both with and without diabetes, studies have also demonstrated that stress can also result in diminished glucose values, and patients not given sufficient exogenous glucose can develop hypoglycemia and ketosis in times of stress (9–27). Stress hyperglycemia should theoretically result in higher glycemic excursions in diabetic individuals than in nondiabetic individuals

Table 1—The distribution of hyperglycemic patients by medical service

| Service | Glucose >200 mg/dl | Diabetes recognized before or at admission | Diabetes unrecognized at admission | Unrecognized at admission (%) |
|------------|----------------------|--|------------------------------------|-------------------------------|
| n | 115 | 74 | 41 | — |
| Medicine | 66 | 40 | 25 | 37.5 |
| Surgery | 48 | 32 | 16 | 33 |
| Gynecology | 2 | 1 | 1 | 50 |
| Podiatry | 1 | 1 | 0 | 0 |

Data are n or %.

Table 2—Characteristics of hyperglycemic patients without a prior diagnosis of diabetes

| | |
|---|-----------------|
| Sex (%) (M/F) | 56.1/43.9 |
| Age (years) | 61 ± 18 (27–92) |
| Mean peak glucose (mg/dl) | 299 (202–503) |
| Patients with two or more glucose values >200 mg/dl (%) | 65.9 |

because of a relative or absolute deficiency of endogenous insulin.

Husband et al. (28) followed a group of hospitalized patients with acute myocardial infarction with newly recognized hyperglycemia. Two months after hospital discharge, patients were evaluated with glucose tolerance testing with findings that an admission glucose value of ≥ 180 mg/dl predicted undiagnosed diabetes rather than stress hyperglycemia (28). Although stress can account for hyperglycemia, our patient population had glucose values in a range that warrants the consideration for a diagnosis of diabetes rather than the assumption of stress hyperglycemia.

The Diabetes Insulin-Glucose in Acute Myocardial Infarction (DIGAMI) series underscores the importance of aggressive early intervention in the management of hyperglycemic patients who have been admitted with suspected myocardial infarction, including those without a previous history of diabetes. Significantly lower mor-

tality rates were seen during the 3-year follow-up period for patients in whom intravenous insulin, followed by multiple daily injections, was initiated in the hospital and continued as an outpatient (29–31).

Although dozens of medications have been described as inducing diabetes, there has been little documented about the glycemic status of patients before the initiation of the offending drugs. Corticosteroids have been evaluated as a means of unmasking impaired glucose tolerance, but among normal controls, only 3% had a positive glucose tolerance test when pretreated with corticosteroids (32). Studies of corticosteroid-treated individuals have found that <20% of steroid-treated individuals develop diabetes (33,34).

It is critical that diabetes be diagnosed as early as possible, since diabetic complications frequently occur before the diagnosis of type 2 diabetes. More than 20% of the individuals with diabetes have retinopathy at the time of diagnosis, and it is often their retinopathy eye findings that lead to the subsequent diagnosis of diabetes (4,5). The prevalence of unrecognized diabetes remains extremely high. By 65 years of age, 18.7% of the population has diabetes, yet half remain undiagnosed (35). Regardless of the etiology, persistent hyperglycemia leads to the acceleration of both microvascular and macrovascular disease (36).

In the U.S., expenditures resulting from the morbidity of patients with diabetes exceed \$100 billion per year and account for 15% of the entire health care

Table 3—Therapeutic intervention for patients with hyperglycemia and without a prior history of diabetes

| Intervention | Percentage |
|---|------------|
| Received insulin treatment and bedside glucose monitoring | 53.6 |
| Received bedside monitoring without insulin treatment | 4.9 |
| No medical orders for hyperglycemia | 41.5 |

budget (37). Delays in the diagnosis of diabetes carry with it substantial financial and health ramifications. Further study is needed to define differences between the hospitalized patient population with transient stress-induced hyperglycemia and those with unrecognized diabetes.

We believe that the failure to consider the possibility of diabetes represents a missed window of opportunity for making an earlier diagnosis and for initiating interventions that may delay the devastating complications of this disease. We recommend that physicians assume that hyperglycemia is diabetes until they prove otherwise.

References

1. U.S. Department of Health and Human Services: *Diabetes in the United States: A Strategy for Prevention 1993*. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Division of Diabetes Translation, 1993
2. Harris MI, Hadden WC, Knowler WC, Bennett PH: Prevalence of diabetes and impaired glucose tolerance and plasma glucose levels in the U.S. population aged 20–74 yr. *Diabetes* 36:523–534, 1987
3. Harris MI: Undiagnosed NIDDM: clinical and public health issues. *Diabetes Care* 16:642–652, 1993
4. Harris MI: Classification, diagnostic criteria, and screening for diabetes. In *Diabetes in America*. 2nd ed. Harris MI, Cowie CC, Stern MP, Boyko EJ, Reiber GE, Bennett PH, Eds. Washington, DC, U.S. Govt. Printing Office, 1995, p. 15–35 (NIH publ. no. 95-1468)
5. Harris MI, Klein R, Welborn TA, Knuiaman MW: Onset of NIDDM occurs at least 4 to 7 years before clinical diagnosis. *Diabetes Care* 15:815–819, 1992
6. U.S. National Center for Health Statistics: *1995 National Hospital Discharge Survey (Public Use Data Tapes)*. U.S. Department




Figure 2—Documentation of diabetes or hyperglycemia in the physicians' daily progress notes.

- of Health and Human Services, Washington, DC, 1995
7. Lustman PJ, Gavard JA: Psychosocial aspects of diabetes adult populations. In *Diabetes in America*. 2nd ed. Harris MI, Cowie CC, Stern MP, Boyko EJ, Reiber GE, Bennett PH, Eds. Washington, DC, U.S. Govt. Printing Office, 1995, p. 507-517 (NIH publ. no. 95-1468)
 8. World Health Organization: *Diabetes Mellitus: Report of a WHO Study Group*. Geneva, World Health Org., 1985, p. 1-113 (Tech. Rep. Ser., no. 727)
 9. World Health Organization: *Report of the Expert Committee on Diabetes*. Geneva, World Health Org., 1985 (Tech. Rep. Ser., no. 646)
 10. National Diabetes Data Group: Classification and diagnosis of diabetes mellitus and other categories of glucose intolerance. *Diabetes* 28:1039-1057, 1979
 11. The Expert Committee on the Diagnosis and Classification of Diabetes Mellitus: Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care* 20:1183-1197, 1997
 12. Gavin LA: Management of diabetes mellitus during surgery. *West J Med* 151:525-529, 1989
 13. Gavin LA: Perioperative management of the diabetic patient. *Endocrinol Metab Clin North Am* 21:457-475, 1992
 14. Hirsch IB, McGill JB: Role of insulin management of surgical patients with diabetes mellitus. *Diabetes Care* 13:980-991, 1990
 15. Hirsch IB, Paauw DS, Brunzell J: Inpatient management of adults with diabetes. *Diabetes Care* 18:870-878, 1995
 16. Gonder-Frederick LA, Carter WR, Cox DJ, Clarke WL: Environmental stress and blood glucose change in insulin-dependent diabetes mellitus. *Health Psychol* 9:503-515, 1990
 17. Hanson CL, Henggeler SW, Burghen GA: Social competence and parental support as mediators of the link between stress and metabolic control in adolescents with insulin-dependent diabetes mellitus. *J Consult Clin Psychol* 55:529-533, 1987
 18. Chase HP, Jackson GG: Stress and sugar control in children with insulin-dependent diabetes mellitus. *J Pediatr* 98:1011-1013, 1981
 19. Hanson CL, Pichert JW: Perceived stress and diabetes control in adolescents. *Health Psychol* 5:439-452, 1986
 20. Vandenberg RL, Sussman KE, Titus CC: Effects of hypnotically induced acute emotional stress on carbohydrate-lipid metabolism in patients with diabetes mellitus. *Psychosom Med* 28:382-390, 1966
 21. Vandenberg RL, Sussman KE, Vaughan GD: Effects of combined physical-anticipatory stress on carbohydrate-lipid metabolism in patients with diabetes mellitus. *Psychosomatics* 8:16-19, 1967
 22. Greydanus DE, Hofmann AD: Psychological factors in diabetes mellitus: a review of the literature with emphasis on adolescence. *Am J Dis Child* 133:1061-1088, 1979
 23. Delamater AM, Bubb J, Kurtz SM, Kuntze J, Smith JA, White NH, Santiago JV: Physiologic responses to acute psychological stress in adolescents with type I diabetes mellitus. *J Pediatr Psychol* 13:69-86, 1988
 24. Gilbert BO, Johnson SB, Silverstein J, Malone J: Psychological and physiological responses to acute laboratory stressors in insulin-dependent diabetes mellitus adolescents and non-diabetic controls. *J Pediatr Psychol* 14:577-591, 1989
 25. Kemmer FW, Bisping R, Steingruber HJ, Baar H, Hardtmann F, Schlaghecke R, Berger M: Psychological stress and metabolic control in patients with type I diabetes mellitus. *N Engl J Med* 314:1078-1084, 1986
 26. Goetsch VL, Wiebe DJ, Vetum LG, Van Dorsten B: Stress and blood glucose in type II diabetes mellitus. *Behav Res Ther* 28:531-537, 1990
 27. Fletcher J, Langman MJS, Kellock RK: Effect of surgery on blood-sugar levels in diabetes mellitus. *Lancet* ii:52-55, 1965
 28. Husband DJ, Alberti KG, Julian DG: Stress hyperglycaemia during acute myocardial infarction: an indicator of pre-existing diabetes? *Lancet* ii:179-181, 1983
 29. Malmberg K, Ryden I, Iamsten A, Herlitz J, Walatdenstrom A, Wedel H: Effect of insulin treatment on cause-specific one-year mortality and morbidity in diabetic patients with acute myocardial infarction: DIGAMI Study Group. *Diabetes Insulin-Glucose in Acute Myocardial Infarction. Eur Heart J* 17:1337-1344, 1996
 30. Malmberg K for the DIGAMI Study Group: Prospective randomized study of intensive insulin treatment on long-term survival after acute myocardial infarction in patients with diabetes mellitus. *BMJ* 314:1512-1515, 1997
 31. Malmberg K, Ryden L, Efendic S, Herlitz J, Nicol P, Waldenstrom A, Wedel H, Welin L: Randomized trial of insulin-glucose infusion followed by subcutaneous insulin treatment in diabetic patients with acute myocardial infarction (DIGAMI Study): effects on mortality at 1 year. *J Am Coll Cardiol* 26:57-64, 1995
 32. Fajans SS, Conn JW: An approach to the prediction of diabetes mellitus by modification of the glucose tolerance test with cortisone. *Diabetes* 3:296-304, 1954
 33. Weisz A, Ratner R: Factors in renal transplant-induced diabetes mellitus (Abstract). *Diabetes* 35:72A, 1986
 34. Landy HJ, Isada NB, McGinnis J, Ratner R, Grossman JH: The effect of chronic steroid therapy on glucose tolerance in pregnancy. *Am J Obstet Gynecol* 159:612-615, 1988
 35. Harris MI, Hadden WC, Knowler WC, Bennett PH: Prevalence of diabetes and impaired glucose tolerance and plasma glucose levels in the U.S. population aged 20-74 yr. *Diabetes* 36:523-534, 1987
 36. Haffner SM, Stern MP, Hazuda HP, Mitchell BD, Patterson JK: Cardiovascular risk factors in confirmed prediabetic individuals: does the clock for coronary heart disease start ticking before the onset of clinical diabetes? *JAMA* 263:2892-2898, 1990
 37. Rubin RJ, Altman WM, Mendelson DN: Health care expenditures for people with diabetes mellitus, 1992. *J Clin Endocrinol Metab* 78:809A-809F, 1994