

Rates of river incision and scarp retreat in eastern and central Grand Canyon over the past half million years: Evidence for passage of a transient knickzone: COMMENT

Ryan Crow¹, Karl Karlstrom², Laura Crossey², Richard Young³, Michael H. Ort⁴, Yemane Asmerom², Victor Polyak², and Andrew Darling⁵

¹U.S. Geological Survey, 2255 N. Gemini Drive, Flagstaff, Arizona 86001, USA

²Department of Earth & Planetary Sciences, Northrop Hall, University of New Mexico, Albuquerque, New Mexico 87131, USA

³Department of Geological Sciences, 1 College Circle, State University of New York, Geneseo, New York 14454, USA

⁴School of Earth Sciences and Environmental Sustainability, Box 4099, Northern Arizona University, Flagstaff, Arizona 86011, USA

⁵School of Earth and Space Exploration, Arizona State University, Tempe, Arizona 85287, USA

Abbott et al. (2015) and Crow et al. (2014), two Grand Canyon incision studies, come to very different conclusions despite apparent methodological similarities. Crow et al. (2014) used U-Th, Ar-Ar, and cosmogenic burial dating of material associated with Colorado River (CR) strath terrace sequences at 6 sites throughout Grand Canyon and concluded that average bedrock incision rates have been temporally steady at each site over at least the last 650 ka, but vary spatially from 100 to 160 m/Ma due to differential mantle-driven uplift. Abbott et al. (2015) used our unpublished Ar-Ar dating on a dike in western Grand Canyon, near river mile 159, plus U-Th dating of travertine-cement in sidestream alluvium, near Hermit Rapid, to suggest that incision rates were 1–4 km/Ma from 500 to 400 ka and <200 m/Ma after 400 ka, a difference they ascribe to a migrating knickpoint. The conclusions of these studies are contradictory.

The simplest criticism of the Abbott et al. (2015) interpretation is that neither of their sites contains unequivocal CR deposits. We visited both sites repeatedly but rejected them as CR incision points because datable materials could not be unambiguously related to the paleoposition of the CR. The ca. 520 ka, 159-mile dikes intrude bedrock, along en echelon joints, on both sides of the river to near a height of 400–450 m, where welded pyroclastic tuff is found (Wenrich et al., 1997). The paleo-position of the CR at the time of dike propagation is unknown because the dikes could have intruded into an extant canyon or one filled with lacustrine deposits. Abbott et al.'s argument that the canyon would act as a “stress concentrator” is complicated by joint control that created preexisting anisotropy for magma movement. Numerous modern examples show that dikes can propagate across significant relief without concentrating eruptions at the lowest point (or in some cases, even breaking out there). For example, during the 1973 Mauna Ulu eruption, en echelon dikes crossed the 100-m-deep Pauahi Crater with significant fissure eruptions on the flanks and rims of the crater but not at its bottom (Tilling, 1987). Thus, we reject the 159-mile dikes as a robust incision constraint for the CR.

At the Hermit site, travertines cemented a ~70-m-thick aggradational sequence of sidestream and colluvial deposits on the Tonto platform, interpreted

by Abbott et al. (2015) as a tributary fan that was at the same elevation as the CR. However, the lack of verified CR gravels suggests the CR could have been much lower. Additionally, the age of the deposits is uncertain, as only 30% of the analyzed samples gave U-Th ages. The rest of the samples show evidence for open-system behavior, are plausibly outside of U-Th dating range (>600–700 ka), and would give ²³⁴U model ages of up to ca. 1.6 Ma. Their successful ca. 500 ka U-Th ages come from the base of deposits and are plausibly younger infillings, similar to well-documented examples throughout Grand Canyon (Crow et al., 2014). Such infillings are expected at a deposit's base where carbonic groundwater can migrate through permeable hill-slope deposits. Careful dating of both travertine clasts and infillings (Crow et al., 2014) is needed to bracket the age of gravel deposition.

We suggest two possible models that could explain all existing data. 1) The ~500 ka U-Th ages reported by Abbott et al. (2015) may be secondary travertine infillings, and gravel deposition took place at ca. 2 Ma. 2) The deposits at the Hermit site are due to sidestream aggradation with local base level controlled by the Tapeats sandstone lithologic bench or a lake behind a ca. 500 ka downstream lava dams (Crow et al., 2015). As an analog for the lake possibility, the Spencer Canyon sidestream debris fan has aggraded ~30 m since the 1930s due to construction of Hoover Dam. Like the Hermit location, it is dominated by sidestream alluvium, it largely lacks CR gravel or appreciable lake deposits except directly at its confluence with the CR/Lake Mead, respectively, and its elevation is well above the bedrock channel.

Figure 1 shows CR paleoprofiles with a transient knickpoint as suggested by Abbott et al. (2015) as well as our CR gravel-constrained incision rates (Crow et al., 2014; Karlstrom et al., 2007). Most of our data would have to be wrong for their model to work. In contrast, the alternate models of their data presented above honor the abundant CR gravel-constrained incision rates, postulate subparallel CR paleoprofiles (Fig. 1), and do not require 1000–4000 m/Ma incision rates that are 2–8 times greater than the highest knickpoint-controlled rates known for bedrock incision in the southwestern U.S. (Donahue et al., 2013).




Figure 1. Longitudinal Colorado River profile showing paleo profiles implied by Abbott et al. (2015) and those suggested by Karlstrom et al. (2007) and Crow et al. (2014). Our preferred profiles honor incision points constrained by unambiguous mainstream-river-gravel and do not support a ~400 m high knickpoint, which we consider unreasonable. Data from Crow et al. (2014), Karlstrom et al. (2007), and Abbott et al. (2015). Numbers next to incision points are mean ages in ka.

REFERENCES CITED

Abbott, L.D., Lundstrom, C., and Traub, C., 2015, Rates of river incision and scarp retreat in eastern and central Grand Canyon over the past half million years: Evidence for passage of a transient knickzone: *Geosphere*, v. 11, p. 638–659, doi:10.1130/GES00978.1.

Crow, R.S., Karlstrom, K.E., Darling, A., Crossey, L.J., Polyak, V.J., Granger, D., Asmerom, Y., and Schmandt, B., 2014, Steady incision of Grand Canyon at the million year timeframe: A case for mantle-driven differential uplift: *Earth and Planetary Science Letters*, v. 397, p. 159–173, doi:10.1016/j.epsl.2014.04.020.

Crow, R.S., Karlstrom, K.E., McIntosh, W.C., Peters, L., Crossey, L.J., and Eyster, A., 2015, A new model for Quaternary lava dams in Grand Canyon based on ⁴⁰Ar/³⁹Ar dating, basalt geochemistry, and field mapping: *Geosphere*, v. 11, p. 1305–1342, doi:10.1130/GES01128.1.

Donahue, M.S., Karlstrom, K.E., Aslan, A., Darling, A., Granger, D., Wan, E., Dickinson, R.G., and Kirby, E., 2013, Incision history of the Black Canyon of Gunnison, Colorado, over the past

~1 Ma inferred from dating of fluvial gravel deposits: *Geosphere*, v. 9, p. 815–826, doi: 10.1130/GES00847.1.

Karlstrom, K.E., Crow, R.S., Peters, L., McIntosh, W.C., Raucci, J., Crossey, L.J., Umhoefer, P., and Dunbar, N., 2007, ⁴⁰Ar/³⁹Ar and field studies of Quaternary basalts in Grand Canyon and model for carving Grand Canyon: Quantifying the interaction of river incision and normal faulting across the western edge of the Colorado Plateau: *Geological Society of America Bulletin*, v. 119, p. 1283–1312, doi:10.1130/0016-7606(2007)119[1283:AAFSSQ]2.0.CO;2.

Tilling, R.I., 1987, Fluctuations in surface height of active lava lakes during 1972–1974 Mauna Ulu Eruption, Kilauea Volcano, Hawaii: *Journal of Geophysical Research*, v. 92, p. 13721, doi:10.1029/JB092iB13p13721.

Wenrich, K.J., Billingsley, G., and Huntoon, P.W., 1997, Breccia pipe and geologic map of the north-eastern Hualapai Indian Reservation and vicinity, Arizona: U.S. Geologic Survey, Miscellaneous Investigations Series Map I-2440, scale 1:48,000.