

Are Circulating Immune Cells a Determinant of Pancreatic Cancer Risk? A Prospective Study Using Epigenetic Cell Count Measures



Verena A. Katzke¹, Charlotte Le Cornet¹, Rayaan Mahfouz², Bianca Brauer², Theron Johnson¹, Federico Canzian³, Vinciane Rebours^{4,5}, Marie-Christine Boutron-Ruault⁶, Gianluca Severi^{6,7}, Matthias B. Schulze^{8,9}, Anja Olsen^{10,11}, Anne Tjønneland^{10,12}, Kim Overvad¹¹, Marta Crous-Bou^{13,14}, Esther Molina-Montes^{15,16,17,18}, Pilar Amiano^{17,19}, José María Huerta^{17,20}, Eva Ardanaz^{17,21,22}, Aurora Perez-Cornago²³, Giovanna Masala²⁴, Valeria Pala²⁵, Rosario Tumino²⁶, Carlotta Sacerdote²⁷, Salvatore Panico²⁸, Bas Bueno-de-Mesquita²⁹, Roel Vermeulen^{30,31,32}, Malin Sund³³, Oskar Franklin³³, Sofia Christakoudi^{32,34,35}, Laure Dossus³⁶, Elisabete Weiderpass³⁷, Sven Olek², and Rudolf Kaaks^{1,38}

ABSTRACT

Background: Evidence is accumulating that immune cells play a prominent role in pancreatic cancer etiology but prospective investigations are missing.

Methods: We conducted a nested case-control study within the European Prospective Investigation into Cancer and Nutrition (EPIC) study with 502 pairs of incident pancreatic cancer cases and matched controls. Relative counts of circulating immune cells (neutrophils and lymphocyte sublineages: total CD3⁺, CD8⁺, CD4⁺, and FOXP3⁺ regulatory T cells (Tregs) relative to nucleated cells, (white blood cells) were measured by qRT-PCR. ORs with 95% confidence intervals were estimated using logistic regressions, modeling relative counts of immune cells on a continuous scale.

Results: Neither relative counts of immune cell types taken individually, nor mutually adjusted for each other were associated

with pancreatic cancer risks. However, in subgroup analyses by strata of lag-time, higher relative counts of Tregs and lower relative counts of CD8⁺ were significantly associated with an increased pancreatic cancer risks in participants diagnosed within the first 5 years of follow-up.

Conclusions: These results might reflect reverse causation, due to higher relative counts of Tregs and lower counts of CD8⁺ cells among individuals with more advanced stages of latent pancreatic cancer, who are closer to the point of developing clinical manifest disease.

Impact: We have shown, for the first time, that increased relative counts of regulatory T cells and lower relative counts of CD8⁺, cytotoxic T cells may be associated with pancreatic cancer risk or relatively late-stage tumor development.

See related commentary by Michaud and Kelsey, p. 2176

¹Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany. ²Ivana Türbachova Laboratory for Epigenetics, Epiontis GmbH, Berlin, Germany, Precision for Medicine Group. ³Genomic Epidemiology Group, German Cancer Research Center (DKFZ), Heidelberg, Germany. ⁴Pancreatology Department, Beaujon Hospital, AP-HP, Clichy, France. ⁵Inserm UMR1149, DHU Unit, Paris-Diderot University, Paris, France. ⁶Paris-Saclay University, UVSQ, Inserm, Gustave Roussy, “Exposome and Heredity” team, CESP, F-94805, Villejuif, France. ⁷Department of Statistics, Computer Science and Applications “G. Parenti,” University of Florence, Florence, Italy. ⁸Department of Molecular Epidemiology, German Institute of Human Nutrition Potsdam-Rehbruecke, Nuthetal, Germany. ⁹Institute of Nutritional Science, University of Potsdam, Nuthetal, Germany. ¹⁰Danish Cancer Society Research Center, Copenhagen, Denmark. ¹¹Department of Public Health, University of Århus, Århus, Denmark. ¹²Department of Public Health, University of Copenhagen, Copenhagen, Denmark. ¹³Unit of Nutrition and Cancer, Cancer Epidemiology Research Program, Catalan Institute of Oncology (ICO) - Bellvitge Biomedical Research Institute (IDIBELL), L’Hospitalet de Llobregat, Barcelona 08908, Spain. ¹⁴Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts. ¹⁵Department of Nutrition and Food Sciences, Faculty of Pharmacy, University of Granada, Granada, Spain. ¹⁶Instituto de Investigación Biosanitaria (ibs.GRANADA), Granada, Spain. ¹⁷CIBERESP, Madrid, Spain. ¹⁸Institute of Nutrition and Food Technology “José Mataix,” Center of Biomedical Research, University of Granada, Granada, Spain. ¹⁹Public Health Division of Gipuzkoa, BioDonostia Research Institute, Donostia-San Sebastian, Spain. ²⁰Department of Epidemiology, Murcia Regional Health Council, IMIB-Arixaca, Murcia, Spain. ²¹Navarra Public Health Institute, Pamplona, Spain. ²²IdiSNA, Navarra Institute for Health Research, Pamplona, Spain. ²³Cancer Epidemiology Unit, Nuffield Department of Population Health, University of Oxford, Oxford UK. ²⁴Cancer Risk Factors and Life-Style Epidemiology Unit, Institute for Cancer Research, Prevention and Clinical Network - ISPRO, Florence, Italy. ²⁵Epidemiology and Prevention Unit, Fondazione IRCCS Istituto Nazionale dei

Tumori di Milano, Milano, Italy. ²⁶Cancer Registry and Histopathology Department, Provincial Health Authority (ASP 7) Ragusa, Italy. ²⁷Unit of Cancer Epidemiology, Città della Salute e della Scienza University-Hospital, Turin, Italy. ²⁸Dipartimento di Medicina Clinica e Chirurgia, Federico II University, Naples, Italy. ²⁹Former senior scientist, Dept. for Determinants of Chronic Diseases (DCD), National Institute for Public Health and the Environment (RIVM), PO Box 1, 3720 BA Bilthoven, the Netherlands. ³⁰Institute for Risk Assessment Sciences, Utrecht University, Utrecht, the Netherlands. ³¹Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht, the Netherlands. ³²School of Public Health, Imperial College London, London, UK. ³³Department of Surgical and Perioperative Sciences, Umeå University, Sweden. ³⁴Department of Epidemiology and Biostatistics, Imperial College London, Norfolk Place, St Mary’s Campus, London, United Kingdom. ³⁵MRC Centre for Transplantation, King’s College London, Great Maze Pond, London, United Kingdom. ³⁶International Agency for Research on Cancer, World Health Organization, Lyon, France. ³⁷Director, International Agency for Research on Cancer, World Health Organization, Lyon, France. ³⁸Translational Lung Research Center (TLRC) Heidelberg, Member of the German Center for Lung Research (DZL), Heidelberg, Germany.

Note: Supplementary data for this article are available at Cancer Epidemiology, Biomarkers & Prevention Online (<http://cebp.aacrjournals.org/>).

V.A. Katzke and C. Le Cornet contributed equally to this article.

Corresponding Author: Verena A. Katzke, Division of Cancer Epidemiology, German Cancer Research Center, Heidelberg 69120, Germany. E-mail: v.katzke@dkfz.de

Cancer Epidemiol Biomarkers Prev 2021;30:2179-77

doi: 10.1158/1055-9965.EPI-21-0169

©2021 American Association for Cancer Research

Introduction

Animal experiments and clinical studies have shown that innate and adaptive immune responses (immuno-surveillance) play a key role in pancreatic cancer development (1–3). The formation of pancreatic precursor lesions and the further development into an invasive tumor is accompanied by progressive infiltration of various types of immune cells (4). In patients with pancreatic cancer, the total or relative counts of various types of immune cells in tumor tissue (5, 6), but also in peripheral blood (7, 8), have been found to correlate with clinical outcomes. In general, higher total or relative counts of CD8⁺ cytotoxic T cells, CD4⁺ T-helper 1 cells, and natural killer cells have been associated with more favorable patient outcomes, whereas higher counts of CD4⁺ T-helper 2 cells and of FOXP3⁺ regulatory T cells have been associated with greater immunosuppression, accelerated cancer development and worse prognosis (9). In addition, neutrophil counts have often been found to be higher among patients with cancer, including patients with pancreatic cancer (10), as compared with cancer-free control subjects, and elevated pretreatment ratios of circulating neutrophil-to-lymphocyte (NLR) have been associated with lower overall patient survival (11, 12). However, although substantial evidence now documents the prognostic significance of immune defense among patients, so far no studies have examined whether precisely quantified immune cell homeostasis in initially healthy individuals also determines future pancreatic cancer risk. The major reason for this latter deficit is that blood samples stored in large-scale population cohort studies usually do not contain intact blood cells, prohibiting flow cytometry analyses of immune cell composition.

Recently, we have developed and validated methods that allow the determination of immune cell composition relative to total nucleated cells by qPCR of epigenetic markers specific for different immune cell lineages (13). This type of assay can be applied to DNA extracted from frozen, nonintact white blood cells as well as other tissues. By applying this method to stored buffy coat samples of the European Prospective Investigation into Cancer and Nutrition (EPIC)-Heidelberg cohort, we found that higher counts of FOXP3⁺ (regulatory) T-cell relative to total nucleated cells were associated with higher risks of cancers of the lung, colorectum and breast, whereas lower relative CD8⁺ (cytotoxic) T cells counts were inversely associated with lung and breast cancer risks (14). In contrast, this first study showed no associations of cancer risk with relative counts of monocytes, B cells, or natural killer cells, and only marginally suggestive associations (nonsignificant) for neutrophils.

To explore whether similar prospective relationships exist between counts of circulating immune cells relative to total nucleated cells and risk of pancreatic cancer, overall or depending on the prospective lag-time between blood sampling (time of immune cell counts) and cancer diagnosis, we performed a case-control study nested within the Europe-wide EPIC cohort (15), with a focus on estimated relative counts for neutrophils and lymphocyte T-cell sublineages.

Materials and Methods

Study population

The current investigation was based on a case-control study nested within the European EPIC cohort—a multicenter prospective study in Europe described in detail previously (15). Briefly, 519,978 healthy men and women, mostly aged between 35 and 70 years, were recruited by 23 collaborating centers in 10 European countries (Denmark, France, Germany, Greece, Italy, the Netherlands, Norway, Spain, Sweden, and the United Kingdom) between 1992 and 2000. At

baseline, participants filled in comprehensive questionnaires on lifestyle, nutrition, and diseases and their height, weight, and body circumferences (waist and hip) were measured. In addition, study participants provided blood samples, which were processed into serum, plasma, and red blood cells, as well as a buffy coat containing all nucleated cells. Processed blood cell fractions were frozen and stored centrally at the International Agency for Research on Cancer (IARC), as well as locally in each recruitment center. The EPIC centers in Malmö (Sweden) and Greece did not contribute to this study.

Incident cancer cases were identified by population cancer registries (Denmark, Italy, the Netherlands, Spain, Sweden, and the United Kingdom) or by a combination of methods including linkage with cancer and pathology registries and systemic verification of active follow-up (France, Germany). In all EPIC centers, data on vital status were collected through population registries, in combination with national health insurance data (France).

Nested case-control design

The present project is based on incident pancreatic cancer cases diagnosed between 1993 and 2008.

Pancreatic cancer incidence data were coded according to ICD-10 and included all invasive exocrine pancreatic cancers that were coded as C25 (25.0–25.3, 25.7–25.9). Exclusion criteria were the occurrence of other malignant tumors preceding the diagnosis of pancreatic cancer, except for nonmelanoma skin cancer, and nonavailability of blood specimens. Seventy-three percent were adenocarcinomas whereas the other 27% were not otherwise specified. Of these 145 unspecified cases, 11% were of advanced and 74% of unknown stage. Microscopically confirmed were 391 cases (78%), whereas the diagnosis of the remaining 22% was based on clinical observations, imaging, or autopsy.

For each case subject, one control subject was selected using an incidence density sampling protocol. Case and control subjects were matched on study recruitment center, sex, age at blood collection (± 6 months), date of blood collection (± 2 month), time of blood collection (± 2 hours), length of follow-up, and time between blood sampling and time of last consumption of any foods or drinks (< 3 , 3 – 6 , ≥ 6 hours). We included initially 530 incident pancreatic cancer cases and 533 controls, building 502 matched case-control pairs and 59 single subjects (31 controls, 28 cases) with no matching partner due to depleted blood samples in the biorepositories.

Laboratory methods: quantification of immune cells

Relative counts of neutrophils and lymphocyte subpopulations were measured by quantitative epigenetic real-time PCR at EPIONTIS GmbH (13).

DNA was extracted at the IARC, from frozen pellets of nucleated blood cells (buffy coats) using the Autopure LS DNA preparation platform (Autogen; ref. 16). Relative counts of immune cells in genomic DNA extracted from buffy coats were assessed by qRT-PCR assisted cell counting (qPACC). Descriptions of the laboratory methods are given on the Epiontis web site (<http://www.epiontis.com>) have been published previously (13, 14, 17–19), and are detailed in the Supplementary Methods and Materials M1 to M4. In brief, for each subject, 1.5 μ g of genomic DNA bisulfite treated prior to qPCR analysis. Relative counts of neutrophils, and lymphocyte sublineages (Tregs, CD4⁺, and CD8⁺ T cells) were assessed by qPACC. The assays are based on the measurement of genetic loci that have been shown to be specifically unmethylated in the target cell type, whereas methylated in all other nucleated cells of the blood. Specific gene loci (amplicon regions) used for the epigenetic cell count quantification include

CD3G/CD3D (all CD3⁺ T cells), CD8B (CD3⁺/CD8⁺ T cells), CD4 (CD4⁺ T cells), FOXP3 (CD3⁺/CD4⁺/FOXP3⁺ Tregs), and LCN2 (CD15⁺ neutrophils). The qPCR signals for each of the immune cell types were normalized relative to the number of nucleated (white blood) cells by using a constitutively unmethylated locus in the GAPDH gene. Each of the assays has been extensively validated in various studies by comparison against absolute cell counts by flow cytometry and show very high correlations (0.85 and higher), both for absolute cells counts (per mm³ of whole blood) and for relative counts as a proportion of total nucleated cells (in whole blood; refs. 13, 20).

Statistical analyses

Each individual cell lineage measured was expressed as a relative percentage of total nucleated cells in the circulation. In addition, for each study participant, the percentages for the CD4⁺ and CD8⁺ T-cell subfractions were further normalized (recalibrated) so as to add up to the fraction of total CD3⁺ T cells, and the percentages of FOXP3⁺ and FOXP3⁻ T helper cells were calculated as fractions of the recalibrated percentage of total CD4⁺ cells (see also Supplementary Fig. S1, additional description in Supplementary Methods and Materials M1–M4 and in ref. 14). The latter, hierarchical recalibration of T-cell subfractions allows an evaluation of improvement in model fit upon stepwise decomposition of the total CD3⁺ T cells into CD4⁺ and CD8⁺ sublineages, and of CD4⁺ into FOXP3⁺ and FOXP3⁻ sublineages, as discussed below.

Multiple imputation was used to impute relative counts of immune cells missing at random in complete matched case sets, that is, neutrophils (proportion of missing 11.1%), CD3⁺ (1.7%), CD8⁺ (1.8%), CD4⁺ (2.7%), FOXP3⁺ (5.5%), adjusted for smoking status (ever vs. never), body mass index (BMI, kg/m²), diabetes (yes, no, unknown), and case/control status. This allowed us to perform analyses in 502 matched case–control sets instead of 416 sets with originally complete immune cell data. The Markov Chain Monte Carlo method for arbitrary missing patterns was used to generate 25 imputed datasets and results were combined to generate valid statistical inferences (21). Comparable results were obtained between imputed and nonimputed datasets, albeit number of case sets were different in the two analyses and risk estimates not significant (Supplementary Table S2).

To examine cross-sectional relationships between relative counts of immune cells and selected baseline risk covariates in controls, partial correlations were calculated adjusted for age at blood draw and sex, for each of the 25 imputed dataset. Resulting correlation coefficients were combined using Fisher z transformation as explained in the SAS documentation PROC MIANALYZE. In addition, a generalized linear model (GLM) was performed to estimate the association between immune markers and diabetes risk in controls.

Conditional logistic regression models were used to examine associations of relative immune cell counts with pancreatic cancer risk, modeling immune cells as continuous variables. Statistical analyses focused in parallel on: (i) the association of pancreatic cancer risk with relative neutrophil counts; and (ii) the association of pancreatic cancer risk with relative blood counts of total (CD3⁺) T cells and T-cell subfractions (CD4⁺ vs. CD8⁺, and within the CD4⁺ fraction FOXP3⁺ vs. other CD4 helper cells). Risk associations were examined for neutrophils and overall CD3⁺ T cells (Model 1), as well as in a series of nested models. Stepwise decomposition models were fitted within the recalibrated T cells, breaking down total T-lymphocytes (CD3⁺) into its CD8⁺ (cytotoxic) and CD4⁺ (helper cell) subfractions (Model 2), and CD4⁺ into its FOXP3⁺ (regulatory) and FOXP3⁻ (nonregulatory) subfractions (Model 3; ref. 14). In this stepwise decomposition

approach, log-likelihood ratio tests were used to examine improvements in overall model fit. As the variables for subcomponents always add up precisely to those for the total of higher-order T-cell lineages (i.e., CD4⁺ plus CD8⁺ equals total CD3⁺, and FOXP3⁺ plus FOXP3⁻ equals CD4⁺), models within this two-step decomposition hierarchy can be considered nested, and stepwise improvements in model fit indicate whether, or not, subcomponent lineages have identical associations with cancer risk as compared with their higher-order sum. For example, if the model including separate variables for CD4⁺ and CD8⁺ cells shows better fit than a model including a variable only for CD3⁺ cells (the sum of CD4⁺ and CD8⁺) this implies that pancreatic risk has significantly different associations with CD4⁺ and with CD8⁺, and that these two risk associations are not summarized well by association of pancreatic cancer risk with the single variable of total CD3⁺ cells.

Analyses were performed for the full, overall case–control sets as well as by strata of lag-time (i.e., strata of 5-years follow-up) between blood draw and cancer diagnosis (<5, 5–10, and ≥10 years). Heterogeneity of the associations of pancreatic cancer risk with immune cell counts by strata of prospective follow-up time was assessed as statistical interaction effect, using likelihood ratio tests. Additional adjustments for potential confounders including smoking status (never, former, current), BMI (kg/m²), self-reported preexisting diabetes at baseline (yes/no), and alcohol lifetime intake (g/day) were examined separately in each model. Although some potential confounders were associated with pancreatic cancer risk (BMI, smoking, diabetes), none were related to immune cell counts and none of the potential confounders changed risk estimates by more than 10% and, therefore, we discarded them from our main models. ORs adjusted for lifestyle factors are very similar and shown in Supplementary Table S3. Sensitivity analyses were performed in microscopically confirmed cases only (*n* = 391, 78% of cases) and by modeling lag-time in shorter intervals close to baseline (i.e., ≤2 years and 2–5 years).

All statistical analyses and imputations were conducted using the Statistical Analysis System (SAS) software package, version 9.4 (SAS Institute Inc.).

Ethics approval and consent to participate

This research project has been performed in accordance with the Declaration of Helsinki and has been approved by the Ethical Committee at IARC (IEC 14-01) and the Medical Faculty Heidelberg (S-103/2015).

Data availability

The EPIC project was launched in the 1990s. Unlike in new studies that we run today, public access to data from the EPIC population was not part of the study protocol at that time. Thus, the data protection statement and informed consent of the EPIC participants do not cover the provision of data in public repositories. Nevertheless, we are open to providing our dataset upon request for (i) statistical validation by reviewers and (ii) pooling projects under clearly defined and secure conditions and based on valid data transfer agreements.

Results

Baseline characteristics of the nested case–control participants are displayed in **Table 1**. Pancreatic cancer cases were on average 64 years old at diagnosis (range: 37–87) with a median follow-up of 8 years (range: 0–16) after baseline blood sampling. More than half of case and control subjects were overweight at recruitment (62% of cases and 55% of controls). Compared with control subjects, cases more often reported current smoking (32% vs. 23%) and preexisting diabetes

Table 1. Baseline characteristics of the study subjects in the nested case-control study within EPIC [median (min-max) or *n* (%)].

Variable	Cases (<i>n</i> = 502)	Controls (<i>n</i> = 502)
Women	248 (49)	248 (49)
Age at recruitment (years)	57 (30-76)	57 (30-76)
Age at diagnosis (years)	64 (37-87)	-
Follow-up (years)	8 (0-16)	
BMI (kg/m ²)		
<25	193 (38)	224 (45)
≥25	309 (62)	278 (55)
Alcohol intake at recruitment (g/d) ^a		
Men	15 (0-138)	14 (0-147)
Women	3 (0-81)	3 (0-59)
Smoking status ^a		
Never	187 (37)	222 (44)
Former	148 (30)	164 (33)
Current	162 (32)	114 (23)
Pack-years	5.0 (0-86)	0.4 (0-81)
Diabetes status ^a		
Self-reported diabetes at recruitment	29 (6)	19 (4)
No diabetes	418 (83)	434 (87)
Counts of immune markers relative to total nucleated cells (%) ^{a,b}		
Neutrophils	55 (3-107)	54 (4-97)
CD3 ⁺	22 (4-60)	22 (6-54)
CD8 ⁺	7 (1-28)	7 (0-27)
CD4 ⁺	15 (2-40)	15 (1-42)
FOXP3 ⁺	1 (0-4)	1 (0-4)
FOXP3 ⁻	14 (1-37)	14 (0-39)

^aMissing (*n* case/*n* control): alcohol intake at recruitment (2/3), smoking status (5/2), pack year (72/55), diabetes (55/49), neutrophils (62/49), CD3⁺ (7/10), CD8⁺ (9/9), CD4⁺ (15/12), FOXP3⁺ (26/29), FOXP3⁻ is the difference between CD4 and FOXP3⁺.

^bImmune marker averages calculated after multiple imputation and recalibration. Neutrophils were not recalibrated; therefore individual percentage can exceed 100.

(6% vs. 4%) at the time of recruitment. Cases and controls showed no differences in alcohol intake (g/day) at recruitment in either men (15% of cases and 14% of controls) or women (3% in cases and controls).

Among the control subjects, on average the neutrophils constituted 54% (range: 4-97) of all nucleated cells, whereas the CD3⁺ cells represented about 22% (6-54). Within the CD3⁺ T-cell compartment, 15% and 7% of the total nucleated cells were CD4⁺ and CD8⁺ cells, respectively, and within the CD4⁺ compartment 1% of total nucleated cells were FOXP3⁺. Percentages were similar among the cases (Table 1). Adjusting for age at blood draw and sex, the relative counts of total CD3⁺ T-lymphocytes and all T-cell subfractions (CD8⁺, CD4⁺, FOXP3⁺) were inversely correlated with neutrophils ($r \leq -0.40$) in controls. Within the T-cell lineage, there was a moderate correlation ($r = 0.51$) between the relative counts of CD8⁺ T and CD4⁺ T cells, and a positive correlation ($r = 0.65$) was also observed between relative counts of FOXP3⁺ and total CD4⁺ cells. All cell types showed only weak and nonsignificant correlations ($r < 0.16$) with BMI, waist circumference, smoking (pack years), time from quitting smoking or alcohol consumption (Fig. 1). In addition to using GLM, the average of relative counts of immune cells were not statistically different between diabetes status (e.g., neutrophils $P = 0.21$, CD3⁺ $P = 0.83$, CD8⁺ $P = 0.78$, CD4⁺ $P = 0.90$, FOXP3⁺ $P = 0.89$).

Regarding associations with pancreatic cancer risk, conditional logistic regression showed no significant associations per percent unit increase of neutrophils or total CD3⁺ T-lymphocytes with pancreatic cancer, irrespectively of the lag time between blood donation and cancer diagnosis (Table 2). Likewise, in the full case-control study set there were no significant improvements in the overall fit for multi-variable models when the counts for overall CD3⁺ T cells (model M1) were broken down into the constituent counts for cytotoxic (CD8⁺) T-lymphocytes and CD4⁺ (helper) T cells (model M2), or when counts of the CD4⁺ compartment were further broken down into regulatory (FOXP3⁺) and nonregulatory (FOXP3⁻) T-lymphocytes (model M3). In analyses by strata of lag-time, however, the hierarchical decomposition into T-cell constituents resulted in borderline significant improvement of model fit when focusing on pancreatic cancer cases whose blood samples were collected less than 5 years prior to cancer diagnosis ($P_{M2M3} = 0.05$ or $P_{M1M3} = 0.03$). Within this 0- to 5-year lag-time interval, the decomposed, mutually adjusted model showed higher pancreatic cancer risk for higher relative counts of regulatory (FOXP3⁺) T-lymphocytes [for a percent unit: OR = 1.80; 95% confidence interval (CI), 1.01-3.20], and for lower counts of cytotoxic CD8⁺ lymphocytes (OR = 0.92; 95% CI, 0.85-1.00; Table 2). For lag-times between blood donation and cancer diagnosis longer than 5 years, these associations and improvements in overall model fit were not observed. Above observed risks were stronger in analyses restricted to microscopically confirmed cases, that is, OR = 2.22 (95% CI, 1.12-4.40) for higher FOXP3 and OR = 0.92 (95% CI, 0.83-1.02) for higher CD8⁺ within the first 5 years of follow-up ($P_{M2M3} = 0.01$ or $P_{M1M3} = 0.02$; not tabled). Further detailed lag-time analyses showed step-wise higher nonsignificant associations for FOXP3 the shorter the follow-up, that is, the closer to baseline (≤ 2 years OR = 2.15; 95% CI, 0.55-8.40 with $n_{cases} = 52$, 2-5 years OR = 1.74; 95% CI, 0.92-3.31 with $n_{cases} = 95$, and 5-10 years OR = 1.06; 95% CI, 0.69-1.63; data not in tables). Significant Likelihood ratio tests were observed for the interaction between length of follow-up and relative immune cell counts of FOXP3⁺ across follow-up categories (<5 years vs. 5-10 years, $P = 0.04$).

No significant associations were observed in analyses stratified by smoking status (ever/never; not tabled). Further stratification by follow-up, however, did show higher pancreatic cancer risk with higher FOXP3⁺ in ever smokers (for a percent unit: OR = 1.87; 95% CI, 1.05-3.33) with follow-up <5 years, although not significant in fully adjusted models.

Additional adjustments for relative counts of neutrophils did not materially alter the fit of any of the above models or alter any of the estimated associations (logistic regression coefficients) between relative cell counts and pancreatic cancer risk.

Discussion

Using a nested case-control design within the European EPIC cohort, we examined the prospective relationship of relative immune cell counts in blood of initially healthy individuals with pancreatic cancer risk, focusing on neutrophils and total (CD3⁺), cytotoxic (CD8⁺), and regulatory (FOXP3⁺) and nonregulatory (FOXP3⁻) CD4⁺ (helper) T-lymphocytes. To determine the relative quantities of these cell types, we used quantitative real-time PCR assays of DNA methylation markers, which provide highly accurate measurements as validated by comparison to absolute to relative cell counts by flow cytometry (13).

The adaptive immune response among healthy individuals depends on the balance between cytotoxic CD8⁺ effector T cells, which drive




Figure 1. Heatmap representing correlations between immune markers and selected baseline factors in 502 controls, adjusted for age and sex, and combined using Fisher transformation. *, Denotes significance $P < 0.05$. Missing (*n* case/*n* control): alcohol average lifetime intake (87/82), waist circumference (56/56), Lifetime smoking in pack-year (72/55), time quitting smoking in former smokers (16/15), neutrophils (62/49), T-lymphocytes after normalization.

the elimination of abnormal cells, and FOXP3⁺ regulatory T lymphocytes (Tregs), which modulate the aggressiveness of the cellular immune response (22, 23). In patients with pancreatic cancer, studies have found greater tumor aggressiveness and worse survival when tumor tissue (especially the tumor center) showed higher FOXP3⁺ T-cell infiltration, whereas high infiltration of particularly CD8⁺ T cells generally corresponded with better outcomes (24, 25). Similarly, studies have shown worse oncologic outcomes, for example, cancer-specific survival including pancreatic cancer (9), among patients who present with higher Tregs and lower CD8⁺ T-lymphocyte counts in blood (26–28). It thus appears plausible that, also among initially healthy individuals, lower relative blood cell counts of CD8⁺ effector T-lymphocytes and higher counts of Tregs within the CD4⁺ T helper cell compartment may be related to higher risk of future cancer development.

Contrary to our expectations, our present study shows no long-term associations of pancreatic cancer risk with the relative quantities of

CD8⁺, or FOXP3⁺ and FOXP3⁻ (CD4⁺) T-cell components, when we used statistical models with stepwise decomposition of total (CD3⁺) T cells into cytotoxic (CD8⁺), regulatory (FOXP3⁺), and nonregulatory (FOXP3⁻) helper cells. These overall null findings stand in contrast to our earlier results from the EPIC-Heidelberg cohort, where over prospective follow-up times up to 15 years (median 6.7 years) this step-wise modeling approach revealed a higher long-term risk of cancers (lung, breast) among initially healthy individuals who had lower proportions of CD8⁺ T cells within the overall T-cell compartment, or who had higher proportions of FOXP3⁺ regulatory T cells among the total circulating CD4⁺ helper T cells (cancers of the lung, breast, and colorectum; ref. 14). However, in subgroup analyses, considering blood samples collected no more than 5 years prior to cancer diagnosis, our present findings did suggest an association of pancreatic cancer risk with a T-lymphocyte signature similar to that identified in our previous EPIC-Heidelberg study, namely higher pancreatic cancer risk among those who had higher relative counts

Table 2. ORs for the association between circulating immune cell composition (relative counts) and pancreatic cancer risk.

	Models ^a	Neutrophils	Total CD3 ⁺	T cells			Model fit improvement P value ^b
				CD8 ⁺	CD4 ⁺	FOXP3 ⁺	
All (502 cases/502 controls)							
Cells modeled individually	M1	1.01 (1.00,1.02)	0.99 (0.97,1.01)				
Cells modeled with mutual adjustments across T-lymphocyte cells	M2		0.99 (0.95,1.03)	0.99 (0.96,1.02)			$P_{M1M2} = 0.89$
	M3		0.99 (0.95,1.03)		1.18 (0.89,1.57)	0.98 (0.94,1.01)	$P_{M2M3} = 0.22$ $P_{M1M3} = 0.45$
Lag time < 5 years (147 cases/147 controls)							
Cells modeled individually	M1	1.01 (0.99,1.03)	0.99 (0.96,1.03)				
Cells modeled with mutual adjustments across T-lymphocyte cells	M2		0.93 (0.86,1.01)	1.03 (0.98,1.08)			$P_{M1M2} = 0.08$
	M3		0.93 (0.85,1.00)		1.80 (1.01,3.23)	0.99 (0.94,1.06)	$P_{M2M3} = 0.05$ $P_{M1M3} = 0.03$
Lag time 5 to 10 years (221 cases/221 controls)							
Cells modeled individually	M1	1.00 (0.99,1.02)	0.99 (0.96,1.02)				
Cells modeled with mutual adjustments across T-lymphocyte cells	M2		1.04 (0.98,1.10)	0.96 (0.92,1.00)			$P_{M1M2} = 0.06$
	M3		1.04 (0.98,1.10)		1.08 (0.70,1.66)	0.95 (0.90,1.00)	$P_{M2M3} = 0.58$ $P_{M1M3} = 0.14$
Lag time 10 years or more (134 cases/134 controls)							
Cells modeled individually	M1	1.00 (0.98,1.03)	0.98 (0.95,1.02)				
Cells modeled with mutual adjustments across T-lymphocyte cells	M2		0.97 (0.90,1.04)	1.00 (0.93,1.07)			$P_{M1M2} = 0.60$
	M3		0.97 (0.90,1.05)		0.92 (0.53,1.62)	1.00 (0.93,1.08)	$P_{M2M3} = 0.77$ $P_{M1M3} = 0.81$

Note: $n = 502$ controls matched to 502 cases on study recruitment centre, sex, age at blood collection, date and time of blood collection, length of follow-up, and fasting status.

^aM1: model including CD3⁺ only, M2: model including CD8⁺ and CD4⁺, M3: model including CD8⁺, FOXP3⁺, FOXP3⁻.

^bImprovement in fit between M1, M2, and M3.

Conditional logistic regression, modelling immune cells continuously as percent counts. Foxp3⁺ with unit 0.1.

Fractions of CD4⁺ and CD8⁺ cells were recalibrated so as to add up to the renormalized fraction of total (CD3⁺) T cells; fractions of FOXP3⁺ and FOXP3⁻ cells were recalibrated so as to add up to fraction of renormalized CD4⁺ cells.

for FOXP3⁺ regulatory T cells among the total CD4⁺ helper T cells, and lower proportions of cytotoxic (CD8⁺) T-lymphocytes. For pancreatic cancer patients whose blood samples had been collected more than 5 years prior to diagnosis, and their matched control subjects, this association pattern was not observed.

The most plausible explanation for seeing associations between relative immune cell counts and pancreatic cancer risk only in the first five years of follow-up is reverse causation, the associations reflecting progressive increases in circulating Tregs, and decreases in CD8⁺ cells, as future pancreatic cancer patients gradually develop more advanced-stage and eventually symptomatic tumors. The well-described model of pancreatic cancer development via precursor lesions (29) and massive infiltration and later shedding of immune cells to the circulation (3) may also support this latter interpretation. Unfortunately, we could not examine whether the associations of immune cell composition with pancreatic cancer within the first 5 years of prospective follow-up varied according to stage at diagnosis, due to incomplete information on tumor stage. Most pancreatic cancers, however, are diagnosed in advanced stage (30).

Neutrophils were originally considered to have pro-inflammatory functions as part of the innate immune responses and as effectors of acute inflammation, but are increasingly being recognized to also exert a broader array of specialized functions in adaptive immune reactions and chronic inflammatory responses to cancer and other diseases (31, 32). In blood, patients with various types of solid tumors, including pancreatic cancer (10, 33, 34), often show increased neutrophil counts compared with cancer-free control subjects, and higher ratios of circulating NLR have been associated with more advanced disease stage and poorer cancer-specific survival rates (35, 36). Again, however, contrary to our initial expectations our present data show no significant association of prediagnosis relative counts of peripheral neutrophils with risk of developing pancreatic cancer. It is worth noting that, in our previous study in the EPIC-Heidelberg cohort, we also did not identify any prospective association of relative neutrophil counts with risks of cancers of the lung, breast, colorectum, or prostate (14).

To our knowledge, our analyses are the first to relate pancreatic cancer risk to relative measures of T-lymphocyte composition using a biologically validated set of cell lineage-specific epigenetic markers, in individuals initially free of known cancer. Cell proportions estimated from epigenetic cell lineage markers—derived from methylation arrays—are nowadays more frequently used. Although this chip array technology is functionally not associated and the role and association of individual CpGs is mostly unknown, these approaches do confirm our currently used approach. Michaud and colleagues recently published on DNA methylation derived immune cell profiles and pancreatic cancer risk but did not observe any association with CD3⁺, CD4⁺, and neutrophils, among others, irrespective of lag-time (37). Their method differs from ours, and used deconvolution algorithms to estimate relative immune cell counts from a broader series of epigenome-wide methylation markers, whereas our more precise and more powerful method uses not only cell type associated but also comethylated regions that were biologically valid and extensively validated against flow cytometry (13).

A limitation of our study is that the epigenetic assays for buffy coat samples allowed quantification only of relative immune cell counts, relative either to total nucleated cells or to a higher-order cell lineage for T-cell compartments, but not of absolute cell counts relative to blood volume. Nonetheless, relative counts by our epigenetic assays show excellent correlations ($r \geq 0.85$) with counts obtained by flow cytometry (13). Another limitation is that only a single, baseline blood

sample is available for the EPIC cohort participants. In our previous study in the EPIC-Heidelberg cohort, however, we found that there was a relatively high intra-individual stability of relative immune cell counts between repeat blood samples collected over time for a sub-set of study participants, with age- and sex-adjusted partial Spearman correlations over a 15-year time interval of about 0.50 for neutrophils, total CD3⁺, total CD4⁺ and FOXP3⁺ cells, and of 0.67 for CD8⁺ cells (14).

In summary, this study in initially cancer-free individuals showed no long-term relationship between blood counts of neutrophils or T-lymphocyte sublineages with later risk of developing pancreatic cancer. However, within prospective follow-up times of less than 5 years between blood sampling and cancer diagnosis, our data suggest a possible higher pancreatic cancer risk in relation to circulating immune cell signature characterized by higher Treg-mediated immune tolerance, and lower CD8⁺ mediated cytotoxicity—a signature that we previously found to be associated with higher risks of cancers of the lung, breast, and colorectum. These time-restricted associations may reflect reverse causation—that is, alterations in circulating immune cell composition induced that result from progressive pancreatic cancer development.

To our knowledge, these analyses are the first to relate pancreatic cancer risk to specific and well-validated epigenetic markers as quantitative measures of circulating T-lymphocyte composition and neutrophil counts in individuals initially free of known cancer. Further studies will be needed to confirm our observations and to assess whether, at all, increased Treg-mediated immune tolerance or reduced CD8⁺ mediated cytotoxicity have any impact on pancreatic cancer risk, and if so, whether this impact is mostly on relatively late-stage tumor development, that is, within relatively short lag-times between blood donation and cancer diagnosis. If further epidemiologic studies confirm findings and show a consistent pattern of associations between circulating immune cell composition and cancer risk, we hope this might open up novel avenues for cancer prevention, focusing on ways to optimize individuals' general immune defense.

Authors' Disclosures

M.-C. Boutron-Ruault reports personal fees from MAYOLI-SPINDLER and GILEAD outside the submitted work. S. Olek reports a patent for US2012107810A1 issued, a patent for WO2010069499A2 issued, a patent for WO2013014122A1 issued, and a patent for WO2011135088A1 issued. No disclosures were reported by the other authors.

Disclaimer

Where authors are identified as personnel of the International Agency for Research on Cancer/World Health Organization, the authors alone are responsible for the views expressed in this article and they do not necessarily represent the decisions, policy or views of the International Agency for Research on Cancer/World Health Organization.

Authors' Contributions

V.A. Katzke: Conceptualization, resources, data curation, formal analysis, supervision, validation, investigation, visualization, methodology, writing—original draft, project administration. **C. Le Cornet:** Conceptualization, resources, data curation, formal analysis, supervision, validation, investigation, visualization, methodology, writing—original draft, project administration. **R. Mahfouz:** Resources. **B. Brauer:** Resources. **T. Johnson:** Resources, laboratory analyses. **F. Canzian:** Resources, writing—review and editing. **V. Rebours:** Resources, writing—review and editing. **M.-C. Boutron-Ruault:** Resources, writing—review and editing. **G. Severi:** Resources, writing—review and editing. **M.B. Schulze:** Resources, writing—review and editing. **A. Olsen:** Resources, writing—review and editing. **A. Tjønneland:** Resources, writing—review and editing. **K. Overvad:** Resources, writing—review and editing. **M. Crous-Bou:** Resources, writing—review and editing. **E. Molina-Montes:** Resources, writing—review and editing. **P. Amiano:** Resources, writing—review and editing. **J.M. Huerta:** Resources, writing—review and

editing. **E. Ardanaz**: Resources, writing–review and editing. **A. Perez-Cornago**: Resources, writing–original draft. **G. Masala**: Resources, writing–review and editing. **V. Pala**: Resources, writing–review and editing. **R. Tumino**: Resources, writing–review and editing. **C. Sacerdote**: Resources, writing–review and editing. **S. Panico**: Resources, writing–review and editing. **B. Bueno-de-Mesquita**: Resources, writing–review and editing. **R. Vermeulen**: Resources, writing–review and editing. **M. Sund**: Resources, writing–review and editing. **O. Franklin**: Resources, writing–review and editing. **S. Christakoudi**: Resources, writing–review and editing. **L. Dossus**: Resources, writing–review and editing. **E. Weiderpass**: Resources, writing–review and editing. **S. Olek**: Resources, writing–review and editing, coordination of laboratory analyses. **R. Kaaks**: Conceptualization, resources, formal analysis, supervision, funding acquisition, validation, investigation, methodology, writing–original draft, project administration, writing–review and editing.

Acknowledgments

We acknowledge the use of data and biological samples from the EPIC-Asturias cohort, PI Ramón Quirós, and the EPIC-Norfolk cohort, PI Nick Wareham. We are grateful for the continuous participation of our cohort participants—without their commitment, this work would not have been possible. We would also like to thank Ms. Julia Oppatt for performing the laboratory work and Ms. Suzanne M Jeurnink (RIVM, the Netherlands) and Daniele Campa (Pisa University, Italy) for intellectual input.

The coordination of EPIC is financially supported by the European Commission (DG-SANCO) and the International Agency for Research on Cancer (IARC) and also by the Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, which has additional infrastructure support provided by the NIHR Imperial Biomedical Research Centre (BRC).

References

- Clark CE, Hingorani SR, Mick R, Combs C, Tuveson DA, Vonderheide RH. Dynamics of the immune reaction to pancreatic cancer from inception to invasion. *Cancer Res* 2007;67:9518–27.
- Huber M, Brehm CU, Gress TM, Buchholz M, Alashkar Alhamwe B, von Strandmann EP, et al. The immune microenvironment in pancreatic cancer. *Int J Mol Sci* 2020;21:7307.
- Inman KS, Francis AA, Murray NR. Complex role for the immune system in initiation and progression of pancreatic cancer. *World J Gastroenterol* 2014;20:11160–81.
- Karamitopoulou E. Tumour microenvironment of pancreatic cancer: immune landscape is dictated by molecular and histopathological features. *Br J Cancer* 2019;121:5–14.
- Fukunaga A, Miyamoto M, Cho Y, Murakami S, Kawarada Y, Oshikiri T, et al. CD8+ tumor-infiltrating lymphocytes together with CD4+ tumor-infiltrating lymphocytes and dendritic cells improve the prognosis of patients with pancreatic adenocarcinoma. *Pancreas* 2004;28:e26–31.
- Ino Y, Yamazaki-Itoh R, Shimada K, Iwasaki M, Kosuge T, Kanai Y, et al. Immune cell infiltration as an indicator of the immune microenvironment of pancreatic cancer. *Br J Cancer* 2013;108:914–23.
- Ikemoto T, Yamaguchi T, Morine Y, Imura S, Soejima Y, Fujii M, et al. Clinical roles of increased populations of Foxp3+CD4+ T cells in peripheral blood from advanced pancreatic cancer patients. *Pancreas* 2006;33:386–90.
- Yamamoto T, Yanagimoto H, Satoi S, Toyokawa H, Hirooka S, Yamaki S, et al. Circulating CD4+CD25+ regulatory T cells in patients with pancreatic cancer. *Pancreas* 2012;41:409–15.
- Karakhanova S, Ryschich E, Mosl B, Harig S, Jager D, Schmidt J, et al. Prognostic and predictive value of immunological parameters for chemoradioimmunotherapy in patients with pancreatic adenocarcinoma. *Br J Cancer* 2015;112:1027–36.
- Arima K, Okabe H, Hashimoto D, Chikamoto A, Tsuji A, Yamamura K, et al. The diagnostic role of the neutrophil-to-lymphocyte ratio in predicting pancreatic ductal adenocarcinoma in patients with pancreatic diseases. *Int J Clin Oncol* 2016;21:940–5.
- Guo W, Lu X, Liu Q, Zhang T, Li P, Qiao W, et al. Prognostic value of neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio for breast cancer patients: an updated meta-analysis of 17079 individuals. *Cancer Med* 2019;8:4135–48.
- Mei Z, Shi L, Wang B, Yang J, Xiao Z, Du P, et al. Prognostic role of pretreatment blood neutrophil-to-lymphocyte ratio in advanced cancer survivors: a systematic review and meta-analysis of 66 cohort studies. *Cancer Treat Rev* 2017;58:1–13.
- Baron U, Werner J, Schildknecht K, Schulze JJ, Mulu A, Liebert UG, et al. Epigenetic immune cell counting in human blood samples for immunodiagnoses. *Sci Transl Med* 2018;10:eaan3508.
- Le Cornet C, Schildknecht K, Rossello Chornet A, Fortner RT, Gonzalez Maldonado S, Katzke VA, et al. Circulating immune cell composition and cancer risk: a prospective study using epigenetic cell count measures. *Cancer Res* 2020;80:1885–92.
- Riboli E, Hunt KJ, Slimani N, Ferrari P, Norat T, Fahey M, et al. European Prospective Investigation into Cancer and Nutrition (EPIC): study populations and data collection. *Public Health Nutr* 2002;5:1113–24.
- Caboux E, Lallemand C, Ferro G, Hemon B, Mendy M, Biessy C, et al. Sources of pre-analytical variations in yield of DNA extracted from blood samples: analysis of 50,000 DNA samples in EPIC. *PLoS One* 2012;7:e39821.
- Baron U, Floess S, Wiczorek G, Baumann K, Grutzkau A, Dong J, et al. DNA demethylation in the human FOXP3 locus discriminates regulatory T cells from activated FOXP3(+) conventional T cells. *Eur J Immunol* 2007;37:2378–89.
- Sehoulji J, Loddenkemper C, Cornu T, Schwachula T, Hoffmuller U, Grutzkau A, et al. Epigenetic quantification of tumor-infiltrating T-lymphocytes. *Epigenetics* 2011;6:236–46.
- Singh A, Yamamoto M, Ruan J, Choi JY, Gauvreau GM, Olek S, et al. Th17/Treg ratio derived using DNA methylation analysis is associated with the late phase asthmatic response. *Allergy Asthma Clin Immunol* 2014;10:32.
- Baron U, Turbachova I, Hellweg A, Eckhardt F, Berlin K, Hoffmuller U, et al. DNA methylation analysis as a tool for cell typing. *Epigenetics* 2006;1:55–60.
- Schlittgen R. Analysis of incomplete multivariate data. *Comput Stat Data Anal* 1999;30:478–9.
- Sakaguchi S, Wing K, Yamaguchi T. Dynamics of peripheral tolerance and immune regulation mediated by Treg. *Eur J Immunol* 2009;39:2331–6.
- Sakaguchi S, Yamaguchi T, Nomura T, Ono M. Regulatory T cells and immune tolerance. *Cell* 2008;133:775–87.
- Orhan A, Vogelsang RP, Andersen MB, Madsen MT, Holmich ER, Raskov H, et al. The prognostic value of tumour-infiltrating lymphocytes in pancreatic cancer: a systematic review and meta-analysis. *Eur J Cancer* 2020;132:71–84.
- Martinez-Bosch N, Vinaixa J, Navarro P. Immune evasion in pancreatic cancer: from mechanisms to therapy. *Cancers* 2018;10:6.
- Schnell A, Schmid C, Herr W, Siska PJ. The peripheral and intratumoral immune cell landscape in cancer patients: a proxy for tumor biology and a tool for outcome prediction. *Biomedicines* 2018;6:25.
- Turbachova I, Schwachula T, Vasconcelos I, Mustea A, Baldinger T, Jones KA, et al. The cellular ratio of immune tolerance (immunoCRIT) is a definite marker for aggressiveness of solid tumors and may explain tumor dissemination patterns. *Epigenetics* 2013;8:1226–35.

The national cohorts are supported by: Danish Cancer Society (Denmark); Ligue Contre le Cancer, Institut Gustave Roussy, Mutuelle Générale de l'Éducation Nationale, Institut National de la Santé et de la Recherche Médicale (INSERM; France); German Cancer Aid, German Cancer Research Center (DKFZ), German Institute of Human Nutrition Potsdam-Rehbruecke (DIfE), Federal Ministry of Education and Research (BMBF; Germany), Associazione Italiana per la Ricerca sul Cancro-AIRC-Italy, Compagnia di SanPaolo and National Research Council (Italy), Dutch Ministry of Public Health, Welfare and Sports (VWS), Netherlands Cancer Registry (NKCR), LK Research Funds, Dutch Prevention Funds, Dutch ZON (Zorg Onderzoek Nederland), World Cancer Research Fund (WCRF), Statistics Netherlands (the Netherlands); Health Research Fund (FIS) - Instituto de Salud Carlos III (ISCIII), PI13/00061 to Granada; PI13/01162 to EPIC-Murcia, Regional Governments of Andalucía, Asturias, Basque Country, Murcia and Navarra, ISCIII RETIC (RD06/0020) and the Catalan Institute of Oncology - ICO (Spain), Swedish Cancer Society, Swedish Research Council and County Councils of Skåne and Västerbotten (Sweden), Cancer Research UK (14136 to EPIC-Norfolk; C570/A16491 and C8221/A29017 to EPIC-Oxford), Medical Research Council (1000143 to EPIC-Norfolk; MR/M012190/1 to EPIC-Oxford; United Kingdom).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received February 9, 2021; revised May 17, 2021; accepted September 3, 2021; published first September 20, 2021.

28. Schreiber RD, Old LJ, Smyth MJ. Cancer immunoediting: integrating immunity's roles in cancer suppression and promotion. *Science* 2011;331:1565–70.
29. Maitra A, Hruban RH. Pancreatic cancer. *Annu Rev Pathol* 2008;3:157–88.
30. Ryan DP, Hong TS, Bardeesy N. Pancreatic adenocarcinoma. *N Engl J Med* 2014; 371:1039–49.
31. Carnevale S, Ghasemi S, Rigatelli A, Jaillon S. The complexity of neutrophils in health and disease: focus on cancer. *Semin Immunol* 2020;48:101409.
32. Liew PX, Kubes P. The neutrophil's role during health and disease. *Physiol Rev* 2019;99:1223–48.
33. Tao L, Zhang L, Peng Y, Tao M, Li G, Xiu D, et al. Preoperative neutrophil-to-lymphocyte ratio and tumor-related factors to predict lymph node metastasis in patients with pancreatic ductal adenocarcinoma (PDAC). *Oncotarget* 2016;7: 74314–24.
34. Ben Q, An W, Wang L, Wang W, Yu L, Yuan Y. Validation of the pretreatment neutrophil-lymphocyte ratio as a predictor of overall survival in a cohort of patients with pancreatic ductal adenocarcinoma. *Pancreas* 2015;44:471–7.
35. Cupp MA, Cariolou M, Tzoulaki I, Aune D, Evangelou E, Berlanga-Taylor AJ. Neutrophil to lymphocyte ratio and cancer prognosis: an umbrella review of systematic reviews and meta-analyses of observational studies. *BMC Med* 2020; 18:360.
36. Yang JJ, Hu ZG, Shi WX, Deng T, He SQ, Yuan SG. Prognostic significance of neutrophil to lymphocyte ratio in pancreatic cancer: a meta-analysis. *World J Gastroenterol* 2015;21:2807–15.
37. Michaud DS, Ruan M, Koestler DC, Alonso L, Molina-Montes E, Pei D, et al. DNA methylation-derived immune cell profiles, CpG markers of inflammation, and pancreatic cancer risk. *Cancer Epidemiol Biomarkers Prev* 2020;29:1577–85.