

Research Article

See related editorial by Mayne, p. 2145, and article by Kantor et al., p. 2312

Serum Antioxidant Nutrients, Vitamin A, and Mortality in U.S. Adults

Abhishek Goyal¹, Mary Beth Terry¹, and Abby B. Siegel²

Abstract

Background: Observational studies have suggested that antioxidant nutrients may reduce cancer and overall mortality risks. However, most randomized trials have failed to show survival benefits. Examining nonlinear associations between antioxidant levels and health outcomes may help to explain these discrepant findings.

Methods: We evaluated all-cause, cancer, and cardiovascular mortality risks associated with quintiles (Q1–Q5) of serum antioxidant (vitamins C and E, β -carotene, and selenium) and vitamin A levels, in 16,008 adult participants of The Third National Health and Nutrition Examination Survey (NHANES III; 1988–1994).

Results: Over a median follow-up period of 14.2 years, there were 4,225 deaths, including 891 from cancer and 1,891 from cardiovascular disease. We observed a dose–response decrease in cancer and overall mortality risks with higher vitamin C levels. In contrast, for vitamin A, risk of cancer death decreased from Q1–Q2, with no further decline in risk at higher levels. For vitamin E, having levels in Q4 was associated with the lowest cancer mortality risk. Both vitamin A and E had U-shaped associations with all-cause mortality. Cancer mortality risks decreased from Q1–Q2 for β -carotene and from Q1–Q4 for selenium. However, for β -carotene and selenium, overall mortality risks decreased from Q1–Q2 but then did not change significantly with higher levels.

Conclusions: Antioxidant supplement use should be studied in the context of overall mortality and other competing mortality risks.

Impact: These data suggest the need for novel intervention studies where doses of these agents are individualized based on their serum levels, and possibly, markers of oxidative stress and systemic inflammatory response. *Cancer Epidemiol Biomarkers Prev*; 22(12); 2202–11. ©2013 AACR.

Introduction

Recent data show that between 28% and 30% of U.S. adults use supplements containing vitamins A, C, and E, and 18% to 19% report using selenium (1, 2). Higher serum antioxidant levels have been associated with lower overall mortality risk (3–15), and fruit and vegetable intake (a rich source of antioxidants) has also been shown to predict better health outcomes in observational studies (16–20). In contrast, most randomized controlled trials have either failed to show significant health benefits from taking these supplements or reported possible harm. A recent systematic review of 78 primary and secondary

prevention trials by the Cochrane Collaboration concluded that use of β -carotene, vitamin E, or high doses of vitamin A supplements was associated with increased all-cause mortality, while the role of vitamin C or selenium supplementation was not clear (21). Although the Physicians' Health Study II (22) recently showed a small reduction in total cancer incidence with multivitamin supplement use, most large clinical trials assessing cancer incidence or mortality have failed to show any beneficial effects of taking these supplements (23–29).

Many authors have suggested that the reason for these discrepant findings is that antioxidants have different effects on various disease processes, and that beyond a certain "threshold" level, they can be potentially toxic (30–40). Surprising little literature exists, however, examining nonlinear effects between serum antioxidant nutrient levels and all-cause and cause-specific mortality outcomes. There are also few studies examining competing risks of mortality in the context of antioxidant nutrient use.

Using data from The Third National Health and Nutrition Examination Survey (NHANES; 1988–1994), a large nationally representative cohort of U.S. adults, we examined whether serum levels of micronutrients with antioxidant properties (vitamins C and E, β -carotene, and

Authors' Affiliations: ¹Department of Epidemiology, Columbia University Mailman School of Public Health and ²Departments of Medicine and Surgery, Columbia University College of Physicians and Surgeons, New York, New York

Note: Supplementary data for this article are available at Cancer Epidemiology, Biomarkers & Prevention Online (<http://cebp.aacrjournals.org/>).

Corresponding Author: Abby B. Siegel, Departments of Medicine and Surgery, Columbia University College of Physicians and Surgeons, 622 West 168th St., PH 14, 105C, New York, NY 10032. Phone: 212-305-9781; Fax: 212-305-4343; E-mail: aas54@columbia.edu

doi: 10.1158/1055-9965.EPI-13-0381

©2013 American Association for Cancer Research.

selenium) and vitamin A, predicted the risks of all-cause, cancer, and cardiovascular disease mortality outcomes. Furthermore, by assessing nonlinear associations, we explored some of the reasons behind the inconsistency in findings between observational studies and randomized trials about the role of these agents on health outcomes.

Materials and Methods

Study population

NHANES III was conducted from October 1988 through October 1994 by the Centers for Disease Control and Prevention (CDC) to provide national health estimates of the United States' civilian population (41). The overall sample size was 39,695; interview and examination response rates were 86% (33,974 participants) and 78% (30,818 participants), respectively (42). Follow-up was from the date of survey participation (1988–1994) to December 31, 2006. Of the 16,573 NHANES III participants (ages 20 years and above) who underwent medical examination, our study included 16,008 (97%) participants for whom data on serum antioxidant nutrient levels and vital status were available. Missing data on covariates ranged from less than 1% to 8% of the study participants. In the analysis, we included only those individuals with complete information available for these covariates. NHANES III was approved by the Institutional Review Board at the CDC. Written informed consent was obtained from all study participants (43).

Measures

Our primary exposures of interest were serum levels of vitamin A, and micronutrients with antioxidant properties. Levels of vitamin A (retinol), vitamin C, vitamin E (α -tocopherol), and β -carotene were measured by isocratic high-performance liquid chromatography (44). Serum selenium levels were measured with atomic absorption spectrometry (44). NHANES III laboratory procedures including quality control systems have been described elsewhere (45).

Our primary outcomes of interest were cause-specific and all-cause mortality. Mortality data were primarily obtained by probabilistic matching to the National Death Index records using the 2010 public release version of the NHANES III Linked Mortality File (46). NHANES III also used various other sources of information to determine the final mortality status and causes of death for survey participants including death certificates, Social Security Administration data, and records from the Centers for Medicare and Medicaid Services. The 9th and 10th revisions of the International Statistical Classification of Diseases (ICD-9 and ICD-10) were used to classify deaths due to cancer (C00–C97) and cardiovascular disease (I00–I78; refs. 46–48). Final mortality status was determined for more than 99% of the study participants (46).

Statistical analysis

To accommodate the complex survey design of NHANES III, we applied appropriate statistical weights

in our analyses (42). Cox Proportional Hazards Models were used to estimate the HRs, and the proportional-hazards assumption was tested using the Kolmogorov-type supremum test (49). To examine whether the associations had a dose–response relationship, we modeled serum levels of antioxidants as quintiles, using the first quintile as the reference group.

Covariates used in the analysis were selected *a priori* based on their suspected roles as confounders. We first fit a basic model adjusted for age and gender (Model 1). For multivariable analysis, Model 2 additionally included race–ethnicity, education, income, body mass index (BMI), smoking status, serum cotinine levels, alcohol consumption, fruit and vegetable intake, physical activity, serum total cholesterol levels, hypertension status (systolic or diastolic blood pressure ≥ 140 or ≥ 90 mm Hg, respectively, use of antihypertensive drugs, or hypertension medical history), diabetes mellitus status (glycosylated hemoglobin $\geq 6.5\%$, use of antidiabetic drugs, or diabetes medical history), history of heart attack, congestive heart failure, stroke or cancer, hormone use among women (use of any estrogen or progesterone including oral contraceptive pills in the past one month), and use of vitamin or mineral supplements (in the past one month). Categories of variables used in the analysis were consistent with the NHANES III survey design (42).

Additional analyses examined these associations in the first 5 or 10 years of follow-up, after excluding deaths within the first 3 years of the survey, and after excluding current smokers. We also assessed the independent effects of these agents by adjusting for other micronutrients in the multivariable models. Because systemic inflammatory response has been shown to affect plasma micronutrient measurements, in an additional analysis, we further adjusted for C-reactive protein levels (50). As participants with self-reported history of comorbidities may have changed their dietary habits and supplement usage, we conducted separate analyses after excluding those with known history of heart attack, congestive heart failure, stroke, or cancer. Finally, to investigate dose–response associations using vitamin A and antioxidant nutrient levels as continuous variables, we created restricted cubic spline functions for all-cause, cancer, and cardiovascular disease mortality outcomes (51). All analyses were done using SAS (Version 9.3; SAS Institute Inc.).

Results

Of the 16,008 study participants, 4,225 died over a median follow-up period of 14.2 years. Eight hundred and ninety-one deaths were due to cancer and 1,891 were due to cardiovascular disease. Table 1 shows the mean baseline serum levels subdivided by sociodemographic, lifestyle, and health-related variables. There were significant differences in serum micronutrient levels for different participant characteristics, especially for BMI categories and smoking status, with current smokers and those with BMI ≥ 30 kg/m² having

Table 1. Serum antioxidant nutrient levels in NHANES III participants

Characteristic	n	Mean serum levels (95% CI)				
		Vitamin A ($\mu\text{mol/L}$)	Vitamin C (mmol/L)	Vitamin E ($\mu\text{mol/L}$)	β -Carotene ($\mu\text{mol/L}$)	Selenium (nmol/L)
Age						
20–39 years	6,425	1.93 (1.90–1.96)	39.77 (37.95–41.60)	22.84 (22.45–23.22)	0.29 (0.28–0.31)	1.58 (1.55–1.60)
40–59 years	4,252	2.08 (2.05–2.11)	41.94 (39.90–43.97)	28.80 (27.89–29.72)	0.39 (0.37–0.41)	1.60 (1.57–1.62)
≥ 60 years	5,331	2.24 (2.20–2.28)	49.52 (47.35–51.69)	32.85 (32.21–33.49)	0.49 (0.47–0.52)	1.59 (1.56–1.61)
Gender						
Male	7,510	2.18 (2.15–2.21)	38.24 (36.58–39.90)	26.38 (25.82–26.93)	0.31 (0.30–0.33)	1.61 (1.58–1.63)
Female	8,498	1.93 (1.90–1.95)	46.59 (44.60–48.58)	27.52 (26.95–28.09)	0.42 (0.41–0.43)	1.57 (1.54–1.59)
Race-ethnicity						
Non-Hispanic White	6,783	2.09 (2.07–2.12)	43.80 (41.70–45.89)	27.77 (27.19–28.35)	0.37 (0.36–0.39)	1.60 (1.57–1.63)
Non-Hispanic Black	4,270	1.90 (1.88–1.93)	34.27 (33.27–35.26)	22.91 (22.51–23.30)	0.36 (0.33–0.38)	1.50 (1.49–1.52)
Mexican-American	4,325	1.86 (1.83–1.89)	39.34 (37.61–41.08)	25.06 (24.64–25.49)	0.32 (0.29–0.34)	1.57 (1.55–1.59)
Other	630	1.92 (1.84–2.00)	43.68 (39.63–47.73)	25.77 (24.48–27.06)	0.39 (0.36–0.43)	1.57 (1.54–1.60)
Level of education						
Less than high school	6,556	2.06 (2.02–2.09)	37.85 (34.73–40.97)	26.77 (26.36–27.18)	0.35 (0.33–0.37)	1.56 (1.54–1.58)
High school or more	9,344	2.05 (2.02–2.07)	44.08 (42.57–45.59)	27.03 (26.47–27.60)	0.37 (0.36–0.39)	1.60 (1.57–1.62)
Annual family income						
<\$20,000	7,730	2.02 (1.98–2.05)	38.54 (36.67–40.40)	25.71 (25.31–26.11)	0.34 (0.33–0.36)	1.56 (1.54–1.58)
\geq \$20,000	8,018	2.06 (2.04–2.09)	44.53 (42.75–46.30)	27.59 (27.02–28.17)	0.38 (0.37–0.40)	1.60 (1.57–1.62)
BMI						
<25 kg/m^2	6,314	1.98 (1.95–2.01)	45.07 (43.27–46.87)	25.37 (24.78–25.96)	0.41 (0.40–0.43)	1.60 (1.57–1.62)
25–29.9 kg/m^2	5,609	2.13 (2.10–2.15)	42.44 (40.51–44.37)	28.27 (27.70–28.85)	0.36 (0.35–0.37)	1.58 (1.56–1.61)
≥ 30 kg/m^2	4,042	2.06 (2.03–2.10)	37.79 (35.60–39.97)	28.27 (27.50–29.03)	0.29 (0.27–0.31)	1.57 (1.54–1.60)
Smoking status						
Current smoker	4,086	1.99 (1.96–2.03)	31.67 (29.28–34.05)	24.27 (23.82–24.72)	0.25 (0.23–0.26)	1.56 (1.53–1.58)
Former smoker	4,018	2.18 (2.15–2.21)	45.99 (44.31–47.66)	29.68 (28.91–30.44)	0.41 (0.38–0.43)	1.61 (1.59–1.64)
Never smoker	7,903	2.01 (1.98–2.03)	47.45 (45.79–49.10)	27.12 (26.54–27.70)	0.42 (0.41–0.44)	1.59 (1.57–1.61)
Alcohol consumption						
Yes	7,598	2.09 (2.06–2.12)	41.50 (39.94–43.05)	26.03 (25.54–26.53)	0.34 (0.32–0.35)	1.60 (1.57–1.62)
No	8,410	1.99 (1.97–2.02)	43.95 (41.62–46.29)	28.15 (27.45–28.86)	0.41 (0.39–0.43)	1.57 (1.55–1.60)
Physical activity ^a						
More active	4,958	2.11 (2.08–2.14)	46.11 (44.26–47.96)	28.52 (27.71–29.34)	0.44 (0.42–0.46)	1.60 (1.57–1.62)
Less active	3,518	2.00 (1.96–2.03)	39.04 (37.59–40.49)	25.66 (25.12–26.20)	0.30 (0.29–0.32)	1.57 (1.54–1.60)
About the same	7,217	2.03 (2.00–2.05)	41.44 (39.27–43.61)	26.48 (25.96–26.99)	0.35 (0.33–0.36)	1.59 (1.56–1.61)
Hypertension status						
Yes	5,484	2.23 (2.20–2.26)	42.90 (40.71–45.09)	30.88 (30.29–31.46)	0.39 (0.37–0.42)	1.59 (1.57–1.62)
No	10,522	1.98 (1.96–2.00)	42.45 (40.78–44.12)	25.47 (25.01–25.94)	0.36 (0.35–0.37)	1.58 (1.56–1.61)
Diabetes mellitus status						
Yes	1,770	2.13 (2.07–2.19)	38.37 (36.00–40.74)	32.24 (30.79–33.68)	0.37 (0.35–0.40)	1.60 (1.57–1.62)
No	14,238	2.04 (2.02–2.07)	42.89 (41.12–44.66)	26.57 (26.10–27.04)	0.37 (0.36–0.38)	1.59 (1.56–1.61)
Hypercholesterolemia						
Yes	4,875	2.25 (2.23–2.28)	44.53 (42.28–46.78)	33.91 (33.12–34.69)	0.44 (0.41–0.46)	1.61 (1.59–1.64)
No	11,124	1.95 (1.93–1.98)	41.69 (39.99–43.38)	23.81 (23.40–24.22)	0.34 (0.32–0.35)	1.57 (1.55–1.60)
Hormone use in women						
Yes	818	2.30 (2.25–2.36)	51.16 (47.51–54.80)	30.93 (29.14–32.72)	0.38 (0.34–0.41)	1.65 (1.62–1.68)
No	7,494	2.03 (2.01–2.05)	41.89 (40.17–43.60)	26.66 (26.23–27.10)	0.37 (0.36–0.38)	1.58 (1.56–1.60)
Supplement use						
Yes	6,050	2.13 (2.10–2.16)	52.43 (50.31–54.55)	31.22 (30.46–31.98)	0.46 (0.44–0.48)	1.59 (1.56–1.61)
No	9,948	1.99 (1.96–2.01)	35.33 (33.56–37.11)	23.85 (23.50–24.21)	0.30 (0.29–0.31)	1.59 (1.56–1.61)

^aCompared with others of same age.

significantly lower vitamin C and β -carotene levels. NHANES III Analytic and Reporting Guidelines provide further details about the characteristics of the study population (42).

Table 2 shows the HRs and 95% confidence intervals (CI) for all-cause mortality by quintiles (Q1–Q5) of micronutrient levels. Table 2 also provides the cutoff points used for dividing serum levels into quintiles. HRs for cancer and cardiovascular disease mortality are shown in Table 3.

We observed U-shaped associations between serum levels of vitamins A and E, and all-cause mortality (Fig. 1) with those with levels in Q1 or Q5 having higher mortality risks compared with those with levels in Q2–Q4. For vitamin A, risk of cancer death decreased from Q1 to Q2, with no further decline in risk at higher levels, whereas for vitamin E, having levels ≥ 26.08 $\mu\text{mol/L}$ (Q4–Q5) were associated with the lowest cancer mortality risk. The increased all-cause mortality risk for those with levels in Q5 for vitamin A was mainly driven by higher cardio-

vascular disease mortality (Fig. 2), and for vitamin E, by higher stroke mortality.

For vitamin C (Model 2), all-cause mortality risk decreased with increases in serum levels from Q1 to Q4 ($P_{\text{trend}} < 0.001$) with no further decline in risk with higher levels (≥ 60.19 mmol/L). For cancer mortality, we observed a dose-response relationship between higher levels and reduced risk ($P_{\text{trend}} < 0.001$). Cardiovascular disease mortality decreased with higher vitamin C levels, except for those with levels in Q5.

For β -carotene and selenium, there was a significant decrease in the overall mortality risk from Q1 to Q2 (HR for Q2 vs. Q1 for β -carotene: 0.75; 95% CI, 0.61–0.93; for selenium: 0.74; 95% CI, 0.63–0.87); however, higher levels (≥ 0.15 $\mu\text{mol/L}$ for β -carotene; ≥ 1.40 nmol/L for selenium) did not significantly change the risk. Cancer mortality risks decreased from Q1 to Q2 for β -carotene and from Q1 to Q4 for selenium. Beyond Q1, higher levels seemed to increase the cardiovascular disease mortality risk for both β -carotene and selenium.

Table 2. HRs for all-cause mortality by quintiles of serum levels

	Quintiles of serum levels					P_{trend}
	Q1	Q2	Q3	Q4	Q5	
Vitamin A levels, $\mu\text{mol/L}$	≤ 1.50	1.54–1.78	1.82–2.06	2.09–2.41	≥ 2.44	
All-cause mortality, no. of events	641	629	804	916	1218	
Model 1 ^a	1 (Reference)	0.80 (0.67–0.96)	0.73 (0.61–0.86)	0.70 (0.59–0.82)	0.83 (0.70–0.98)	0.16
Model 2 ^b	1 (Reference)	0.89 (0.73–1.09)	0.82 (0.68–0.98)	0.82 (0.69–0.97)	0.95 (0.78–1.15)	0.93
Vitamin C levels, mmol/L	≤ 15.33	15.90–31.80	32.36–45.42	45.99–59.62	≥ 60.19	
All-cause mortality, no. of events	949	691	651	662	838	
Model 1	1 (Reference)	0.78 (0.65–0.93)	0.63 (0.51–0.78)	0.53 (0.45–0.62)	0.55 (0.47–0.65)	<0.01
Model 2	1 (Reference)	0.90 (0.77–1.06)	0.80 (0.63–1.01)	0.75 (0.61–0.91)	0.75 (0.62–0.91)	<0.01
Vitamin E levels, $\mu\text{mol/L}$	≤ 18.65	18.67–22.08	22.11–26.05	26.08–32.16	≥ 32.18	
All-cause mortality, no. of events	555	620	803	979	1251	
Model 1	1 (Reference)	0.80 (0.66–0.97)	0.69 (0.57–0.84)	0.69 (0.56–0.84)	0.71 (0.60–0.84)	<0.01
Model 2	1 (Reference)	0.84 (0.67–1.04)	0.81 (0.68–0.97)	0.84 (0.66–1.07)	0.89 (0.70–1.13)	0.77
β -Carotene levels, $\mu\text{mol/L}$	≤ 0.13	0.15–0.20	0.22–0.32	0.34–0.50	≥ 0.52	
All-cause mortality, no. of events	693	662	814	912	1127	
Model 1	1 (Reference)	0.72 (0.61–0.85)	0.65 (0.56–0.76)	0.60 (0.51–0.72)	0.58 (0.49–0.69)	<0.01
Model 2	1 (Reference)	0.75 (0.61–0.93)	0.76 (0.65–0.89)	0.75 (0.63–0.89)	0.75 (0.63–0.90)	0.01
Selenium levels, nmol/L	≤ 1.38	1.40–1.50	1.51–1.60	1.61–1.71	≥ 1.73	
All-cause mortality, no. of events	930	858	763	679	856	
Model 1	1 (Reference)	0.70 (0.60–0.82)	0.67 (0.57–0.79)	0.58 (0.48–0.70)	0.66 (0.57–0.76)	<0.01
Model 2	1 (Reference)	0.74 (0.63–0.87)	0.77 (0.66–0.91)	0.69 (0.56–0.85)	0.79 (0.68–0.92)	0.03

NOTE: Data are given as: HR (95% CI).

^aModel 1: adjusted for age and gender.

^bModel 2: adjusted for variables in Model 1 and race-ethnicity, level of education, annual family income, BMI, smoking status, serum cotinine level, alcohol consumption, fruit and vegetable intake, physical activity, serum total cholesterol levels, hypertension status, diabetes status, history of heart attack, congestive heart failure, stroke or cancer, hormone use in women, and supplement use.

Table 3. HRs for cancer and cardiovascular disease mortality by quintiles of serum levels

	Quintiles of serum levels					<i>P</i> _{trend}
	Q1	Q2	Q3	Q4	Q5	
Vitamin A levels, $\mu\text{mol/L}$	≤ 1.50	1.54–1.78	1.82–2.06	2.09–2.41	≥ 2.44	
Cancer, no. of deaths	150	137	171	215	216	
Model 1 ^a	1 (Reference)	0.83 (0.58–1.18)	0.79 (0.61–1.02)	0.80 (0.61–1.06)	0.72 (0.53–0.98)	0.08
Model 2 ^b	1 (Reference)	0.89 (0.59–1.34)	0.87 (0.65–1.15)	0.96 (0.72–1.28)	0.91 (0.64–1.28)	0.89
Cardiovascular disease, no. of deaths	250	265	365	404	596	
Model 1	1 (Reference)	0.87 (0.71–1.07)	0.90 (0.71–1.13)	0.78 (0.64–0.96)	1.08 (0.86–1.36)	0.24
Model 2	1 (Reference)	0.97 (0.78–1.21)	0.93 (0.72–1.20)	0.84 (0.66–1.06)	1.09 (0.83–1.43)	0.45
Vitamin C levels, mmol/L	≤ 15.33	15.90–31.80	32.36–45.42	45.99–59.62	≥ 60.19	
Cancer, no. of deaths	234	161	147	141	140	
Model 1	1 (Reference)	0.79 (0.61–1.04)	0.56 (0.37–0.85)	0.49 (0.37–0.64)	0.40 (0.29–0.57)	<0.01
Model 2	1 (Reference)	0.86 (0.64–1.15)	0.70 (0.45–1.09)	0.69 (0.51–0.91)	0.55 (0.39–0.79)	<0.01
Cardiovascular disease, no. of deaths	409	288	267	281	416	
Model 1	1 (Reference)	0.73 (0.56–0.94)	0.62 (0.48–0.81)	0.49 (0.39–0.61)	0.60 (0.49–0.73)	<0.01
Model 2	1 (Reference)	0.83 (0.64–1.08)	0.76 (0.58–1.01)	0.66 (0.49–0.89)	0.77 (0.60–1.00)	0.04
Vitamin E levels, $\mu\text{mol/L}$	≤ 18.65	18.67–22.08	22.11–26.05	26.08–32.16	≥ 32.18	
Cancer, no. of deaths	117	155	192	195	230	
Model 1	1 (Reference)	0.98 (0.64–1.51)	0.85 (0.59–1.22)	0.68 (0.42–1.09)	0.68 (0.49–0.94)	<0.01
Model 2	1 (Reference)	1.04 (0.64–1.68)	1.01 (0.67–1.55)	0.88 (0.50–1.53)	0.93 (0.57–1.50)	0.52
Cardiovascular disease, no. of deaths	210	249	356	469	596	
Model 1	1 (Reference)	1.06 (0.78–1.44)	0.91 (0.64–1.30)	1.00 (0.75–1.32)	1.03 (0.78–1.35)	0.73
Model 2	1 (Reference)	1.05 (0.76–1.45)	1.01 (0.69–1.48)	1.10 (0.83–1.47)	1.16 (0.84–1.59)	0.29
β -Carotene levels, $\mu\text{mol/L}$	≤ 0.13	0.15–0.20	0.22–0.32	0.34–0.50	≥ 0.52	
Cancer, no. of deaths	173	150	180	181	205	
Model 1	1 (Reference)	0.67 (0.49–0.93)	0.63 (0.47–0.86)	0.60 (0.46–0.77)	0.51 (0.39–0.67)	<0.01
Model 2	1 (Reference)	0.74 (0.50–1.09)	0.76 (0.53–1.08)	0.83 (0.59–1.16)	0.74 (0.54–1.01)	0.25
Cardiovascular disease, no. of deaths	263	265	354	437	561	
Model 1	1 (Reference)	0.73 (0.57–0.93)	0.71 (0.55–0.92)	0.70 (0.54–0.89)	0.69 (0.53–0.90)	0.03
Model 2	1 (Reference)	0.75 (0.56–1.01)	0.84 (0.64–1.10)	0.86 (0.67–1.12)	0.91 (0.68–1.21)	0.82
Selenium levels, nmol/L	≤ 1.38	1.40–1.50	1.51–1.60	1.61–1.71	≥ 1.73	
Cancer, no. of deaths	194	191	169	128	182	
Model 1	1 (Reference)	0.79 (0.56–1.12)	0.77 (0.58–1.02)	0.49 (0.34–0.69)	0.73 (0.54–0.97)	0.01
Model 2	1 (Reference)	0.76 (0.52–1.12)	0.81 (0.62–1.08)	0.53 (0.36–0.79)	0.86 (0.62–1.20)	0.20
Cardiovascular disease, no. of deaths	400	387	335	320	383	
Model 1	1 (Reference)	0.71 (0.57–0.90)	0.67 (0.53–0.85)	0.73 (0.58–0.93)	0.71 (0.58–0.88)	0.04
Model 2	1 (Reference)	0.76 (0.61–0.95)	0.77 (0.63–0.96)	0.89 (0.71–1.12)	0.83 (0.67–1.04)	0.58

NOTE: Data are given as: HR (95% CI).

^aModel 1: adjusted for age and gender.

^bModel 2: adjusted for variables in Model 1 and race–ethnicity, level of education, annual family income, BMI, smoking status, serum cotinine level, alcohol consumption, fruit and vegetable intake, physical activity, serum total cholesterol levels, hypertension status, diabetes status, history of heart attack, congestive heart failure, stroke or cancer, hormone use in women, and supplement use.

In our analysis, in multivariable models, HRs did not change significantly after further adjusting for serum C-reactive protein levels (Supplementary Table S1) or after adjusting for other micronutrients in the study (Supplementary Table S2). Excluding NHANES III participants

who are current smokers (Supplementary Table S3) or those with history of heart attack, congestive heart failure, stroke, or cancer (Supplementary Table S4) also did not materially change the findings. We observed similar patterns for mortality risks after excluding deaths within the




Figure 1. HRs for all-cause mortality. HRs are adjusted for variables in Model 2: age, gender, race–ethnicity, level of education, annual family income, BMI, smoking status, serum cotinine level, alcohol consumption, fruit and vegetable intake, physical activity, serum total cholesterol levels, hypertension status, diabetes status, history of heart attack, congestive heart failure, stroke or cancer, hormone use in women, and supplement use. The y-axis is shown on a log scale.

first 3 years of the survey (Supplementary Table S5) or after limiting follow-up to 5 or 10 years (Supplementary Table S6). We also examined HRs for all-cause mortality stratified by age, gender, race–ethnicity, BMI, and supplement use (Supplementary Table S7). More detailed results for cancer and cardiovascular disease mortality outcomes are provided in Supplementary Tables S8 and

S9, respectively. Restricted cubic splines for all-cause mortality, cancer mortality, and cardiovascular mortality, are presented in Supplementary Figures S1, S2, and S3, respectively.

Discussion

Oxidative stress has been implicated in the pathogenesis of several chronic diseases, including various cancers (52–55). Vitamin A and antioxidant nutrients such as vitamins C and E, β-carotene, and selenium have been hypothesized to prevent or delay this damage (56–61). Consequently, use of these agents is rising, and almost 40% of U.S. adults take them for their perceived health benefits (1, 2).

In this study, using data from a large, nationally representative cohort of 16,008 U.S. adults, we observed threshold effects and nonlinear associations between serum levels of these micronutrients, and all-cause and cause-specific mortality outcomes. These results help to explain some of the discrepant findings between the observational studies and randomized trials about the role of these agents on health outcomes.

We found that higher levels were generally associated with a modest decrease in all-cancer mortality risk. However, the most significant decline in risk was generally from the first to the second quintile. A number of high profile trials assessing antioxidants as cancer prevention agents have failed to show significant mortality benefits with taking these supplements (23–27) and some have reported possible harm (28, 29). A recent meta-analysis of 22 primary and secondary prevention trials concluded that there was no evidence to support preventive effects of any of the antioxidant supplements on any cancer type




Figure 2. HRs for cancer and cardiovascular disease mortality. HRs are adjusted for variables in Model 2: age, gender, race–ethnicity, level of education, annual family income, BMI, smoking status, serum cotinine level, alcohol consumption, fruit and vegetable intake, physical activity, serum total cholesterol levels, hypertension status, diabetes status, history of heart attack, congestive heart failure, stroke or cancer, hormone use in women, and supplement use. The y-axes are shown on a log scale.

(62). The Physicians' Health Study II also recently showed only a small reduction in total cancer incidence (HR = 0.92; 95% CI, 0.86–0.998) and a nonsignificant reduction in mortality (HR = 0.88; 95% CI, 0.77–1.01) with more than a decade of daily multivitamin use and follow-up (22). Furthermore, most of these large prevention trials did not consider baseline nutrition level in their inclusion criteria (63, 64).

The importance of assessing baseline nutritional status is underscored by the Linxian Study in China, which specifically targeted a geographic area where subjects had chronically low blood levels of multiple micronutrients. It is the only prevention trial to our knowledge so far that has reported statistically significant reductions in all-cancer (RR = 0.87; 95% CI, 0.75–1.00) as well as overall mortality (RR = 0.91; 95% CI, 0.84–0.99) risks with antioxidant supplementation (65).

Interestingly, in the Physicians' Health Study I, there was no decrease in total cancer incidence with β -carotene supplementation. However, in subgroup analyses, lower baseline β -carotene levels were associated with alcohol consumption and higher BMI, and overall cancer incidence was modestly reduced with supplementation in these subgroups (among daily alcohol drinkers, RR = 0.9; 95% CI, 0.8–1.0; among those in the highest BMI quartile, RR = 0.9; 95% CI, 0.7–1.0; ref. 23). Similarly, in the SU.VI. MAX trial, lower total cancer incidence (RR = 0.69; 95% CI, 0.53–0.91) and all-cause mortality (RR = 0.63; 95% CI, 0.42–0.93) with multivitamin use in men was attributable to their lower baseline antioxidant status (27). Although some of these results could be due to testing multiple hypotheses, the findings presented in our study as well as the results from others examining a nutritionally deficient population further support the hypothesis that supplementation might only be useful for those who have low serum antioxidant levels, and that beyond a threshold, higher levels do not lead to additional benefit, and may potentially be toxic.

Our results also have broader implications for assessing overall mortality benefits from the use of dietary supplements. For example, using antioxidant levels as continuous variables, a study in the British National Diet and Nutrition Survey found that higher levels of vitamin C, selenium, and β -carotene were associated with lower overall mortality risk (HR per SD for vitamin C = 0.81, $P < 0.001$; for selenium = 0.76, $P < 0.001$; and for β -carotene = 0.92, $P = 0.08$), whereas the results for vitamins A and E were not significant (HR for vitamin A = 0.96, $P = 0.4$; α -tocopherol = 0.96, $P = 0.4$; ref. 6). If we had used antioxidant levels as continuous variables, we would have obtained similar results. However, by assessing nonlinear associations, we were able to show that for selenium and β -carotene, beyond the first quintile, there was no apparent decrease in the mortality risk with higher levels (Fig. 1). Moreover, we showed a U-shaped association between levels of vitamins A and E, and all-cause mortality.

These results for overall mortality are also consistent with the findings of the recent systematic review by the

Cochrane Collaboration, which concluded that supplementation with β -carotene, vitamin E, or high doses of vitamin A was associated with increased mortality risk, whereas the role of vitamin C or selenium supplement use was not clear (21). Figure 1 in our study shows that except for vitamin C, having antioxidant levels beyond the first quintile did not lead to any further decrease in all-cause mortality risk. Moreover, for vitamins A and E, higher levels increased the overall risk of death.

Our study has a number of strengths compared with previous studies. First, this is the largest study to date assessing serum vitamin A and antioxidant nutrient levels, and cause-specific and all-cause mortality outcomes. Therefore, instead of examining serum levels as continuous variables, we were able to divide them into quintiles, and assess threshold and nonlinear relationships with mortality. Because of small sample sizes and/or short follow-up durations, most observational studies have been unable to show such associations (3–15). Furthermore, NHANES III used a nationally representative sample of the U.S. population, and final mortality status was available for more than 99% of the participants, minimizing the possibility of selection bias. Assessing serum levels of these agents instead of using dietary history, along with standardized and validated laboratory methods, reduced the potential for information bias. Finally, use of appropriate sampling weights in the analysis helped to obtain statistical estimates similar to those if the entire sampling frame (the United States) had been surveyed.

Our study has several limitations. We tested multiple hypotheses. Therefore, results must be interpreted with caution. We limited our analysis to vitamin A and antioxidant nutrients that are commonly used as dietary supplements. Other potential agents with antioxidant properties such as zinc, lycopene, and other carotenoids need further evaluation. As with any observational study, residual confounding by socioeconomic status, lifestyle variables, and other factors cannot be excluded. However, NHANES III assessed a large number of health-related variables, which enabled us to control for many potential confounders. Furthermore, we obtained different results for different micronutrients as well as for different health outcomes. These findings would be difficult to explain entirely on the basis of residual confounding. In addition to assessing cardiovascular mortality, there are many other possible competing risks for mortality that we could have assessed. We assessed cardiovascular mortality as an example of potential different effects of antioxidants on different disease processes.

Finally, in this study, we used a single measurement of serum levels to assess long-term nutritional status. In our analysis (Supplementary Table S6), we observed similar HRs for 5-year, 10-year, and for the entire duration of follow-up. This suggests that at the population level, a single measurement of serum levels may provide a reliable estimate of long-term antioxidant status. A major limitation of randomized trials assessing use of

supplements for primary prevention is that the participants typically have to be kept on intervention for long periods to significantly affect health outcomes. Therefore, observational studies such as this may facilitate the assessment of the relationship between long-term nutritional status and health outcomes.

In summary, using data from a nationally representative cohort of more than 16,000 U.S. adults, we were able to identify specific plasma levels of vitamin A and various antioxidant nutrients, which were associated with maximum cancer-related and overall survival. We found that beyond a certain threshold, there was generally no additional benefit with higher serum levels with respect to overall mortality. Specifically, for vitamins A and E, higher levels increased the overall mortality risk. These data support the findings of recent randomized trials that have generally failed to show health benefits with taking these supplements (21, 62, 66). We also showed that having low serum antioxidant nutrient levels was associated with higher mortality risk, suggesting that supplementation might still be useful for those who are nutritionally deficient.

Our findings underscore the need to assess safety of these agents like other drugs, rather than classifying them as "dietary supplements," which affects their regulatory oversight (67). Although the current Institute of Medicine guidelines provide recommended dietary allowances and tolerable upper intake levels for these agents, we also highlight the potential significance of measuring serum levels to guide their use as supplements (68). Novel intervention studies might then be planned where doses of these agents are individualized on the basis of serum levels, lifestyle behaviors such as smoking which affect

levels, and possibly, markers of oxidative stress and systemic inflammatory response (50, 69, 70). Such a strategy would help ensure that those who are deficient get the required micronutrients, while also preventing toxic levels of antioxidants that could potentially lead to worse health outcomes.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: A. Goyal, M.B. Terry, A.B. Siegel

Development of methodology: A. Goyal

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): A. Goyal, A.B. Siegel

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): A. Goyal, M.B. Terry, A.B. Siegel

Writing, review, and/or revision of the manuscript: A. Goyal, M.B. Terry, A.B. Siegel

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): A. Goyal, A.B. Siegel

Study supervision: A. Goyal, M.B. Terry, A.B. Siegel

Acknowledgments

The authors thank Dr. Zhezhen Jin from the Department of Biostatistics, Mailman School of Public Health, Columbia University, for his help with restricted cubic spline functions.

Grant Support

This work was supported by the Steven J. Levinson Medical Research Foundation and NIH K23 CA149084-01A1 (to A.B. Siegel).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received April 12, 2013; revised June 26, 2013; accepted July 23, 2013; published OnlineFirst August 8, 2013.

References

- Gahche J, Bailey R, Burt V, Hughes J, Yetley E, Dwyer J, et al. Dietary supplement use among U.S. adults has increased since NHANES III (1988-1994). *NCHS Data Brief* 2011;1-8.
- Bailey RL, Gahche JJ, Lentino CV, Dwyer JT, Engel JS, Thomas PR, et al. Dietary supplement use in the United States, 2003-2006. *J Nutr* 2011;141:261-6.
- Buijsse B, Feskens EJ, Schlettwein-Gsell D, Ferry M, Kok FJ, Kromhout D, et al. Plasma carotene and alpha-tocopherol in relation to 10-y all-cause and cause-specific mortality in European elderly: the Survey in Europe on Nutrition and the Elderly, a Concerted Action (SENECA). *Am J Clin Nutr* 2005;82:879-86.
- Buijsse B, Feskens EJ, Kwape L, Kok FJ, Kromhout D. Both alpha- and beta-carotene, but not tocopherols and vitamin C, are inversely related to 15-year cardiovascular mortality in Dutch elderly men. *J Nutr* 2008;138:344-50.
- Bleys J, Navas-Acien A, Guallar E. Serum selenium levels and all-cause, cancer, and cardiovascular mortality among US adults. *Arch Intern Med* 2008;168:404-10.
- Bates CJ, Hamer M, Mishra GD. Redox-modulatory vitamins and minerals that prospectively predict mortality in older British people: the National Diet and Nutrition Survey of people aged 65 years and over. *Br J Nutr* 2011;105:123-32.
- Akbaraly NT, Arnaud J, Hininger-Favier I, Gourlet V, Rousset AM, Berr C. Selenium and mortality in the elderly: results from the EVA study. *Clin Chem* 2005;51:2117-23.
- Akbaraly TN, Favier A, Berr C. Total plasma carotenoids and mortality in the elderly: results of the Epidemiology of Vascular Ageing (EVA) study. *Br J Nutr* 2009;101:86-92.
- Gale CR, Martyn CN, Winter PD, Cooper C. Vitamin C and risk of death from stroke and coronary heart disease in cohort of elderly people. *BMJ* 1995;310:1563-6.
- Hu P, Reuben DB, Crimmins EM, Harris TB, Huang MH, Seeman TE. The effects of serum beta-carotene concentration and burden of inflammation on all-cause mortality risk in high-functioning older persons: MacArthur studies of successful aging. *J Gerontol A Biol Sci Med Sci* 2004;59:849-54.
- Ito Y, Suzuki K, Ishii J, Hishida H, Tamakoshi A, Hamajima N, et al. A population-based follow-up study on mortality from cancer or cardiovascular disease and serum carotenoids, retinol and tocopherols in Japanese inhabitants. *Asian Pac J Cancer Prev* 2006;7:533-46.
- Khaw KT, Bingham S, Welch A, Luben R, Wareham N, Oakes S, et al. Relation between plasma ascorbic acid and mortality in men and women in EPIC-Norfolk prospective study: a prospective population study. *European Prospective Investigation into Cancer and Nutrition. Lancet* 2001;357:657-63.
- Loria CM, Klag MJ, Caulfield LE, Whelton PK. Vitamin C status and mortality in US adults. *Am J Clin Nutr* 2000;72:139-45.
- Ray AL, Semba RD, Walston J, Ferrucci L, Cappola AR, Ricks MO, et al. Low serum selenium and total carotenoids predict mortality among

- older women living in the community: the women's health and aging studies. *J Nutr* 2006;136:172–6.
15. Sahyoun NR, Jacques PF, Russell RM. Carotenoids, vitamins C and E, and mortality in an elderly population. *Am J Epidemiol* 1996;144:501–11.
 16. Agudo A, Cabrera L, Amiano P, Ardanaz E, Barricarte A, Berenguer T, et al. Fruit and vegetable intakes, dietary antioxidant nutrients, and total mortality in Spanish adults: findings from the Spanish cohort of the European Prospective Investigation into Cancer and Nutrition (EPIC-Spain). *Am J Clin Nutr* 2007;85:1634–42.
 17. Genkinger JM, Platz EA, Hoffman SC, Comstock GW, Helzlsouer KJ. Fruit, vegetable, and antioxidant intake and all-cause, cancer, and cardiovascular disease mortality in a community-dwelling population in Washington County, Maryland. *Am J Epidemiol* 2004;160:1223–33.
 18. Nothlings U, Schulze MB, Weikert C, Boeing H, van der Schouw YT, Bamia C, et al. Intake of vegetables, legumes, and fruit, and risk for all-cause, cardiovascular, and cancer mortality in a European diabetic population. *J Nutr* 2008;138:775–81.
 19. Dauchet L, Amouyel P, Hercberg S, Dallongeville J. Fruit and vegetable consumption and risk of coronary heart disease: a meta-analysis of cohort studies. *J Nutr* 2006;136:2588–93.
 20. Hercberg S, Galan P, Preziosi P, Alfarez MJ, Vazquez C. The potential role of antioxidant vitamins in preventing cardiovascular diseases and cancers. *Nutrition* 1998;14:513–20.
 21. Bjelakovic G, Nikolova D, Gluud LL, Simonetti RG, Gluud C. Antioxidant supplements for prevention of mortality in healthy participants and patients with various diseases. *Cochrane Database Syst Rev* 2012;3:CD007176.
 22. Gaziano JM, Sesso HD, Christen WG, Bubes V, Smith JP, MacFadyen J, et al. Multivitamins in the prevention of cancer in men: the Physicians' Health Study II randomized controlled trial. *JAMA* 2012;308:1871–80.
 23. Cook NR, Le IM, Manson JE, Buring JE, Hennekens CH. Effects of beta-carotene supplementation on cancer incidence by baseline characteristics in the Physicians' Health Study (United States). *Cancer Causes Control* 2000;11:617–26.
 24. Lippman SM, Klein EA, Goodman PJ, Lucia MS, Thompson IM, Ford LG, et al. Effect of selenium and vitamin E on risk of prostate cancer and other cancers: the Selenium and Vitamin E Cancer Prevention Trial (SELECT). *JAMA* 2009;301:39–51.
 25. Lonn E, Bosch J, Yusuf S, Sheridan P, Pogue J, Arnold JM, et al. Effects of long-term vitamin E supplementation on cardiovascular events and cancer: a randomized controlled trial. *JAMA* 2005;293:1338–47.
 26. Lee IM, Cook NR, Manson JE, Buring JE, Hennekens CH. Beta-carotene supplementation and incidence of cancer and cardiovascular disease: the Women's Health Study. *J Natl Cancer Inst* 1999;91:2102–6.
 27. Hercberg S, Galan P, Preziosi P, Bertrais S, Mennen L, Malvy D, et al. The SU.VI.MAX Study: a randomized, placebo-controlled trial of the health effects of antioxidant vitamins and minerals. *Arch Intern Med* 2004;164:2335–42.
 28. The effect of vitamin E and beta carotene on the incidence of lung cancer and other cancers in male smokers. The Alpha-Tocopherol, Beta Carotene Cancer Prevention Study Group. *N Engl J Med* 1994;330:1029–35.
 29. Omenn GS, Goodman GE, Thornquist MD, Balmes J, Cullen MR, Glass A, et al. Risk factors for lung cancer and for intervention effects in CARET, the Beta-Carotene and Retinol Efficacy Trial. *J Natl Cancer Inst* 1996;88:1550–9.
 30. Rock CL, Jacob RA, Bowen PE. Update on the biological characteristics of the antioxidant micronutrients: vitamin C, vitamin E, and the carotenoids. *J Am Diet Assoc* 1996;96:693–702.
 31. Padayatty SJ, Katz A, Wang Y, Eck P, Kwon O, Lee JH, et al. Vitamin C as an antioxidant: evaluation of its role in disease prevention. *J Am Coll Nutr* 2003;22:18–35.
 32. Burton GW, Traber MG. Vitamin E: antioxidant activity, biokinetics, and bioavailability. *Annu Rev Nutr* 1990;10:357–82.
 33. Tapiero H, Townsend DM, Tew KD. The antioxidant role of selenium and seleno-compounds. *Biomed Pharmacother* 2003;57:134–44.
 34. Burton GW, Ingold KU. beta-Carotene: an unusual type of lipid antioxidant. *Science* 1984;224:569–73.
 35. Hathcock JN, Hattan DG, Jenkins MY, McDonald JT, Sundaresan PR, Wilkening VL. Evaluation of vitamin A toxicity. *Am J Clin Nutr* 1990;52:183–202.
 36. Galli F, Azzi A. Present trends in vitamin E research. *Biofactors* 2010;36:33–42.
 37. Omaye ST, Krinsky NI, Kagan VE, Mayne ST, Liebler DC, Bidlack WR. beta-carotene: friend or foe? *Fundam Appl Toxicol* 1997;40:163–74.
 38. Zwolak I, Zaporowska H. Selenium interactions and toxicity: a review. *Selenium interactions and toxicity. Cell Biol Toxicol* 2012;28:31–46.
 39. Rutkowski M, Grzegorzczak K. Adverse effects of antioxidative vitamins. *Int J Occup Med Environ Health* 2012;25:105–21.
 40. Rogovik AL, Vohra S, Goldman RD. Safety considerations and potential interactions of vitamins: should vitamins be considered drugs? *Ann Pharmacother* 2010;44:311–24.
 41. National Center for Health Statistics. The National Health and Nutrition Examination Survey (NHANES). [cited 2013 April 1]. Available from: <http://www.cdc.gov/nchs/nhanes.htm>
 42. National Center for Health Statistics. The Third National Health and Nutrition Examination Survey (NHANES III), 1988-94. Analytic and Reporting Guidelines. [cited 2013 April 1]. Available from: <http://www.cdc.gov/nchs/data/nhanes/nhanes3/nh3gui.pdf>
 43. National Center for Health Statistics. Research Ethics Review Board (ERB) Approval. [cited 2013 April 1]. <http://www.cdc.gov/nchs/nhanes/irba98.htm>
 44. National Center for Health Statistics. Laboratory Procedures Used for the Third National Health and Nutrition Examination Survey (NHANES III), 1988-1994 [cited 2013 April 1]. Available from: <http://www.cdc.gov/nchs/data/nhanes/nhanes3/cdrom/nchs/manuals/labman.pdf>
 45. Gunter EW, McQuillan G. Quality control in planning and operating the laboratory component for the Third National Health and Nutrition Examination Survey. *J Nutr* 1990;120 Suppl 11:1451–4.
 46. National Center for Health Statistics. The Third National Health and Nutrition Examination Survey (NHANES III) Linked Mortality File, Mortality follow-up through 2006: Matching Methodology May 2009. [cited 2013 April 1]. http://www.cdc.gov/nchs/data/datainkage/matching_methodology_nhanes3_final.pdf
 47. National Center for Health Statistics. The Third National Health and Nutrition Examination Survey (NHANES III) Linked Mortality File. [cited 2013 April 1]. http://www.cdc.gov/nchs/data/datainkage/nh3_file_layout_public_2010.pdf
 48. World Health Organization. International statistical classification of diseases and related health problems, 10th revision. Geneva, Switzerland: World Health Organization; 1992.
 49. Lin DY, Wei LJ, Ying Z. Checking the Cox model with cumulative sums of martingale-based residuals. *Biometrika* 1993;80:557–72.
 50. Duncan A, Talwar D, McMillan DC, Stefanowicz F, O'Reilly DS. Quantitative data on the magnitude of the systemic inflammatory response and its effect on micronutrient status based on plasma measurements. *Am J Clin Nutr* 2012;95:64–71.
 51. Desquilbet L, Mariotti F. Dose-response analyses using restricted cubic spline functions in public health research. *Stat Med* 2010;29:1037–57.
 52. Finkel T, Holbrook NJ. Oxidants, oxidative stress and the biology of ageing. *Nature* 2000;408:239–47.
 53. Griendling KK, Alexander RW. Oxidative stress and cardiovascular disease. *Circulation* 1997;96:3264–5.
 54. Klaunig JE, Kamendulis LM. The role of oxidative stress in carcinogenesis. *Annu Rev Pharmacol Toxicol* 2004;44:239–67.
 55. Halliwell B. Free radicals, antioxidants, and human disease: curiosity, cause, or consequence? *Lancet* 1994;344:721–4.
 56. Willcox JK, Ash SL, Catignani GL. Antioxidants and prevention of chronic disease. *Crit Rev Food Sci Nutr* 2004;44:275–95.
 57. Halliwell B. Antioxidants in human health and disease. *Annu Rev Nutr* 1996;16:33–50.
 58. Halliwell B. Antioxidant defence mechanisms: from the beginning to the end (of the beginning). *Free Radic Res* 1999;31:261–72.

59. Tomita Y, Himeno K, Nomoto K, Endo H, Hirohata T. Augmentation of tumor immunity against syngeneic tumors in mice by beta-carotene. *J Natl Cancer Inst* 1987;78:679–81.
60. Glatthaar BE, Hornig DH, Moser U. The role of ascorbic acid in carcinogenesis. *Adv Exp Med Biol* 1986;206:357–77.
61. Sandhu JK, Haqqani AS, Birnboim HC. Effect of dietary vitamin E on spontaneous or nitric oxide donor-induced mutations in a mouse tumor model. *J Natl Cancer Inst* 2000;92:1429–33.
62. Myung SK, Kim Y, Ju W, Choi HJ, Bae WK. Effects of antioxidant supplements on cancer prevention: meta-analysis of randomized controlled trials. *Ann Oncol* 2010;21:166–79.
63. Morris MC, Tangney CC. A potential design flaw of randomized trials of vitamin supplements. *JAMA* 2011;305:1348–9.
64. Mayne ST, Ferrucci LM, Cartmel B. Lessons learned from randomized clinical trials of micronutrient supplementation for cancer prevention. *Annu Rev Nutr* 2012;32:369–90.
65. Blot WJ, Li JY, Taylor PR, Guo W, Dawsey S, Wang GQ, et al. Nutrition intervention trials in Linxian, China: supplementation with specific vitamin/mineral combinations, cancer incidence, and disease-specific mortality in the general population. *J Natl Cancer Inst* 1993;85:1483–92.
66. Myung SK, Ju W, Cho B, Oh SW, Park SM, Koo BK, et al. Efficacy of vitamin and antioxidant supplements in prevention of cardiovascular disease: systematic review and meta-analysis of randomised controlled trials. *BMJ* 2013;346:f10.
67. Dietary Supplement Health and Education Act of 1994. Public Law 103-417. 103rd Congress [cited 2013 April 1]. http://ods.od.nih.gov/About/DSHEA_Wording.aspx
68. Institute of Medicine. Nutrient Recommendations: Dietary Reference Intakes. [cited 2013 April 1]. www.nap.edu/catalog/9810.htmlhttp://ods.od.nih.gov/Health_Information/Dietary_Reference_Intakes.aspx
69. Mayne ST. Antioxidant nutrients and chronic disease: use of biomarkers of exposure and oxidative stress status in epidemiologic research. *J Nutr* 2003;133 Suppl 3:933S–40S.
70. Therond P, Bonnefont-Rousselot D, Davit-Spraul A, Conti M, Legrand A. Biomarkers of oxidative stress: an analytical approach. *Curr Opin Clin Nutr Metab Care* 2000;3:373–84.