

OBSERVATIONS

Characteristics of 98 Children and Adolescents Diagnosed With Type 2 Diabetes by Their Health Care Provider at Initial Presentation

Although the number of children and youth with type 2 diabetes is increasing, a clear case definition that describes children with type 2 diabetes at presentation remains elusive. Most initial diagnoses are decided on the clinical picture at presentation (1). Characteristics and risk factors have been outlined in several review and clinical articles (2–4). The purpose of this study was to describe the characteristics of youth presenting for an initial visit to the outpatient clinic of a large tertiary children's care center and diagnosed with type 2 diabetes.

For this retrospective study, data were abstracted from a consecutive sample of 98 patients' medical records at Texas Children's Hospital starting 1 January 1998 and ending 31 October 2001. The sample's mean age at diagnosis was 13.6 years (SD 2.33; range 8.7–18.4 years). Fifty-one percent of the children were female and 49% were male (female:male ratio 1:1). For 43% race/ethnicity was not specified; the remaining participants were 28.6% African American, 22.4% Hispanic, 3.1% non-Hispanic white, and 3.1% Asian. Of those for whom data were available, a maternal history of type 2 diabetes was reported by 32.7% (18/55) and an unspecified type of diabetes by 12.7% (7/55). Twenty-seven percent (13/47) reported a father with type 2 diabetes and 21% (10/47) an unspecified type of diabetes.

Mean BMI was 34.67 kg/m² (SD 6.91). Ninety-three percent had a BMI \geq 95th percentile. All but three of the individuals had BMIs \geq 85th percentile. Of those for whom data were recorded, acanthosis nigricans was identified in 94%

(48/51). A Tanner stage of 3, 4, or 5 was identified in 73.2% (49/67).

Blood pressure readings indicated that 49.4% (41/83) had a systolic (SBP) and 10.8% (9/83) a diastolic (DBP) \geq 95th percentile for age, sex, and height ($n = 83$). Fifty-five percent (46/83) had SBP and 19.3% (16/83) DBP readings \geq 90th percentile for blood pressure. Of 72 pulse rates recorded, 2.6% were \geq 95th percentile for age. Average HbA_{1c} was 10.38 (SD 3.52) ($n = 95$).

Of those who had symptoms documented in the medical record, 83.6% (56/67) reported polyuria, 83.9% (52/62) polydipsia, and 61% (36/59) polyphagia. Seventy-five percent reported both polyuria and polydipsia (46/61). Of the cases available, 46.2% (24/52) reported all three of the polys at initial presentation, 46.8% (29/62) had weight loss, and 62.5% (30/48) had ketones. Of those for whom islet cell antibody data were recorded (50/98), 49 had JDF units <5 . Fifty-three percent were started on insulin, 46.3% on oral agents, and 13.7% on both insulin and oral agent ($n = 96$). Initial mean insulin dose was 0.6 units/kg.

Our sample is similar to those described in previous reports except for a more even ratio of female to male subjects, a greater percent with elevated SBP and/or DBP, and more individuals reporting weight loss. We are the first to report blood pressure by the 95th and 90th percentiles and the first to report pulse rate. These data contribute to the growing body of clinical evidence defining the characteristics of youth with type 2 diabetes.

SANDRA L. UPCHURCH, PHD, RN, CDE¹

CHRISTINE A. BROSNAN, DRPH, RN¹

JANET C. MEININGER, PHD, RN, FAAN¹

DORIS E. WRIGHT, PHD, RD²

JILL A. CAMPBELL, MS, RD³

SIRIPOOM V. MCKAY, MD⁴

BARBARA SCHREINER, MN, RN, CDE, BCADM⁴

From the ¹School of Nursing, University of Texas Health Science Center at Houston, Houston, Texas; the ²Department of Nutrition and Food Sciences, Texas Woman's University, Houston, Texas; the ³Department of Pediatrics Endocrinology and Metabolism, Texas Children's Hospital, Houston, Texas; and the ⁴Department of Pediatrics, Endocrinology and Metabolism, Baylor College of Medicine, Houston, Texas.

Address correspondence to Sandra L. Upchurch, The University of Texas Health Science Center at Houston, School of Nursing, 1100

Holcombe #5.518, Houston, TX 77025. E-mail: sandra.l.upchurch@uth.tmc.edu.

© 2003 by the American Diabetes Association.

References

1. American Diabetes Association: Type 2 diabetes in children and adolescents. *Pediatr* 105:671–680, 2000
2. Fagot-Campagna A, Pettitt DJ, Engelgau MM, Burrows NR, Geiss LS, Valdez R, Beckles GL, Saaddine J, Gregg EW, Williamson DE, Narayan KM: Type 2 diabetes among North American children and adolescents: a epidemiologic review and a public health perspective. *J Pediatr* 136: 664–672, 2000
3. Maculoso CJ, Bauer UE, Deeb LC, Malone LC, Chaudhari M, Silverstein J, Eidson M, Goldberg RB, Gaughan-Bailey B, Brooks RG, Rosenbloom AL: Type 2 diabetes mellitus among Florida children and adolescents, 1994 through 1998. *Public Health Reports* 117:373–379, 2002
4. Zuhri-Yafi MI, Brosnan PG, Hardin DS: Treatment of type 2 diabetes mellitus in children and adolescents. *J Pediatr Endocrinol Metab* 15:541–546, 2002

Asymptomatic Bacteriuria and Leukocyturia in Type 1 Diabetic Children and Young Adults

In the study of Geerlings et al. (1), one of five type 1 diabetic women had asymptomatic bacteriuria (ASB). In the few studies of diabetic children a low prevalence of \sim 1% was found (2–4). Our clinical experience suggested a much higher prevalence; therefore, we decided to estimate the prevalence and possible risk factors of ASB in type 1 diabetic children.

There were 178 (86 male) type 1 diabetic children and young adults (age 15.1 ± 5.9 years) with diabetes duration of 6.2 (3.0–10.1) [median (interquartile range)] years who participated in this study.

The control group consisted of 194 (103 male) school children/medical students (14.4 ± 5.1 years). After careful cleaning, midstream voiding morning urine samples were collected and immediately cultured on 2 consecutive days.

ASB was defined as the presence of $\geq 10^5$ colony-forming units/ml of one and

weights <2,500, 2,501–3,699, and >3,700 g were defined as low, normal, and high, respectively. In the occupational cohort, subjects with known type 2 diabetes and with HbA_{1c} >6.5% were defined as diabetic patients. In the hospital cohort, current use of antihypertensive agents was defined as hypertension. In both cohorts, subjects who had a diabetic parent, sibling, and/or offspring were considered to have a family history of diabetes. The significance of difference of the results in the different groups was tested by χ^2 analysis or Student's *t* test.

The prevalence of low birth weight in the 301 diabetic subjects, including those in the occupational cohort and the 1,823 nondiabetic subjects in the occupational cohort, was 18.6% (56 of 301) and 9.8% (178 of 1,823), respectively ($P < 0.001$), whereas the prevalence of high birth weight in the same groups was 9.3 and 11.6%, respectively (NS). Of the 56 diabetic subjects with low birth weight, 32 had a family history of diabetes (57.1%), and of the 178 corresponding nondiabetic subjects, only 25 had a family history of diabetes (14.0%) ($P < 0.0001$). Mean BMI at the age of 20 was 21.8 ± 3.3 and 20.6 ± 2.2 kg/m² ($P < 0.01$), and mean maximum BMI was 26.1 ± 3.3 and 24.8 ± 3.0 kg/m² ($P < 0.01$) in the same groups. The prevalence of hypertension in diabetic subjects with low and normal birth weight in the hospital cohort was 46.2% (18 of 39) and 23.4% (22 of 94) ($P < 0.01$), respectively, and the prevalence in those with high birth weight was 29.0% (9 of 31) (NS compared with low and normal weight groups), while mean BMI did not differ significantly among the three groups (23.0 ± 2.3 , 23.9 ± 3.2 , and 25.4 ± 4.2 kg/m²).

Our results showed that low birth weight was also associated with the development of type 2 diabetes in Japanese subjects and that not only genetic influences but also higher BMI in adulthood seemed to be important in the development of type 2 diabetes in individuals with low birth weight. The significantly higher incidence of hypertension in diabetic subjects with low birth weight compared with diabetic subjects with normal birth weight suggests that insulin resistance might be stronger in the former. Further prospective studies with a large number of participants using more appropriate indices of insulin resistance are required to clarify this association be-

tween birth weight and insulin resistance in individuals with type 2 diabetes.

SONOKO ANAZAWA, MD
YOSHIHITO ATSUMI, MD
KEMPEI MATSUOKA, MD

From Saiseikai Central Hospital, Tokyo, Japan.

Address correspondence to Sonoko Anazawa, MD, 1-4-17 Mita, Minato-Ku, Tokyo 108-0073, Japan. E-mail: sonoko-a@fa2.so-net.ne.jp.

© 2003 by the American Diabetes Association.

References

1. Hales CN, Barker DJP, Clark PMS Cox LJ, Fall C, Osmond C, Winter PD: Fetal and infant growth and impaired glucose tolerance at age 64. *BMJ* 303:1019–1022, 1991
2. Barker DJP, Hales CN, Fall CHD, Osmond C, Phillips K, Clark PMS: Type 2 (non-insulin dependent) diabetes mellitus, hypertension and hyperlipidemia (syndrome X): relation to reduced fetal growth. *Diabetologia* 36:62–67, 1993
3. Valdez R, Athens MA, Thompson GH, Bradshaw BS, Stern MP: Birth weight and adult health outcomes in a biethnic population in the USA. *Diabetologia* 37:624–631, 1994
4. Lithell HO, McKeigue PM, Berglund L, Mohsen R, Lithell UB, Leon DA: Relation of size at birth to non-insulin dependent diabetes and insulin concentrations in men aged 50–60 years. *BMJ* 312:406–410, 1996
5. Rich-Edwards JW, Colditz GA, Stampfer MJ, Willet WC, Gillman MW, Hennekens CH, Speizer FE, Manson JA: Birth weight and the risk of type 2 diabetes in adult women. *Ann Intern Med* 130:278–284, 1999
6. McCane DR, Pettit DJ, Hanson RL, Jacobsen LTH, Knowler WC, Bennett PH: Birth weight and non-insulin dependent diabetes: thrifty genotype, thrifty phenotype, or surviving small baby genotype? *BMJ* 308:942–945, 1994
7. Wei JN, Lin RS, Sung FC, Lin CC, Li CY, Chiang CC, Chang CH, Chaung LM: Low birth weight and high birth weight infants are both at an increased risk to have type 2 diabetes among schoolchildren in Taiwan. *Diabetes Care* 26:343–348, 2003

Coffee Consumption and the Incidence of Type 2 Diabetes

There have been few reports of associations between coffee consumption and diabetes. In healthy subjects given an oral glucose tolerance test (OGTT), the 1-h glucose concentra-

tion was lower when the glucose load was given with coffee (1). There were no significant differences in the other time points in the test, nor in serum insulin concentration at any time point. In contrast, others have reported a deterioration of glucose tolerance after coffee ingestion (2,3). In healthy subjects a rise in blood glucose levels after caffeine intake was detected at the 2nd, 3rd, and 4th hours in comparison to those taking a placebo (4). Blood insulin levels were comparable after caffeine or placebo ingestion during the entire OGTT. Ingestion of coffee or the injection of caffeine is associated with elevated plasma catecholamines and free fatty acids, as well as decreased insulin sensitivity (5), all of which might be expected to increase the incidence of diabetes. Recently, coffee consumption was reported to decrease the incidence of type 2 diabetes in the Netherlands (6). Prompted by these inconsistent findings, we attempted to replicate the Dutch finding in the longitudinal population-based study of diabetes among the Pima Indians (7).

The average daily use of coffee was assessed by questionnaire from 1978 to 1992 in 2,680 nondiabetic individuals aged ≥ 15 years (mean age 27 years; 60% women) who had at least one follow-up examination by September 2002. Consumption was recorded as never, occasionally (less than one a day), one or two a day, three or more a day, or occasionally heavy. Diabetes was diagnosed by OGTT using the 1985 World Health Organization criteria (8).

During an average follow-up time of 11 years (range 1–23), 824 individuals developed diabetes. After adjustment for age, sex, and BMI in a time-dependent proportional hazards model, the risk of developing diabetes for those who reported drinking coffee compared with those who never drink coffee (referent category) was as follows: occasionally 1.09 (95% CI: 0.89–1.34), one or two a day 0.92 (0.74–1.13), three or more a day 1.01 (0.82–1.26), and occasionally heavy 0.81 (0.55–1.03). Overall, adjusted for age, sex, and BMI, there was no significant association between consumption categories and incidence of diabetes ($P = 0.6$). Tea consumption was also unrelated to the incidence of diabetes (not shown).

One strength of our study was the periodic use of an OGTT, as compared with the subjective self-reporting of diabetes

used in the Dutch study (6). In that study, coffee consumption was associated with lower socioeconomic status and less healthy behaviors, factors that might be associated with a lower likelihood of being tested for diabetes. Our study included nearly three times as many incident cases of diabetes as the Dutch study (824 vs. 306), resulting in a narrow 95% CI (0.84–1.18) around our estimate of the hazard rate ratio of 0.99 for any coffee consumption. In conclusion, our data provide no evidence for a relationship of coffee consumption and risk of type 2 diabetes.

ARAMESH SAREMI, MD
MARSHALL TULLOCH-REID, MBBS, MPHIL
WILLIAM C. KNOWLER, MD, DRPH

From the Diabetes and Arthritis Epidemiology Section, National Institute of Diabetes and Digestive and Kidney Diseases, Phoenix, Arizona.

Address correspondence to Aramesh Saremi, MD, 1550 East Indian School Rd., Phoenix, Arizona 85014. E-mail: asaremi@mail.nih.gov.

© 2003 by the American Diabetes Association.

References

1. Feinberg LJ, Sandberg H, De Castro O, Bellet S: Effects of coffee ingestion on oral glucose tolerance curves in normal human subjects. *Metabolism* 17:916–922, 1968
2. Jankelson OM, Beaser SB, Howard FM, Mayer J: Effect of coffee on glucose tolerance and circulating insulin in men with maturity-onset diabetes. *Lancet* 1:527–529, 1967
3. Wachmann A, Hattner RS, George B, Bernstein DS: Effects of decaffeinated and nondecaffeinated coffee ingestion on blood glucose and plasma radioimmuno-reactive insulin responses to rapid intravenous infusion of glucose in normal man. *Metabolism* 19:539–546, 1970
4. Pizzol A, Tikhonoff V, Paleari CD, Russo E, Mazza A, Ginocchio G, Onesto C, Pavan L, Casiglia E, Pessina AC: Effects of caffeine on glucose tolerance: a placebo-controlled study. *Eur J Clin Nutr* 52:846–849, 1998
5. Keijzers GB, De Galan BE, Tack CJ, Smits P: Caffeine can decrease insulin sensitivity in humans. *Diabetes Care* 25:364–369, 2002
6. Van Dam RM, Feskens EJM: Coffee consumption and risk of type 2 diabetes mellitus. *Lancet* 360:1477–1478, 2002
7. Knowler WC, Pettitt DJ, Saad MF, Bennett PH: Diabetes mellitus in the Pima Indians: incidence, risk factors and pathogenesis. *Diabetes Metab Rev* 6:1–27, 1990

8. World Health Organization: *Diabetes Mellitus: Report of a WHO Study Group*. Geneva, World Health Org. (Tech. Rep. Ser., no. 727), 1985

Impairment of Glucose Tolerance Over 10 Years in Middle-Aged Normal Glucose Tolerant Indians

We followed 191 normal glucose tolerant (NGT; 1985 World Health Organization criteria) nondiabetic subjects (115 men) as control subjects in the Wellcome Diabetes Study (1). Their mean age was 41 years (SD 11.2), BMI 23.6 kg/m² (34% >25 kg/m²), and 31% had a first-degree relative with diabetes.

During the next 10 years, 8 (7 men) died, 40 were lost to follow up, 14 men and 8 women became impaired glucose tolerant (IGT), and 2 men and 4 women developed diabetes. Men whose glucose tolerance deteriorated were heavier at entry (71.8 vs. 62.3 kg, $P < 0.001$), more obese (BMI 25.3 vs. 22.6 kg/m², $P < 0.01$), and more centrally obese (waist circumferences 85.9 vs. 78.9 cm, $P < 0.01$) than those who remained NGT, all adjusted for age. They also had higher 2-h glycemia (oral glucose tolerance test, 6.6 vs. 5.9 mmol/l, $P < 0.05$), fasting triglyceridemia (1.6 vs. 1.1 mmol/l, $P < 0.01$), and fasting and 2-h insulinemia (95.1 vs. 47.9 and 929 vs. 515 pmol/l, $P < 0.05$ for both), which was reflected in insulin resistance (homeostasis model assessment [HOMA] 2.7 vs. 1.4, $P < 0.05$). Among women, triglyceridemia (1.5 vs. 0.9 mmol/l, $P < 0.01$) and higher systolic blood pressure (137 vs. 122 mmHg, $P < 0.05$) were predictive.

On multivariate analysis, after forcing in age, sex, and family history of diabetes, glucose tolerance deterioration (both sexes) was predicted by initial HOMA (odds ratio 1.38, 95% CI 1.01–1.85), 2-h plasma glucose (1.04, 1.00–1.08), fasting plasma triglyceride concentration (1.01, 1.00–1.02), and weight gain (1.2, 1.02–1.32).

These results, from a first prospective study of such duration among Indians in India, confirm studies from elsewhere (2) in associating deterioration of glucose tol-

erance in the NGT with obesity, weight gain, insulin resistance, higher circulating triglycerides, and 2-h glucose concentrations. Clearly, there is an excess of insulin resistance over B-cell deficiency markers. Finally, we wish to emphasize the relative thinness at which these effects were seen. The relative risk of deterioration of glucose tolerance during 10 years among the whole group was 2.4 (1.1–5.3) with BMI above and below 23 kg/m². This may reflect both the higher body fat percentage for a given BMI among Indians and their marked central adiposity (3). This has already prompted a reduction in the target BMI for obesity-related action among Asian Indians to 23 kg/m² (4).

Therefore, among Indians reduction in adiposity must be a prime target for diabetes prevention. This will have to start at levels that are accepted in the west without demur. This is necessary at all ages, but will be made difficult by our recent observation that central obesity and hyperinsulinemia are present in Indians at birth (5).

CHITTARANJAN S. YAJNIK, MD¹
KISHORE M. SHELGIKAR, MD¹
SADANAND S. NAIK, PHD¹
MEHMOOD G. SAYYAD, MSC¹
KONDIRAM N. RAUT, MSW¹
DATTATRAYA S. BHAT, MSC¹
JYOTI A. DESHPANDE, MSC¹
SHAILAJA D. KALE, MD¹
DEREK HOCKADAY, FRCP²

From the ¹Diabetes Unit, KEM Hospital and Research Centre, Pune, India; and the ²Oxford Lipid Metabolism Unit, Sheikh Rashid Laboratories, Radcliffe Infirmary, Oxford, U.K.

Address correspondence to Chittaranjan S. Yajnik, MD, Diabetes Unit, KEM Hospital and Research Centre, Rasta Peth, Pune 411011, India. E-mail: diabetes@vsnl.com

© 2003 by the American Diabetes Association.

References

1. Shelgikar KM, Hockaday TDR, Yajnik CS: Central rather than generalized obesity is associated with hyperglycemia in Asian Indian subjects. *Diabet Med* 8:712–717, 1991
2. Haffner SM, Stern MP, Mitchell BD, Hazuda HP, Patterson JK: Incidence of type II diabetes in Mexican Americans predicted by fasting insulin and glucose level, obesity and body fat distribution. *Diabetes* 39:283–288, 1990
3. Yajnik CS: The insulin resistance epidemic in India: fetal origins, later lifestyle, or both? *Nutr Reviews* 59:1–9, 2001

4. James WP, Chunming C, Inoue S: Appropriate Asian body mass indices? (Editorial). *Obes Rev* 3:139, 2002
5. Yajnik CS, Lubree HG, Rege SS, Naik SS, Deshpande JA, Deshpande SS, Joglekar CV, Yudkin JS: Adiposity and hyperinsulinemia in Indians are present at birth. *J Clin Endocrinol Metab* 87:5575–5580, 2002

Risk Factors of Autonomic and Sensory Nerve Dysfunction in Patients With Newly Diagnosed Type 1 Diabetes

Autonomic neuropathy (AN) in patients with newly diagnosed type 1 diabetes was first described by Fraser et al. (1). Of the six patients examined, two had evidence of AN. In this study, autonomic dysfunction detected during the initial metabolic derangement in newly diagnosed diabetic patients was not reversible after a prolonged period of improved control, indicating that established alterations may occur from the time of diagnosis onward. Other authors (2,3) have also shown that standard cardiovascular reflex tests are able to detect AN in newly diagnosed diabetic patients. A relationship between the severity of AN and prolongation of the corrected QT interval has also been noted (4).

According to the results of the EURO-DIAB IDDM Complications Study, the development of neuropathy is related to cardiovascular risk factors (5). The EURO-DIAB Prospective Complications Study (6) also confirms this finding.

However, there are no data regarding potential risk factors of nerve dysfunction in patients with newly diagnosed type 1 diabetes. We examined 40 patients with newly diagnosed type 1 diabetes with a mean (\pm SD) age of 34.7 ± 11.3 years. The control group comprised 25 healthy subjects (age 38.3 ± 14.8 years). The five standard tests of cardiovascular autonomic function were applied (7). Heart rate tests (heart rate responses to deep breathing, the 30:15 ratio, and the Valsalva ratio) mainly reflect parasympathetic function, while blood pressure responses to sustained handgrip and standing primarily allow the assessment

of sympathetic integrity. The results of each of the five tests were scored as 0 (normal), 1 (borderline), or 2 (abnormal). A final score was calculated (range 0–10) to express the severity of the overall autonomic disorder. Patients with at least one abnormal or two borderline cardiovascular tests (score ≥ 2) were considered to have autonomic neuropathy. Peripheral sensory function was characterized by the evaluation of the current perception threshold (CPT), with a neuroselective diagnostic stimulator (Neurotron, Baltimore, MD), which permits transcutaneous testing (8) at three sinusoidal frequencies (2 kHz, 250 Hz, and 5 Hz). Median and peroneal nerves (digital branches) were studied. All tests were performed after 9 days (range 3–34) of insulin therapy.

As multiple comparisons increase the risk of the error of first kind, we considered results at $P \leq 0.01$ as statistically proven, while those at $P \leq 0.05$ were regarded as marginally significant.

Twelve diabetic patients (30%) had at least one abnormal autonomic function test. Parasympathetic neuropathy was found in six patients, sympathetic nerve dysfunction was observed in three patients, and three subjects had both parasympathetic and sympathetic damage. A significant decrease of the 30:15 ratio (mean \pm SE) was found in diabetic patients compared with control subjects (1.28 ± 0.03 vs. 1.42 ± 0.03 , $P = 0.003$). The autonomic score was higher in diabetic patients (1.08 ± 0.24) than in control subjects (0.17 ± 0.08 , $P = 0.005$).

At least one abnormal sensory parameter was observed in 10 patients (25%). Higher CPT values indicating hypesthesia were found in the diabetic group compared with control subjects at peroneal nerve testing at 250 Hz (1.6 ± 0.1 vs. 1.1 ± 0.07 mA, $P = 0.03$) and 5 Hz (1.1 ± 0.09 vs. 0.6 ± 0.06 , $P = 0.007$), just as at median nerve testing at 5 Hz (0.6 ± 0.03 vs. 0.49 ± 0.05 , $P = 0.048$).

Analyzing the relationship between blood pressure and autonomic function in diabetic patients, the 30:15 ratio correlated significantly negatively with the diastolic blood pressure values ($r = 0.3240$, $P = 0.044$). There was a significant positive relationship between systolic blood pressure and the CPT values testing median nerve at 5 Hz ($r = 0.3988$, $P = 0.012$). The decrease of systolic blood pressure after standing correlated significantly

cantly negatively with CPT values at the peroneal nerve at 2 kHz ($r = -0.3436$, $P = 0.032$), 250 Hz ($r = -0.3893$, $P = 0.014$), and 5 Hz ($r = -0.3273$, $P = 0.042$).

Assessing the relationship between smoking and autonomic function, a significant negative correlation was found between the duration of smoking and the deep breathing test ($r = -0.3452$, $P = 0.006$). The duration of smoking correlated significantly positively with the parasympathetic score ($r = 0.3817$, $P = 0.002$), just as with the autonomic score ($r = 0.3398$, $P = 0.006$). There was a significant correlation between plasma cholesterol and the parasympathetic score ($r = 0.3937$, $P = 0.047$).

A significant negative correlation was observed between the deep breathing test and the CPT values testing the median nerve at 2 kHz ($r = -0.4452$, $P = 0.005$) as well as at 250 Hz ($r = -0.4048$, $P = 0.01$).

In conclusion, autonomic and sensory nerve dysfunction are quite frequent complications in newly diagnosed type 1 diabetic patients and seem to be related to each other. Our data suggest that traditional cardiovascular risk factors (smoking, hypertension, and serum cholesterol) should be considered as potential risk factors for the development of neuropathy, even in newly diagnosed type 1 diabetic patients. These observations may confirm the role of vascular factors in the pathogenesis of neuropathy and may be important for the development of risk reduction strategies.

KATALIN KERESZTES, MD
 ILDIKO ISTENES, MD
 ZSOLT HERMANYI, MD
 PETER VARGHA, MD, PHD
 ISTVAN BARNA, MD, PHD
 PETER KEMPLER MD, PHD, DSC

From the 1st Department of Medicine, Semmelweis University, Budapest, Hungary.

Address correspondence to Dr. Peter Kempler, 1st Department of Medicine, Semmelweis University Budapest, H-1083, Budapest, Koranyi S.u.2/a, Hungary. E-mail: eva.kempet@ediport.hu.

© 2003 by the American Diabetes Association.

References

1. Fraser DM, Campbell IW, Ewing DJ, Murray A, Neilson JMM, Clarke BF: Peripheral and autonomic nerve function in newly diagnosed diabetes mellitus. *Diabetes* 26: 546–550, 1977

- Ziegler D, Dannehl K, Mühlen H, Spüler M, Gries FA: Prevalence of cardiovascular autonomic dysfunction assessed by spectral analysis and standard tests of heart-rate variation in newly diagnosed IDDM patients. *Diabetes Care* 15:908–911, 1992
- Kempler P, Váradi A, Tamás GY: Autonomic neuropathy in newly diagnosed diabetes mellitus (Letter). *Diabetes Care* 16: 848–849, 1993
- Kempler P, Keresztes K, Hermányi ZS, Marton A: Studies must establish whether prolonged QTc interval in newly diagnosed type 1 diabetes is reversible (Letter). *BMJ* 317:678–679, 1998
- Kempler P, Tesfaye S, Chaturvedi N, Stevens LK, Webb DJ, Eaton S, Kerényi Zs, Tamás Gy, Ward JD, Fuller JH, the EURODIAB IDDM Complications Study Group: Autonomic neuropathy is associated with increased cardiovascular risk factors: the EURODIAB IDDM Complications Study. *Diabet Med* 19:900–909, 2002
- The EURODIAB Prospective Complications Study (PSC) Group: Cardiovascular risk factors predict diabetic peripheral neuropathy in type 1 subjects in Europe (Abstract). *Diabetologia* 42 (Suppl. 1):A50, 1999
- Ewing DJ, Clarke BF: Diagnosis and management of diabetic autonomic neuropathy. *BMJ* 28:5916–918, 1982
- Barkai L, Kempler P: Puberty as a risk factor for diabetic neuropathy (Letter). *Diabetes Care* 23:1044–1045, 2000

LDL Electronegativity Is Enhanced in Type 1 Diabetes

LDL particles exhibit heterogeneity in density, size, chemical composition, and charge (1). Lipoperoxidation, oxidation, and glycosylation increase the net negative charge and may enhance LDL atherogenicity with important metabolic consequences. A relevant role of more electronegative LDL in atherogenesis is supported by the observation that it is elevated in subjects at high risk, such as familial hypercholesterolemic and type 1 diabetic patients (2).

We reported the precise measurement of the electrophoretic mobility of LDL as an indicator of modification by capillary electrophoresis and the UV absorption at 234 nm that results from the formation of conjugated dienes in constituent polyenoic fatty acids in 14 type 1 diabetic patients (7 normoalbuminuric and 7 microalbuminuric patients) and in

6 nondiabetic subjects. In type 1 diabetic patients with normoalbuminuria (six men and one woman; mean age 38 ± 12 years) the mean duration of diabetes was 25 ± 7 years, and they were in stable glycemic control ($HbA_{1c} = 7.1 \pm 0.6\%$). The seven diabetic patients with microalbuminuria (six men and one woman; mean age 52 ± 9 years, $P < 0.01$ vs. normoalbuminuric patients) had a mean duration of diabetes of 22 ± 14 years and a mean HbA_{1c} value of $8.8 \pm 1\%$ ($P < 0.01$ vs. normoalbuminuric patients). Diabetic patients had significantly higher BMI (25 ± 2 kg/m²) ($P < 0.01$ for normoalbuminuric subjects, 25 ± 3 kg/m²; $P < 0.05$ for microalbuminuric vs. control group, 21 ± 2 kg/m²) and fasting glucose levels (215 ± 83 mg/dl) ($P < 0.01$ for normoalbuminuric subjects, 197 ± 91 mg/dl; $P < 0.01$ for microalbuminuric vs. control group, 99 ± 18 mg/dl) than control subjects. There was no difference in triglycerides, total cholesterol, LDL cholesterol, and HDL cholesterol levels between diabetic subjects and the control group.

LDL was isolated by preparative sequential ultracentrifugation at the density of 1.063 g/ml. Dialyses, capillary electrophoresis (CE), and the electrophoretic mobility (μ) of LDL were performed as described by Stock and Miller (3). Migration of LDL particles was monitored at 200 and 234 nm. The amount of conjugated dienes is obtained from the percentage of the height of LDL peak at 234 nm related to the height of LDL peak at 200 nm. Student's *t* test and Pearson's correlation were used to assess statistical significance.

The electrophoretic mobility (mean \pm SD) for the diabetic LDL was $-1.249 \pm 0.065 \cdot 10^{-4} \cdot \text{cm}^2 \cdot \text{vol}^{-1} \cdot \text{s}^{-1}$, while that for the control LDL was -1.032 ± 0.121 ($P = 0.0001$). The diabetic group, subdivided into normoalbuminuric and microalbuminuric subjects, presented an electrophoretic mobility mean of -1.234 ± 0.068 and $-1.263 \pm 0.064 \cdot 10^{-4} \cdot \text{cm}^2 \cdot \text{vol}^{-1} \cdot \text{s}^{-1}$, respectively. When each group was compared with the control, the differences were always statistically significant in both cases ($P = 0.0032$ for normoalbuminuric patients vs. control subjects; $P = 0.0001$ for microalbuminuric patients vs. control subjects). Diabetic subjects have LDL with significantly higher migration rates, which were independent from microalbuminuria.

In LDL obtained from the diabetic patients the content of diene conjugates was not statistically different from the control group ($6.22 \pm 1.199\%$ for diabetic subjects vs. $5.509 \pm 0.219\%$ for control subjects).

The difference between diabetic and control subjects was still not statistically significant when the content of diene conjugates in normoalbuminuric ($6.235 \pm 1.544\%$) and microalbuminuric ($6.214 \pm 0.854\%$) subjects was individually compared with that of the control group. In the diabetic group, the electrophoretic mobility was not significantly correlated with HbA_{1c} , duration of diabetes, the subjects' age, or fasting glucose levels.

The finding of electronegative LDL in type 1 diabetic subjects could be related to the increase of the so-called LDL(-), which is also detectable in normal subjects, although in small amounts (4). Capillary electrophoresis cannot separate the fraction LDL(-) from the bulk of plasma LDL. It gives an estimate of the algebraic sum of the electronegative charges distributed on the surface of LDL particles.

Nonenzymatic glycosylation should, surprisingly, be excluded as a cause of higher LDL electronegativity. In this regard, we found no significant correlation between electrophoretic mobility and HbA_{1c} and the fasting plasma glucose levels in the diabetic group. Furthermore, neither the duration of diabetes nor subject age had effects on LDL mobility. Thus, the increased negative charge could be related to compositional abnormalities or other modifications not evaluated in this report, such as an enrichment in sialic acid. Desialylated LDL is more resistant to copper oxidation than native LDL (5).

In conclusion, the finding of more electronegative LDL in diabetic subjects could be an additional risk factor for atherosclerosis in diabetes. Investigations are under way to assess if electrophoretic mobility of LDL in type 1 diabetes can be decreased by further lowering HbA_{1c} levels.

ROBERTO GAMBINO, PHD
SARA GIUNTI, MD
BARBARA UBERTI, PHD
PAOLO CAVALLO PERIN, MD
GIANFRANCO PAGANO, MD
MAURIZIO CASSADER, PHD

From the Dipartimento di Medicina Interna, Università di Torino, Torino, Italy.

Address correspondence to Dr. Maurizio Casader, Dipartimento di Medicina Interna, Università di Torino, Corso AM Dogliotti 14, Torino 10126, Italy.

© 2003 by the American Diabetes Association.

References

1. La Belle M, Blanche PJ, Krauss M: Charge properties of low density lipoprotein subclasses. *J Lipid Res* 38:690–700, 1997
2. Sánchez-Quesada JL, Pérez A, Caixàs A, Ordóñez-Llanos J, Carreras G, Payés A: Electronegative low density lipoprotein subform is increased in patients with short-duration IDDM and is closely related to glycaemic control. *Diabetologia* 39:1469–1476, 1996
3. Stock J, Miller NE: Capillary electrophoresis to monitor the oxidative modification of LDL. *J Lipid Res* 39:1305–1309, 1998
4. Demuth K, Myara I, Chappey B, Vedie B, Pech-Amsellem MA, Haberland ME, Moatti N: A cytotoxic electronegative LDL subfraction is present in human plasma. *Arterioscl Thromb Vasc Biol* 16:773–783, 1996
5. Myara I, Haberland ME, Demuth K, Chappey B, Moatti N: Susceptibility to copper oxidation of neuraminidase-treated LDL. *Clin Chem Acta* 240:221–223, 1995

Effect of Glimepiride on Serum Adiponectin Level in Subjects With Type 2 Diabetes

Sulfonylurea is known to lower glucose levels by stimulating pancreatic insulin secretion. Glimepiride, a new agent of sulfonylurea, is unique in that the glucose-lowering efficacy is similar but the ability to stimulate insulin secretion is lower in comparison with conventional sulfonylureas such as glibenclamide, glipizide, and gliclazide (1). Thus, glimepiride is hypothesized to have greater extrapancreatic effect, such as an improvement in insulin resistance (1). The previous report by Muller et al. (1) supports the hypothesis that insulin-resistant diabetic KK-Ay mice can be well controlled by glimepiride, but not by glibenclamide and gliclazide. Glimepiride is reported to increase insulin-stimulated glycogen synthesis in cultured human skeletal muscle cells. Very recently,

Tsunekawa et al. (2) clearly demonstrated that glimepiride actually increases insulin sensitivity in type 2 diabetic patients. They also proposed that the increase in insulin sensitivity might be associated with increased adiponectinemia. Here we report our data regarding the effects of glimepiride on insulinemia, insulin sensitivity, and serum adiponectin levels in type 2 diabetic subjects. In addition, the effects of glimepiride are compared with those achieved by metformin, which has been proven to have little effect on body weight during glycemic control.

A total of 28 Japanese patients with type 2 diabetes (19 men and 9 women, aged 59 ± 2 years [mean \pm SE], BMI 26.5 ± 0.8 kg/m²) were investigated before and after treatment with glimepiride. The treatment duration was 3 months, and the daily dose of glimepiride was 1.9 ± 0.2 mg (range 1.0–3.0). Changes in indices were analyzed by Wilcoxon's sign-rank test. After the treatment, fasting plasma glucose (166 ± 7 vs. 147 ± 7 mg/dl, $P = 0.009$) and HbA_{1c} (7.9 ± 0.3 vs. $7.4 \pm 0.2\%$, $P = 0.006$) levels fell significantly. Both fasting insulin (11.7 ± 1.5 vs. 9.4 ± 1.0 μ U/ml, $P = 0.007$) and homeostasis model assessment for insulin resistance (HOMA-IR) (3) (5.0 ± 0.8 vs. 3.8 ± 0.6 , $P = 0.005$) decreased, suggesting an amelioration of insulin resistance. Serum adiponectin concentration, measured by Linco RIA kits (St. Charles, MO), increased significantly (22.1 ± 2.7 vs. 28.5 ± 2.8 μ g/ml, +29%, $P = 0.015$), whereas no significant change was observed in BMI (26.5 ± 0.9 vs. 26.5 ± 0.8 kg/m², $P = 0.748$). There was also no change in serum concentrations of total (214 ± 6 vs. 210 ± 6 mg/dl, $P = 0.125$) and HDL (54 ± 3 vs. 53 ± 3 mg/dl, $P = 0.438$) cholesterol and triglyceride (123 ± 10 vs. 120 ± 10 mg/dl, $P = 0.387$) before and after the treatment.

In a separate group of type 2 diabetic patients matched with the glimepiride group for sex, age, BMI, glycemia, and insulinemia (seven men and five women, aged 58 ± 3 years, and BMI 25.7 ± 0.7 kg/m²), the effect of metformin (daily dose 750 mg) was evaluated. Three months of the metformin treatment also decreased both fasting glucose (159 ± 4 to 135 ± 4 mg/dl, $P = 0.006$) and HbA_{1c} (7.9 ± 0.2 to $7.1 \pm 0.2\%$, $P = 0.013$) levels. In contrast to the glimepiride treatment, fasting insulin (12.4 ± 2.0 vs. 13.8 ± 4.3 μ U/ml, $P = 0.689$) and

HOMA-IR (4.8 ± 0.7 vs. 4.1 ± 1.0 , $P = 0.695$) remained unchanged, whereas serum adiponectin concentration was increased slightly but significantly (18.7 ± 3.0 vs. 20.6 ± 3.3 μ g/ml, +10%, $P = 0.034$). No significant change was observed in BMI and serum lipid concentrations (data not shown).

Our present finding supports the notion that one of the glucose-lowering mechanisms of glimepiride is to improve insulin resistance. In accordance with a recent article (2), the glimepiride treatment increased serum adiponectin levels without affecting BMI. In the present study, serum adiponectin levels were also increased by the treatment of metformin, which, unlike insulin-sensitizing thiazolidinediones, is known to not affect circulating adiponectin concentration (4). Therefore, it seems possible that the increase in adiponectinemia by the glimepiride treatment could be, in part, due to an effect of glycemic control per se. Another difference between the glimepiride and metformin groups is the change in insulinemia; fasting insulin was decreased in the former group and unchanged in the latter. Since insulin seems to suppress expression and secretion of adiponectin in both in vitro and in vivo studies (5,6), the decrease in insulinemia by glimepiride may conversely increase circulating adiponectin concentration.

We agree that the improvement in glycemic control, insulinemia, and adiponectinemia by glimepiride is of potential benefit to decrease risk factors of atherosclerosis in type 2 diabetic patients. The mechanisms of the increased adiponectinemia by glimepiride may be complex and multifactorial. It also remains to be elucidated whether conventional sulfonylureas would increase adiponectinemia in subjects with type 2 diabetes.

SHOICHIRO NAGASAKA, MD¹
 ATARU TANIGUCHI, MD²
 YOSHITAKA AISO, MD³
 TOSHIMITSU YATAGAI, MD¹
 TOMOATSU NAKAMURA, MD¹
 YOSHIKATSU NAKAI, MD⁴
 MITSUO FUKUSHIMA, MD⁵
 AKIRA KUROE, MD²
 SHUN ISHIBASHI, MD¹

From the ¹Division of Endocrinology and Metabolism, Jichi Medical School, Tochigi, Japan; the ²Division of Diabetes, Kansai-Denryoku Hospital, Osaka, Japan; the ³Aiso Clinic, Tokyo, Japan; the ⁴College of Medical Technology, Kyoto University,

Kyoto, Japan; and the ⁵Department of Diabetes and Clinical Nutrition, Graduate School of Medicine, Kyoto University, Kyoto, Japan.

Address correspondence to Shoichiro Nagasaka, MD, Division of Endocrinology and Metabolism, Jichi Medical School, Yakushiji 3311-1, Minami-kawachi, Tochigi 329-0498, Japan. E-mail: sngsk@jichi.ac.jp.

© 2003 by the American Diabetes Association.

References

- Muller G, Satoh Y, Geisen K: Extrapancratic effects of sulfonyleureas: a comparison between glimepiride and conventional sulfonyleureas (Review). *Diabetes Res Clin Pract* 28:S115–S137, 1995
- Tsunekawa T, Hayashi T, Suzuki Y, Matsui-Hirai H, Kano H, Fukatsu A, Nomura N, Miyazaki A, Iguchi A: Plasma adiponectin plays an important role in improving insulin resistance with glimepiride in elderly type 2 diabetic patients. *Diabetes Care* 26:285–289, 2003
- Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC: Homeostasis model assessment: insulin resistance and β -cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 28:412–419, 1985
- Combs TP, Wagner JA, Berger J, Doebber T, Wang W-J, Zhang BB, Tanen M, Berg AH, O'Rahilly S, Savage DB, Chatterjee K, Weiss S, Larson PJ, Gottesdiener KM, Gertz BJ, Charron MJ, Scherer PE, Moller DE: Induction of adipocyte complement-related protein of 30 kilodaltons by PPAR γ agonists: a potential mechanism of insulin sensitization. *Endocrinology* 143:998–1007, 2002
- Fasshauer M, Klein J, Neumann S, Eszlinger M, Paschke R: Hormonal regulation of adiponectin gene expression in 3T3-L1 adipocytes. *Biochem Biophys Res Commun* 290:1084–1089, 2002
- Yu JG, Javorschi S, Hevener AL, Kruszynska YT, Norman RA, Sinha M, Olefsky JM: The effect of thiazolidinediones on plasma adiponectin levels in normal, obese, and type 2 diabetic subjects. *Diabetes* 51:2968–2974, 2002

Glargine Insulin Is Not an Alternative in Insulin Allergy

Allergy to insulin is rare with human recombinant insulin and is now reported for <1% of diabetic patients. Clinic symptoms are usually local and appear a few minutes after the injection

(red blotch, induration, pruritus, and burning sensation at insulin injection sites) and are rarely general (from urticaria to anaphylactic shock). A decrease of the efficiency of the insulin is usually associated with these symptoms. Different methods have been proposed for the treatment of insulin allergy including the use of oral antihistaminics, the addition of glucocorticoids to insulin, and the change to human insulin analogues.

To our knowledge, we report the first case of allergy to a new long-acting insulin analogue, insulin glargine

An 81-year-old man with type 2 diabetes was admitted for uncontrolled diabetes and insulin initiation. He had a coronary artery bypass 2 years previous and recently had a critical limb ischemia. The patient had no history of any allergy. He was first treated by Mixtard 30 twice daily (Innolet; Novo Nordisk). The patient presented local induration and pruritus at insulin injection site and general urticarian lesions from 10 to 15 min after the injection. An allergy to insulin was then suspected.

Skin-prick tests (5 UI/ml) were positive for human and porcine insulin and negative for all additives (protamine, paraben, metacresol, phenol, zinc, and isophane) using the Novo Insulin Allergy Kit (Novo Nordisk). These tests confirmed the allergy to insulin.

To test the possibility of treating the patient with rapid-acting insulin analogs, we examined skin reactions to aspart and lispro insulin. We have therefore performed additive skin-prick tests with aspart and lispro insulins. They were positive for lispro and negative for aspart insulin. Similar results with insulin analogues have been previously reported (1).

A treatment with subcutaneous continuous aspart insulin infusion was then initiated. No local reaction was observed, and glycemic control gradually improved. However, a prolonged treatment with an insulin pump was very difficult for this older patient. We therefore decided to test glargine insulin, a new long-acting human insulin analogue. Unfortunately, skin tests were very positive with glargine insulin. To our knowledge, we report the first case of allergy to this new insulin.

Human insulin analogues, lispro or aspart, have been proposed for the treatment of insulin allergy (1,2). Allergy to lispro, aspart, or both has been recently reported (1–3). In our observation, we es-

tablish a similar case with lispro but not with aspart insulins. However, to our knowledge, we report the first case of allergy with glargine insulin.

KARINE-NOELLE DURAND-GONZALEZ, MD¹
NICOLAS GUILLAUSSEAU, MD¹
CATHERINE PECQUET, MD²
JEAN-PIERRE GAYNO, MD¹

From the ¹Department of Endocrinology, Poissy-Saint Germain en Laye Hospital, Laye, France; and the ²Department of Allergology, Tenon Hospital, Tenon, France.

Address correspondence to Dr. Durand-Gonzalez, Service de Diabétologie et Endocrinologie, 18 rue Armagis, 78100 Saint Germain en Laye, France. E-mail: durander@club-internet.fr.

© 2003 by the American Diabetes Association.

References

- Airaghi L, Lorini M, Tedeschi A: The insulin analog aspart: a safe alternative in insulin allergy (Letter). *Diabetes Care* 24:2000, 2001
- Abraham MR, Al-Sharafi BA, Saavedra GA, Khardori R: Lispro in the treatment of insulin allergy. *Diabetes Care* 22:1916–1917, 1999
- Takata H, Kumon Y, Osaki F, Kumagai C, Arai K, Ikeda Y, Suehiro T, Hashimoto K: The human insulin analogue aspart is not the almighty solution for insulin allergy. *Diabetes Care* 26:253–254, 2003

Association Between Endothelial Nitric Oxide Synthase Glu298Asp Polymorphism and Postchallenge Insulin Levels in Nondiabetic Japanese Subjects

Endothelial nitric oxide synthase (eNOS) catalyzes NO production in vascular endothelial cells, and NO regulates local blood flow by inducing vasodilation (1). Enhancement of skeletal muscle glucose uptake occurs during elevation of muscle blood flow, which is induced by increased eNOS expression and activated by insulin stimulation (2–4). It was previously reported that low production of NO in eNOS knockout mice causes reduction of insulin-induced

blood flow and glucose uptake in whole body (5). These findings suggest that eNOS plays an important role in the regulation of insulin-induced glucose uptake in whole body.

Polymorphism in eNOS exon 7 with G→T conversion at nucleotide position 894 results in amino acid substitution of glutamic acid for aspartic acid in amino acid residue 298 (Glu298Asp). Structural alteration in this variant affects the susceptibility to cleavage and reduces activity of this enzyme (6,7). Dysfunction of eNOS by this polymorphism may cause reduction of insulin-induced blood flow and glucose uptake. It was recently demonstrated that insulin secretion and sensitivity could be assessed by 75-g oral glucose tolerance test (8,9). Therefore, we examined the association of this polymorphism with fasting and postchallenge glucose and insulin levels in nondiabetic Japanese subjects by the 75-g oral glucose tolerance test.

This study comprised 247 Japanese nondiabetic volunteers (69 men and 178 women). Written informed consent was obtained from all subjects enrolled in this study. A 75-g glucose tolerance test was performed early in the morning after fasting overnight. Venous sampling was obtained before loading (0 min), at 30 min, and 120 min after glucose loading, and blood glucose and insulin levels were measured. The serum insulin levels were measured with an EIA kit (Eiken insulin kit; EIA, Tokyo, Japan). All subjects were nondiabetic according to American Diabetes Association criteria (10). For assessing the substitution of G→T at position 894 (Glu298Asp), genomic DNA isolated from peripheral blood leukocytes was amplified by PCR and digested with *Ban*II restriction enzyme as previously described (11,12). Data are expressed as means ± SEM. Differences between each group were tested by two-tailed unpaired Student's *t* test. A *P* < 0.05 was considered as statistically significant.

The allele frequency was 0.927 for G and 0.073 for T in all subjects. Genotype distribution was 86.6% (214 of 247) for Glu/Glu, 12.2% (30 of 247) for Glu/Asp, and 1.2% (3 of 247) for Asp/Asp. The frequency and distribution are compatible with previous data (11,12). Because the number of homozygous mutants was too small, the combined data from homozygous and heterozygous individuals were used in the following analysis. There was

no significant difference in age (53.8 ± 0.5 vs. 56.0 ± 1.6 years), BMI (23.4 ± 0.2 vs. 24.4 ± 0.7 kg/m²), waist-to-hip ratio (1.01 ± 0.01 vs. 0.98 ± 0.04), systolic blood pressure (130.0 ± 1.3 vs. 130.1 ± 3.4 mmHg), diastolic blood pressure (77.6 ± 0.8 vs. 77.2 ± 2.0 mmHg), total cholesterol (5.46 ± 0.10 vs. 5.58 ± 0.21 mmol/l), triglyceride (1.28 ± 0.05 vs. 1.16 ± 0.10 mmol/l), HDL cholesterol (1.62 ± 0.03 vs. 1.63 ± 0.09 mmol/l), or HbA_{1c} (5.0 ± 0.1 vs. $5.0 \pm 0.2\%$) between Glu/Glu and Glu/Asp + Asp/Asp. The results of the glucose tolerance test were as follows: plasma glucose in Glu/Glu was not significantly different from Glu/Asp + Asp/Asp at 0 min (5.07 ± 0.05 vs. 5.12 ± 0.12 mmol/l), 30 min (8.36 ± 0.16 vs. 8.94 ± 0.30 mmol/l), and 120 min (6.36 ± 0.18 vs. 6.93 ± 0.35 mmol/l). However, serum insulin levels were significantly increased in Glu/Asp + Asp/Asp compared with Glu/Glu at 30 min (309.6 ± 40.8 vs. 236.4 ± 9.6 pmol/l, *P* < 0.02) and 120 min (342.0 ± 36.0 vs. 220.2 ± 10.8 pmol/l, *P* < 0.0005). There was no significant difference in serum insulin levels at 0 min (35.2 ± 1.2 vs. 41.8 ± 3.0 pmol/l), homeostasis model assessment for insulin resistance (HOMA-IR) (1.36 ± 0.06 vs. 1.59 ± 0.12), and insulinogenic index ($\Delta I_{30}/\Delta G_{30}$, 1.09 ± 0.22 vs. 0.74 ± 0.12) between Glu/Glu and Glu/Asp + Asp/Asp.

In the present study, elevation of insulin levels at 30 and 120 min after glucose loading test was observed in subjects with Glu/Asp + Asp/Asp polymorphism compared with wild-type. However, the blood levels of glucose were not significantly different between these two groups. These data showed that there is a remarkable difference in postchallenge insulin levels between Glu/Glu and Glu/Asp + Asp/Asp groups. Subjects with Glu/Asp + Asp/Asp require more insulin to maintain the same glucose levels than subjects with Glu/Glu during glucose loading test. It was reported (8,9) that insulin level during postchallenge (120 min) is correlated with insulin sensitivity as measured by the glucose clamp method in nondiabetic subjects. Thus, one explanation for the elevated postchallenge (120 min) insulin levels may be reduced insulin sensitivity due to impaired insulin-mediated local blood flow in subjects with Glu/Asp + Asp/Asp polymorphism.

However, no significant difference was observed in HOMA-IR, a marker of

insulin sensitivity, between subjects with Glu/Asp + Asp/Asp and those with Glu/Glu, suggesting that another mechanism in addition to insulin sensitivity may affect postchallenge insulin levels. It was previously observed (13) that decreased insulin-mediated blood flow in muscles is associated with reduction of insulin clearance in obese subjects with insulin resistance. Also, remarkable difference in blood flow and insulin clearance has been observed between lean and obese subjects in hyperinsulinemic conditions (13). These observations suggest that decreased insulin-mediated blood flow reduces insulin clearance, which leads to increased circulating insulin levels. Therefore, impairment of insulin-mediated vasodilation with subsequent reduction of insulin clearance may be another explanation for the changes of postchallenge insulin levels in subjects with the Glu/Asp + Asp/Asp polymorphism.

In conclusion, the present study showed for the first time that eNOS Glu289Asp polymorphism affects postchallenge insulin levels in nondiabetic Japanese subjects.

NORIKO MARUYAMA, MD¹

YUTAKA YANO, MD¹

ESTEBAN C. GABAZZA, MD¹

RIKA ARAKI, MD¹

AKIRA KATSUKI, MD¹

YASUKO HORI, MD¹

KANAME NAKATANI, MD²

YASUHIRO SUMIDA, MD¹

YUKIHIKO ADACHI, MD¹

From the ¹Third Department of Internal Medicine, Mie University School of Medicine, Tsu, Mie, Japan; and the ²Department of Laboratory Medicine, Mie University School of Medicine, Tsu, Mie, Japan.

Address correspondence to Dr. Yutaka Yano, Third Department of Internal Medicine, Mie University School of Medicine, Edobashi 2-174, Tsu, Mie 514-8507, Japan. E-mail: yanoyuta@clin.medic.mie-u.ac.jp.

© 2003 by the American Diabetes Association.

References

1. Ignarro LJ, Cirino G, Casini A, Napoli C: Nitric oxide as a signaling molecule in the vascular system: an overview. *J Cardiovasc Pharmacol* 34:879–886, 1999
2. Kuboki K, Jiang ZY, Takahara N, Ha SW, Igarashi M, Yamauchi T, Feener EP, Herbert TP, Rhodes CJ, King GL: Regulation of endothelial constitutive nitric oxide synthase gene expression in endothelial cells and in vivo: a specific vascular action of insulin. *Circulation* 101:676–681, 2000

3. Steinberg HO, Brechtel G, Johnson A, Fineberg N, Baron AD: Insulin-mediated skeletal muscle vasodilation is nitric oxide dependent: a novel action of insulin to increase nitric oxide release. *J Clin Invest* 94:1172–1179, 1994
4. Baron AD, Brechtel-Hook G, Johnson A, Cronin J, Leaming R, Steinberg HO: Effect of perfusion rate on the time course of insulin-mediated skeletal muscle glucose uptake. *Am J Physiol* 271:E1067–E1072, 1996
5. Duplain H, Burcelin R, Sartori C, Cook S, Egli M, Lepori M, Vollenweider P, Pedrazzini T, Nicod P, Thorens B, Scherrer U: Insulin resistance, hyperlipidemia, and hypertension in mice lacking endothelial nitric oxide synthase. *Circulation* 104:342–345, 2001
6. Tesouro M, Thompson WC, Rogliani P, Qi L, Chaudhary PP, Moss J: Intracellular processing of endothelial nitric oxide synthase isoforms associated with differences in severity of cardiopulmonary diseases: cleavage of proteins with aspartate vs. glutamate at position 298. *Proc Natl Acad Sci U S A* 97:2832–2835, 2000
7. Wang XL, Sim AS, Wang MX, Murrell GA, Trudinger B, Wang J: Genotype dependent and cigarette specific effects on endothelial nitric oxide synthase gene expression and enzyme activity. *FEBS Lett* 471:45–50, 2000
8. Stumvoll M, Mitrakou A, Pimenta W, Jenssen T, Yki-Jarvinen H, Van Haeflén T, Renn W, Gerich J: Use of the oral glucose tolerance test to assess insulin release and insulin sensitivity. *Diabetes Care* 23:295–301, 2000
9. Hanson RL, Pratley RE, Bogardus C, Narayan KM, Roumain JM, Imperatore G, Fagot-Campagna A, Pettitt DJ, Bennett PH, Knowler WC: Evaluation of simple indices of insulin sensitivity and insulin secretion for use in epidemiologic studies. *Am J Epidemiol* 151:190–198, 2000
10. The Expert Committee on the Diagnosis and Classification of Diabetes Mellitus: Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care* 20:1183–1197, 1997
11. Ohtoshi K, Yamasaki Y, Gorogawa S, Hayaishi-Okano R, Node K, Matsuhisa M, Kajimoto Y, Hori M: Association of (-)786T-C mutation of endothelial nitric oxide synthase gene with insulin resistance. *Diabetologia* 45:1594–1601, 2002
12. Miyamoto Y, Saito Y, Kajiyama N, Yoshimura M, Shimasaki Y, Nakayama M, Kamitani S, Harada M, Ishikawa M, Kuwahara K, Ogawa E, Hamanaka I, Takahashi N, Kaneshige T, Teraoka H, Akamizu T, Azuma N, Yoshimasa Y, Yoshimasa T, Itoh H, Masuda I, Yasue H, Nakao K: Endothelial nitric oxide synthase gene is positively associated with essential hypertension. *Hypertension* 32:3–8, 1998
13. Sjöstrand M, Gudbjörnsdóttir S, Holmäng A, Lönn L, Strindberg L, Lönnroth P: Delayed transcapillary transport of insulin to muscle interstitial fluid in obese subjects. *Diabetes* 51:2742–2748, 2002

A Diabetic Subject With MELAS and Antiphospholipid Syndrome

Mitochondrial encephalopathy, lactic acidosis, and stroke-like episodes (MELAS) syndrome has been reported to coexist with autoimmune type 1 diabetes (1) and Graves' disease (2). We described, for the first time, a diabetic patient with MELAS syndrome, autoimmune hemolytic anemia, and antiphospholipid syndrome.

During routine health examination, a 38-year-old man was diagnosed with diabetes. There was no family history of diabetes. On 22 January 2001, he was admitted due to right hemiparesis, slurred speech, and headache. Magnetic resonance imaging of the brain revealed increased signal intensity on diffusion scan with decreased apparent diffusion coefficient confined to left middle cerebral arterial territory, which was compatible with acute ischemic infarct. For his young stroke, we checked carotid duplex, cardiac sonography, anti-nuclear antibodies, rheumatic factors, protein C, and protein S, which were all negative. The level of antiphospholipid antibodies was 14.7 phospholipid units/ml (normal <5, positive >15) and that of anticardiolipin antibodies was 17.5 phospholipid units/ml (normal <16, positive >21). He was then discharged in stable condition. Serum markers were repeated on 16 July 2001 and showed positivity for antiphospholipid antibodies (19.5 phospholipid units/ml) and anticardiolipin antibodies (22.4 phospholipid units/ml). Antiphospholipid syndrome was favored due to a history of vascular thrombosis and the presence of circulating antibodies. However, gradual loss of cognition and muscle power developed progressively. Bilateral hearing impairment was found, and the patient was once again admitted for hyperglycemia with metabolic acidosis on

11 July 2001. Serum lactic acid was high (6.95 mmol/l). The presence of lactic acidosis, bilateral hearing loss, progressive muscle weakness, young stroke, and diabetes prompted us to examine him for mitochondrial disease. MELAS syndrome was then diagnosed with a demonstration of an A-to-G point mutation at position 3243 of mitochondrial DNA. No such a mutation was found in his mother or siblings, indicating that a de novo mutation occurred in this subject. Besides, autoimmune hemolytic anemia was also found during admission (hemoglobin 9.0 g/dl, mean corpuscular volume 91.6 fl, reticulocyte count 10.81%, haptoglobin <5.83 mg/dl, direct Coombs test 2+, antinuclear antibodies [–] 1:160, and rheumatoid factor <20 IU/ml). His condition deteriorated progressively, and he died 5 months later of pneumonia.

It is difficult to distinguish patients with antiphospholipid syndrome from MELAS syndrome based on brain image studies. Patients with antiphospholipid syndrome may suffer from oxidant-mediated injury (3). Since the mitochondrial genome lacks a DNA repairing system and protecting proteins, it is susceptible to oxidative stress. Thus, the presence of antiphospholipid syndrome might be one of the causes of de novo mutation or may accelerate accumulation of mutated DNA, which may result in a rapidly deteriorating course such as that seen in this patient. We therefore suggest that patients with MELAS syndrome be examined for the presence of antiphospholipid syndrome, especially those presenting with vascular thrombosis.

HUNG-YUAN LI, MD¹

WEI-SHIUNG YANG, MD, PHD^{1,2}

TONG-YUAN TAI, MD, PHD¹

LEE-MING CHUANG, MD, PHD^{1,2}

From the ¹Department of Internal Medicine, National Taiwan University Hospital, Taipei, Taiwan; and the ²Graduate Institute of Clinical Medicine, National Taiwan University Medical College, Taipei, Taiwan.

Address correspondence to Lee-Ming Chuang, MD, PhD, Department of Internal Medicine, National Taiwan University Hospital, 7 Chung Shan S. Rd., Taipei, Taiwan, R.O.C. E-mail: leeming@ha.mc.ntu.edu.tw.

© 2003 by the American Diabetes Association.

References

1. Huang CN, Jee SH, Hwang JJ, Kuo YF, Chuang LM: Autoimmune IDDM in a

- sporadic MELAS patient with mitochondrial tRNA Leu(UUR) mutation. *Clin Endocrinol* 49:265–270, 1998
2. Yang CY, Lam HC, Lee HC, Wei YH, Lu CC, Han TM, Tsai JL, Chuang YH, Lee JK: MELAS syndrome associated with diabetes mellitus and hyperthyroidism: a case report from Taiwan. *Clin Endocrinol* 43: 235–239, 1995
 3. Levine JS, Branch DW, Rauch J: The antiphospholipid syndrome. *N Engl J Med* 346:752–763, 2002

High Degree of Mitochondrial 3243 Mutation in Gastric Biopsy Specimen in a Patient With MELAS and Diabetes Complicated by Marked Gastrointestinal Abnormalities

A point mutation of mitochondrial DNA at nucleotide position 3243 has been shown to cause mitochondrial encephalomyopathy with lactic acidosis and stroke-like episodes (MELAS) (1). This mutation, however, is also found in maternally inherited diabetes and deafness (MIDD) (2), which accounts for ~1% of the diabetic population in Japan (3). The same point mutation of mitochondrial DNA causes a wide range of symptoms that have been suggested to be due to the difference in the degree of heteroplasmy; thus, the proportion of the mutant mitochondrial DNA is divergent among different tissues (4). Little evidence, however, is available due to difficulty in obtaining viable samples from active lesions associated with complications causing symptoms. In a diabetic patient with MELAS and severe gastrointestinal disease, including functional ileus, duodenal ulcer, and acute gastric mucosal lesions, which were resistant to treatment, we had a rare opportunity to investigate the degree of heteroplasmy of the 3243 mutation in a biopsy specimen of gastric mucosa, tissue with a major lesion that causes gastrointestinal complications, compared with peripheral white blood cells.

A 21-year-old woman had diabetes (HbA_{1c} 7.3%), bilateral hearing loss,

muscle weakness, and several stroke-like episodes with high serum lactate and pyruvate levels (lactate 7.4 mmol/l and pyruvate 261 μmol/l). Neurological findings showed bilateral sensory hearing loss, bilateral external ophthalmoplegia and droopy eyelids, muscle weakness (proximal > distal), and cerebellar ataxia. Magnetic resonance imaging scans showed cerebellar atrophy and mild cerebral atrophy. Her mother, grandmother, and mother's brother also had hearing loss, but there was no obvious family history of diabetes. She also had functional ileus, duodenal ulcer, and acute gastric mucosal lesions, which were resistant to treatment. She was diagnosed as having MELAS with diabetes and gastrointestinal disease. DNA extracted from peripheral blood cells from the patient, her mother, and her elder sister was positive for the 3243 mutation. The proportion of the mutated allele in the proband (39%) was much higher than that in her sister (19%) and mother (10%). We also analyzed mitochondrial DNA of biopsy specimens of her gastric mucosa, which exhibited a higher proportion (57%) of the mutated allele than her peripheral white blood cells (39%), suggesting that her gastrointestinal complications were attributable to a high proportion of the mitochondrial variant in the gastrointestinal tract.

A high degree of mutated mitochondria in the gastrointestinal tract was previously reported in a case without typical clinical features of MELAS but with diabetes and gastrointestinal symptoms (5). There was 70% heteroplasmy of the mutation in his gastrointestinal tract, while that in peripheral white blood cells was 37% (5). Taking these findings together, it is likely that gastrointestinal symptoms in patients with the 3243 mutation, with either MELAS or MIDD, correlate with the degree of heteroplasmy in the gastrointestinal tract. These observations provide further evidence that the clinical diversity of symptoms related to mitochondrial 3243 mutation may be due to diversity in the proportion of the mutation in each organ.

KAORI INOUE, MD
HIROSHI IKEGAMI, MD, PHD
TOMOMI FUJISAWA, MD, PHD
YUMIKO KAWABATA, MD, PHD
MAKI SHINTANI, MD, PHD
KOJI NIJIMA, MD
ONO MASAYA, MD

MASANORI NISHINO, MD
MICHIKO ITOI-BABAYA, MD
NARU BABAYA, MD, PHD
TOSHIO OGIHARA, MD, PHD

From the Department of Geriatric Medicine, Osaka University Graduate School of Medicine, Osaka, Japan.

Address correspondence to Hiroshi Ikegami, Department of Geriatric Medicine, Osaka University Graduate School of Medicine 2-2, Yamadaoka, Suita, Osaka, 565-0871, Japan. E-mail: ikegami@geriat.med.osaka-u.ac.jp.

© 2003 by the American Diabetes Association.

References

1. Kari M, Jacca SM, Seija U, Anne MR: Epidemiology of A3243G, the mutation for mitochondrial encephalomyopathy, lactic acidosis and stroke-like episodes (MELAS): prevalence of the mutation in adult population. *Am J Hum Genet* 63: 447–454, 1998
2. van-den-Oueland JM, Lemkes HH, Trembath RC, Ross R, Velho G, Massen JA: Maternally inherited diabetes and deafness is a distinct subtype of diabetes and associates with a single point mutation in the mitochondrial tRNA^{Leu (UUR)} gene. *Diabetes* 43:746–751, 1994
3. Kadowaki T, Kadowaki H, Mori Y, Tobe K, Sakuta R, Yazaki Y: A subtype of diabetes mellitus associated with a mutation of mitochondrial DNA. *N Engl J Med* 330: 962–968, 1994
4. Matsuo O, Ikuya N, Yuichi G: Single muscle fiber analysis in patients with 3243 mutation in mitochondrial DNA: comparison with the phenotype and the proportion of mutant gene. *J Neurol Sci* 159:170–175, 1998
5. Kishimoto M, Hashiramoto M, Kanda F, Tanaka M: Mitochondrial mutation in diabetic patient with gastrointestinal symptoms (Letter). *Lancet* 345:452, 1995

Frequency of Diabetes Transmission From Two Type 1 Diabetic Parents to Their Children

There is little information in the literature about the risk of diabetes in children of type 1 diabetic parents (mother and father affected), and evidence is based on small numbers (1). Analysis of a larger number of such trios could importantly contribute to the clar-

ification of type 1 diabetes inheritance. In the Karlsburg Clinic for Diabetes and Metabolic Diseases, 77 offspring of 61 pairs with type 1 diabetes have registered since 1955. Until 1989, the parents and their children (diabetic and nondiabetic) were repeatedly treated or checked for diabetes in our clinic; since 1990 some of the patients have visited the clinic or have been contacted as outpatients. The diagnosis of type 1 diabetes in the affected individuals is proofed by clinical and in many cases laboratory data (C-peptide, antibodies, and HLA), as is obvious from our previous study (2).

Here we present data of 58 offspring from 46 pairs with type 1 diabetes that were last contacted in December of 2001. There are 35 daughters and 23 sons born between 1955 and 1993. Until now, 25 of 58 (43%) descendants developed diabetes at ages 1–42 years (mean \pm SD 11 \pm 10). The diabetes incidence did not increase during the observation time in 10-year intervals [54 (7 of 13), 42 (5 of 12), and 50 (11 of 22), respectively, and so far 18% (2 of 11) in those born 1986–1995]. The difference in the diabetes affection of sons (12 of 23) and daughters (13 of 35) is not significant. A strong preponderance of female descendants (18 female and 7 male) was seen in children born before 1975, whereas the sex ratio is nearly one (18 female and 15 male) for those born after 1975.

We observed a dramatic decrease in the age of diabetes onset. Children born during 1955–1975 developed the disease at 1–42 years (mean 15.1 \pm 12.1), and children born during 1975–1995 developed the disease at 2–13 years (7.3 \pm 5.2) ($P = 0.0449$). The mean age of mothers when they gave birth to their children was 24 years in both periods.

In children born before 1975 the mean age of diabetes manifestation (15.1 \pm 12.1 years) is not different from that of their parents (fathers 16 \pm 4.9 years and mothers 16.2 \pm 7.4 years), but in children born after 1976 (7.3 \pm 5.2) it is significantly ($P = 0.014$) younger than that of their fathers (13.6 \pm 5.9 years) but not of their mothers (11 \pm 6.8 years).

The constant diabetes incidence of >40% over >40 years of observation time implies the definite role of a genetic background for type 1 diabetes development. The high diabetes frequency of 43% (near half) in a representative number of children of two type 1 diabetic par-

ents points, according to Mendelian laws, to a small number of responsible genes. On the other hand, we believe that these data demonstrate the influence of the altered environment over time. We previously described (3) a disturbed proportion of male to female (1:2) in children of diabetic mothers, but not of fathers. We now show that this shifted sex ratio does not exist in offspring of diabetic mothers (and fathers) born in the last 20 years. As most children were born in the Karlsburg hospital, we know that the therapeutical principles of diabetes treatment during pregnancy changed to normoglycemia in the late 1970s (4,5). As described in the Swedish study (6), we also confirm the dramatic shift to younger age at diagnosis for this subgroup of German diabetic children.

IIONA RJASANOWSKI, MD¹
INGRID KÖLTING, PHD²
WOLFGANG KERNER, MD¹

From the ¹Heart and Diabetes Center Mecklenburg-Vorpommern, Department of Diabetes and Diseases of Metabolism, Karlsburg, Germany; and the ²University of Greifswald, Department of Laboratory Animal Science, Medical Faculty, Karlsburg, Germany.

Address correspondence to Wolfgang Kerner, Klinikum Karlsburg, Klinik fuer Diabetes, Greifswalder Str. 11, Karlsburg D-17495, Germany. E-mail: wkerner@mail.uni-greifswald.de.

© 2003 by the American Diabetes Association.



References

1. Köbberling J, Tillil H: Risk to Family members of Becoming Diabetic. A Study of the Genetics of Type 1 Diabetes. In *Future Trends in Juvenile Diabetes, Therapy and Research*. Laron Z, Karp M, Tikva P, Eds. Basel, Karger, 1986, p. 26–38
2. Rjasanowski I, Vogt L, Michaelis D, Keilacker H, Kohnert K: The frequency of diabetes in children of type 1 diabetic parents. *Diabete Metab* 19:173–177, 1993
3. Rjasanowski I, Klötting I, Kovacs P: Altered sex ratio in offspring of mothers with insulin-dependent diabetes mellitus. *Lancet* 351:497–498, 1998
4. Gödel E, Amendt P, Amendt U, Festge B, Fett KD, Glöckner E, Jutzi E, Kublun G, Bibergeil H, Bruns W: Diabetes in pregnancy: a clinical and statistical analysis of 1800 pregnancies and deliveries from 1952–1971. 1. Perinatal mortality in relation to the classification of White and delivery date. *Fortschr Geburtshilfe Gynakol* 54:33–44, 1975
5. Fuhrmann K, Reiher H, Semmler K, Fischer F, Fischer M, Glockner E: Prevention of congenital malformations in

- infants of insulin-dependent diabetic mothers. *Diabetes Care* 6:219–223, 1983
6. Pundziute-Lyckå A, Dahlquist G, Nyström L, Amquist H, Björk E, Blohmé G, Bolinder J, Eriksson JW, Sundquist G, Östmann J: The incidence of type 1 diabetes has not increased but shifted to a younger age at diagnosis in the 0–34 years group in Sweden 1983 to 1998. *Diabetologia* 45:783–791, 2002

Putting Diabetes to the Test

Analyzing glycemic control based on patients' diabetes knowledge

Diabetes has reached epidemic proportions. Most of the morbidity, mortality, and cost of type 2 diabetes is related to cardiovascular complications (1). These complications can be decreased by improving glycemic control (2,3). However, one problem in diabetes care is the poor translation of knowledge derived from research to clinical practice (4). A lack of patients understanding the long-term ramifications of poor glycemic control may play a role in this. To further test this hypothesis, we performed the following study to determine whether patient diabetes knowledge was related to glycemic control.

Diabetes knowledge was measured using the Michigan Diabetes Knowledge Test (J.T. Fitzgerald, University of Michigan, Ann Arbor, MI) (5). Only questions pertaining to type 2 diabetes were used. HbA_{1c} values were determined. Data were analyzed using SAS software version 6.1 (SAS, Cary, NC). Logistic regression analysis was done using Pearson's correlation coefficient.

Seventy-seven subjects completed the study. The average number of questions answered correctly was 8.5 \pm 2.3. The mean HbA_{1c} value was 8.05 \pm 1.6%. Regression analysis demonstrated an inverse correlation between the number of correct responses and HbA_{1c} values. For each increase in the number of questions answered correctly, HbA_{1c} decreased by 0.239 ($r = -0.337$, $P < 0.003$).

These results suggest that improved diabetes knowledge improves glycemic control. Other studies have shown that patient educational interventions lower HbA_{1c} values (6). Patient education

should be part of the diabetes treatment plan as this provides another modality for reaching glycemic targets. Patient empowerment is a strong tool that can be attained through knowledge of disease processes.

In conclusion, we have shown that improved diabetes knowledge is associated with better glycemic control. Empowering diabetic patients with knowledge of their disease may help combat this financially draining epidemic.

KATHLEEN M. COLLERAN, MD¹
 BRIAN STARR, BS²
 MARK R. BURGE, MD¹

From the ¹Department of Medicine, University of New Mexico, Albuquerque, New Mexico; and the ²School of Medicine, University of New Mexico, Albuquerque, New Mexico.

Address correspondence to Kathleen Colleran, MD, University of New Mexico HSC, Department of Medicine ACC-5th, 2211 Lomas Blvd. NE, Albuquerque, NM 87131. E-mail: kcolleran@salud.unm.edu.

© 2003 by the American Diabetes Association.

Acknowledgments—We thank the Michigan Diabetes Research and Training Center for use of the Michigan Diabetes Knowledge Test.

References

1. Rubin RJ, Altman WM, Mendelson DN: Health expenditures for people with diabetes. *J Clin Endo and Metab* 78:809A–809F, 1992
2. U.K. Prospective Diabetes Study Group: Effect of intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 34). *Lancet* 352:837–853, 1998
3. Stratton IM, Adler AI, Neil HA, Matthews DR, Manley SE, Cull CA, Hadden D, Turner RC, Holman RR: Association of glycemia with macrovascular and microvascular complications of type 2 diabetes (UKPDS 35): prospective observational study. *BMJ* 321:405–412, 2000
4. George PB, Tobin KJ, Corpus RA, Devlin WH, O'Neill WW: Treatment of cardiac risk factors in diabetic patients. How well do we follow the guidelines? *Am Heart J* 142:857–863, 2001
5. Fitzgerald JT, Anderson RM, Funnell MM, Hiss RG, Hess GE, Davis WK, Barr PA: The reliability and validity of a brief diabetes knowledge test. *Diabetes Care* 21: 706–710, 1998
6. Van den Arend IJ, Stolk RP, Rutten GE, Schrijvers GJ: Education integrated into

structured general practice care for type 2 diabetic patients results in sustained disease knowledge and self-care. *Diabet Med* 17:190–197, 2000

Ethnic Differences in Diabetes Symptoms Among People Without Known Diabetes in New Zealand

National diabetes awareness programs often emphasize the importance of symptoms in undiagnosed diabetes. Here we describe the frequency of diabetes symptoms in a multiethnic community in New Zealand.

Randomly selected nondiabetic European, Maori, and Pacific residents were asked if any of six cardinal symptoms of hyperglycemia (thirst, weight loss, polyuria, boils, tiredness, and blurred vision) were present. Subjects were screened using random venous blood sampling as previously described (6). Those with a random glucose ≥ 6.5 mmol/l within 2 h of a meal, or ≥ 6.0 2 h after a meal, and a random 20% of others were invited to attend a 75-g, 2-h oral glucose tolerance test (OGTT) (1999 World Health Organization criteria for diabetes, impaired glucose tolerance [IGT], and impaired fasting glucose [IFG]). Those diagnosed elsewhere, or with no OGTT but a random glucose of ≥ 11.1 mmol/l, have been included in the undiagnosed diabetes group. Odds ratios (ORs) and 95% CIs are shown.

Of the 2,423 people invited to participate, 1,585 (65.4%) were interviewed. Overall (and after removing those with new diabetes), Maori and Pacific individuals were significantly more likely to have more than two symptoms (Europeans 4.2% [3.8%] vs. 14.3% [13.5%] and 9.4% [9.0%], both $P < 0.001$, respectively). Each of the individual symptoms was present significantly less frequently among Europeans.

Of the 786 invited to the 75-g OGTT, 534 (67.9%) attended. Symptoms were uncommon among those with new diabetes, but still 2.6-fold (1.4–4.8) as likely when compared with IGT/IFG or normal subjects (e.g., more than two symptoms present in 16.9% vs. 6.5% and 7.5%, respectively, $P < 0.001$). This is a similar

frequency to that found in the Australian Diabetes Screening Study (2) but less than in the Hoorn study (3). IGT/IFG and normal subjects had very similar symptom frequencies. A logistic regression for each symptom revealed that after adjusting for ethnicity, age, sex, and BMI, only thirst was significantly associated with hyperglycemia, and then only at ≥ 17.0 mmol/l vs. < 5.0 mmol/l, OR = 9.2 (3.2–26.6). In a similar analysis, more than two symptoms were 3.5-fold (1.2–10.1) more common among those with a random glucose ≥ 11.0 mmol/l. When diabetes replaced random glucose in each regression, thirst (2.0 [1.0–4.0]), polyuria (1.8 [1.0–3.2]), tiredness (1.9 [1.0–3.4]), and having two symptoms (1.9 [1.1–3.4]) were significant entrants. Obesity was not associated with excess symptoms.

We conclude that few individuals with undiagnosed diabetes had symptoms, and that symptoms were only marginally increased in undiagnosed diabetes. Furthermore, symptoms were not increased in IGT/IFG. Ethnic differences in symptoms exist, confounding their interpretation in some settings. Our data suggest that unless a patient presents acutely, using prompted symptomatology to decide who should be screened for dysglycemia using blood testing is futile.

DAVID SIMMONS, FRACP, MD¹
 COLIN F. THOMPSON, FRACP²
 MICHAEL M. ENGELGAU, MD, MP³

From the ¹Waikato Clinical School, University of Auckland, Hamilton, New Zealand; the ²South Auckland Diabetes Project, Middlemore Hospital, New Zealand; and the ³Division of Diabetes Translation, Centers for Disease Control and Prevention, Atlanta, Georgia.

Address correspondence to David Simmons, FRACP, MD, Waikato Clinical School, Waikato Hospital, Hamilton, New Zealand. E-mail: simmonsdc@waikatodhb.govt.nz.

© 2003 by the American Diabetes Association.

Acknowledgments—We are grateful to Servier for funding a research fellow (C.F.T), Auckland Medical Research Foundation, Maurice and Phyllis Paykel Trust, Hector Trust, SmithKline Beecham, and the members of the Diabetes Projects Trust.

References

1. Simmons D, Thompson CF, Volklander D: Polynesians: prone to obesity and type 2 diabetes but not hyperinsulinaemia. *Diabet Med* 18:193–198, 2001

2. Welborn TA, Reid CM, Marriott G: Australian Diabetes Screening Study: impaired glucose tolerance and non-insulin dependent diabetes mellitus. *Metabolism* 46 (Suppl. 1):35–39, 1997
3. Ruige JB, de Neeling JN, Kostense PJ, Bouter LM, Heine RJ: Performance of an NIDDM screening questionnaire based on symptoms and risk factors. *Diabetes Care* 20:491–416, 1997

COMMENTS AND RESPONSES

Hyperglycemia After Myocardial Infarction

In their article on hyperglycemia in subjects admitted with myocardial infarction, Dandona, Aljada, and Bandyopadhyay (1) summarize much of the current knowledge about the anti-inflammatory effects of insulin. They propose mechanisms to explain the decreased morbidity and mortality seen in subjects on insulin infusions with tight blood glucose control. This may explain the results seen in another trial (2) in the intensive care population.

However, the authors fail to mention another of the possible effects of insulin. One of the systemic responses to critical illness is acute protein breakdown. This is thought to permit release of amino acids from skeletal muscle for high-priority use in threatened tissues. This protein breakdown may be due to the catabolic actions of the counter regulatory hormones and/or through the actions of a variety of cytokines (3). The insulin resistance that occurs as a result of these excess hormones and cytokines may reduce the inhibitory effect that insulin has on the ATP-ubiquitin proteasome proteolytic pathway, thus leading to an increase in skeletal muscle protein loss (4). This breakdown occurs despite the provision of adequate enteral or parenteral nutrition (5).

The protein breakdown seen in critical illness is analogous to the situation seen during prolonged insulin deprivation in subjects with type 1 diabetes. Insulin has been shown to prevent this breakdown from occurring (6,7). Thus, one of the reasons for the improved outcomes in the intensive care population on insulin may be that they lose less protein.

The anti-catabolic action of insulin in these patients results in fewer complications due to the maintenance of immunocompetence, reduced incidence of infection, normalized wound healing, less muscle weakness, and lower mortality seen in the hyperglycemic critically ill (8).

KETAN DHATARIYA, MBBS, MRCP, MSc

From the Endocrine Research Unit, Mayo Clinic and Foundation, Rochester, Minnesota.

Address correspondence to Ketan Dhatariya, MBBS, MRCP, MSc, Department of Endocrine Research Fellow, Mayo Clinic, Joseph 5-194, Rochester, MN 55905. E-mail: dhatariya.ketan@mayo.edu.
© 2003 by the American Diabetes Association.

References

1. Dandona P, Aljada A, Bandyopadhyay A: The potential therapeutic role of insulin in acute myocardial infarction in patients admitted to intensive care and in those with unspecified hyperglycemia. *Diabetes Care* 26:516–519, 2003
2. Van den Berghe G, Wouters P, Weekers F, Verwaest C, Bruyninckx F, Schetz M, Vlasselaers D, Ferdinande P, Bouillon R, Lauwers P: Intensive insulin therapy in the surgical intensive care unit. *N Engl J Med* 345:1359–1367, 2001
3. Vary TC: Regulation of skeletal muscle protein turnover during sepsis. *Curr Opin Clin Nutr Metab* 1:217–224, 1998
4. Bailey JL, Wang X, Price SR: The balance between glucocorticoids and insulin regulates muscle proteolysis via the ubiquitin-proteasome pathway. *Miner Electrolyte Metab* 25:220–223, 1999
5. Streat SJ, Beddoe AH, Hill GL: Aggressive nutritional support does not prevent protein loss despite fat gain in septic intensive care patients. *J Trauma* 27:262–266, 1987
6. Nair KS, Ford GC, Ekberg K, Fernqvist-Forbes E, Wahren J: Protein dynamics in whole body and in splanchnic and leg tissues in type I diabetic patients. *J Clin Invest* 95:2926–2937, 1995
7. Nair KS, Ford GC, Halliday D: Effect of intravenous insulin treatment on in vivo whole body leucine kinetics and oxygen consumption in insulin-deprived type I diabetic patients. *Metabolism* 36:491–495, 1987
8. Cuthbertson DP: The metabolic response to injury and its nutritional implications: retrospect and prospect. *J Parenter Enteral Nutr* 3:108–129, 1979

Hyperglycemia After Myocardial Infarction

Response to Dhatariya

We appreciate the comments of Dhatariya (1) in this issue of *Diabetes Care*. Clearly, insulin is the ultimate anabolic hormone that may not only keep inflammation at bay, but also regulate the appropriate utilization of metabolites such that it conserves protein and fat and prevents their breakdown. Its usefulness in preventing protein catabolism in the clinical setting was demonstrated in the 1980s. The key studies of Nair and colleagues (2,3) referred to by Dhatariya provide the scientific basis for this important insulin action. The next challenge is to determine how inflammation induces a state of protein catabolism and exactly how insulin exerts its beneficial effects against the background of inflammation.

It is also worth mentioning two other key actions of insulin described recently: 1) apo E^{-/-} mice that develop atherosclerosis suppress this process when given insulin (4), and 2) insulin suppresses reperfusion-induced myocardial damage following ischemia in isolated rat heart, as well as reduces myocardial apoptosis (5).

We believe this is just the beginning of a new era in understanding insulin action beyond the conventional biochemical/metabolic paradigm that we have been accustomed to for the first 80 years of its life. As discussed in our commentary, as we understand more about these novel actions of insulin, its clinical application will expand.

PARESH DANDONA, MD
AHMAD ALJADA, PHD
ARINDAM BANDYOPADHYAY, MD

From the Division of Endocrinology, Diabetes and Metabolism, State University of New York at Buffalo, Buffalo, New York; and Kaleida Health, Buffalo, New York.

Address correspondence to Paresh Dandona, MD, PhD, Diabetes-Endocrinology Center of WNY, 3 Gates Circle, Buffalo, NY 14209. E-mail: pdandona@kaleidhealth.org.

© 2003 by the American Diabetes Association.



References

1. Dhatariya K: Hyperglycemia after myocardial infarction (Letter). *Diabetes Care* 26:2222, 2003
2. Nair KS, Ford GC, Ekberg K, Fernqvist-Forbes E, Wahren J: Protein dynamics in whole body and in splanchnic and leg tissues in type I diabetic patients. *J Clin Invest* 95:2926–2937, 1995
3. Nair KS, Ford GC, Halliday D: Effect of intravenous insulin treatment on in vivo whole body leucine kinetics and oxygen consumption in insulin-deprived type I diabetic patients. *Metabolism* 36:491–495, 1987
4. Shamir R, Shehadeh N, Rosenblat M, Eschach-Adiv O, Coleman R, Kaplan M, Hamoud S, Lischinsky S, Hayek T: Oral insulin supplementation attenuates atherosclerosis progression in apolipoprotein E-deficient mice. *Arterioscler Thromb Vasc Biol* 23:104–110, 2003
5. Jonassen AK, Sack MN, Mjos OD, Yellon DM: Myocardial protection by insulin at reperfusion requires early administration and is mediated via Akt and p70s6 kinase cell-survival signaling. *Circ Res* 89: 1191–1198, 2001

Clinical and Genetic Heterogeneity of Latent Autoimmune Diabetes in Adults

We read with great interest the recent article by Hosszufalusi et al. (1) reporting that latent autoimmune diabetes in adults (LADA) manifests clinical features similar to those of adult-onset type 1 diabetes with rapid progression, and that prevalences of predisposing alleles and haplotypes did not differ between these two forms of autoimmune diabetes. Patients with LADA have been found to be heterogeneous in their clinical attributes (2). Some patients develop insulin deficiency within a few years (LADA-type 1), while others show clinical and metabolic markers similar to patterns in antibody-negative type 2 diabetes (LADA-type 2). As the authors noted, selection bias may have been present among the patients with LADA in their study because they tested for antibodies against islet cell cytoplasm (ICA) only in patients clinically suspected to have LADA. Therefore, patients with

LADA in their study were not representative of all LADA patients, but represented only a subgroup, namely LADA-type 1. As a result, patient characteristics and prevalences of predisposing alleles and haplotypes were similar to those seen in adult-onset type 1 diabetes. We previously reported that HLA-DRB1 alleles influenced the prognosis of Japanese patients with diabetes who were positive for antibodies to GAD (3). Patients with LADA who later developed insulin deficiency showed increased prevalence of one of the predisposing alleles. Notably, patients with LADA who did not develop insulin deficiency were more likely to have protective alleles and less likely to have predisposing alleles than patients with type 1 diabetes showing rapid progression.

In addition, the authors diagnosed type 1 diabetes when patients had typical diabetes symptoms, were ketosis-prone, and required prompt insulin treatment at the time of diagnosis. The median and upper interquartile fasting concentrations of C-peptide (0.46 and 1.05 nmol/l, respectively) in the subjects with type 1 diabetes were somewhat higher than we expected, given their tendency toward ketosis and immediate need for insulin treatment. One suspects that patients who required insulin within 6 months of diagnosis as opposed to the time of diagnosis may have been included in the type 1 diabetes category. We propose that those patients should be placed in another subgroup, given their clinical differences from typical type 1 diabetic patients.

Demonstrating that ICA detected at diagnosis disappeared in six LADA patients with a relatively long disease course, the authors speculated that the tendency of ICA to disappear with increasing disease duration was similar in LADA and type 1 diabetes. In fact, the meaning of this disappearance may differ between LADA and type 1 diabetes. ICA positivity has been reported to persist in LADA (4), in contrast to the disappearance during the course of type 1 diabetes. Low levels of antibodies to GAD in patients with LADA declined to undetectable levels in our study (5), which suggests the possibility of pseudopositivity of antibodies to GAD.

In conclusion, patients with LADA constitute a clinically and genetically heterogeneous group. A more precise classification of autoimmune diabetes in adults

is needed to define differences between forms of autoimmune diabetes in adults.

MICHIAKI FUKUI, MD¹
YOSHIHIRO KITAGAWA, MD¹
NAOTO NAKAMURA, MD²
TOSHIKAZU YOSHIKAWA, MD²

From the ¹Department of Endocrinology and Hematology, Osaka General Hospital of West Japan Railway Company, Osaka, Japan; and the ²First Department of Internal Medicine, Kyoto Prefectural University of Medicine, Kyoto, Japan.

Address correspondence to Michiaki Fukui, MD, The Department of Endocrinology and Hematology, Osaka General Hospital of West Japan Railway Company, 1-2-22 Matsuzaki-cho, Abeno-ku, Osaka 545-0053, Japan. E-mail: sayarinapm@hotmail.com.

© 2003 by the American Diabetes Association.



References

1. Hosszufalusi N, Vatay A, Rajczy K, Prohaszka Z, Pozsonyi E, Horvath L, Grosz A, Gero L, Madacsy L, Romics L, Karadi I, Fust G, Panczel P: Similar genetic features and different islet cell autoantibody pattern of latent autoimmune diabetes in adults (LADA) compared with adult-onset type 1 diabetes with rapid progression. *Diabetes Care* 26:452–457, 2003
2. Lohmann T, Kellner K, Verlohren HJ, Krug J, Steindorf J, Scherbaum WA, Seissler J: Titre and combination of ICA and autoantibodies to glutamic acid decarboxylase discriminate two clinically distinct types of latent autoimmune diabetes in adults (LADA). *Diabetologia* 44: 1005–1010, 2001
3. Fukui M, Nakano K, Nakamura N, Maruya E, Saji H, Obayashi H, Ohta K, Ohta M, Mori H, Kajiyama S, Wada S, Kida Y, Kosaka K, Deguchi M, Shigeta H, Kitagawa Y, Kondo M: HLA-DRB1 alleles contribute to determining the prognosis of Japanese diabetes mellitus positive for antibodies to glutamate decarboxylase. *J Clin Immunol* 18:89–92, 1998
4. Kobayashi T, Nakanishi K, Okubo M, Murase T, Kosaka K: GAD antibodies seldom disappear in slowly progressive IDDM (Letter). *Diabetes Care* 19:1031, 1996
5. Fukui M, Nakamura N, Nakano K, Kajiyama S, Matsuo S, Obayashi H, Ohta M, Shigeta M, Shigeta H, Kitagawa Y, Kondo M: HLA-associated cellular response to GAD in type 2 diabetes with antibodies to GAD. *Endocrine J* 47:753–761, 2000

Clinical and Genetic Heterogeneity of Latent Autoimmune Diabetes in Adults

Response to Fukui et al.

We read with interest the comments of Fukui et al. (1) in this issue of *Diabetes Care* regarding our recent article (2). They claim that patients with latent autoimmune diabetes in adults (LADA) are heterogeneous in their clinical attributes and that the LADA patients in our study represented the LADA type 1 subgroup. Because of this, we observed that their clinical and genetic characteristics resembled patients having rapidly progressive adult-onset type 1 diabetes. We agree that LADA patients may show heterogeneous clinical features, but we feel that our patients with LADA are representative of the whole LADA group: 25% of our LADA patients belonged to the overweight category based on their BMI, and the same percentage of patients had lipid abnormalities. The median of the insulin-free period after the diagnosis of diabetes was 3.0 years (1.0–6.0), despite the fact that we initiated insulin therapy in 23 of our 54 patients with LADA during the first year after diagnosis. The indication of insulin treatment in these patients was their autoantibody positivity and not their metabolic status. We suspect that LADA type 1 and LADA type 2 subgroups may not represent different clinical entities. Instead we think that a certain proportion of patients with LADA, mainly with aging, develop clinical features of metabolic syndrome beside their autoimmune diabetes.

We agree with Palmer and Hirsch (3) that phenotypically there are at least three separate populations of autoimmune diabetes in adults: adult-onset type 1 diabetes, LADA, and obese phenotypic type 2 diabetes with autoantibody positivity. In the study of Fukui et al. (4) anti-GAD-positive type 2 diabetic patients with secondary failure of sulfonylurea therapy ($n = 44$, we think that these are the LADA patients) showed an increased prevalence of one of the predisposing alleles, while the anti-GAD-positive and well-controlled

type 2 diabetic patients ($n = 22$, we think that they represent type 2 diabetes with autoantibody positivity) were more likely to have protective alleles and less likely to have predisposing alleles compared with type 1 diabetes showing rapid progression. Notably, the type 1 diabetic group from their study instead represents childhood-onset diabetes (age at onset was 14.5 ± 12.9 years). Another explanation could be the ethnic differences between the Japanese and Hungarian populations.

The unexpected high level of fasting C-peptide at onset in the type 1 diabetic group, (median 0.46 nmol/l [range 0.24–1.05]) was also surprising. However, Mallone et al. (5) also reported a wide range of fasting plasma C-peptide levels in newly diagnosed type 1 diabetes, even with childhood onset (median 0.44 ng/ml [0–5.70]). Since the diagnosis of type 1 diabetes was established according to the World Health Organization criteria, and the decision of prompt insulin therapy was based on the clinical picture (presence of ketonuria or ketoacidosis) in our study, we do not think that another subgroup of type 1 diabetic patients should have been formed on the basis of the fasting C-peptide level. As a result of the comments by Fukui et al., we noticed a regrettable typing error in Table 1 of our report (1): fasting C-peptide in adult-onset type 1 diabetes 1–10 years after onset is 0.40 nmol/l (0.24–0.62) instead of 0.40 nmol/l (0.24–1.05).

We reported that the islet cell antibody (ICA) positivity documented earlier disappeared in six patients having LADA with longer disease course. There was a considerable interval between the positive and the negative ICA tests (6–11 years). The data regarding persistence of ICA in LADA are controversial. In the cited study (6), ICA either persisted ($n = 18$) or disappeared ($n = 9$) and anti-GAD antibody persisted ($n = 10$) in patients having type 2 diabetes with further insulin requirement. Further studies are necessary to evaluate the long-term characteristics of ICA in patients with LADA.

Regarding the classification of autoimmune diabetes, we would divide it into two or three subtypes. One subtype would be the childhood-onset type 1 diabetes (age at onset <20 years), showing the highest prevalence of the predisposing genotypes and the most aggressive β -cell destruction. Another subtype

could be adult-onset type 1 diabetes, which has two forms: rapidly and slowly progressive. The latter should be called LADA without age restriction. The problem of obese phenotypic type 2 diabetes with autoantibody positivity remains unsolved; it needs to be decided whether this group belongs to type 1 or type 2 diabetes, represents a mixture of type 1 and type 2 diabetes, represents a distinct clinical entity, or is merely a laboratory bias.

NÓRA HOSSZÚFALUSI, MD, PHD
PÁL PÁNCZÉL, MD, PHD

From the 3rd Department of Internal Medicine, Semmelweis University, Budapest, Hungary.

Address correspondence to Nóra Hosszúfalusi, MD, PhD, Budapest Kútvolgyi út 4. H-1125 Hungary. E-mail: hono@kut.sote.hu.

© 2003 by the American Diabetes Association.

References

1. Fukui M, Kitagawa Y, Nakamura N, Yoshikawa T: Clinical and genetic heterogeneity of latent autoimmune diabetes in adults (Letter). *Diabetes Care* 26:2223, 2003
2. Hosszúfalusi N, Vatay A, Rajczy K, Prohaszka Z, Pozsonyi E, Horvath L, Grosz A, Gero L, Madacsy L, Romics L, Karadi I, Fust G, Panczel P: Similar genetic features and different islet cell autoantibody pattern of latent autoimmune diabetes in adults (LADA) compared with adult-onset type 1 diabetes with rapid progression. *Diabetes Care* 26:452–457, 2003
3. Palmer JP, Hirsch IR: What's in a name: latent autoimmune diabetes of adults, type 1.5, adult onset, and type 1 diabetes. *Diabetes Care* 26:536–538, 2003
4. Fukui M, Nakano K, Maruya E, Saji H, Obayashi H, Ohta K, Ohta M, Mori H, Kajiyama S, Wada S, Kida Y, Kosaka K, Deguchi M, Shigeta H, Kitagawa Y, Kondo M: HLA-DRB1 alleles contribute to determining the prognosis of Japanese diabetes mellitus positive for antibodies to glutamate decarboxylase. *J Clin Immunol* 18: 89–92, 1998
5. Mallone R, Ortolan E, Pinach S, Volante M, Zanone MM, Bruno G, Baj G, Lohmann T, Cavallo-Perin P, Malavasi F: Characterisation in new-onset type 1 diabetes and latent autoimmune diabetes of the adult (LADA) and comparison with other islet autoantibodies. *Diabetologia* 45:1667–1677, 2002
6. Kobayashi T, Nakanishi K, Okubo M, Murase T, Kosaka K: GAD antibodies seldom disappear in slowly progressive IDDM (Letter). *Diabetes Care* 19:1031, 1996