

A new methodological protocol for the use of dendrogeomorphological data in flood risk analysis

A. Díez-Herrero, J. A. Ballesteros-Cánovas, J. M. Bodoque
and V. Ruiz-Villanueva

ABSTRACT

Dendrogeomorphology uses information sources recorded in the roots, trunks and branches of trees and bushes located in the fluvial system to complement (or sometimes even replace) systematic and palaeohydrological records of past floods. The application of dendrogeomorphic data sources and methods to palaeoflood analysis over nearly 40 years has allowed improvements to be made in frequency and magnitude estimations of past floods. Nevertheless, research carried out so far has shown that the dendrogeomorphic indicators traditionally used (mainly scar evidence), and their use to infer frequency and magnitude, have been restricted to a small, limited set of applications. New possibilities with enormous potential remain unexplored. New insights in future research of palaeoflood frequency and magnitude using dendrogeomorphic data sources should: (1) test the application of isotopic indicators ($^{16}\text{O}/^{18}\text{O}$ ratio) to discover the meteorological origin of past floods; (2) use different dendrogeomorphic indicators to estimate peak flows with 2D (and 3D) hydraulic models and study how they relate to other palaeostage indicators; (3) investigate improved calibration of 2D hydraulic model parameters (roughness); and (4) apply statistics-based cost-benefit analysis to select optimal mitigation measures. This paper presents an overview of these innovative methodologies, with a focus on their capabilities and limitations in the reconstruction of recent floods and palaeofloods.

Key words | dendrogeomorphology, flood hazard, natural risk, palaeoflood, roughness, tree-rings

A. Díez-Herrero (corresponding author)
J. A. Ballesteros-Cánovas

V. Ruiz-Villanueva
Geological Survey of Spain (IGME),
Ríos Rosas 23,
E-28003 Madrid,
Spain
E-mail: andres.diez@igme.es

J. M. Bodoque
University of Castilla-La Mancha (UCLM),
Avenida de Carlos III,
45004 Toledo,
Spain

Associated researchers to IMDEA-Water

INTRODUCTION

Floods are among the natural catastrophes and disasters that cause the highest number of victims, injuries and displacements as well as economic losses in goods and infrastructure. This is the case both in Spain and worldwide, as shown by the results of the floods affecting many countries over recent decades; in Spain alone, flood damage totalling 1% of GDP has caused more than 215 deaths in the last 20 years. The wide-scale flooding that took place in central Europe in 2002 was the turning point for the EU institutions concerned with this problem, leading to the new EU Directive initiative. In Europe, lowland floods are rarely associated with fatalities; in contrast, flash floods often result in loss of life (Gaume *et al.* 2009).

In mountain torrents the traditional methods used for flood hazard analysis (hydrological-hydraulic, palaeohydrological and geological-geomorphological) present enormous scientific uncertainties because of the lack of availability of input data. Normally, in these areas there are very few meteorological and flow gauge stations, or even none at all; their space-time validity may be heterogeneous due to altitude distribution (concentrated in lower level urban areas), and the time series could be discontinued or statistically not representative when fewer than 30 data samples are available. Historic and palaeohydrological data are often used to complete the systematic data log (Benito & Thorndycraft 2004); however, many of the

analysed areas do not have available documentation on historic flooding as they are a considerable distance from administrative centres and document production centres. There are often no records of flood deposits either, as there are no debris materials in the source area, or no appropriate deposit-prone or retention areas (because of the high steep dynamic velocities). As a result, in the absence of input data, it is very difficult to apply reliable statistical analyses or hydrometeorological models that are adequately calibrated and validated. In the upper basins, conditions for stable human activities such as agriculture and industry are less favourable, significant forest areas have survived and in almost all high-gradient river beds, trees and riparian or other species can be easily found on or near river banks. The growth distribution pattern and single trees of these plant species are affected by the interference of flash floods occurring in these river courses and on their banks.

Dendrogeomorphology (Alestalo 1971) uses information sources recorded in the roots, trunks and branches of trees and bushes located at specific geomorphological points (river banks, bars, flood plains, etc.), to complement (or even replace) the record of flash floods that have occurred in a water stream (Díez Herrero *et al.* 2007). This scientific analysis involves the following research: finding examples with ideal dendro-evidence (exposed roots, adventitious branches, impact scarring, etc.); species selection with homogeneous growth; planned sampling of ring sequences; study of plant anatomy searching for characteristic signatures of flash flood events; synchronization of ring sequences for event dating; including events to better fit the frequency distribution function; hydraulic modelling of the site to estimate flow rates; assigning magnitudes to the estimated frequencies. This allows systematic record data series to be extended, completed or even replaced using proxy data evidencing physical events, not statistical models.

The key aim of this paper is therefore to propose a methodological scheme or protocol for the use of dendrogeomorphological data in flood risk analysis. The objective of this methodological protocol is to improve frequency and magnitude estimates for flash floods and enable the implementation of more effective preventive measures for mitigation of related risks. To achieve this aim the specific objectives are to: (1) review the application of different data sources and scientific methods resulting from

dendromorphological analysis; and (2) analyse the preliminary results from case studies in which this protocol is followed.

BACKGROUND AND STATE OF THE ART

Dendrogeomorphology is a relatively recent branch of science, especially compared with other closely related sub-disciplines such as dendroclimatology. In fact, both derive from dendrochronology, the science of dating tree-rings to characterize and synchronize environmental changes that have taken place, or for dating archaeological remains. The origins can be traced back to Leonardo da Vinci's observations in the late 15th and early 16th centuries (France manuscripts M 78v and 79r), but it was not until the early 20th century that the theoretical bases were established (Douglass 1909, 1914). However, its laws and principles, although originating in the 18th century with theories such as uniformitarianism, were not explicitly formulated until the 1970s (Fritts 1976): uniformitarianism, limiting factors, ecological range, aggregate tree-ring growth (Cook 1990), site selection, replicate measurement and cross-dating.

The early dendrogeomorphological studies on the relationship between tree-rings and geomorphological processes date from the 1960s (La Marche 1963), but it was not until a few years later that the first publications appeared using this term (Alestalo 1971). In general, these dealt with the application of dendrochronological techniques to the study of erosive processes of soils, snow avalanches, slope mass movements (rockfalls, rock avalanches, landslides, etc.) and surface dating, among others.

The application of dendrogeomorphological analyses to torrent dynamic-related events such as debris flows and flash floods dates back to the same time. The earliest papers on these applications also date from the 1960s (Sigafos 1964; Harrison & Ried 1967), but it was not until the 1980s that these became standard, with the papers by Yanosky (1982a, 1982b, 1983, 1984), Hupp (1984, 1986, 1988), Hupp *et al.* (1987), Gottesfeld & Gottesfeld (1990), McCord (1990) and Gottesfeld (1996). These techniques and data sources have since become more widely used, although largely limited in geographical terms to

North America and Central Europe (Astrade & Bégin 1997; St. George *et al.* 1999, 2002; St. George 2010; Stoffel & Wilford 2012). These include the methodological contributions of C. R. Hupp, particularly the book chapter by Hupp (1988). In this contribution he includes the only classification of flood dendrogeomorphological evidence (FDE) (limited to six types) prior to this article. Other book chapters by Yanosky & Jarrett (2002) and by St. George (2010) also provide an overview of the main contributions of dendrogeomorphology to hydrological studies, but focus on aspects of the frequency of extreme events, and on one type of magnitude indicator, respectively. The other publications are simply case studies, and do not include a methodological proposal apart from short sections on materials and methods.

With an accumulation of more than 40 years' experience in these studies, the current state of the art in terms of methods and knowledge is the result of the evolution and recent fine-tuning of techniques in the following three aspects:

- Botanical features: in early studies only external evidence was used, such as corrosion scarring, bifurcations at the base of the trunk, adventitious branches and eccentric growth; nowadays, however, many more tree-ring pattern growth anomalies are used; these are called growth disturbances (GD), such as the lines formed by traumatic resin ducts (TRDs) or sudden variations in tree-ring width (Bollschweiler *et al.* 2008b; Ruiz-Villanueva *et al.* 2010a).
- Detail of the information extracted: formerly, tree-rings were only studied as indivisible elements, whereas nowadays, plant histology and plant cytology techniques are applied (Ballesteros *et al.* 2010a, 2010b) that allow zone rescaling within the tree-ring and identification of even the season and month when the disturbance occurred (Stoffel *et al.* 2006) and differentiation between various processes in mountain areas (avalanches, flash floods, debris flows, etc.).
- New data from dendrogeomorphological analysis, from the classical estimation of flood discharges, from simple equations (Yanosky & Jarrett 2002) to the calibration of interesting flow parameters such as roughness (Ballesteros *et al.* 2010b).

This means that in the current state of the art, dendrogeomorphological analysis techniques require the study of a wide range of elements, the use of synchronization and the application of microscopic techniques to extract the maximum information on the event, its origin and dating. Good examples of some of the most recent studies including advances in scientific knowledge are: Gottesfeld (1996), Kozłowski (1997), Yanosky & Jarrett (2002), Coriell (2002), Stoffel, Bollschweiler and Ehmiş (Stoffel *et al.* 2003, 2005, 2006, 2008; Bollschweiler *et al.* 2006, 2008a, 2008b; Stoffel & Wilford 2012) and Zielonka *et al.* (2008). A summary on recent state-of-the art research in dendrogeomorphology is given in Stoffel & Bollschweiler (2009) and Stoffel *et al.* (2010).

FIRST METHODOLOGICAL PROPOSAL: CLASSIFICATION OF FLOOD DENDROGEOMORPHOLOGICAL EVIDENCE

As mentioned above, dendrogeomorphological techniques have been applied to the analysis of flood risks for more than four decades. As there is no classification or any commonly accepted nomenclature of dendrogeomorphological evidence in use today, and our review of many studies revealed inconsistencies in the description of dendrogeomorphic data, we elaborated a simplified classification of evidence used in flood analysis. Therefore, this system would facilitate realization of cross studies and spatial correlation of the results, and it would also be useful as a guide to novice researchers in this field, providing a basic doctrinal corpus.

The conductive thread of this classification is the spatial scale of the evidence (from macro- to microscopic, from kilometres to microns) and the scale of studied elements (from forest communities and vegetal formations to changes in cell structure). More than 29 types of disturbance signs have been distinguished and hierarchically arranged, with 15 of them referring to entire individuals (tree or bush) or their parts (trunk, branches or roots). The classification has a cross-tabular shape (Table 1) and is complemented with graphics of the evidence, both in the form of sketches (Figure 1) and pictures from field examples (Figure 2).

Table 1 | Proposal for classification of the main types of flood dendrogeomorphological evidence (FDE) useful in the study of past floods, depending on the scale of the studied element (modified from Díez-Herrero *et al.* (2009))

Spacial scale	Studied element		Type of dendrogeomorphological evidence		No.	
km	Macroscopic	Bottomland vegetation patterns		Species distribution pattern	FDE 1	
		Individual (tree or bush)	Complete tree or bush	Coverage distribution pattern		FDE 2
Ages distribution pattern				FDE 3		
Candelabra growth ('Sigafos' trees)				FDE 4		
Tilted and overturned trees				FDE 5		
Decapitated trees (tops missing)				FDE 6		
dm	Part of tree or bush			Trunk	Stripped bark with callus marks	Scars by sediment load impact
			Branches torn off		FDE 8	
			Scraping from other falling trees		FDE 9	
			M	Bark erosion		FDE 10
Elbows and angles (sharp changes in the trunk growth direction)				FDE 11		
Sudden narrowings in trunk		FDE 12				
Bifurcations		FDE 13				
Sprouts from buried trunks and roots		FDE 14				
dm	Branches	Sharp changes in the branch growth direction		FDE 15		
		Sprouting from burried trunk		FDE 16		
		Roots	Exposed roots	Stripped bark and erosion	FDE 17	
Float roots without substrate contact			FDE 18			
cm	Mesoscopic	Tissues, wedges and slices	Rings	New roots from buried trunks	FDE 19	
				Eccentric growths (reaction wood)		FDE 20
			False or absent tree rings		FDE 21	
			mm	Tissues	Discontinuities, erosion, and internal scars	
Changes in tree-ring parameters (width, % early wood, late wood, etc.)		FDE 23				
mm	Microscopic	Thin slice (microslices)	Ratio parenchyma-lignification tissue		FDE 24	
			Size and density of vessels		FDE 25	
			Cells	Changes in cell parameters	Size and morphometry of cell lumens	FDE 26
				Cell wall thickness		FDE 27
				Appearance and/or abundance of special types of cells	Traumatic resin ducts (TRDs)	FDE 28
			mm	Fibre-tracheid		FDE 29
				Traumatic structures in cell wall		FDE 30
Å	Atomic	Cell wall	Cellulose	Isotopic fractionation	$^{18}\text{O}/^{16}\text{O}$ ratio	FDE 31




Figure 1 | Types of the main individual macroscopic dendrogeomorphological evidence useful in the study of past floods: FDE 4, 'candelabrum' growths (Sigafos trees); FDE 5, tilted tree with feet tipped over; FDE 6, decapitation; FDE 7, scars caused by sediment load impact; FDE 8, branches torn off; FDE 9, grazes caused by falling boulders; FDE 10, erosion on the trunk; FDE 11, trunk bends and angles; FDE 12, sudden narrowing of trunk; FDE 13, bifurcations; FDE 14/16, re-growth from buried trunks; FDE 15, bends and angles in branches; FDE 17, exposed roots with stripped bark and eroded surface; FDE 18, exposed roots without contact with substratum; FDE 19, new roots from buried trunks.

However, not all signs appear to be equally useful for the reconstruction of magnitudes or frequencies of past flood events. Using a cross table (Table 2), the most significant signs are shown in the sense that they provide the date as well as the area affected by an event, the minimum levels reached that can later be used for the estimation of peak flows with hydraulic modelling and simplified equations (e.g., critical flow, super-elevation) (e.g., Elleder *et al.* 2013).

SECOND METHODOLOGICAL PROPOSAL: A 15-TASK WORKING PLAN WITH ADVANCED TECHNIQUES

A working plan has been drawn up structured into 15 activities or tasks and its relationships and links as shown in Figure 3. To carry out the activities and tasks envisaged in the working plan, simultaneous and complementary analysis and synthesis-based methodological approaches may be adopted with a sequential combination of field data acquisition methods, in-laboratory study protocols, and desk-based processing and analysis methods.

The first steps of the field data acquisition, i.e., geomorphological analysis (task 1) and flora characterization (task 2) should be carried out using classical methods that do not require specific adaptations or additional explanations. Species sampling (task 3), comprising sampling of individual examples of trees and bushes, should start with a rigorous and objective selection, based on external characteristics which can be used to provide evidence for interference between the sample tree and the river dynamics (see the previous section). The collection of detailed topographical data (task 4) also uses conventional methods and standardized techniques (GPS receivers in differential mode, total stations, 3D terrestrial laser scanner).

Neither is there anything new in laboratory work methodologies, as they are standardized procedures in dendrochronological analysis (Picher 1990; Wilford *et al.* 2005). This is the case of sample preparation (task 5) and tree-ring counting and measurement (task 6). However, some new methodological techniques are required for the anatomical and histological study (task 7) focused on the detection of past flood events. Anomalies in tissues and cells associated with flash flood events should be observed, with special emphasis on: cell dimensions (radial length and tangential width of parenchymal cells, vessels, fibres for broad leaves and tracheids for conifers; e.g., Stoffel & Hitz 2008), cell lumen, cell wall thickness, percentage of late-wood, resin ducts (lines caused by traumatism, TRDs; see Bollschweiler *et al.* 2008a, 2008b; Schneuwly *et al.* 2009a, 2009b), existing anatomical elements of traumatic origin (e.g., tyloses, sticky deposits) or the existence of intra-annual density fluctuation. Another source of 3D anatomical data




Figure 2 | Examples of the types of the main dendrogeomorphological evidence useful in the study of past floods.

could be the tomographic analysis of portions of tree trunks, using sophisticated medical devices. Finally, variance analysis (ADEVA) should be performed on the different variables considered in anatomical analysis to verify (or disprove) statistically significant differences between different mean groups. Variogram analysis should be applied to explore 3D intrinsic structure of cell data sampled at different distance intervals and directions (Bianchi & Zheng 2009; Remy *et al.* 2009).

Most desk-based data processing and analysis methods are classical and standard from dendrochronological or hydrological research; for example, series synchronization and tree-ring dating (task 9), hydrometeorological modelling of river basins (task 11), hydraulic modelling of reaches (task 12; Lang *et al.* 2004) or mapping flash flood hazard areas and risk maps (task 14; Díez-Herrero *et al.* 2009). Others methods do have peculiarities for their application to the analysis of past floods.

Table 2 | Use of flood dendrogeomorphological evidence (FDE) types and their combinations for the study of the frequency and magnitude of past floods. Type of information conveyed and degree of utility for each sign or combination in the study of flood hazard: black, high utility; grey, medium; white, none (modified from Díez-Herrero *et al.* (2009))

FDE/FDEs	Temporal considerations							
	Unique events			Magnitude considerations				
	Dating	Seasonality	Frequency	Extent	Flow depth	Sediment load	Energy	Duration
FDE 1	Grey		Grey	Black	Grey	Grey	Grey	Grey
FDE 2	Grey		Grey	Black	Grey	Grey	Grey	Grey
FDE 10				Black	Black	Black	Grey	Grey
FDE 23	Black		Black	Black	Grey	Grey	Grey	Grey
	FDE 3		Black	Black	Grey	Grey	Grey	Grey
	FDE 4	Grey	Black	Black	Grey	Grey	Grey	Grey
	FDE 5		White	Black	Grey	Grey	Grey	Grey
	FDE 6		Black	Black	Black	Black	Black	Black
	FDE 7		Black	Black	Black	Black	Black	Black
FDE 8	Black		Black	Black	Grey	Grey	Grey	Grey
FDE 9	Grey		Grey	Black	White	White	White	White
FDE 11	Black		Black	Black	Grey	Grey	Grey	Grey
FDE 12	Black		Black	Black	Grey	Grey	Grey	Grey
FDE 13	Black		Black	Black	Grey	Grey	Grey	Grey
FDE 14	Black		Black	Black	White	Black	Black	Black
FDE 15	Black		Black	Black	Grey	Grey	Grey	Grey
FDE 16	Black		Black	Black	White	Black	Black	Black
FDE 19	Grey		Grey	Grey	White	Black	Black	Black
FDE 20	Black		Black	Black	Grey	Grey	Grey	Grey
	FDE 17	Grey	Black	Black	Grey	Grey	Grey	Grey
	FDE 18	Grey	Black	Black	Grey	Grey	Grey	Grey
	FDE 5		Grey	Grey	Black	Black	Black	Black
FDE 7/FDE 21	Black		Black	Black	Black	Black	Black	Black
FDE 22 28-30	Black		Black	Black	White	Black	Black	Black
	FDE 17	Grey	Black	Black	White	Black	Black	Black
	FDE 7		Black	Black	Black	Black	Black	Black
FDE 24	Black		Black	Black	White	White	White	White
FDE 25	Black	Grey	Black	Black	White	White	White	White
FDE 24–28 27–29	Black		Black	Black	Black	Black	Black	Black
	FDE 7		Black	Black	Black	Black	Black	Black
FDE 17	Black		Black	Black	White	White	White	White
FDE 31	Black		Black	Black	White	White	White	White
FDE 32	Black		Black	Black	White	White	White	White

This is the case for task 10, flood event detection and dating in tree-ring series. From the synchronized tree-ring series we can proceed to the dating of the anatomical elements associated with torrent phenomena (GDs, such as TRDs, wounds, callus tissue, reaction wood, reduced or increased growth); and obtain a list or database of flash flood events with assigned dates, indicating whether these are definite or approximate dates, and the type of evidence marking maximum, minimum or exact level. Similarly,

depending on what part of the tree-ring the GDs (late or earlywood) are situated, it is even possible to infer the season of the year (at a seasonal level and sometimes to the precise month) in which the event took place. The possible identification of flash flood events that caused unearthing and exposure of the tree root system (e.g., by erosion of river banks or pedestal effect on bars and islands), and which may have led to reduced leaf system production (crown) should also be investigated, observing possible




Figure 3 | Chart summarizing the 15 activities or tasks of the methodological proposal, from the data sources to the final expected results, and their mutual relations.

changes in the tree-ring growth pattern at the standard height (1.30 m). These dates, especially those that are repeated and common, should be contrasted with recent historic records of intense precipitation and flooding where hydrometeorological information exists. This information should be used to calculate similarity indices and conditional probabilities. To calibrate the elements used for detecting and dating events, the same methods and data sources should be used as applied to recent events studied in detail, from which many effects on riverbank vegetation are conserved.

Also flood frequency analysis (task 13), using the combination of systematic and palaeoflood data from dendrochronological sources, has some particularities. When the dates of the flood events and their estimated flows are available, these can be used to produce a statistical analysis to allow a frequency distribution function adjusted to these flows. For this, different formulas can be used to assign sample probability to different events, depending on whether they are exact, minimum or maximum levels, and on their degree of certainty (classified as EX, LB, UB and

DB type); and specific frequency distribution functions should be used (for both stationary and non-stationary models) based on setting thresholds or ranges, which have already been successfully used for including historic and palaeohydrological data (Francés 2004; Macdonald 2013). To facilitate these analyses, various computer applications that implement these formulas can be used comparatively, with programs including FRESH (University of Québec; Ouarda et al. 2004), AFINS (Grupo de Investigación de Hidráulica e Hidrología; UPV) and MAX (Stedinger et al. 1988). Finally, the same analyses should be carried out including the data from the dendrogeomorphology in the systematic record flow series from the nearby measuring stations, observing the possible improved fit by applying different statistical tests (Kolmogorov-Smirnov, PPCC, L-moments ratio, etc.).

Finally, our methodological proposal also includes improvements and new methods of cost-benefit analysis for risk mitigation measures. From the results of the maps and risk analysis, a detailed study should be made of all the

mitigation measures that can reasonably be applied in each case (predictive, preventive and corrective), evaluating the cost of applying each individually or in combination. When this battery of measures and possible logical combinations is available, their effects should be input into the hydrometeorological models, to study to what extent and in what conditions a reduction of risk levels could occur. Geostatistics and geographic information systems should be integrated for assessment of socioeconomic factors related to flood risk management. From these two estimates, the cost of the measures and the benefits of their application, complex cost–benefit analyses should be produced, using stochastic and geostatistic methods to take into account their spatial dimension (Sánchez-Silva 2004; Zhai & Ikeda 2008); to do this, computer-based tools should be used as specific developments for the MATLAB program package. In short, the aim of this analysis should be to integrate the concept of sustainability of these preventive measures with essential aspects related to risk calculation, definition of acceptability criteria and the optimization of investment. The results indicate optimum measures for minimizing risks of flash floods in terms of cost–benefit analysis (Ballesteros *et al.* submitted).

INITIAL RESULTS AND DISCUSSION OF ADVANTAGES AND LIMITATIONS

Both methodological proposals described and explained above are not only theoretical exercises. They have been

applied in different projects and study areas (Figure 4), mainly in Spain, but also in other areas of Europe such as the United Kingdom (Ruiz Villanueva *et al.* 2010b) or Poland.

The first results have been published over the last 3 years, or will be published soon, in some of the most prestigious journals, both in the botanic thematic category (*Tree Physiology*, *Tree-Ring Research*, etc.) and in hydrologic–geomorphological scope (*Journal of Hydrology*, *Hydrological Processes*, *Water Resources Research*, *Water Resources Management*, *Geomorphology*, *Catena*, etc.); i.e., see the publications of Ballesteros *et al.* (2010a, 2010b, 2011a, 2011b, 2012, submitted), Ruiz-Villanueva *et al.* (2010a) and Genova *et al.* (2011).

Nevertheless, this is the first time that well-developed methodological proposals have been published. The necessarily brief methodology section of the papers has prevented proper development of specific aspects of the analysis carried out (St. George *et al.* 1999). Such detailed explanation is necessary for two reasons: to allow other research teams to follow these techniques; and more importantly, to allow other scientists to replicate the analysis outlined in this work, to ensure they achieve the same results. For this reason, we consider it important to explain all the details of our advanced methods, and bring them to the scientific community so they can be corrected and completed.

Some of the methodologies presented are classical techniques that require no further discussion, such as tasks 1, 2, 4 and 11 to 15. Another group is pertinent to the




Figure 4 | Location of study areas where advanced dendrogeomorphological methods are being applied to the study of past floods: (a) in the world, Spain, Switzerland, United Kingdom, El Salvador and Poland; (b) in Spain, Navaluenga, Arenas de San Pedro, Guisando Segovia, Pajares de Pedraza, Taburiente and Valsain.

dendrochronological field and these tasks (3, 5, 6, 7 and 9) are well developed in the scientific literature (Picher 1990; Wilford et al. 2005) but for other applications (palaeoclimatology, vegetation dynamics, etc.). Finally, for another set of tasks (8 and 10), the relationship of linked tasks and the classification of evidence are described in detail here for the first time and require some discussion of their advantages and limitations.

One of the most innovative tasks that could be included in the second methodological proposal is isotope analysis in alpha-cellulose (Figure 3, task 8). This task is still at the development stage, and has not yet been tested, calibrated and validated. It should therefore be considered as a working hypothesis, rather than a methodological step. To analyse the isotope contents of the tree-ring wood for specific stable isotopes (^{16}O , ^{18}O , ^{13}C , etc.), the following procedures should be carried out or requested from specialized laboratories: samples of late and/or early wood from specific tree-rings; extraction of alpha-cellulose from the

wood; measurement of the $^{16}\text{O}/^{18}\text{O}$ isotope ratio and interpretation of the results. The aim of this novel technique is to study the relationship between the above ratio in the tree-ring wood corresponding to a particular year with flash flood events and the ratios measured in rainfall recorded in the main precipitation events of that year or season (taken from the CEDEX national measurement network). The aim of this comparison is to establish possible relationships between both ratios to study the meteorological origin (of frontal, of convective, of orographic, etc.) of the precipitation that generated the flash flood from tree-ring wood analysis, allowing two or more past event populations to be differentiated and their frequency and magnitude to be examined separately (Figure 5).

We are beginning to study the $^{16}\text{O}/^{18}\text{O}$ ratio patterns of different types of precipitation in central Spain corresponding to recent well-known flood events. These singular events will be selected looking for vegetative periods in which only one type of precipitation has been described in the




Figure 5 | Methodological procedure framework for the analysis of isotopes in tree-ring alpha-cellulose and its use for improving flood frequency analysis.

meteorological reports. Later, the alpha-cellulose $^{16}\text{O}/^{18}\text{O}$ ratio from the wood sequence used for dating the flood event by dendrochronological techniques will be analysed. Both isotopic ratios will be compared looking for statistical relations. These results will be used to extrapolate this relation to the entire tree-ring sequence, completing the past flood record and their classification in two or three event populations. The last step (equivalent to task 13) will be frequency analysis of each single population, and comparison with traditional frequency analysis of the entire flood dataset.

The proposed classification of dendrogeomorphological evidence applied to flood risk analysis is an important advance with respect to earlier classifications (Hupp 1988), because it is not a simple list of evidence without order and criteria. On the contrary, this proposal uses a clear criterion for the type of evidence, its spatial dimension and the element upon which the analysis is performed. This allows the potential user to easily locate its study elements, and the designations are internationally comparable. Furthermore, it has expanded from four or five evidence types of earlier classifications, to more than 30 different types, increasing the range of possibilities for study. Among the limitations is that the study of plant communities as FDE is not just a dendrogeomorphological analysis using its own techniques. Other aspects of population dynamics, species distributions and densities, etc., which are not studied exclusively in tree-ring dendrochronology, must also be taken into account.

The assignment of data obtained for each type of evidence (Table 2), although based on scientific literature and our own experience, remains hypothetical at present as very little work has been done to demonstrate validity. It would therefore be interesting to explore new areas with suitable studies, for example, using the angle of tilt of trees to obtain previous flood energy levels and thus estimate the water depth of palaeofloods.

CONCLUSIONS

Although dendrogeomorphological techniques have been used for the study of floods for more than 40 years, proposals for methodological approaches can still be advanced

to improve risk analysis. Two of these proposals are classification of dendrogeomorphological evidence and a working schema comprising 15 tasks, with some additional innovative techniques. The first proposal enables studies to normalize and standardize nomenclature improving the possibility of obtaining data on past floods. The advanced methods of the working schema, such as the isotope analysis of tree-rings, can improve flood frequency analysis and reduce uncertainties in risk analysis. Thus it is possible to obtain very accurate and reliable information from past floods and help prevent future flooding using dendrogeomorphological techniques.

ACKNOWLEDGEMENTS

This study was funded by the *MAS Dendro-Avenidas* research project (CGL2010-19274; www.dendro-avenidas.es) of the Spanish Ministry of Economy and Competitiveness, and by the *IDEA-GESPPNN* project (163/2010; www.idea-gesppnn.es) funded by the Spanish Ministry of Agriculture, Food and the Environment (National Parks Research Program). The authors wish to acknowledge the collaboration of: the other members of research teams of both projects; the Tagus and Duero River Water Authorities; the Environmental Authority, technicians and rangers of the Junta de Castilla y León; the village, city and town councils of Navalunga, Arenas de San Pedro, Guisando, Segovia and Pajares de Pedraza; and the Social and Cultural Division of Caja Avila (Bankia).

REFERENCES

- Alestalo, J. 1971 Dendrochronological interpretation of geomorphic processes. *Fennia* **105**, 1–140.
- Astrade, L. & Bégin, Y. 1997 Tree-ring response of *Populus tremula* L. and *Quercus robur* L. to recent spring floods of the Saône River, France. *Ecoscience* **4**, 232–239.
- Ballesteros, J. A., Stoffel, M., Bodoque, J. M., Bollschweiler, M., Hitz, O. & Díez-Herrero, A. 2010a Wood anatomy of *Pinus pinaster* Ait. following wounding by flash floods. *Tree-Ring Res.* **66** (2), 93–103.
- Ballesteros, J. A., Stoffel, M., Bollschweiler, M., Bodoque, J. M. & Díez-Herrero, A. 2010b Flash-flood impacts cause changes in wood anatomy of *Alnus glutinosa*, *Fraxinus angustifolia* and *Quercus pyrenaica*. *Tree Physiol.* **30**, 773–781.

- Ballesteros Cánovas, J. A., Eguibar, M., Bodoque, J. M., Díez-Herrero, A., Stoffel, M. & Gutiérrez-Pérez, I. 2011a [Estimating flash flood discharge in an ungauged mountain catchment with 2D hydraulic models and dendrogeomorphic paleostage indicators](#). *Hydrol. Process.* **25**, 970–979.
- Ballesteros Cánovas, J. A., Bodoque, J. M., Díez-Herrero, A., Sanchez-Silva, M. & Stoffel, M. 2011b [Calibration of floodplain roughness and estimation of palaeoflood discharge based on tree-ring evidence and hydraulic modelling](#). *J. Hydrol.* **403**, 103–115.
- Ballesteros-Cánovas, J. A., Díez-Herrero, A. & Bodoque, J. M. 2012 [Searching for useful non-systematic tree-ring data sources for flood hazard analysis using GIS tools](#). *Catena* **92**, 130–138.
- Ballesteros-Cánovas, J. A., Sanchez-Silva, M., Bodoque, J. M. & Díez-Herrero, A. (submitted) [An example of integrated approach to flood risk management: the case of Navalunga \(Central Spain\)](#). *Water Resour. Manage.* (submitted 2011-09-14).
- Benito, G. & Thorndyraft, V. 2004 [Use of systematic, palaeoflood and historical data for the improvement of flood risk estimation: an introduction](#). In: *Systematic, Palaeoflood and Historical Data for the Improvement of Flood Risk Estimation: Methodological Guidelines* (G. Benito & V. R. Thorndyraft, eds). CSIC – Centro de Ciencias Medioambientales, Madrid, pp. 5–14.
- Bianchi, M. & Zheng, C. M. 2009 [SGeMS: a free and versatile tool for three-dimensional geostatistical applications](#). *Groundwater* **47** (1), 8–12.
- Bollschweiler, M., Stoffel, M. & Ehmsich, M. 2006 [Channel activity of debris flows on forested cones – a case study using dendrogeomorphology](#). *Geophys. Res. Abstr.* **8**, 06463.
- Bollschweiler, M., Stoffel, M. & Schneuwly, D. M. 2008a [Dynamics in debris-flow activity on a forested cone – a case study using different dendroecological approaches](#). *Catena* **72**, 67–78.
- Bollschweiler, M., Stoffel, M., Schneuwly, D. M. & Bourqui, K. 2008b [Traumatic resin ducts in *Larix decidua* trees impacted by debris flows](#). *Tree Physiol.* **28**, 255–263.
- Cook, E. R. 1990 [A conceptual linear aggregate model for tree rings](#). In: *Methods of Dendrochronology: Applications in the Environmental Sciences* (E. R. Cook & L. A. Kairiukstis, eds). Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 98–104.
- Coriell, F. 2002 [Reconstruction of a Paleoflood Chronology for the Middlebury River Gorge using Tree Scars as Flood Stage Indicators](#). Unpublished Senior Thesis, Middlebury College, Department of Geology, Middlebury, Vermont, USA, 45 pp.
- Díez-Herrero, A., Ballesteros, J. A., Bodoque, J. M., Eguibar, M. A., Fernández, J. A., Génova, M., Laín, L., Llorente, M., Rubiales, J. M. & Stoffel, M. 2007 [Mejoras en la estimación de la frecuencia y magnitud de avenidas torrenciales mediante técnicas dendrogeomorfológicas](#). *Bol. Geológico Minero* **118** (4), 789–802.
- Díez-Herrero, A., Lain-Huerta, L. & Llorente-Isidro, M. 2009 [A Handbook on Flood Hazard Mapping Methodologies](#). Publications of the Geological Survey of Spain, Series Geological Hazards/Geotechnics No. 2, Madrid, Spain, 190 pp.
- Douglass, A. E. 1909 [Weather cycles in the growth of big trees](#). *Month. Weather Rev.* **37** (5), 225–237.
- Douglass, A. E. 1914 [A method for estimating rainfall by the growth of trees](#). In: *The Climatic Factor as Illustrated in Arid America* (E. Huntington, ed.). Publication 192, Carnegie Institute of Washington, Washington, DC, pp. 101–121.
- Elleder, L., Herget, J., Roggenkamp, T. & Nießen, A. 2013 [Historic floods in the city of Prague – a reconstruction of peak discharges for 1481–1825 based on documentary sources](#). *Hydrol. Res.* **44** (2), 202–214.
- Francés, F. 2004 [Flood frequency analysis using systematic and non-systematic information](#). In: *Systematic, Palaeoflood and Historical Data for the Improvement of Flood Risk Estimation: Methodological Guidelines* (G. Benito & V. R. Thorndyraft, eds). CSIC – Centro de Ciencias Medioambientales, Madrid, pp. 55–70.
- Fritts, H. C. 1976 [Tree Rings and Climate](#). Academic Press, New York, NY.
- Gaume, E., Bain, V., Bernardara, P., Newinger, O., Barbuc, M., Bateman, A., Blaskovicova, L., Blöschl, G., Borga, M., Dumitrescu, A., Daliakopoulos, I., Garcia, J., Irimescu, A., Kohnova, S., Koutroulis, A., Marchi, L., Matreata, S., Medina, V., Preciso, E., Sempere-Torres, D., Stancalie, G., Szolgay, J., Tsanis, I., Velasco, D. & Viglione, A. 2009 [A collation of data on European flash floods](#). *J. Hydrol.* **367**, 70–78.
- Génova, M., Ballesteros-Cánovas, J. A., Díez-Herrero, A. & Martínez-Callejo, B. 2011 [Correlating documentary records of historical floods with the dendrochronological dating of a wooden deck in the old Mint of Segovia, Spain](#). *Geoarchaeology* **26** (5), 786–808.
- Gottesfeld, S. 1996 [British Columbia flood scars: maximum flood-stage indicators](#). *Geomorphology* **14** (4), 319–325.
- Gottesfeld, A. S. & Gottesfeld, L. M. J. 1990 [Floodplain dynamics of a wandering river, dendrochronology of the Morice River, British Columbia, Canada](#). *Geomorphology* **3**, 159–179.
- Harrison, S. S. & Ried, J. R. 1967 [A flood-frequency graph based on tree-scar data](#). *Proc. N. D. Acad. Sci.* **21**, 23–33.
- Hupp, C. R. 1984 [Dendrogeomorphic evidence of debris flow frequency and magnitude at Mount Shasta, California](#). *Environ. Geol. Water Sci.* **6** (2), 121–128.
- Hupp, C. R. 1986 [Botanical evidence of floods and paleoflood frequency](#). Paper presented at the International Symposium on Flood Frequency and Risk Analysis, Baton Rouge, LA, USA.
- Hupp, C. R. 1988 [Plant ecological aspects of flood geomorphology and paleoflood history](#). In: *Flood Geomorphology* (V. R. Baker, R. C. Kochel & P. C. Patton, eds), Chapter 20. John Wiley and Sons, Chichester, UK, pp. 335–356.
- Hupp, C. R., Osterkamp, W. R. & Thornton, J. L. 1987 [Dendrogeomorphic evidence and dating of recent debris flows on Mount Shasta, northern California](#). U.S. Geol. Surv. Prof. Paper 1396-B, B1-B39.

- Kozłowski, T. T. 1997 Responses of woody plants to flooding and salinity. *Tree Physiol. Monogr.* **1**, 1–29.
- La Marche Jr., V. C. 1963 Origin and geologic significance of buttress roots of bristlecone pines, White Mountains, California. U.S. Geol. Surv. Prof. Paper 475-C, C149–C150.
- Lang, M., Fernández, J. F., Recking, A., Naulet, R. & Grau, P. 2004 Methodological guide for palaeoflood and historical peak discharge estimation. In: *Systematic, Palaeoflood and Historical Data for the Improvement of Flood Risk Estimation: Methodological Guidelines* (G. Benito & V. R. Thorndycraft, eds). CSIC – Centro de Ciencias Medioambientales, Madrid, pp. 43–53.
- Macdonald, N. 2013 Reassessing flood frequency for the River Trent, Central England, since AD 1320. *Hydrol. Res.* **44** (2), 215–233.
- McCord, V. A. S. 1990 Augmenting Flood Frequency Estimates using Flood-Scarred Trees. Unpublished PhD Dissertation, Department of Geosciences, University of Arizona, Tucson, Arizona, USA, 182 pp.
- Ouarda, T. B. M. J., Hamdi, Y. & Bobée, B. 2004 A general system for frequency estimation in hydrology (FRESH) with historical data. In: *Systematic, Palaeoflood and Historical Data for the Improvement of Flood Risk Estimation: Methodological Guidelines* (G. Benito & V. R. Thorndycraft, eds). CSIC – Centro de Ciencias Medioambientales, Madrid, pp. 71–74.
- Picher, J. R. 1990 Sample preparation, cross-dating, and measurement. In: *Methods of Dendrochronology: Applications in the Environmental Sciences* (E. R. Cook & L. A. Kairiuksis, eds). Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 40–51.
- Remy, N., Boucher, A. & Wu, J. 2009 *Applied Geostatistics with SGeMS: A User's Guide*. Cambridge University Press, Cambridge, UK, 284 pp.
- Ruiz-Villanueva, V., Díez-Herrero, A., Stoffel, M., Bollschweiler, M., Bodoque, J. M. & Ballesteros, J. A. 2010a Dendrogeomorphic analysis of flash floods in a small ungauged mountain catchment (Central Spain). *Geomorphology* **118**, 383–392.
- Ruiz-Villanueva, V., Thorndycraft, V. & Díez-Herrero, A. 2010b Análisis dendrogeomorfológico y registro de paleoinundaciones en la cuenca del río Till (NE Inglaterra): resultados preliminares. In: *Avances de la Geomorfología en España, 2008–2010* (X. Úbeda, D. Vericat & R. J. Batalla, eds). XI Reunión Nacional de Geomorfología, 20–24 Septiembre de 2010. UB, CTFC, ULI y SEG, Solsona, Lérida, pp. 521–524.
- Sánchez-Silva, M. 2004 Risk analysis and the decision making process in engineering. In: *Intelligent Knowledge-Based Systems: Business and Technology in the New Millennium* (C. T. Leondes, ed.). Kluwer Press, Dordrecht, The Netherlands, Vol. 4, Chapter 7.
- Schneuwly, D. M., Stoffel, M. & Bollschweiler, M. 2009a Formation and spread of callus tissue and tangential rows of resin ducts in *Larix decidua* and *Picea abies* following rockfall impacts. *Tree Physiol.* **29**, 281–289.
- Schneuwly, D. M., Stoffel, M., Dorren, L. K. A. & Berger, F. 2009b Three-dimensional analysis of the anatomical growth response of European conifers to mechanical disturbance. *Tree Physiology* **29**, 1247–1257.
- Sigafoos, R. S. 1964 Botanical evidence of floods and flood-plain deposition. Geol. Surv. Prof. Paper (U.S.) 485-A, 1–35.
- St. George, S. 2010 Tree rings as paleoflood and paleostage indicators. In: *Tree-Rings and Natural Hazards. A State-of-the-Art* (M. Stoffel, M. Bollschweiler, D. R. Butler & B. H. Luckman, eds). Springer, Dordrecht, Heidelberg, London, New York, pp. 233–239.
- St. George, R. S., Nielsen, E. & Brooks, G. 1999 Tree rings into the next Millennium! In: Report of Activities, 1999. Manitoba Industry, Trade and Mines, Geological Services, pp. 126–129.
- St. George, R. S., Nielsen, E., Conciatori, F. & Tardif, J. 2002 Trends in *Quercus macrocarpa* vessel areas and their implications for tree-ring paleoflood studies. *Tree-Ring Res.* **58** (1/2), 3–10.
- Stedinger, J., Surani, R. & Therivel, R. 1988 *MAX Manual*. Cornell University, Ithaca, NY, 51 pp.
- Stoffel, M. & Bollschweiler, M. 2009 What tree rings can tell about earth-surface processes. *Teaching the principles of dendrogeomorphology. Geogr. Compass* **3**, 1013–1037.
- Stoffel, M., Bollschweiler, M., Butler, D. R. & Luckman, B. H. 2010 *Tree Rings and Natural Hazards: A State of the Art*. Springer, Heidelberg, New York.
- Stoffel, M., Bollschweiler, M. & Hassler, G. R. 2006 Differentiating past events on a cone influenced by debris-flow and snow avalanche activity – a dendrogeomorphological approach. *Earth Surf. Proc. Land.* **31** (11), 1424–1437.
- Stoffel, M., Conus, D., Grichting, M. A., Lièvre, I. & Maître, G. 2008 Unraveling the patterns of late Holocene debris-flow activity on a cone in the Swiss Alps: chronology, environment and implications for the future. *Glob. Planet. Change* **60**, 222–234.
- Stoffel, M., Gärtner, H., Lièvre, I. & Monbaron, M. 2003 Comparison of reconstructed debris flow event years (Ritigraben, Switzerland) and existing flooding data in neighboring rivers. In: *Debris Flow Hazard Mitigation: Mechanics, Prediction, and Assessment* (D. Rickenmann & C. Chen, eds). Vol. 1, pp. 243–253. Millpress, Davos, Switzerland.
- Stoffel, M. & Hitz, O. M. 2008 Snow avalanche and rockfall impacts leave different anatomical signatures in tree rings of *Larix decidua*. *Tree Physiol.* **28**, 1713–1720.
- Stoffel, M., Lièvre, I., Conus, D., Grichting, M. A., Rietzo, H., Gärtner, H. W. & Monbaron, M. 2005 400 years of debris flow activity and triggering weather conditions: Ritigraben VS, Switzerland. *Arct. Antarct. Alp. Res.* **37** (3), 387–395.
- Stoffel, M. & Wilford, D. J. 2012 Hydrogeomorphic processes and vegetation: disturbance, process histories, dependencies and interactions. *Earth Surf. Proc. Land.* **37**, 9–22.
- Yanosky, T. M. 1982a Effects of flooding upon woody vegetation along parts of the Potomac River floodplain. Geol. Surv. Prof. Paper (U.S.) **1206**, 1–21.

- Yanosky, T. M. 1982b Hydrologic inferences from ring widths of flood-damaged trees, Potomac River, Maryland. *Environ. Geol. Water Sci.* **4**, 43–52.
- Yanosky, T. M. 1983 Evidence of floods on the Potomac River from anatomical abnormalities in the wood of flood-plain trees. *Geol. Surv. Prof. Paper (U.S.)* **1296**, 1–51.
- Yanosky, T. M. 1984 Documentation of high summer flows on the Potomac River from the wood anatomy of ash trees. *Water Resour. Bull.* **20**, 241–250.
- Yanosky, T. M. & Jarrett, R. D. 2002 Dendrochronologic evidence for the frequency and magnitude of paleofloods. Ancient floods, modern hazards: principles and application of paleoflood hydrology. *Water Sci. Appl.* **5**, 77–89.
- Wilford, D. J., Cherubini, P. & Sakals, M. E. 2005 *Dendroecology: A Guide for Using Trees to Date Geomorphic and Hydrologic Events*. Land Management Handbook No. 58, British Columbia Ministry of Forests, Canada.
- Zhai, G. & Ikeda, S. 2008 Empirical analysis of Japanese flood risk acceptability within multi-risk contest. *Nat. Hazard Earth Syst. Sci.* **8** (5), 1049–1066.
- Zielonka, T., Holeksa, J. & Ciapala, S. 2008 A reconstruction of flood events using scarred trees in the Tatra Mountains, Poland. *Dendrochronologia* **26**, 173–183.

First received 28 October 2011; accepted in revised form 6 July 2012. Available online 7 November 2012