

The National Cancer Institute Early Detection Research Network: Two Decades of Progress **AACR**

Robert C. Bast Jr¹ and Sudhir Srivastava²



Introduction

Over the last two decades, substantial progress has been made in the development of cancer biomarkers and their application to unmet needs across the spectrum of cancer care. NCI's Early Detection Research Network (EDRN) has had an important role in facilitating this progress (1). Having been formally established in 2000, the EDRN now includes more than 300 investigators from academic institutions and the private sectors who have discovered, developed, or evaluated more than a thousand cancer biomarkers that have been reported in more than 3,400 peer-reviewed articles, with 20% in high-impact journals. Possibly of greater importance, the EDRN has provided support, infrastructure, standards, quality control, and a process to evaluate and validate candidate biomarkers, facilitating "go" or "no go" decisions regarding their further development. Collaborative research has been encouraged with monthly meetings of working groups and semiannual gatherings of all EDRN investigators. Through the efforts of the EDRN and its investigators, seven biomarkers and devices have attained full FDA approval and an additional assay has received Breakthrough Device designation.

To recognize the progress that has been made over the last 20 years, as well as the contribution of the EDRN, this issue of *Cancer Epidemiology, Biomarkers & Prevention* includes a series of reviews prepared by members of the EDRN that captures the current state and paths to further development of cancer biomarkers, which permits not only early detection, but more effective triage, prognostication, and prediction and monitoring of response to treatment across a wide range of cancers from different sites.

Improving Established Screening Strategies

For some sites including lung, colon, prostate, breast, esophagus, and liver, screening is already a part of clinical practice. Here, biomarkers are being developed to improve specificity, identify individuals who would benefit from screening, decrease overdiagnosis, make screening more convenient, and increase the prevalence of screening. Biomarkers have generally been detected in blood, but have also been measured in more proximate sources, including sputum for lung cancer, stool for colorectal cancer, and urine for prostate cancer after digital rectal examination.

¹Department of Experimental Therapeutics, The University of Texas MD Anderson Cancer Center, Houston, Texas. ²Division of Cancer Prevention, National Cancer Institute, National Institutes of Health, Bethesda, Maryland.

See all articles in this *CEBP Focus* section, "NCI Early Detection Research Network: Making Cancer Detection Possible."

Corresponding Author: Robert C. Bast Jr, The University of Texas MD Anderson Cancer Center, 1400 Pressler Street, FCT 8.5066, Houston, TX 77030. Phone: 713-792-7743; Fax: 713-792-7864; E-mail: rbast@mdanderson.org

Cancer Epidemiol Biomarkers Prev 2020;29:2396-400

doi: 10.1158/1055-9965.EPI-20-1158

©2020 American Association for Cancer Research.

Lung cancer

One of the most important advances of the last two decades has been early detection of lung cancer with spiral "low-dose" CT (LDCT). Screening individuals with a heavy smoking history of 30 pack-years has detected enough early-stage diseases to improve survival among the screened by as much as 20%. LDCT screening, however, has a high false-positive rate and detects large numbers of small indeterminate nodules up to 30 mm in diameter (2). Indeterminate nodules have also been found in as many as a quarter of chest CT scans performed for other medical indications. To date, screening has been limited to heavy smokers, who represent only 27% of individuals destined to get lung cancer in our country. Discovery and development of biomarkers to estimate lung cancer risk with greater precision, to reduce false positives, and to categorize indeterminate pulmonary nodules from CT scans have great potential to improve early detection of lung cancer. A proteomic signature measured by mass spectroscopy (Nodify XL2; ref. 3) and a panel of seven autoantibodies (EarlyCDT-Lung; refs. 4, 5) with high negative predictive value are now commercially available to aid in identifying indeterminate lesions with a low risk of malignancy. The Percepta gene expression mRNA signature of bronchial epithelial washings has identified patients with suspicious lung lesions and negative bronchoscopy who are unlikely to have cancer and who can be spared additional invasive diagnostic procedures with a >90% negative predictive value (6).

Other alterations of nucleic acids in blood and sputum have been developed as biomarkers for lung cancer (7). Liquid biopsy has been used to detect EGFR DNA mutations that predict sensitivity or resistance to different tyrosine kinase inhibitors (8). Detection of stage I-II lung cancer has been reported by detecting mutations in peripheral blood circulating tumor DNA (ctDNA; ref. 9). Profiles of miRNA have been used to identify candidates for LDCT and to stratify risk of indeterminate nodules (10, 11). DNA methylation of CpG islands in the promoter regions of particular genes has been measured in plasma and sputum. Methylated DNA biomarkers have been used to identify individuals who might benefit from LDCT and to stratify indeterminate nodules. The most impressive results have been obtained with models based on whole-methylome sequencing, rather than with panels of individual biomarkers. One methylome-based study yielded 75% sensitivity for stage Ia lung cancer and 86% for stage Ib lung cancer, testing plasma from patients with indeterminate nodules (12).

Inflammation contributes to the initiation, progression, and metastasis of lung cancer (13). Inflammatory cytokines, including IL6, IL8, and C-reactive protein (CRP), are elevated in sera from patients with lung cancer, years prior to diagnosis (14). Development of better risk models that include inflammatory and noninflammatory biomarkers and that do not depend solely upon smoking history should extend LDCT to a larger population with a greater impact on lung cancer-related mortality.

Colorectal cancer

Colorectal cancer is the second leading cause of cancer-related death in the United States, second only to lung cancer. Despite the availability of multiple screening options, 40% of eligible patients are not screened

in our country, necessitating the need for additional accessible, patient acceptable, and cost-effective screening methods. In the United States, optical colonoscopy has been the standard of care, but in other developing and developed countries, two-stage strategies are in place where stool biomarkers prompt colonoscopy (15). The fecal immunochemical test (FIT) for blood in stool has generally provided the initial stage. DNA biomarkers, both genetic and epigenetic, have been evaluated alone and in combination with FIT (16). In 2014, the FDA approved the Cologuard (FIT-DNA) stool test that measures aberrantly methylated *BMP3* and *NDRG4* DNA, mutant *K-RAS* DNA, β -actin, and hemoglobin by FIT. While FIT-DNA detected 92.3% of all colorectal cancers and 94% of early-stage cancers, FIT detected 73.8% and 70% (17). With regard to the polyp precursors of colorectal cancer that are excised during colonoscopy, FIT-DNA detected 42.4% of advanced precancerous lesions, whereas FIT identified 23.8%. Thus, FIT-DNA offers significantly greater sensitivity as an initial phase of a two-stage strategy where a positive test prompts colonoscopy, but there is still room for improvement in detecting premalignant lesions. In addition, a blood test could provide greater convenience and possibly wider adoption by the community. A single blood test has been approved by the FDA for early detection of colorectal carcinoma (Epi proColon) that detects methylated *SEPT9* DNA in plasma. A meta-analysis of 25 studies indicated that the overall sensitivity for detecting colorectal cancer was 71% at 92% specificity and the sensitivity for detecting adenomas and polyps >1 cm was 23% at 91% specificity (18). Panels of blood protein biomarkers show greater promise for detecting premalignant disease. A combination of galectin-3 ligand, galectin-3, *CYFRA21*, and carcinoembryonic antigen (CEA) in serum has detected 56% of advanced polyps at 90% specificity (15). The combination of galectin-3 ligand and CEA has comparable sensitivity and is being tested prospectively in an EDRN-sponsored study.

Prostate cancer

Since implementing PSA screening in the early 1990s, prostate cancer-related mortality has decreased by nearly 50% in the United States (19). PSA, however, detects both aggressive, lethal cancers and more indolent disease, which does not shorten survival. The challenge of overdiagnosis and consequent overtreatment has prompted the U.S. Preventive Services Task Force to discourage routine screening and to individualize the use of PSA based on discussions between patients and their physicians. Over the last two decades, the EDRN has contributed to the development of tests that distinguish aggressive from indolent disease, reducing the number of unnecessary biopsies in men unlikely to have prostate cancer. The Prostate Health Index (PHI) includes total PSA, free PSA, and [−2]pro PSA. The PHI test has been approved by the FDA to identify men with a total PSA between 4 and 10 ng/mL who are likely to benefit from biopsies. Elevation of PCA3 long noncoding RNA in urine after prostate examination provides a highly specific biomarker for prostate cancer. The PROGENSA PCA3 assay has been FDA approved to identify men suspected of having prostate cancer who should have a repeat prostate biopsy. Combining detection of the *TMPRSS2:ERG* fusion gene and PCA3 in urine provides greater sensitivity than PCA3 alone (19). EDRN investigators are prospectively evaluating men with low-grade prostate cancers who have been followed with active surveillance and ultimately undergo prostatectomy to identify biomarkers that predict upgrading to more aggressive disease that needs treatment. EDRN studies are also underway to integrate changes in MRI with multiple biomarkers to detect more aggressive disease.

Breast cancer

Mammography has contributed to the 30% decline in breast cancer-related mortality observed over the last three decades, but screening has also detected substantial amounts of premalignant disease that might never have progressed. Overdiagnosis and consequent overtreatment can be reduced by developing biomarkers to identify indolent disease, incorporating biology and risk assessment in screening strategies, changing the pathology guideline for tumor classification, and refining the classification of precancerous lesions (20). Improvements in diagnostics and in biomarker-driven classification must be linked to a willingness to avoid unnecessary treatment in well-defined groups of patients.

Despite improvements in breast imaging, around 15% to 20% of breast cancers still go undetected (21). Many developing nations cannot afford routine mammography screening. Blood-based assays that detect early-stage breast cancers could be quite valuable as a cost-effective alternative to mammography or to detect cases missed by conventional imaging. While small cancers may not shed sufficient protein or DNA to permit detection, tumor-specific autoantibodies could be evoked by very early lesions. Autoantibodies have been detected against TP53, MUC1, HER2, and a variety of other tumor-associated antigens. Panels of protein targets have detected autoantibodies with 66% to 67% sensitivity at 82% to 84% specificity (21). The EDRN has encouraged further development of this approach.

Esophageal cancer

Most esophageal adenocarcinomas originate from metaplastic small intestinal epithelial cells at the gastroesophageal junction, termed Barrett esophagus (22). Detection and monitoring of Barrett esophagus are generally performed with upper endoscopy. EDRN investigators identified a two-marker panel of methylated *VIM* and methylated *CCNA1* DNA that detected 95% of Barrett esophagus, dysplasia, and esophageal adenocarcinoma with 91% specificity in brushings obtained at endoscopy (23). To make screening more convenient, a swallowable balloon-based device has been developed, called EsoCheck, that can sample cells at the gastroesophageal junction in 5 minutes without endoscopy. Detection of methylated *VIM* and methylated *CCNA1* DNA in cells sampled with the balloon device detected nondysplastic Barrett esophagus with a sensitivity of 90% and specificity of 92% (23). This strategy was approved by the FDA in 2019 and promises to enhance detection of Barrett esophagus and esophageal adenocarcinoma.

Hepatocellular cancer

Hepatocellular cancer (HCC) is the leading cause of cancer-related death worldwide related to endemic hepatitis B infection (24). HCC is also the fastest growing cause of cancer-related mortality in the United States due to increasing prevalence of nonalcoholic fatty liver disease, alcohol-related liver disease, and hepatitis C infection (24). HCC generally arises in the context of chronic liver disease where screening has been recommended every 6 months with abdominal ultrasound. With this approach, only 45% of cases are detected in early stage where surgery can be curative. Addition of alpha-fetoprotein (AFP) increases early detection to 63% (25). As 40% to 50% of HCCs do not elevate levels of AFP, additional biomarkers are being developed to aid in early detection. The EDRN provided independent validation of des-gamma carboxyprothrombin (DCP) and lectin-bound AFP (AFPL3) prior to FDA approval in 2011 (26). Using a GALAD score that integrates gender, age, AFP, AFPL3, and DCP, a sensitivity of 80% to 91% and specificity of 81% to 90% were achieved for detecting early-stage (<3 cm) disease across several international cohorts (27), but in a

German study of patients with nonalcoholic fatty liver disease, sensitivity was only 68% at 95% specificity (28). EDRN investigators are currently evaluating more than a dozen protein, glycomics, and nucleic acid biomarkers that could further improve sensitivity for early detection of HCC.

Developing New Screening Strategies

Effective screening strategies have not yet been established for several malignancies, including ovarian cancer, pancreatic cancer, and mesothelioma.

Ovarian cancer

Over the last three decades, 5-year survival for patients with ovarian cancer has improved from 37% to 46% through more widespread use of cytoreductive surgery and combination chemotherapy, but the cure rate of <40% has changed little due, in part, to detection at a late stage (III–IV) in 75% to 80% of cases. Currently, there is no established screening test for ovarian cancer, but a two-stage strategy where rising serum CA125 triggers transvaginal sonography shows promise (29). In the United Kingdom Collaborative Trial for Ovarian Cancer Screening that included around 200,000 postmenopausal women, a prespecified subset that excluded patients with prevalent disease had a 20% reduction in ovarian cancer–related mortality (30). Statistical bounds around this estimate are broad and reanalysis will be required later this year before it can be accepted. Whatever the outcome, there is room for improvement in the sensitivity for both stages of the strategy. The EDRN has supported studies of biomarker panels that can detect early-stage (I–II) disease at 98% specificity. A four-institution EDRN collaboration is studying different autoantibodies and antigen–autoantibody complexes. Notably, anti-TP53 autoantibodies are associated with 22% of ovarian cancers and can be elevated 8 months prior to CA125 and 22 months prior to clinical diagnosis in patients who do not experience a rise in CA125 (31). Multiple autoantibodies are being evaluated with standard panels of sera from patients with early-stage ovarian cancer and from controls to identify biomarkers that can detect cases missed by CA125.

Currently, only about half of women with ovarian cancer in the United States are referred at diagnosis to gynecologic oncologists who are specially trained to perform complete excision of all visible cancer, improving overall survival. Most generally present with a pelvic mass that can either be malignant or benign. The EDRN has supported development of three FDA-approved biomarker panels [ROMA (CA125 and HE4; ref. 32), OVA1 (CA125, transferrin, transthyretin, apolipoprotein AI, and beta-2 microglobulin; ref. 33), and OVERA (CA125, HE4, FSH, transferrin, apolipoprotein A1, transferrin, HE4, and FSH; ref. 34)] that identify women with pelvic masses who are most likely to have ovarian cancer, rather than benign disease, and who would benefit from referral to a specially trained surgeon.

Pancreatic cancer

Pancreatic ductal adenocarcinoma (PDAC) is another malignancy with poor prognosis that is diagnosed at late stage in 85% of cases (35). For the 15% of patients with disease limited to the pancreas, approximately 50% will survive 4.5 years, compared with 5% at 5 years with metastatic disease. PDAC is generally detected with imaging, but image-based screening is recommended only for individuals with a strong family history and/or predisposing germ line mutation (*ATM*, *BRCA2*, *BRCA1*, *PALB2*, *p16/CDKN2A*, *hMLH1*, *hMSH2*, and *hPMS6*). Blood biomarkers are required to identify individuals at

average genetic risk who would benefit from imaging. Other risk factors include recent onset of diabetes and the presence of intraductal papillary mucinous neoplasms (IPMN) and mucinous cystic neoplasms (MCN) that can progress to PDAC in 3% to 6% of cases. As the mortality of pancreatic surgery can be as high as 2% to 4% in some centers, biomarkers in cyst fluid that can distinguish premalignant from benign lesions would aid in appropriate management.

Consortia of EDRN investigators have established reference sets of sera from early-stage (I–II) patients and fluids from pancreatic cysts to evaluate conventional and novel biomarkers. A panel of serum CA19-9, sTRA, thrombospondin, DCD, TIMP1, MUC4, angiostatin, and PMS2 detected 79% of patients with early-stage PDAC at 95% specificity (35). Abnormalities in cyst fluids are being evaluated with next-generation sequencing of DNA, assays for methylated DNA, and measurement of telomerase, mucins, and DAS-1. Several other candidate biomarkers are being evaluated for biomarkers in blood and cyst fluids.

Mesothelioma

Malignant pleural mesothelioma (MPM) is a rare disease that occurs in asbestos-exposed individuals after 20 to 40 years (36). Less than 5% of patients are detected in stage I. Soluble mesothelin-related peptide is the one FDA-approved biomarker for aiding in diagnosis of MPM. EDRN resources and investigators have contributed to the validation of the MESOMARK assay for diagnosing and monitoring mesothelioma. In different meta-analyses, MPM exhibited a sensitivity of 32% to 64% at 89% to 95% specificity (37, 38). In longitudinal screening studies, MPM could be elevated up to a year before diagnosis. Other biomarkers are being evaluated including proteins (osteopontin, fibulin-3, IL6, vimentin, HMGB1, calretinin, ENOX-2, thioredoxin-1, and VEGF), SOMAmers, glycopeptide signatures, miRNA, ctDNA, and methylated DNA. Promising panels will need to be validated (39).

Universal screening strategies

Five EDRN investigators have helped to develop CancerSEEK that has attempted to detect eight types of cancer in stage I–III by assaying 61 mutations/amplicons in 16 genes and eight protein biomarkers (16). A case–control study was performed with 1,005 patients with cancer (ovary, liver, stomach, pancreas, esophagus, colorectal, lung, and breast cancer) and 812 controls (40). The median sensitivity across the eight cancer types was 70% at >99% specificity. A machine learning algorithm was able to localize the source of the cancer to one of two anatomic sites in a median of 83% of patients (40). In the DETECT-A study of 10,006 women not previously known to have cancer, DNA mutations and protein biomarkers detected 26 cancers and 15 were confirmed and localized by PET-CT (41). While this is a work in progress, there is great potential to detect cancer from multiple organs.

Novel Types of Biomarkers and Methods of Analysis

With EDRN support, new types of biomarkers have been developed over the last two decades, including circular RNA (circRNA). Artificial intelligence has been applied to the analysis of imaging data.

circRNA

circRNA constitutes a class of single-stranded RNAs that are covalently closed in a loop structure without polarity or polyadenylated tails (42). circRNAs are abundant and resist degradation

by exoribonucleases, providing stability in blood and body fluids. Both cancer-associated and tissue-associated circRNAs have been described. Expression of different circRNAs has been linked to prognosis across at least a dozen of different cancer types. circRNAs can predict response or toxicity after radiotherapy, cytotoxic and targeted drugs, endocrine treatment, and immunotherapy. Studies in patients with non-small cell lung, breast, colorectal, gastric, pancreatic, HCC, genitourinary (GU), and head and neck cancers suggest that circRNAs also have potential for early detection.

Radiomics

Early detection of cancers utilizes not only biomarkers, but imaging techniques, including LDCT for lung cancer, X-ray mammography, ultrasound for HCC, or optical imaging for esophageal and colorectal cancers (43). Traditionally, imaging information has been assessed qualitatively, but over the last decade, use of computerized tools has permitted conversion of images into quantitative data (radiomics), with subsequent analysis of these data using artificial intelligence, including convolutional neural networks and generative adversarial networks. This type of analysis requires large amounts of well-annotated data. Radiomics is improving diagnostic accuracy of early lesions, defining risk, and distinguishing malignant or aggressive cancers from benign disease. These techniques have been used to evaluate dysplastic nevi, malignant melanomas, indeterminate pulmonary nodules, malignant breast lesions, pancreatic IPMNs, and prostate cancers during active surveillance and liver stiffness in patients at risk for HCC.

Principles of Biomarker Development Moving Forward

Pitfalls

The number of useful new biomarkers introduced into the clinic remains limited. Despite hundreds of candidates, few maintain their sensitivity, specificity, and predictive values when tested in the clinic (44). False discovery relates to limited datasets, reproducibility, and tumor heterogeneity. Each of these challenges is addressed by the EDRN. ctDNA is a recent example with limitation in specificity, sensitivity, fragmentation, lead time, mutant allele fraction, and clin-

ical relevance. Even when ctDNA is analyzed in 10 mL of blood, it will be difficult to detect cancers <1 cm in diameter. Analysis of ctDNA with artificial intelligence algorithms may improve sensitivity.

Rigor

Since 2000, the EDRN has introduced rigor into biomarker development with guidelines, study design standards, biomarker reference sets, and the most rigorous blinding policy in the biomarker field (45). Importantly, the EDRN has developed an array of statistical and computational tools for early detection biomarker evaluation and developed a multidisciplinary team-science approach. Moving forward, the EDRN can strengthen its discovery pipeline, conduct better and more efficient validation studies, and develop even more effective statistical and computational tools.

Conclusion

This edition of *Cancer Epidemiology, Biomarkers & Prevention* should provide a useful assessment of the current contribution of biomarkers to early detection and the unmet needs that still must be addressed. We are grateful to all members of the EDRN who have contributed to this issue and the many achievements of the Network.

Disclosure of Potential Conflicts of Interest

R.C. Bast Jr reports grants from the NCI Early Detection Research Network Clinical Validation Center (early detection of ovarian cancer) during the conduct of the study and other from Fujirebio Diagnostics Inc. (royalties for the discovery of CA125) outside the submitted work. No potential conflicts of interest were disclosed by the other author.

Acknowledgments

The exceptional administrative assistance of Ruby Robinson in assembling EDRN reviews is gratefully acknowledged. This work was supported by funds from the NCI Early Detection Research Network (5 U01 CA200462-04) and generous donations from the Ann and Henry Zarrow Foundation, the Mossy Foundation, the Roberson Endowment, Stuart and Gaye Lynn Zarrow, Barry Elson, and Arthur and Sandra Williams.

Received August 3, 2020; revised August 29, 2020; accepted August 31, 2020; published first December 1, 2020.

References

1. Srivastava S, Wagner PD. The Early Detection Research Network: a national infrastructure to support the discovery, development, and validation of cancer biomarkers. *Cancer Epidemiol Biomarkers Prev* 2020;29:2401–10.
2. Ostrin EJ, Sidransky D, Spira A, Hanash SM. Biomarkers for lung cancer screening and detection. *Cancer Epidemiol Biomarkers Prev* 2020;29:2411–5.
3. Silvestri GA, Tanner NT, Kearney P, Vachani A, Massion PP, Porter A, et al. Assessment of plasma proteomics biomarker's ability to distinguish benign from malignant lung nodules: results of the PANOPTIC (Pulmonary Nodule Plasma Proteomic Classifier) trial. *Chest* 2018;154:491–500.
4. Massion PP, Healey GF, Peek LJ, Fredericks L, Sewell HF, Murray A, et al. Autoantibody signature enhances the positive predictive power of computed tomography and nodule-based risk models for detection of lung cancer. *J Thoracic Oncol* 2017;12:578–84.
5. Edelsberg J, Weycker D, Atwood M, Hamilton-Fairley G, Jett JR. Cost-effectiveness of an autoantibody test (EarlyCDT-Lung) as an aid to early diagnosis of lung cancer in patients with incidentally detected pulmonary nodules. *PLoS One* 2018;13:e0197826.
6. Silvestri GA, Vachani A, Whitney D, Elashoff M, Porta Smith K, Ferguson JS, et al. A bronchial genomic classifier for the diagnostic evaluation of lung cancer. *N Engl J Med* 2015;373:243–51.
7. Farooq M, Herman JG. Noninvasive diagnostics for early detection of lung cancer: challenges and potential with a focus on changes in DNA methylation. *Cancer Epidemiol Biomarkers Prev* 2020;29:2416–22.
8. Malapelle U, Sirera R, Jantus-Lewintre E, Reclusa P, Calabuig-Fariñas S, Blasco A, et al. Profile of the Roche Cobas® EGFR mutation test v2 for non-small cell lung cancer. *Expert Rev Mol Diagn* 2017;17:209–15.
9. Bettgowda C, Sausen M, Leary RJ, Kinde I, Wang Y, Agrawal N, et al. Detection of circulating tumor DNA in early- and late-stage human malignancies. *Sci Transl Med* 2014;6:224ra24.
10. Montani F, Marzi MJ, Dezi F, Dama E, Carletti RM, Bonizzi G, et al. MiR-test: a blood test for lung cancer early detection. *J Natl Cancer Inst* 2015;107:djv063.
11. Sozzi G, Boeri M, Rossi M, Verri C, Suatoni P, Bravi F, et al. Clinical utility of a plasma-based miRNA signature classifier within computed tomography lung cancer screening: a correlative MILD trial study. *J Clin Oncol* 2014;32:768–73.
12. Liang W, Zhao Y, Huang W, Gao Y, Xu W, Tao J, et al. Non-invasive diagnosis of early-stage lung cancer using high-throughput targeted DNA methylation sequencing of circulating tumor DNA (ctDNA). *Theranostics* 2019;9:2056–70.
13. Lim RJ, Liu B, Krysan K, Dubinett SM. Lung cancer and immunity markers. *Cancer Epidemiol Biomarkers Prev* 2020;29:2423–30.

14. Pine SR, Mechanic LE, Enewold L, Chaturvedi AK, Katki HA, Zheng YL, et al. Increased levels of circulating interleukin 6, interleukin 8, C-reactive protein, and risk of lung cancer. *J Natl Cancer Inst* 2011;103:1112–22.
15. Bresalier RS, Grady WM, Markowitz SD, Nielsen HJ, Batra SK, Lampe PD. Biomarkers for early detection of colorectal cancer: the Early Detection Research Network, a framework for clinical translation. *Cancer Epidemiol Biomarkers Prev* 2020;29:2431–40.
16. Cohen JD, Diergaarde B, Papadopoulos N, Kinzler KW, Schoen RE. Tumor DNA as a cancer biomarker through the lens of colorectal neoplasia. *Cancer Epidemiol Biomarkers Prev* 2020;29:2441–53.
17. Imperiale TF, Ransohoff DF, Itzkowitz SH, Levin TR, Lavin P, Ligard GP, et al. Multitarget stool DNA testing for colorectal-cancer screening. *N Engl J Med* 2014;370:1287–97.
18. Nian J, Sun X, Ming SY, Yan C, Ma Y, Feng Y, et al. Diagnostic accuracy of methylated SEPT9 for blood-based colorectal cancer detection: a systematic review and meta-analysis. *Clin Transl Gastroenterol* 2017;8:e216.
19. Liss MA, Leach RJ, Sanda MG, Semmes OJ. Prostate cancer biomarker development: National Cancer Institute's Early Detection Research Network Prostate Cancer Collaborative Group review. *Cancer Epidemiol Biomarkers Prev* 2020;29:2454–62.
20. Hewitt K, Son J, Glencer A, Borowsky AD, Cooperberg MR, Esserman LJ. The evolution of our understanding of the biology of cancer is the key to understanding and avoiding overdiagnosis and overtreatment. *Cancer Epidemiol Biomarkers Prev* 2020;29:2463–74.
21. Rauf F, Anderson KS, LaBaer J. Autoantibodies in early detection of breast cancer. *Cancer Epidemiol Biomarkers Prev* 2020;29:2475–85.
22. Grady WM, Yu M, Markowitz SD, Chak A. Barrett's esophagus and esophageal adenocarcinoma biomarkers. *Cancer Epidemiol Biomarkers Prev* 2020;29:2486–94.
23. Moinova HR, LaFramboise T, Lutterbaugh JD, et al. Identifying DNA methylation biomarkers for non-endoscopic detection of Barrett's esophagus. *Sci Transl Med* 2018;10:eaa05848.
24. Parikh ND, Mehta AS, Singal AG, Block T, Marrero JA, Lok AS. Biomarkers for the early detection of hepatocellular carcinoma. *Cancer Epidemiol Biomarkers Prev* 2020;29:2495–503.
25. Tzartzeva K, Obi J, Rich NE, Parikh ND, Marrero JA, Yopp A, et al. Surveillance imaging and alpha fetoprotein for early detection of hepatocellular carcinoma in patients with cirrhosis: a meta-analysis. *Gastroenterology* 2018;154:1706–18.
26. Marrero JA, Feng Z, Wang Y, Nguyen MH, Befeler AS, Roberts LR, et al. Alpha-fetoprotein, des-gamma carboxyprothrombin, and lectin-bound alpha-fetoprotein in early hepatocellular carcinoma. *Gastroenterology* 2009;137:110–8.
27. Berhane S, Toyoda H, Tada T, Kumada T, Kagebayashi C, Satomura S, et al. Role of the GALAD and BALAD-2 serologic models in diagnosis of hepatocellular carcinoma and prediction of survival in patients. *Clin Gastroenterol Hepatol* 2016;14:875–86.
28. Best J, Bechmann LP, Sowa JP, Sydor S, Dechene A, Pflanz K, et al. GALAD score detects early hepatocellular carcinoma in an international cohort of patients with nonalcoholic steatohepatitis. *Clin Gastroenterol Hepatol* 2020;18:728–35.
29. Bast RC Jr, Lu Z, Han CY, Lu KH, Anderson KS, Drescher CW, et al. Biomarkers and strategies for early detection of ovarian cancer. *Cancer Epidemiol Biomarkers Prev* 2020;29:2504–12.
30. Jacobs IJ, Menon U, Ryan A, Gentry-Maharaj A, Burnell M, Kalsi JK, et al. Ovarian cancer screening and mortality in the UK Collaborative Trial of Ovarian Cancer Screening (UKCTOCS): a randomised controlled trial. *Lancet* 2016;387:945–56.
31. Yang WL, Gentry-Maharaj A, Simmons A, Ryan A, Fourkala EO, Lu Z, et al. Elevation of TP53 autoantibody before CA125 in preclinical invasive epithelial ovarian cancer. *Clin Cancer Res* 2017;23:5912–22.
32. Moore RG, Miller MC, Disilvestro P, Landrum LM, Gajewski W, Ball JJ, et al. Evaluation of the diagnostic accuracy of the risk of ovarian malignancy algorithm in women with a pelvic mass. *Obstet Gynecol* 2011;118:280–8.
33. Ueland FR, Desimone CP, Seamon LG, Miller RA, Goodrich S, Podzielinski I, et al. Effectiveness of a multivariate index assay in the preoperative assessment of ovarian tumors. *Obstet Gynecol* 2011;117:1289–97.
34. Coleman R, Herzog TJ, Chan DW, Munroe DG, Pappas TC, Smith A, et al. Validation of a second-generation multivariate index assay for malignancy risk of adnexal masses. *Am J Obstet Gynecol* 2016;215:82.
35. Liu Y, Kaur S, Huang Y, Fahrman JF, Rinaldo JA, Hanash SM, et al. Biomarkers and strategy to detect preinvasive pancreatic cancer: state of the field and the impact of the EDRN. *Cancer Epidemiol Biomarkers Prev* 2020;29:2513–23.
36. Pass HI, Alimi M, Carbone M, Yang H, Goparaju CM. Mesothelioma biomarkers: a review highlighting contributions from the Early Detection Research Network. *Cancer Epidemiol Biomarkers Prev* 2020;29:2524–40.
37. Luo L, Shi HZ, Liang QL, Jiang J, Qin SM, Deng JM. Diagnostic value of soluble mesothelin-related peptides for malignant mesothelioma: a meta-analysis. *Respir Med* 2010;104:149–56.
38. Hollevoet K, Reitsma JB, Creaney J, Grigoriu BD, Robinson BW, Scherpereel A, et al. Serum mesothelin for diagnosing malignant pleural mesothelioma: an individual patient data meta-analysis. *J Clin Oncol* 2012;30:1541–9.
39. Bonotti A, Foddis R, Landi S, Melaiu O, De Santi C, Giusti L, et al. A novel panel of serum biomarkers for MPM diagnosis. *Dis Markers* 2017;2017:3510984.
40. Cohen JD, Li L, Wang Y, Thoburn C, Afsari B, Danilova L, et al. Detection and localization of surgically resectable cancers with a multi-analyte blood test. *Science* 2018;359:926–30.
41. Lennon AM, Buchanan AH, Kinde I, Warren A, Ashley Honushefsky A, Cohain AT, et al. Feasibility of blood testing combined with PET-CT to screen for cancer and guide intervention. *Science* 2020;369:eabb9601.
42. Brown JR, Chinnaiyan AM. The potential of circular RNAs as cancer biomarkers. *Cancer Epidemiol Biomarkers Prev* 2020;29:2541–55.
43. Gillies RJ, Schabath MB. Radiomics improves cancer screening and early detection. *Cancer Epidemiol Biomarkers Prev* 2020;29:2556–67.
44. Ren AH, Fiala CA, Diamandis EP, Kulasingam V. Pitfalls in cancer biomarker discovery and validation with emphasis on circulating tumor DNA. *Cancer Epidemiol Biomarkers Prev* 2020;29:2568–74.
45. Feng Z, Pepe MS. Adding rigor to biomarker evaluations: the EDRN experience. *Cancer Epidemiol Biomarkers Prev* 2020;29:2575–82.