

and glycemic control as measured by A1C became similar in the two original DCCT treatment groups, and yearly follow-up has continued through the present time. The current study includes all participants with available frozen plasma collected at or near the end of the DCCT, excluding seven participants who were pregnant at the time of plasma collection ($N = 1,193$; 83% of randomized DCCT participants) (19).

Circulating vitamin D metabolites

Concentrations of 25(OH)D, 1,25(OH)₂D, and 24,25(OH)₂D were measured in plasma samples that were obtained at or soon before the end of the DCCT (19). The median calendar year of collection was 1992 (range 1988–1993). Vitamin D metabolites are known to be stable during long-term storage. Plasma samples were stored continuously at -70°C until measurements were performed using high-performance liquid chromatography–tandem mass spectrometry. These methods quantify 25(OH)D₂ and 25(OH)D₃ [summed to evaluate total 25(OH)D], 1,25(OH)₂D₂ and 1,25(OH)₂D₃ [summed to evaluate total 1,25(OH)₂D], and 24,25(OH)₂D₃ [referred to throughout as 24,25(OH)₂D] (20–22). The lower limits of detection for 25(OH)D₂, 25(OH)D₃, 1,25(OH)₂D₂, 1,25(OH)₂D₃, and 24,25(OH)₂D₃ were 2.5 nmol/L (965 samples below), 2.5 nmol/L (0 samples below), 12.5 pmol/L (973 samples below), 12.5 pmol/L (0 samples below), and 1.25 nmol/L (13 samples below), respectively. Measurements that were below the lower limits of detection were truncated to the limit. Interassay imprecision was 6.63% at 70.2 nmol/L for 25(OH)D₃, 5.27% at 74.5 nmol/L for 25(OH)D₂, 9.6% at 86.3 pmol/L for 1,25(OH)₂D₂, 12.2% at 94.6 pmol/L for 1,25(OH)₂D₃, and 10.5% at 7.9 nmol/L for 24,25(OH)₂D₃. The calibration for the measurement of 25(OH)D was verified using Standard Reference Material 972 from National Institute of Standards and Technology with accuracy of 91–95% for 25(OH)D₃ and 100–116% for 25(OH)D₂.

Measures of subclinical cardiovascular disease

Computed tomography (CT) was performed by certified technicians between November 2000 and March 2003, 11–20 years after enrollment into the DCCT and 8–15 years after collection of blood for vitamin D metabolite measurement (median 10 years), in 1,205 (86%) of the

surviving 1,404 participants with specific patient consent. CT was performed in 19 scanning sites using a C-150 cardiac-gated electron beam CT scanner ($n = 9$; Imatron, San Francisco, CA), a Lightspeed ($n = 7$; General Electric Medical Systems, Waukesha, WI), or a Volume Zoom (Siemens, Erlanger, Germany) multi-detector CT system; a Lightspeed Marconi MX-8000 (GE); or a Somatom 4 (Siemens) ($n = 3$). All participants were scanned twice over calibration phantoms of known physical calcium concentration. Scans were read centrally at the Harbor-UCLA Research and Education Institute (University of California, Los Angeles, Torrance, CA) to identify and quantify coronary artery calcium (CAC), calibrated according to the readings of the phantom using the method of Agatston et al. (23). The average score from the two scans was used in the analysis. Readers were masked to subject identity and prior treatment assignment.

Carotid IMT was measured by B-mode ultrasonography in 1994 and again in 2000/2001 (24). IMT was measured in both the common and internal carotid arteries. A single longitudinal lateral view of the distal 10 mm of the right and left common carotid arteries and three longitudinal views in different imaging planes of each internal carotid artery were obtained. The internal carotid artery was defined as including both the carotid bulb and the 10-mm segment distal to the tip of the flow divider that separates the internal from the external carotid artery. Studies were performed by certified technicians at the clinical centers, recorded on videotapes, and read in a central unit (Tufts University) by a single reader (24).

Covariates

Covariates were ascertained concurrently with measurement of vitamin D metabolites. Demographic characteristics and smoking were ascertained by questionnaire. Solar irradiation of each DCCT site was quantified as the mean annual ultraviolet index in 1992, as published online by the National Weather Service (<ftp://ftp.cpc.ncep.noaa.gov/long/uv/cities>, accessed October 2011). A1C was measured using high-performance liquid chromatography (25). Glomerular filtration rate (GFR) was estimated using the Chronic Kidney Disease–Epidemiology (CKD-EPI) equation on the basis of serum creatinine concentration, age, sex, and race (26). In order to increase precision,

baseline albumin excretion rate (AER) for this study was calculated as the geometric mean of the AER measured concurrently with vitamin D metabolites and the AER preceding vitamin D measurement, usually within 1 year. Plasma concentrations of parathyroid hormone and fibroblast growth factor-23 were measured concurrently with vitamin D metabolites in a subset of 300 participants using immunoassay and ELISA, respectively (19). Data on vitamin D supplement use were not available.

Statistical methods

Associations of circulating vitamin D metabolites with CAC were assessed with a binomial regression model to estimate the relative risk for presence (vs. absence) of CAC. A study of the Multi-Ethnic Study of Atherosclerosis (MESA) developed the CAC categories: 0, 0–100, 100–300, and >300 (27). Proportional odds models were estimated using these established categories as the outcome. Coefficients from the polytomous model are interpreted as the odds ratio for having higher CAC associated with a unit difference in the exposure (28). This model has the advantage of using all of the available data, while the interpretation of the coefficients remains meaningful. In addition, Tobit regression models were estimated to the combined binary and continuous CAC outcomes (29). Nonzero CAC scores were transformed using the natural logarithm because the distribution is highly skewed. The Tobit regression coefficient represents the log ratio of the geometric mean CAC score per unit increase in the covariate, assuming some true measurable calcification for all subjects, including those with undetectable levels.

Associations of vitamin D metabolites with IMT were estimated using linear regression. Internal carotid IMT was log transformed prior to analysis, yielding estimates reflecting relative differences in internal carotid IMT. Generalized estimating equations were used to determine valid inference while using both the measurements from 1994 and the measurements from 2000/2001 for each subject simultaneously (30). Due to observed differences in IMT between 1994 and 2000/2001, the models were adjusted for year of measurement. No interactions by year were observed.

Plasma 25(OH)D concentrations were examined in clinically relevant categories: ≥ 75 nmol/L (30 ng/mL), 50–75

Table 1—Description of DCCT/EDIC study population at the time of biomarker measurement, except for CAC/IMT measures, which occurred at year 6

| Variable | All measured participants | 25(OH)D (nmol/L) | | |
|--|---------------------------|------------------|--------------|--------------|
| | | ≥75 | 50–75 | <50 |
| N | 1,193 | 314 | 550 | 329 |
| Age (years)* | 32.4 (2.6) | 31.5 (2.6) | 32.5 (2.6) | 32.2 (2.6) |
| Diabetes | | | | |
| duration (years)* | 7.5 (2.1) | 6.8 (2) | 7.5 (2.1) | 7.5 (2.1) |
| Female sex, N (%)* | 564 (47) | 103 (36) | 133 (47) | 149 (49) |
| Nonwhite race, N (%)* | 42 (4) | 3 (1) | 1 (0) | 9 (3) |
| DCCT intensive | | | | |
| diabetes therapy, N (%) | 598 (50) | 160 (55) | 139 (49) | 157 (51) |
| Current smoking, N (%) | 236 (20) | 60 (21) | 54 (19) | 49 (16) |
| BMI (kg/m ²)* | 25.7 (1.9) | 25.1 (1.7) | 25.6 (1.9) | 25.6 (1.8) |
| Systolic blood pressure (mmHg)* | 114.8 (3.4) | 115.2 (3.4) | 113.3 (3.4) | 114.8 (3.4) |
| Diastolic blood pressure (mmHg)* | 74 (2.9) | 74 (2.9) | 72.9 (3) | 74 (2.8) |
| A1C (%) | 8.1 (1.3) | 8.1 (1.2) | 8.2 (1.3) | 8.2 (1.3) |
| Iothalamate GFR (mL/min) | 132.9 (4.8) | 135.1 (4.3) | 136.7 (5.1) | 131.5 (5.1) |
| Serum creatinine (μmol/L) | 65.3 (11.3) | 70.4 (12.2) | 69.1 (12.3) | 54.2 (3.5) |
| Estimated GFR (mL/min/1.73 m ²) | 122.3 (5) | 124.9 (5.1) | 122.7 (5) | 122.5 (4.9) |
| Creatinine clearance (mL/min)* | 133.6 (5.6) | 138.6 (5.7) | 133.8 (5.6) | 132.9 (5.6) |
| HDL cholesterol (mmol/L) | 1.3 (0.6) | 1.3 (0.6) | 1.3 (0.6) | 1.3 (0.6) |
| LDL cholesterol (mmol/L) | 2.8 (0.9) | 2.8 (0.9) | 2.8 (0.9) | 2.9 (0.9) |
| Total 25(OH)D (nmol/L)* | 63.3 (4.7) | 90.8 (4) | 62.2 (2.6) | 38.9 (2.9) |
| 24,25(OH) ₂ D ₃ (nmol/L)* | 9.9 (2.3) | 15.7 (2.2) | 9.6 (1.6) | 4.8 (1.4) |
| Total 1,25(OH) ₂ D (pmol/L)* | 107.5 (5.5) | 117.9 (5.7) | 105.4 (5.4) | 101.1 (5.4) |
| FGF-23 (pg/mL)* | 31.4 (2.9) | 34.8 (3.1) | 30.1 (2.8) | 30.1 (2.7) |
| PTH (ng/L)* | 30.8 (3.7) | 26.3 (3.1) | 28.1 (3.4) | 30.7 (3.3) |
| Nonzero CAC, N (%) | 329 (30) | 90 (34) | 85 (33) | 75 (27) |
| CAC score (Agatston units, among those with CAC >0), geometric mean (geometric SD) | 51.2 (2.5) | 60.9 (2.5) | 49.2 (2.5) | 55.9 (2.4) |
| Common carotid IMT thickness (μm) | 619.8 (10.8) | 615.3 (10.6) | 626.2 (11.2) | 607.8 (10.1) |
| Internal carotid IMT thickness (μm) | 729.1 (18.7) | 744.3 (19.5) | 761 (19.5) | 713.3 (18) |

Data are means (SD) unless otherwise indicated. N = 1,193 subjects underwent measurements of vitamin D metabolites. FGF-23, fibroblast growth factor-23; PTH, parathyroid hormone. *P < 0.05 for a test comparing means, geometric means, or proportions, where appropriate, across 25(OH)D groups.

nmol/L (20–30 ng/mL), and < 50 nmol/L (20 ng/mL). Because no such categories exist for 24,25(OH)₂D and 1,25(OH)₂D, these metabolites were examined in quartiles. Adjusted coefficients and corresponding 95% CIs were generated for each category of vitamin D metabolite, and P values were generated evaluating each vitamin D metabolite as a continuous variable.

Models were adjusted for variables (measured at the time of plasma collection) that could confound associations of circulating vitamin D metabolites with study outcomes: age (continuous), sex, race (white vs. nonwhite), duration of diabetes (categories), DCCT treatment assignment, year of measurement (IMT only), solar irradiation of DCCT/EDIC site (continuous), BMI (continuous), AER

(continuous), and estimated GFR (continuous). P values < 0.05 were considered statistically significant. All analyses were performed using R, version 2.15 (R Foundation for Statistical Computing, Vienna, Austria).

RESULTS—Of the 1,193 subjects included in this study, 47% were women and 96% were white (Table 1). At the time of biomarker measurement (at or near the end of the DCCT), mean age was 32.4 years, mean duration of diabetes was 7.5 years, 20% of the subjects smoked, mean systolic blood pressure was 114.8 mmHg, and mean LDL cholesterol was 2.8 mmol/L. Lower plasma 25(OH)D concentration was associated with greater age, longer duration of diabetes, female sex, nonwhite race, and higher BMI (Table 1). Similar associations were observed for 24,25(OH)₂D₃ (Supplementary Table 1), which was highly correlated with 25(OH)D (Kendall $\tau = 0.7$). Lower plasma 1,25(OH)₂D was associated with longer duration of diabetes and white race (Supplementary Table 2).

Coronary artery calcium

Both plasma 25(OH)D concentration and CAC were measured in 1,081 subjects; 329 subjects had nonzero CAC scores (30%), and among those subjects the mean and median Agatston scores were 215.5 and 48.8, respectively, reflecting the highly skewed distribution of CAC. The prevalence of CAC tended to be lower—not higher—with lower concentrations of each vitamin D metabolite, though none of these trends were statistically significant (Supplementary Table 3). When CAC in multiple categories was examined as the outcome, lower circulating levels of all three vitamin D metabolites were associated with lower mean CAC (Table 2). Similar associations were observed using Tobit regression (Supplementary Table 4). There was a suggestion of a U-shaped relationship between plasma 1,25(OH)₂D and CAC (Fig. 1). In a separate post hoc model not reported in the tables, we used the middle two quartiles as the reference category in a polytomous regression model. Compared with the middle two categories, the lowest quartile was associated with 1.2 times the odds of higher mean CAC (95% CI 0.8–1.7) and the highest quartile was associated with 1.7 times the odds (1.2–1.7). Associations of the vitamin D metabolites with CAC did not differ by sex or DCCT treatment assignment (Supplementary Table 5).

Table 2—Associations of circulating vitamin D metabolites measured at the end of the DCCT with CAC measured during the EDIC study among 1,081 DCCT/EDIC participants

| Vitamin D metabolite | N | Mean CAC (SD) | Model 1** | Model 2*** |
|-------------------------------|-----|----------------|------------------|------------------|
| 25(OH)D | | | | |
| ≥75 nmol/L | 285 | 92.34 (340.84) | 1 (ref.) | 1 (ref.) |
| 50–75 nmol/L | 503 | 63.81 (253.64) | 0.89 (0.63–1.25) | 0.86 (0.61–1.21) |
| <50 nmol/L | 293 | 42.56 (138.74) | 0.79 (0.53–1.18) | 0.7 (0.47–1.05) |
| P | | | 0.51 | 0.223 |
| Per 25 nmol/L lower* | | | 0.86 (0.72–1.01) | 0.8 (0.68–0.96) |
| P* | | | 0.071 | 0.013 |
| 24,25(OH)₂D | | | | |
| ≥12.7 nmol/L | 264 | 86.67 (331.34) | 1 (ref.) | 1 (ref.) |
| 9.4–12.7 nmol/L | 257 | 72.69 (272.61) | 0.82 (0.55–1.22) | 0.79 (0.53–1.17) |
| 6.2–9.4 nmol/L | 283 | 57.39 (229.81) | 0.63 (0.42–0.94) | 0.61 (0.41–0.91) |
| <6.2 nmol/L | 267 | 48.6 (175.74) | 0.66 (0.44–0.99) | 0.59 (0.39–0.89) |
| P | | | 0.093 | 0.037 |
| Per 2.5 nmol/L lower* | | | 0.93 (0.87–1) | 0.91 (0.85–0.98) |
| P* | | | 0.055 | 0.015 |
| 1,25(OH)₂D | | | | |
| ≥122 pmol/L | 247 | 79.83 (261.47) | 1 (ref.) | 1 (ref.) |
| 101–122 pmol/L | 259 | 46.01 (225.62) | 0.59 (0.4–0.89) | 0.6 (0.4–0.9) |
| 88–101 pmol/L | 244 | 46.72 (203.4) | 0.56 (0.36–0.85) | 0.56 (0.37–0.86) |
| <88 pmol/L | 286 | 79.71 (280.2) | 0.72 (0.49–1.08) | 0.69 (0.46–1.03) |
| P | | | 0.024 | 0.029 |
| Per 25 pmol/L lower* | | | 0.9 (0.79–1.01) | 0.88 (0.78–1) |
| P* | | | 0.084 | 0.055 |

Data are adjusted odds ratio (95% CI) unless otherwise indicated. Estimates are odds ratios from polytomous logistic regression models in the CAC categories 0, 0–100, 100–300, and >300. Odds ratio estimates are interpreted as the odds of an increase to the next highest category associated with a unit difference in the exposure. Model 0 is adjusted only for scanner type. **Adjusted for scanner type, age, and duration of diabetes at the time of biomarker measurement as well as sex, race, and DCCT treatment assignment. ***Additionally adjusted for BMI, AER, and estimated GFR at the time of biomarker measurement as well as solar irradiation of DCCT clinic site. *Evaluates each vitamin D metabolite as a continuous variable, scaled as indicated. P value for trend created using this model.

IMT

Both plasma 25(OH) concentrations and carotid IMT were measured among 1,086 participants in 1994 and 991 participants in 2000/2001. The distribution of common IMT was unimodal and approximately symmetric, while the distribution of internal IMT was skewed to the right. The mean (SD) common carotid IMT in 1994 was 581 (86) μm and in 2000/2001 was 620 (117) μm. The mean internal carotid IMT in 1994 was 642 (221) μm and in 2000/2001 was 729 (349) μm. There was a suggestion that the lowest concentrations of 25(OH)D and 1,25(OH)₂D were associated with modestly greater common carotid IMT, but none of the three vitamin D metabolites were significantly associated with either common or internal mean IMT (Table 3). No plasma vitamin D metabolite was associated with mean IMT measured in 1994 only (evaluated using linear regression) or with change in IMT from 1994 to 2000/2001 (evaluated using generalized estimating

equations including time-by-vitamin D metabolite interactions). Associations of the vitamin D metabolites with IMT did not differ by sex or DCCT treatment assignment (Supplementary Table 5).

CONCLUSIONS

—In this study of 1,193 people with type 1 diabetes, we did not observe the hypothesized associations of lower plasma vitamin D metabolite concentrations with increased risks of subclinical cardiovascular disease, as measured by CAC or carotid IMT. To our surprise, we observed trends toward associations of lower concentrations of vitamin D metabolites with reduced CAC prevalence and severity, while risk estimates for IMT were close to null. Our findings contrast with those of previous studies that have shown associations of low serum concentrations of 25(OH)D and 1,25(OH)₂D with increased risk of CAC and cardiovascular mortality.

A number of prior studies have reported associations of low circulating

25(OH)D or 1,25(OH)₂D concentrations with increased risk of coronary heart disease. Among 374 participants in the Coronary Artery Calcification in Type 1 Diabetes (CACTI) study, circulating 25(OH)D <50 nmol/L was associated with a 3.3-fold increased odds of prevalent CAC compared with 25(OH)D ≥75 nmol/L in minimally adjusted analyses (15). Among 200 Steno Clinic patients with type 2 diabetes, plasma 25(OH)D <12.5 nmol/L (N = 19) was associated with increased risk of Agatston coronary calcium score ≥400 compared with 25(OH)D ≥12.5 nmol/L (16). Among 1,370 mostly nondiabetic participants in MESA, each 25 nmol/L decrement in 25(OH)D was associated with a 23% increased risk of developing CAC over 3 years' median follow-up (10). Lower 25(OH)D concentration has also been associated with increased risks of cardiovascular disease events in the general population (30,31) and with increased mortality risk in types 1 and 2 diabetes (12,14), each presumably reflecting in part an increased tendency to develop coronary heart disease. Lower circulating 1,25(OH)₂D concentrations were associated with increased risk of prevalent CAC among a population referred for evaluation of hyperlipidemia and with a trend toward increased risk of incident CAC in MESA (10,17).

Differences in our results compared with prior studies may reflect differences in study population, study design, or covariate adjustment. Regarding study population, associations of vitamin D metabolite concentrations with atherosclerosis could conceivably vary by age, sex, or diabetes status, which differ substantially in the studies noted above. Our study is most similar to the CACTI study, since both evaluated participants with type 1 diabetes, evaluated 25(OH)D using similar assays and categories, and measured CAC at similar ages. However, compared with CACTI, mean plasma 25(OH)D concentration (63.2 vs. 88.1 nmol/L) and the prevalence of CAC (30 vs. 45.5%) were each lower in our study. There were also differences in study design and covariate adjustment. Specifically, we measured CAC a median of 10 years after circulating vitamin D metabolites and thus tested associations of vitamin D metabolites with a combination of prevalent and incident CAC in a staggered cross-sectional analysis. The timing of the vitamin D and CAC measurements is temporally appropriate and clinically

Table 3—Associations of circulating vitamin D metabolites measured at the end of the DCCT with internal carotid IMT measured during the EDIC study among 991 DCCT/EDIC participants

| Vitamin D metabolite | N | Common carotid IMT | | | Internal carotid IMT | | |
|-------------------------------|-----|-------------------------------|-----------------------------------|----------------------|---|--|------------------|
| | | Unadjusted mean (SD) IMT (μm) | Adjusted mean difference (95% CI) | | Unadjusted geometric mean (geometric SD) IMT (μm) | Adjusted geometric mean ratio (95% CI) | |
| | | | Model 1** | Model 2*** | | Model 1** | Model 2*** |
| 25(OH)D | | | | | | | |
| ≥75 nmol/L | 550 | 591 (99) | 0 (ref.) | 0 (ref.) | 648 (1.37) | 1 (ref.) | 1 (ref.) |
| 50–75 nmol/L | 954 | 600 (103) | 4.7 (−6.4 to 15.8) | 2.2 (−8.8 to 13.1) | 655 (1.35) | 1.01 (0.97–1.04) | 1 (0.97–1.04) |
| <50 nmol/L | 573 | 608 (110) | 12.4 (0–24.9) | 7.9 (−4.8 to 20.6) | 640 (1.32) | 0.98 (0.94–1.02) | 0.97 (0.93–1.01) |
| P | | | 0.142 | 0.447 | | 0.405 | 0.227 |
| Per 25 nmol/L lower* | | | 4.7 (−0.7 to 10.1) | 2.4 (−3.1 to 8) | | 1 (0.98–1.01) | 0.99 (0.97–1.01) |
| P* | | | 0.087 | 0.387 | | 0.72 | 0.281 |
| 24,25(OH)₂D | | | | | | | |
| ≥12.7 nmol/L | 507 | 596 (104) | 0 (ref.) | 0 (ref.) | 653 (1.37) | 1 (ref.) | 1 (ref.) |
| 9.4–12.7 nmol/L | 490 | 605 (110) | 5.4 (−8.7 to 19.4) | 3.7 (−10 to 17.4) | 662 (1.38) | 1.01 (0.96–1.05) | 1 (0.96–1.04) |
| 6.2–9.4 nmol/L | 537 | 593 (93) | −4.3 (−16.5 to 8) | −6.9 (−19.2 to 5.3) | 644 (1.32) | 0.99 (0.95–1.03) | 0.98 (0.94–1.02) |
| <6.2 nmol/L | 527 | 603 (106) | 2 (−11.4 to 15.5) | −3.6 (−17 to 9.8) | 634 (1.31) | 0.96 (0.92–1) | 0.94 (0.9–0.99) |
| P | | | 0.52 | 0.42 | | 0.102 | 0.025 |
| Per 2.5 nmol/L lower* | | | 0.1 (−2.1 to 2.3) | −1 (−3.2 to 1.2) | | 0.99 (0.99–1) | 0.99 (0.98–1) |
| P* | | | 0.962 | 0.366 | | 0.088 | 0.015 |
| 1,25(OH)₂D | | | | | | | |
| ≥122 pmol/L | 488 | 586 (97) | 0 (ref.) | 0 (ref.) | 626 (1.34) | 1 (ref.) | 1 (ref.) |
| 101–122 pmol/L | 505 | 597 (102) | −1.5 (−14.8 to 11.9) | −2.7 (−15.8 to 10.3) | 654 (1.33) | 1.01 (0.97–1.05) | 1.01 (0.97–1.05) |
| 88–101 pmol/L | 463 | 600 (95) | 1.3 (−11.6 to 14.3) | 0.2 (−12.7 to 13.1) | 644 (1.27) | 1 (0.96–1.04) | 1 (0.96–1.04) |
| <88 pmol/L | 542 | 613 (118) | 11.4 (−2.2 to 24.9) | 8.3 (−5.3 to 21.9) | 666 (1.41) | 1.03 (0.98–1.07) | 1.01 (0.97–1.06) |
| P | | | 0.243 | 0.422 | | 0.606 | 0.848 |
| Per 25 pmol/L lower* | | | 3.5 (−0.3 to 7.2) | 2.5 (−1.2 to 6.3) | | 1 (0.99–1.02) | 1 (0.99–1.01) |
| P* | | | 0.073 | 0.189 | | 0.488 | 0.945 |

Estimates are mean differences and geometric mean ratios from linear regression models on common carotid IMT and the log-transformed internal carotid IMT, respectively. Models use both year 1 and year 6 measurements with inference performed using generalized estimating equations. **Adjusted for scanner type, age, and duration of diabetes at the time of biomarker measurement as well as sex, race, and DCCT treatment assignment. ***Additionally adjusted for BMI, AER, and estimated GFR at the time of biomarker measurement as well as solar irradiation of DCCT clinic site. *Evaluates each vitamin D metabolite as a continuous variable, scaled as indicated. P value for trend created using this model.

could still lead to misclassification and bias results toward the null. In conclusion, we did not find evidence linking impaired vitamin D metabolism with increased risk of subclinical CVD in type 1 diabetes.

Acknowledgments—This study was supported by grants RC4DK090766, R01DK087726, and R01KD088762 from the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). The DCCT/EDIC study is supported by contracts with the Division of Diabetes, Endocrinology, and Metabolic Diseases of the NIDDK; National Eye Institute; National Institute of Neurological Disorders and Stroke; and the General Clinical Research Centers Program and by funding from the Clinical and Translational Science Awards Program; National Center for Research Resources;

and Genentech through a Cooperative Research and Development Agreement with NIDDK. Free or discounted supplies or equipment were contributed by Abbott; Animas; Aventis; Bayer; Becton, Dickinson, and Company; CanAm; Eli Lilly; LifeScan; Medtronic; MiniMed; Omron; OmniPod; Roche; and Sanofi. No other potential conflicts of interest relevant to this article were reported.

M.C.S. analyzed the data and wrote the manuscript. J.D.B., J.M.L., and M.W.S. contributed to the discussion and edited the manuscript. P.A.C., M.E.M., and B.Z. edited the manuscript. A.N.H. measured the data and edited the manuscript. I.H.d.B. conceived the study, contributed to the discussion, and edited the manuscript. M.S. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

The authors thank W. Sun and J.-Y. Backlund of The George Washington University for their help preparing the data.

References

1. Costacou T, Edmundowicz D, Prince C, Conway B, Orchard TJ. Progression of coronary artery calcium in type 1 diabetes mellitus. *Am J Cardiol* 2007;100:1543–1547
2. Lewis EJ, Hunsicker LG, Bain RP, Rohde RD; The Collaborative Study Group. The effect of angiotensin-converting-enzyme inhibition on diabetic nephropathy. *N Engl J Med* 1993;329:1456–1462
3. Colhoun HM, Betteridge DJ, Durrington PN, et al.; CARDS investigators. Primary prevention of cardiovascular disease with atorvastatin in type 2 diabetes in the Collaborative Atorvastatin Diabetes

- Study (CARDS): multicentre randomised placebo-controlled trial. *Lancet* 2004;364:685–696
4. Nathan DM, Cleary PA, Backlund JYC, et al.; Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications (DCCT/EDIC) Study Research Group. Intensive diabetes treatment and cardiovascular disease in patients with type 1 diabetes. *N Engl J Med* 2005;353:2643–2653
 5. Andress DL. Vitamin D in chronic kidney disease: a systemic role for selective vitamin D receptor activation. *Kidney Int* 2006;69:33–43
 6. Li YC, Kong J, Wei M, Chen ZF, Liu SQ, Cao LP. 1,25-Dihydroxyvitamin D(3) is a negative endocrine regulator of the renin-angiotensin system. *J Clin Invest* 2002;110:229–238
 7. Takeda M, Yamashita T, Sasaki N, et al. Oral administration of an active form of vitamin D3 (calcitriol) decreases atherosclerosis in mice by inducing regulatory T cells and immature dendritic cells with tolerogenic functions. *Arterioscler Thromb Vasc Biol* 2010;30:2495–2503
 8. Szeto FL, Reardon CA, Yoon D, et al. Vitamin D receptor signaling inhibits atherosclerosis in mice. *Mol Endocrinol* 2012;26:1091–1101
 9. Targher G, Bertolini L, Padovani R, et al. Serum 25-hydroxyvitamin D3 concentrations and carotid artery intima-media thickness among type 2 diabetic patients. *Clin Endocrinol (Oxf)* 2006;65:593–597
 10. de Boer IH, Kestenbaum B, Shoben AB, Michos ED, Sarnak MJ, Siscovick DS. 25-hydroxyvitamin D levels inversely associate with risk for developing coronary artery calcification. *J Am Soc Nephrol* 2009;20:1805–1812
 11. Reis JP, von Mühlen D, Michos ED, et al. Serum vitamin D, parathyroid hormone levels, and carotid atherosclerosis. *Atherosclerosis* 2009;207:585–590
 12. Joergensen C, Gall MA, Schmedes A, Tarnow L, Parving HH, Rossing P. Vitamin D levels and mortality in type 2 diabetes. *Diabetes Care* 2010;33:2238–2243
 13. Choi AI, Lo JC, Mulligan K, et al. Association of vitamin D insufficiency with carotid intima-media thickness in HIV-infected persons. *Clin Infect Dis* 2011;52:941–944
 14. Joergensen C, Hovind P, Schmedes A, Parving HH, Rossing P. Vitamin D levels, microvascular complications, and mortality in type 1 diabetes. *Diabetes Care* 2011;34:1081–1085
 15. Young KA, Snell-Bergeon JK, Naik RG, et al. Vitamin D deficiency and coronary artery calcification in subjects with type 1 diabetes. *Diabetes Care* 2011;34:454–458
 16. Joergensen C, Reinhard H, Schmedes A, et al. Vitamin D levels and asymptomatic coronary artery disease in type 2 diabetic patients with elevated urinary albumin excretion rate. *Diabetes Care* 2012;35:168–172
 17. Watson KE, Abrolat ML, Malone LL, et al. Active serum vitamin D levels are inversely correlated with coronary calcification. *Circulation* 1997;96:1755–1760
 18. Shen H, Bielak LF, Ferguson JF, et al. Association of the vitamin D metabolism gene CYP24A1 with coronary artery calcification. *Arterioscler Thromb Vasc Biol* 2010;30:2648–2654
 19. de Boer IH, Sachs MC, Cleary PA, et al.; Diabetes Control and Complication Trial/Epidemiology of Diabetes Interventions and Complications Study Research Group. Circulating vitamin D metabolites and kidney disease in type 1 diabetes. *J Clin Endocrinol Metab* 2012;97:4780–4788 DOI: 10.1210/jc.2012-2852
 20. Bosworth CR, Levin G, Robinson-Cohen C, et al. The serum 24,25-dihydroxyvitamin D concentration, a marker of vitamin D catabolism, is reduced in chronic kidney disease. *Kidney Int* 2012;82:693–700 DOI: 10.1038/ki.2012.193
 21. Strathmann FG, Laha TJ, Hoofnagle AN. Quantification of 1 α ,25-dihydroxy vitamin D by immunoextraction and liquid chromatography-tandem mass spectrometry. *Clin Chem* 2011;57:1279–1285
 22. Hoofnagle AN, Laha TJ, Donaldson TF. A rubber transfer gasket to improve the throughput of liquid-liquid extraction in 96-well plates: application to vitamin D testing. *J Chromatogr B Analyt Technol Biomed Life Sci* 2010;878:1639–1642
 23. Agatston AS, Janowitz WR, Hildner FJ, Zusmer NR, Viamonte M Jr, Detrano R. Quantification of coronary artery calcium using ultrafast computed tomography. *J Am Coll Cardiol* 1990;15:827–832
 24. Nathan DM, Lachin J, Cleary P, et al.; Diabetes Control and Complications Trial; Epidemiology of Diabetes Interventions and Complications Research Group. Intensive diabetes therapy and carotid intima-media thickness in type 1 diabetes mellitus. *N Engl J Med* 2003;348:2294–2303
 25. Steffes M, Cleary P, Goldstein D, et al. Hemoglobin A1c measurements over nearly two decades: sustaining comparable values throughout the Diabetes Control and Complications Trial and the Epidemiology of Diabetes Interventions and Complications study. *Clin Chem* 2005;51:753–758
 26. Levey AS, Stevens LA, Schmid CH, et al. A new equation to estimate glomerular filtration rate. *Ann Intern Med* 2009;150:604–612
 27. Detrano R, Guerci AD, Carr JJ, et al. Coronary calcium as a predictor of coronary events in four racial or ethnic groups. *N Engl J Med* 2008;358:1336–1345
 28. Ananth CV, Kleinbaum DG. Regression models for ordinal responses: a review of methods and applications. *Int J Epidemiol* 1997;26:1323–1333
 29. Tobin J. Estimation of relationships for limited dependent variables. *Econometrica* 1958;26:24–36
 30. Liang KY, Zeger SL. Longitudinal data analysis using generalized linear models. *Biometrika* 1986;73:13–22
 31. Wang TJ, Pencina MJ, Booth SL, et al. Vitamin D deficiency and risk of cardiovascular disease. *Circulation* 2008;117:503–511
 32. Melamed ML, Michos ED, Post W, Astor B. 25-hydroxyvitamin D levels and the risk of mortality in the general population. *Arch Intern Med* 2008;168:1629–1637
 33. Mathew S, Lund RJ, Chaudhary LR, Geurs T, Hruska KA. Vitamin D receptor activators can protect against vascular calcification. *J Am Soc Nephrol* 2008;19:1509–1519
 34. Giovannucci E, Liu Y, Hollis BW, Rimm EB. 25-hydroxyvitamin D and risk of myocardial infarction in men: a prospective study. *Arch Intern Med* 2008;168:1174–1180
 35. Lim S, Shin H, Kim MJ, et al. Vitamin D inadequacy is associated with significant coronary artery stenosis in a community-based elderly cohort: the Korean Longitudinal Study on Health and Aging. *J Clin Endocrinol Metab* 2012;97:169–178
 36. Michos ED, Streeten EA, Ryan KA, et al. Serum 25-hydroxyvitamin d levels are not associated with subclinical vascular disease or C-reactive protein in the old order amish. *Calcif Tissue Int* 2009;84:195–202