

F. Hoffmann-La Roche Ltd./Genentech Inc., Celgene Corporation, Gilead Sciences, Pfizer Inc., Amgen Inc., Janssen-Cilag Ltd., and Lundbeck Ltd., and research funding from F. Hoffmann-La Roche Ltd. M.D. and R.H. are employees of Roche Products Ltd., and G.F.-R. and E.W.-F. are employees of F. Hoffmann-La Roche Ltd. M.D. and G.F.-R. have stock ownership in Roche, which is publicly traded. J.G.G. has received honoraria from Celgene Corporation, Gilead Sciences, F. Hoffmann-La Roche Ltd., Janssen-Cilag Ltd., and Pharmacyclics LLC, and other remuneration (travel, accommodation, and expenses) from Gilead Sciences. M.H. has received consultancy fees and research funding from, and has served on a Speaker's Bureau for, F. Hoffmann-La Roche Ltd. G.S. has received consultancy fees from F. Hoffmann-La Roche Ltd./Genentech Inc., Celgene Corporation, Gilead Sciences, Janssen-Cilag Ltd., and Mundipharma; honoraria from F. Hoffmann-La Roche Ltd./Genentech Inc., Amgen Inc., Mundipharma, and Sanofi; and research funding and other remuneration (travel and expenses) from F. Hoffmann-La Roche Ltd. G.C. has received consultancy fees from F. Hoffmann-La Roche Ltd. and honoraria from F. Hoffmann-La Roche Ltd., GlaxoSmithKline, Gilead Sciences, Janssen-Cilag Ltd., and Sanofi. T.L. declares no competing financial interests.

Correspondence: Ciara Freeman, Barts Cancer Institute, 3rd Floor John Vane Science Centre, Charterhouse Square, London, United Kingdom; e-mail: c.freeman@qmul.ac.uk.

References

- Hallek M. Chronic lymphocytic leukemia: 2015 update on diagnosis, risk stratification, and treatment. *Am J Hematol*. 2015;90(5):446-460.
- Byrd JC, Murphy T, Howard RS, et al. Rituximab using a thrice weekly dosing schedule in B-cell chronic lymphocytic leukemia and small lymphocytic lymphoma demonstrates clinical activity and acceptable toxicity. *J Clin Oncol*. 2001;19(8):2153-2164.
- Byrd JC, Waselenko JK, Maneatis TJ, et al. Rituximab therapy in hematologic malignancy patients with circulating blood tumor cells: association with increased infusion-related side effects and rapid blood tumor clearance. *J Clin Oncol*. 1999;17(3):791-795.
- Winkler U, Jensen M, Manzke O, Schulz H, Diehl V, Engert A. Cytokine-release syndrome in patients with B-cell chronic lymphocytic leukemia and high lymphocyte counts after treatment with an anti-CD20 monoclonal antibody (rituximab, IDEC-C2B8). *Blood*. 1999;94(7):2217-2224.
- Golay J, Da Roit F, Bologna L, et al. Glycoengineered CD20 antibody obinutuzumab activates neutrophils and mediates phagocytosis through CD16B more efficiently than rituximab. *Blood*. 2013;122(20):3482-3491.
- Mössner E, Brünker P, Moser S, et al. Increasing the efficacy of CD20 antibody therapy through the engineering of a new type II anti-CD20 antibody with enhanced direct and immune effector cell-mediated B-cell cytotoxicity. *Blood*. 2010;115(22):4393-4402.
- Patz M, Isaeva P, Forcob N, et al. Comparison of the in vitro effects of the anti-CD20 antibodies rituximab and GA101 on chronic lymphocytic leukaemia cells. *Br J Haematol*. 2011;152(3):295-306.
- Goede V, Fischer K, Busch R, et al. Obinutuzumab plus chlorambucil in patients with CLL and coexisting conditions. *N Engl J Med*. 2014;370(12):1101-1110.
- Goede V, Fischer K, Engelke A, et al. Obinutuzumab as frontline treatment of chronic lymphocytic leukemia: updated results of the CLL11 study. *Leukemia*. 2015;29(7):1602-1604.
- Sehn LH, Assouline SE, Stewart DA, et al. A phase 1 study of obinutuzumab induction followed by 2 years of maintenance in patients with relapsed CD20-positive B-cell malignancies. *Blood*. 2012;119(22):5118-5125.
- Cartron G, de Guibert S, Dillhuyd MS, et al. Obinutuzumab (GA101) in relapsed/refractory chronic lymphocytic leukemia: final data from the phase 1/2 GAUGUIN study. *Blood*. 2014;124(14):2196-2202.
- Nyhlen K, Gautam C, Andersson R, Srinivas U. Modulation of cytokine-induced production of IL-8 in vitro by interferons and glucocorticosteroids. *Inflammation*. 2004;28(2):77-88.
- Kara IO, Sahin B, Gunesacar R. Expression of soluble CD27 and interleukins-8 and -10 in B-cell chronic lymphocytic leukemia: correlation with disease stage and prognosis. *Adv Ther*. 2007;24(1):29-40.
- Walz A, Meloni F, Clark-Lewis I, von Tscherner V, Baggiolini M. [Ca²⁺]_i changes and respiratory burst in human neutrophils and monocytes induced by NAP-1/interleukin-8, NAP-2, and gro/MGSA. *J Leukoc Biol*. 1991;50(3):279-286.
- Laprevotte E, Ysebaert L, Klein C, et al. Endogenous IL-8 acts as a CD16 co-activator for natural killer-mediated anti-CD20 B cell depletion in chronic lymphocytic leukemia. *Leuk Res*. 2013;37(4):440-446.
- Krüttgen A, Rose-John S. Interleukin-6 in sepsis and capillary leakage syndrome. *J Interferon Cytokine Res*. 2012;32(2):60-65.
- Gordon MS, Nemunaitis J, Hoffman R, et al. A phase I trial of recombinant human interleukin-6 in patients with myelodysplastic syndromes and thrombocytopenia. *Blood*. 1995;85(11):3066-3076.
- Trikha M, Corringham R, Klein B, Rossi JF. Targeted anti-interleukin-6 monoclonal antibody therapy for cancer: a review of the rationale and clinical evidence. *Clin Cancer Res*. 2003;9(13):4653-4665.
- Liu F, Jia L, Wang P, Farren T, Agrawal S. Tocilizumab overcomes chemo-resistance of CLL cells [abstract]. *Blood*. 2013;122(21). Abstract 5305.
- Giezen TJ, Mantel-Teeuwisse AK, ten Berg MJ, et al. Rituximab-induced thrombocytopenia: a cohort study. *Eur J Haematol*. 2012;89(3):256-266.
- Freeman CL, Dixon M, Houghton R, et al. Risk factors associated with the development of infusion-related reactions in patients with chronic lymphocytic leukaemia treated with anti-CD20 monoclonal antibodies: analysis of the CLL11 study dataset [abstract]. *Blood*. 2014;124(21). Abstract 3339.
- Xia P, Gamble JR, Rye K-A, et al. Tumor necrosis factor-alpha induces adhesion molecule expression through the sphingosine kinase pathway. *Proc Natl Acad Sci USA*. 1998;95(24):14196-14201.
- Mihara M, Hashizume M, Yoshida H, Suzuki M, Shiina M. IL-6/IL-6 receptor system and its role in physiological and pathological conditions. *Clin Sci (Lond)*. 2012;122(4):143-159.
- Ram R, Bonstein L, Gafter-Gvili A, Ben-Bassat I, Shpilberg O, Raanani P. Rituximab-associated acute thrombocytopenia: an under-diagnosed phenomenon. *Am J Hematol*. 2009;84(4):247-250.

DOI 10.1182/blood-2015-09-670802

© 2015 by The American Society of Hematology

To the editor:

Dramatic transient improvement of metastatic BRAF^{V600E}-mutated Langerhans cell sarcoma under treatment with dabrafenib

Langerhans cell sarcoma (LCS) is a rare histiocytic neoplasm with overt malignant cytological features and an aggressive clinical course.¹ Disseminated LCS carries a poor prognosis.¹ We report a case of a metastatic BRAF^{V600E}-mutated LCS that dramatically improved after administration of the BRAF inhibitor (BRAFi) dabrafenib.

A 58-year-old man was referred in August 2014 with a diagnosis of progressive Langerhans cell histiocytosis (LCH). He was treated in July 2013 by surgery and radiotherapy for left humerus LCH diagnosed by bone biopsy. In February 2014, enlargement of the left axillary, pectoral, and supraclavicular lymph nodes was observed. Histologic examination of a lymph node biopsy indicated LCH

recurrence, although some atypical cells were described. CHOP (cyclophosphamide, doxorubicin, vincristine, and prednisone) chemotherapy was initiated but was discontinued after 3 cycles because of disease progression. The patient's condition deteriorated. The lymph nodes increased in size and multiple lung nodules were observed on lung computed tomography (CT) scan. These lesions were highly hypermetabolic on fluorodeoxyglucose (¹⁸F-FDG) positron emission tomography (PET)-CT (Figure 1). A new lymph node biopsy revealed massive infiltration by very large cells with irregular folded nuclei and necrotic areas. The mitotic rate was >50 per 10 high-power fields. The tumor cells expressed CD1a and langerin




Figure 1. BRAFV600E genotyping. (A) 18 F-FDG PET-CT before the onset of treatment. (a) Fused PET-CT axial view of pulmonary nodules (standardized maximal uptake [SUVmax] = 12). (b) Left axillary mass (SUVmax = 11.9). 18 F-FDG PET-CT after 3 months of dabrafenib treatment. (c) Fused PET-CT showing no significant residual uptake in the lung. (d) Presence of a residual small axillary lymph node with decreased uptake (SUVmax = 4.5). The patient was classified as partial metabolic response according to the EORTC criteria (ie, >25% reduction of the SUVmax in the target lesions), with a decrease in SUVmax of 83% in the target lesions. 18 F-FDG PET-CT after 6 months of dabrafenib treatment at the time of LCS relapse. (e) Fused PET-CT showing a hypermetabolic pulmonary nodule (SUVmax = 6.3) and mediastinal lymph nodes (SUVmax = 8.7); (f) axillary nodal relapse (SUVmax = 9.6). (B) Light optic microscopy of the left axillary lymph node biopsy before and during treatment. (a) Hematoxylin-and-eosin (HE) staining showing a very pleomorphic proliferation composed of large tumor cells with folded nuclei, and numerous mitotic cells. (b) Tumor cells stained with an anti-CD1a antibody (clone O10; Dako); (c) most of the cells are langerin (CD207)-positive (clone 12D6; Novocastra). These features are characteristic of LCS. (d) Anti-BRAF^{V600E} (clone VE1; Spring Biosciences) immunostaining is positive in tumor cells. (e) HE staining of a lymph node biopsy at day 10 of dabrafenib treatment, showing massive necrosis of tumor cells. (f) No cells stained positive with the anti-BRAF^{V600E} mutation-specific antibody. Original magnification, $\times 400$ in all sections. (C) Pyrosequencing of DNA extracted from lymph node biopsies. (a) Before treatment, the BRAF^{V600E} mutation is present (mutated allele = 78%); (b) on day 10 of dabrafenib treatment, only wild-type BRAF^{V600E} is observed (mutated allele = 1.4%); and (c) at the time of LCS recurrence, demonstrating the resurgence of the BRAF^{V600E} mutation (mutated allele = 52%). (D) Expression of MAP3K8/COT protein in the pretreatment and relapsing tumor. HT-29 BRAF^{V600E} colorectal cells that express both the long (1-467) and short (30-467) forms of COT are shown as a positive control. β -actin served as loading control. Extracellular signal-regulated kinase (ERK) expression and activation in relapsed LCS. (E) Tumor cells are intensively stained with an anti-phospho-ERK-1/2 antibody (clone MAPK-YT; Sigma) compared with pretreatment tumor cells. Original magnification, $\times 400$. (F) Western blot analysis for phosphorylated ERK (pERK) and ERK (endogenous total ERK) in the pretreatment and relapsing tumors is shown. A375 BRAF^{V600E} melanoma cells were used as a positive control.

(Figure 1), features characteristic of LCS.^{1,2} A review of previous biopsies showed the presence of small areas of similar tumor cells. All tissue specimens harbored the BRAF^{V600E} mutation, as determined by immunohistochemistry and molecular genotyping (Figure 1). After informed consent, dabrafenib (150 mg twice daily) was initiated. Within a week, the patient improved, and the lymph nodes dramatically decreased in size. A lymph node biopsy performed on treatment day 10 showed massive necrosis of tumor cells and the absence of the BRAF^{V600E} mutation (Figure 1). Serial 18 F-FDG PET-CT scans after 3 months of treatment revealed marked improvement of the lesions, with residual small axillary adenopathy (Figure 1). The patient was in complete response according

to Response Evaluation Criteria In Solid Tumors (RECIST) 1.1 criteria and in partial metabolic response according to European Organisation for Research and Treatment of Cancer (EORTC) criteria. The treatment was well tolerated except for diarrhea (which resolved after the dabrafenib dose was reduced to 100 mg twice daily) and palmoplantar hyperkeratosis. Average values of dabrafenib plasma trough and maximal concentrations, which were monitored monthly, reached 16 ng/mL (15-27 ng/mL) and 604 ng/mL (507-786 ng/mL), respectively, consistent with previous pharmacokinetic studies.³

At 6 months, 18 F-FDG PET-CT revealed an increase in size of the left axillary lymph node, the enlargement of mediastinal lymph nodes, and a left pulmonary nodule (Figure 1). The axillary lymph

node biopsy confirmed the recurrence of BRAF^{V600E}-mutated LCS (Figure 1). To determine the mechanisms by which dabrafenib resistance was acquired, genomic and transcriptomic analyses of the main previously described factors involved in BRAFi resistance were performed on the pretreatment and relapsing tumors, which contained 95% and 90% tumor cells, respectively⁴⁻⁶ (supplemental Methods and supplemental Table 1, see supplemental Data available on the *Blood* Web site). No NRAS^{G12/G13/Q61}, KRAS^{G12/G13}, or MAP2K1 (exons 2 and 3) mutations or BRAF splicing variants were detected in the recurrent lesion.⁴⁻⁶ The MAP3K8/COT messenger RNA level, normalized to the number of tumor cells, was threefold (3.3 ± 0.2) increased in the relapsing specimen as compared with the initial tumor. This increase, which was also observed at the protein level, may explain the observed intense activation of the mitogen-activated protein kinase (MAPK) pathway (Figure 1). Indeed, in melanoma, MAP3K8/COT overexpression drives resistance to BRAFi through MAPK pathway reactivation which may be overcome by the addition of a MAPK kinase inhibitor (MEKi).⁴ The patient was thereafter switched to combined vemurafenib and cobimetinib treatment. Within 3 days, the patient clinically improved. At 10 weeks of treatment, he had an Eastern Cooperative Oncology Group (ECOG) performance status of 1 and the different tumor localizations had decreased in size.

LCS is a rare malignant histiocytic disorder that affects the lymph nodes as well as extranodal sites.¹ LCS is differentiated from LCH based on cytological criteria, a high mitotic index, and more aggressive behavior.^{1,2} Various chemotherapy regimens, primarily CHOP, are used with limited response rates and high mortality.¹ Recently, the BRAF^{V600E} mutation was identified in a variable proportion of dendritic cell neoplasms, including LCS, suggesting that these patients may benefit from BRAFi.^{1,7,8} The dramatic response of the present case to dabrafenib further supports the key role of the BRAF^{V600E} mutation in histiocytic disorders,^{9,10} although secondary resistance to BRAFi treatment may occur in malignant forms.⁸

In melanoma, a first-line therapeutic strategy with the combination of BRAFi and MEKi has been demonstrated to prevent or delay the onset of resistance observed with treatment with BRAFi alone, in addition to increasing progression-free survival.^{11,12} A similar strategy might be more appropriate in progressive BRAF^{V600E}-mutated histiocytic malignancies.

Samia Mourah

Laboratoire de Pharmacologie Biologique, Hôpital Saint-Louis,
Assistance Publique-Hôpitaux de Paris,
Paris, France
Université Paris Diderot, Sorbonne Paris Cité,
Paris, France
INSERM U976,
Paris, France

Gwenaël Lorillon

Centre National de Référence de l'Histiocytose Langerhansienne,
Service de Pneumologie,
Hôpital Saint-Louis, Assistance Publique-Hôpitaux de Paris,
Paris, France

Véronique Meignin

Service de Pathologie, Hôpital Saint-Louis,
Assistance Publique-Hôpitaux de Paris,
INSERM UMR_S1165,
Paris, France

Laetitia Vercellino

Service de Médecine Nucléaire, Hôpital Saint-Louis,
Assistance Publique-Hôpitaux de Paris,
Paris, France

Constance de Margerie-Mellon

Service de Radiologie, Hôpital Saint-Louis,
Assistance Publique-Hôpitaux de Paris,
Paris, France

Cécile Pages

Département de Dermatologie, Hôpital Saint-Louis,
Assistance Publique-Hôpitaux de Paris,
Paris, France

Lauriane Goldwirt

Laboratoire de Pharmacologie Biologique, Hôpital Saint-Louis,
Assistance Publique-Hôpitaux de Paris,
Paris, France

Alexandre How-Kit

Laboratoire de Génomique Fonctionnelle,
Fondation Jean Dausset-Centre d'Etude du Polymorphisme Humain,
Paris, France

Jorg Tost

Laboratoire d'Épigénétique, Centre National de Génotypage,
Commissariat à l'énergie atomique et aux énergies
alternatives-Institut de Génomique,
Evry, France

Céleste Lebbe

Département de Dermatologie, Hôpital Saint-Louis,
Assistance Publique-Hôpitaux de Paris,
Paris, France
Université Paris Diderot, Sorbonne Paris Cité,
Paris, France
INSERM U976,
Paris, France

Abdellatif Tazi

Centre National de Référence de l'Histiocytose Langerhansienne,
Service de Pneumologie,
Hôpital Saint-Louis, Assistance Publique-Hôpitaux de Paris,
Paris, France
Université Paris Diderot, Sorbonne Paris Cité,
Paris, France
Equipe de Recherche en Biostatistiques et Epidémiologie Clinique,
INSERM U1153 Centre de Recherche INSERM Epidémiologie et Statistique
Paris Sorbonne Cité (CRESS),
Paris, France

The online version of this article contains a data supplement.

Acknowledgments: The authors thank Aurélie Sadoux, Coralie Reger de Moura (Laboratoire de Pharmacologie Biologique, Hôpital Saint-Louis, Paris, France), and Maeva Valluci (Centre National de Référence de l'Histiocytose Langerhansienne, Service de Pneumologie, Hôpital Saint-Louis, Paris, France) for technical support, and Elisabeth Savariau (Institut Universitaire d'Hématologie, Service d'Infographie, Hôpital Saint-Louis, Paris, France) for assistance with the figures.

Contribution: S.M. and A.T. designed the research, analyzed and interpreted the data, and wrote the manuscript; G.L. managed sample acquisition and collected clinical data; V.M. analyzed and interpreted histologic data; L.V. analyzed and interpreted ¹⁸F-FDG PET-CT findings; C.d.M.-M. analyzed and interpreted radiologic results; C.P. analyzed and interpreted dermatologic clinical findings; L.G. analyzed and interpreted pharmacologic data; A.H.-K. and J.T. analyzed BRAF, NRAS, and KRAS genotyping and interpreted the data; and C.L. provided expertise in the melanoma field and intellectual content.

Conflict-of-interest disclosure: S.M. declares a consulting role for Roche and Novartis. C.d.M.-M. declares travel accommodation by Guerbet. L.G. declares travel accommodation by Janssen. C.L. declares honoraria from Roche, advisory roles for Roche, GlaxoSmithKline (GSK), Novartis, Bristol-Myers Squibb (BMS), Merck Sharp & Dohme (MSD), and Amgen, and travel accommodation by Roche. The remaining authors declare no competing financial interests.

Correspondence: Abdellatif Tazi, Service de Pneumologie, Hôpital Saint-Louis, 1 Avenue Claude Vellefaux, 75475, Paris cedex 10, France; e-mail: abdellatif.tazi@sls.aphp.fr.

References

- Zwerdling T, Won E, Shane L, Javahara R, Jaffe R. Langerhans cell sarcoma: case report and review of world literature. *J Pediatr Hematol Oncol*. 2014;36(6):419-425.
- Swerdlow SH, Campo E, Harris NL, et al. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues. Lyon, France: IARC; 2008.
- Falchook GS, Long GV, Kurzrock R, et al. Dose selection, pharmacokinetics, and pharmacodynamics of BRAF inhibitor dabrafenib (GSK2118436). *Clin Cancer Res*. 2014;20(17):4449-4458.
- Johannessen CM, Boehm JS, Kim SY, et al. COT drives resistance to RAF inhibition through MAP kinase pathway reactivation. *Nature*. 2010;468(7326):968-972.
- Van Allen EM, Wagle N, Sucker A, et al; Dermatologic Cooperative Oncology Group of Germany (DeCOG). The genetic landscape of clinical resistance to RAF inhibition in metastatic melanoma. *Cancer Discov*. 2014;4(1):94-109.
- Wagle N, Van Allen EM, Treacy DJ, et al. MAP kinase pathway alterations in BRAF-mutant melanoma patients with acquired resistance to combined RAF/MEK inhibition. *Cancer Discov*. 2014;4(1):61-68.
- Go H, Jeon YK, Huh J, et al. Frequent detection of BRAF(V600E) mutations in histiocytic and dendritic cell neoplasms. *Histopathology*. 2014;65(2):261-272.
- Idbaih A, Mokhtari K, Emile JF, et al. Dramatic response of a BRAF V600E-mutated primary CNS histiocytic sarcoma to vemurafenib. *Neurology*. 2014;83(16):1478-1480.
- Haroche J, Cohen-Aubart F, Emile JF, et al. Reproducible and sustained efficacy of targeted therapy with vemurafenib in patients with BRAF(V600E)-mutated Erdheim-Chester disease. *J Clin Oncol*. 2015;33(5):411-418.
- Hyman DM, Puzanov I, Subbiah V, et al. Vemurafenib in multiple nonmelanoma cancers with BRAF V600 mutations. *N Engl J Med*. 2015;373(8):726-736.
- Larkin J, Ascierto PA, Dréno B, et al. Combined vemurafenib and cobimetinib in BRAF-mutated melanoma. *N Engl J Med*. 2014;371(20):1867-1876.
- Long GV, Stroyakovskiy D, Gogas H, et al. Combined BRAF and MEK inhibition versus BRAF inhibition alone in melanoma. *N Engl J Med*. 2014;371(20):1877-1888.

DOI 10.1182/blood-2015-06-650036

© 2015 by The American Society of Hematology

To the editor:

Ferric chloride thrombosis model: unraveling the vascular effects of a highly corrosive oxidant

The topical application of ferric chloride (FeCl_3) to the vasculature is one of the most commonly used experimental approaches to induce thrombosis. The method was first described by Kurz and colleagues¹ and has subsequently been proven to be a highly effective and reliable approach to elucidate the role of platelet receptors, ligands, and activation pathways in promoting thrombosis.²⁻⁷ It has also shed new light on the role of coagulation proteases in regulating thrombin generation and thrombus growth⁸⁻¹¹ and has been used in the context of thrombolysis,¹²⁻¹⁷ although the model appears to have significant limitations in this regard. Despite its widespread use, the precise mechanism by which FeCl_3 induces thrombosis remains controversial. A recent study in *Blood* by Ciciliano and colleagues has provided further insight into the molecular mechanisms underlying FeCl_3 -induced thrombosis, suggesting an important role for charge-dependent aggregation effects of FeCl_3 on blood cells and plasma proteins.¹⁸

It has long been assumed that the major effects of FeCl_3 are limited to the vasculature. FeCl_3 ions have been localized to the endothelium with uptake of iron through endothelial pinocytotic processes. Several studies have described ferric ion-rich membrane-enclosed bodies which transmigrate into the endothelium, followed by exocytosis into the vessel lumen.^{19,20} It was assumed that this accumulation of iron and generation of reactive oxygen species produce endothelial toxicity and denudation, leading to the exposure of subendothelial elements that promote thrombus formation.^{1,21} However, a number of studies have revealed minor endothelial denudation and collagen exposure following FeCl_3 treatment,^{20,22,23} raising the possibility that FeCl_3 has effects beyond the vessel wall.

The first demonstration that red blood cells (RBCs) may play an important role in promoting FeCl_3 -induced thrombosis was derived from our *in vitro* studies using isolated blood cell components perfused through mouse aortae *ex vivo*.²³ Surprisingly, endothelial cells exposed to FeCl_3 alone exhibited minor levels of injury. However, in the presence of whole blood or isolated RBCs, FeCl_3 -induced red cell hemolysis and hemoglobin oxidation promoted extensive vascular injury and thrombosis.²³ Elegant electron microscopy studies by Barr and colleagues confirmed extensive red

cell accumulation on the endothelium following FeCl_3 exposure *in vivo*, with platelets rapidly recruited to accumulated red cell-derived structures.²² Eckly and colleagues revealed surface expression of tissue factor on the ferric ion-rich spherical bodies, which they attributed as the primary mediator of platelet adhesion and fibrin formation.²⁰

The plot thickens further. Ciciliano and colleagues used microfluidic devices coated with endothelial cells to dissect the effects of FeCl_3 on individual blood cell and plasma components.¹⁸ These studies demonstrated concentration-dependent effects of FeCl_3 on protein and blood cell aggregation, independent of effects on the endothelium. This aggregation effect was principally attributed to colloidal chemistry, whereby cells and proteins adhere and aggregate as a result of their charge. The authors have proposed that this physicochemical effect of FeCl_3 on blood cells is the primary instigator driving blood cell adhesion to the endothelium. They argue that this mechanism then facilitates the “secondary” phase of FeCl_3 injury, with red cell aggregates and damaged endothelium providing a reactive surface for the accumulation of platelets and initiation of blood coagulation, necessary for stable thrombus formation.¹⁹⁻²¹ However, it remains to be determined to what extent oxidative damage vs physicochemical effects predominate to induce RBC aggregation and thrombosis.²³ In this context, aluminum chloride (AlCl_3), which carries a similar charge to Fe^{3+} and was used as an additional means of evidence to support the role of colloidal chemistry in cellular and protein aggregation, is also well known to cause oxidative damage, including lipid peroxidation and RBC hemolysis.²⁴ This is in contrast to chromium chloride (CrCl_3), which was also used in this study as a negative control; however, this molecule has antioxidant properties.²⁵ Whether the effects of AlCl_3 can be offset by antioxidants, as was demonstrated for FeCl_3 in our own *ex vivo* studies using isolated aorta,²³ will be important to determine. It is also interesting to note that the oxidative effects of FeCl_3 we observed in isolated aorta were initiated with concentrations of FeCl_3 lower than that observed to induce macroscopic precipitation of plasma proteins.²³

Collectively, the studies highlighted above unequivocally demonstrate that the effects of FeCl_3 on vascular injury and