

Decadal carbon discharge by a mountain stream is dominated by coarse organic matter

Jens M. Turowski^{1*}, Robert G. Hilton², Robert Sparkes^{3,4}

¹Swiss Federal Research Institute WSL, Zürcherstrasse 111, 8903 Birmensdorf, Switzerland

²Department of Geography, Durham University, Science Laboratories, South Road, Durham DH1 3LE, UK

³Department of Materials Science and Metallurgy, University of Cambridge, 27 Charles Babbage Road, Cambridge CB3 0FS, UK

⁴School of Earth Atmospheric and Environmental Sciences, University of Manchester, Williamson Building, Oxford Road, Manchester M13 9PL, UK

ABSTRACT

Rapid erosion in mountain forests results in high rates of biospheric particulate organic carbon (POC) export by rivers, which can contribute to atmospheric carbon dioxide drawdown. However, coarse POC (CPOC) carried by particles >1 mm is rarely quantified. In a forested pre-Alpine catchment, we measured CPOC transport rates and found that they increase more rapidly with water discharge than fine POC (<1 mm) and dissolved organic carbon (DOC). As a result, decadal estimates of CPOC yield of 12.3 ± 1.9 t C km⁻² yr⁻¹ are higher than for fine POC and DOC, even when excluding 4 extreme flood events. When including these floods, CPOC dominates organic carbon discharge (~80%). Most CPOC (69%) was water logged and denser than water, suggesting that CPOC has the potential to contribute to long-term sedimentary burial. Global fluxes remain poorly constrained, but if the transport behavior of CPOC shown here is common to other mountain streams and rivers, then neglecting CPOC discharge could lead to a large underestimation of the global transfer of biospheric POC from land to ocean.

INTRODUCTION

Erosion of particulate organic carbon (POC) from the terrestrial biosphere and its transport by rivers redistributes nutrients and can contribute to atmospheric carbon dioxide drawdown (Bernier, 1982; Stallard, 1998; Battin et al., 2008; Galy et al., 2015). High rates of physical erosion in mountain catchments result in elevated rates of fine POC discharge (FPOC, particles >0.2–0.7 μm and <1 mm), with biospheric FPOC (FPOC_{biosphere}) yields >10 t C km⁻² yr⁻¹ (Hilton et al., 2012; Goñi et al., 2013; Smith et al., 2013; Galy et al., 2015). As a result, mountain rivers can contribute significant amounts of FPOC_{biosphere} to large rivers, lakes, and the oceans (Stallard, 1998; Hilton et al., 2012; Galy et al., 2015). This carbon, recently derived from atmospheric carbon dioxide via photosynthesis, is often transported along with large volumes of clastic sediment (Hilton et al., 2012). High sediment accumulation rates in depositional settings can increase the burial efficiency of POC_{biosphere} and promote the drawdown of atmospheric CO₂ over geological time scales (Bernier, 1982; Kao et al., 2014; Galy et al., 2015).

Despite this recognition, the organic carbon (OC) transported as coarse particulate organic matter (CPOM, particles >1 mm) remains poorly constrained, mainly because it is challenging to measure. CPOM can range in size from leaves to entire trees, and is not captured by typical river water sampling methods (e.g., Goñi et al., 2013; Smith et al., 2013; Hilton et al.,

2015), while it is transported episodically during large floods when it is difficult to work in river channels (West et al., 2011; Wohl, 2013; Kramer and Wohl, 2014). CPOM also contributes to ecosystem functions because it typically contains ~50% carbon by weight and can form the basis of the food chain in many streams (Fisher and Likens, 1973). In addition to contributing to carbon and nutrient transfers in rivers, large wood, consisting of CPOM with lengths >1 m, can affect stream morphology and hydraulics, while providing shelter for in-stream fauna and affecting breeding grounds (Wohl, 2013).

A significant challenge remains to accurately measure coarse POC (CPOC) transport in rivers

across the full size range of CPOM, while linking CPOC transfer to hydrodynamic conditions in rivers; only by doing so can CPOC yields (t C km⁻² yr⁻¹) be accurately quantified. In addition, CPOC eroded from the biosphere is often thought to float (West et al., 2011), suggesting that it could be more susceptible to oxidation upon its delivery to floodplains (Fisher and Likens, 1973), lakes and reservoirs (Seo et al., 2008), and the oceans (West et al., 2011). However, water-logged woody debris, with a density higher than water, is a component of FPOC_{biosphere} in large river systems (Bianchi et al., 2007; Hilton et al., 2015). The amount of water-logged CPOC discharged by mountain rivers remains unknown. Here we use detailed measurements of CPOM transport in the Erlenbach, a 0.7 km² catchment in the Swiss Prealps. Although small, the catchment has geomorphic, climatic, and ecological characteristics that are representative of forested mountain headwater streams in a temperate climate (Schleppi et al., 1999; Smith et al., 2013).

METHODS

The Erlenbach is a steep (11% slope) mountain stream with step-pool morphology and drains 0.7 km² in the Swiss Prealps (47.045707°N, 8.708844°E) (Fig. 1). The mean annual air temperature is ~4.5 °C and the mean annual precipi-




Figure 1. A: Location of the Erlenbach catchment in Switzerland. B: Map of the catchment.

*Current address: Section 5.1 Geomorphology, Helmholtz Centre Potsdam, GFZ German Research Centre for Geosciences, Telegrafenberg 14473 Potsdam, Germany; E-mail: turowski@gfz-potsdam.de.

tation is ~2300 mm. Approximately 40% of the total catchment area is covered by alpine forest, mainly comprising Norway Spruce (*Picea abies*) and European Silver Fir (*Abies alba*) (Schleppi et al., 1999), and a small amount of logging has been done in the upper catchment over the past 10 yr. The remaining 60% of the catchment is covered by wetland and alpine meadows. A well-developed riparian zone is generally lacking and active landslide complexes along the channel lead to strong channel-hillslope coupling typical of many steep mountain catchments. Both DOC and FPOC fluxes have been previously determined (Hagedorn et al., 2000; Smith et al., 2013). The FPOC has been partitioned into that derived from the terrestrial biosphere (FPOC_{biosphere}) and that from rock-derived OC using stable carbon isotopes, nitrogen to carbon ratios, and radiocarbon (Smith et al., 2013).

We use CPOM data sampled with three different methods (see the GSA Data Repository¹), each of which is suitable for a different water discharge range (Turowski et al., 2013). All sampling locations were within 30 m of a permanently installed gauge measuring water discharge (Q_w , L s⁻¹) at 10 min intervals (Rickenmann et al., 2012). At low Q_w (1 L s⁻¹ to 1000 L s⁻¹; most samples <250 L s⁻¹), Bunte traps were used (Bunte et al., 2007). These are metal frames placed on the stream bed, to which a net with 6 mm meshing is attached. At intermediate Q_w (200 L s⁻¹ to 1500 L s⁻¹; most samples >400 L s⁻¹) basket samplers were used (Rickenmann et al., 2012), consisting of metal cubes with 1 m edges and walls and floor made of metal mesh with 10 mm holes. The samplers automatically move into the flow when Q_w exceeds a pre-defined threshold value and when bedload transport is recorded. Both traps and baskets sample the entire flow depth with nearly 100% efficiency (Rickenmann et al., 2012; Turowski et al., 2013).

Woody material in the basket and trap samples was separated from clastic material in the field and weighed. Basket samples from A.D. 2011 to 2013 were separated into floating and sinking fractions in the field by dropping them into a water-filled bucket. Subsequently, the material was dried for 24 h at 80 °C, and the dry mass was obtained.

The diameter and length of large woody debris trapped in a retention basin after two extreme events (1995, 2010) complement the data at high Q_w (>5000 L s⁻¹). Masses were calculated assuming a cylindrical shape and a dry density of 410 kg/m³, which is typical for the Norway Spruce (*Picea Abies*) that is common in the catchment. The three methods were made

comparable by using distributions of particle masses (Turowski et al., 2013). CPOC was calculated from CPOM using the mean OC content of 47.8% ± 3.8% (±standard deviation) measured from 37 randomly drawn subsamples.

RESULTS

The transport rate of CPOC (kg C s⁻¹) was positively correlated with Q_w and well described by a power law rating curve ($r^2 = 0.87$; Fig. 2A). CPOC transport increases much more rapidly with increasing Q_w (rating curve exponent $\beta = 4.14 \pm 0.19$) than DOC ($r^2 = 0.98$, $\beta = 1.17 \pm 0.04$) and FPOC_{biosphere} ($r^2 = 0.88$, $\beta = 1.90 \pm 0.10$). The data confirm that high river power is needed to mobilize and transport CPOC (West et al., 2011; Wohl, 2013). The relationship is consistent with the difference between bedload and suspended load transport rates in the Erlenbach (cf. Turowski et al., 2009; Smith et al., 2013), suggesting that CPOC is traveling as part of the bedload. This interpretation is supported by the observation that large fractions (mean 69%, median 78%) of the CPOM were water logged and denser than water, especially at high Q_w (Fig. 2B). Water logging likely occurs during storage of CPOM in log jams in the stream, or within saturated soil and litter on the hillslopes.

To estimate the decadal rate of CPOC discharge, we fitted a linear regression in double-logarithmic space to obtain a rating curve. The data points obtained from the retention basin material were not included in the regression, but are close to the rating curve at high Q_w . We used additional data from 2013, which resulted in a different rating curve than previously published (cf. Turowski et al., 2013). The rating curve was integrated over 31 yr of Q_w measurements. During this period, 4 exceptional flood events affected the catchment (Turowski et al., 2009), with peak Q_w >9000 L s⁻¹ and return periods >20 yr. Not accounting for these 4 floods, the background CPOC yield was 12.3 ± 1.9 t C km⁻² yr⁻¹. Uncertainties were derived from analytical errors of the rating curve fits. The exceptional floods delivered between 331 and 1066 t C km⁻², with an average of 585 t C km⁻². These values are lower than the 6300–19,100 t C km⁻² of large wood carbon (LWC) delivered to the ocean during typhoon Morakot in Taiwan (West et al., 2011), but higher than the 10–24 t C km⁻² of LWC delivered from the upper Rio Chagres, Panama, in a rain storm (Wohl and Ogden, 2013). In total, the 4 floods delivered 2338 ± 1609 t C km⁻², or 75.4 ± 51.9 t C km⁻² yr⁻¹. When added to the background rate,




Figure 2. A: Coarse particulate organic carbon (CPOC) transport rate (kg C s⁻¹) as a function of water discharge (L s⁻¹) for the Erlenbach study catchment, Switzerland. Also shown are published measurements of dissolved organic carbon (DOC) (Hagedorn et al., 2000) and fine biospheric particulate organic carbon (FPOC_{biosphere}) transfer (Smith et al., 2013). Data are fit with power law rating curves for CPOC (thick solid line, exponent $\beta = 4.14 \pm 0.19$), FPOC_{biosphere} (dashed line, $\beta = 1.90 \pm 0.10$), and DOC (fine solid line, $\beta = 1.17 \pm 0.04$). CPOC data from extreme events in A.D. 1995 and 2010 (box, cross) support the rating curve fit. **B:** Percent of water-logged coarse particulate organic matter (CPOM) at the time of collection. Mean (69%) and median (78%) of 35 basket samples are indicated by the solid and the dashed line, respectively. Low values at water discharges <600 L s⁻¹ arose from autumn samples with small absolute mass consisting mainly of fresh leaves. Summed over all samples, water-logged CPOM contributed 76% to the total dry mass.

¹GSA Data Repository item 2016007, additional method information and Figures DR1–DR3, is available online at www.geosociety.org/pubs/ft2016.htm, or on request from editing@geosociety.org or Documents Secretary, GSA, P.O. Box 9140, Boulder, CO 80301, USA.

the average CPOC discharge estimate is $87.7 \pm 51.9 \text{ t C km}^{-2} \text{ yr}^{-1}$. Exceptional flood events appear to be even more important for CPOC than for $\text{FPOC}_{\text{biosphere}}$ (Hilton et al., 2012), which results from the steep relationship between CPOC transport rate and Q_w (Fig. 2A; Fig. DR1 in the Data Repository).

The background CPOC yield ($12.3 \pm 1.9 \text{ t C km}^{-2} \text{ yr}^{-1}$) from the Erlenbach is a significant catchment-scale carbon transfer (Hilton et al., 2012; Galy et al., 2015) and on its own is comparable to the upper range of estimates of $\text{FPOC}_{\text{biosphere}}$ yields from temperate and tropical active mountain belts (Fig. 3). Other carbon transfers from the Erlenbach, obtained using the same methods on previously collected data (Hagedorn et al., 2000; Smith et al., 2013), are lower than CPOC transfer, with a DOC yield of $11.3 \pm 0.0 \text{ Mg C km}^{-2} \text{ yr}^{-1}$, and an $\text{FPOC}_{\text{biosphere}}$ yield of $10.7 \pm 0.1 \text{ Mg C km}^{-2} \text{ yr}^{-1}$. The background CPOC transfer thus represents $\sim 36\%$ of the decadal biospheric OC discharge by this catchment. Inclusion of the exceptional events raises CPOC transfer to as much as $\sim 80\%$ of the total OC (TOC) discharge (Fig. 3). We can assess the sustainability of OC export by comparing it to the net primary production (NPP) of $\sim 740 \text{ Mg C km}^{-2} \text{ yr}^{-1}$ in the Erlenbach catchment (see the Data Repository). The background rate of CPOC discharge is $\sim 1.7\%$ of this NPP and is sustainable, in agreement with a global compilation of river $\text{FPOC}_{\text{biosphere}}$ yields (Galy et al., 2015). However, extreme events may severely deplete the biosphere stock of carbon. The CPOC discharge during a single event appears to have the potential to exceed the catchment's yearly production; our data suggest that on decadal time scales, exceptional events discharge $\sim 10\%$ of the NPP.

DISCUSSION

The contribution of CPOC to carbon discharge by rivers is not typically quantified, and a direct comparison with data from other catchments remains challenging. Notwithstanding, it has been calculated that LWC alone contributes at least 10% and as much as 35% of the total carbon yields in mountain rivers with catchment areas as large as 2000 km^2 (Fig. DR2) (Seo et al., 2008). CPOM particles smaller than large wood to sizes of 1 mm were not considered in that study, but dominate CPOC in the Erlenbach (cf. Turowski et al., 2013). Based on the Erlenbach's size, its $\text{FPOC}_{\text{biosphere}}$ and LWC yields are similar to those observed in other mountain regions in the world (Fig. DR2). $\text{FPOC}_{\text{biosphere}}$ yields are known to be strongly linked to physical erosion rate (Fig. 3) (Galy et al., 2015), and high yields are observed in active mountain belts in temperate and tropical settings (Hilton et al., 2012). In accord with this, estimates of LWC transfer in Taiwanese catchments are larger than for the Erlenbach (West et al., 2011). Therefore, we pro-




Figure 3. Literature data of biospheric particulate organic carbon ($\text{POC}_{\text{biosphere}}$) yield ($\text{t C km}^{-2} \text{ yr}^{-1}$) plotted against suspended sediment yield ($\text{t km}^{-2} \text{ yr}^{-1}$) for global rivers with fitted relationship (dashed line) (Galy et al., 2015) (Q_w — water discharge). The Erlenbach (Switzerland) catchment does not show exceptional fine $\text{POC}_{\text{biosphere}}$ ($\text{FPOC}_{\text{biosphere}}$) yields for its suspended sediment yield. Inclusion of coarse POC (CPOC) for the Erlenbach, not available for the other catchments, increases carbon export by an order of magnitude (black and white stars).

pose that the often unmeasured CPOC fraction is a significant component of $\text{POC}_{\text{biosphere}}$ export from forested mountain catchments.

To make a tentative first assessment of the global significance of CPOC transport, we assume that the Erlenbach catchment is representative for temperate mountain forests, which cover a total area of $1.2 \times 10^6 \text{ km}^2$ worldwide (Sands, 2005). While the climatic, geomorphic, and ecological characteristics of the Erlenbach support that assumption, its physical erosion rate is high (Fig. 3). Without more measurements of CPOC transport (Fig. 1) and estimation of CPOC yields (Fig. 3), a global CPOC discharge estimate remains poorly constrained. Based on the Erlenbach background CPOC yield over 31 yr ($12.3 \pm 1.9 \text{ t C km}^{-2} \text{ yr}^{-1}$), the global CPOC discharge from temperate mountain forest catchments could be $\sim 15 \text{ Mt C yr}^{-1}$. This is $\sim 10\%$ of the recent estimate of global $\text{FPOC}_{\text{biosphere}}$ discharge to the oceans by rivers of $157 +74/-50 \text{ Mt C yr}^{-1}$ (Galy et al., 2015). If extreme floods are included, CPOC discharge from temperate mountain forests could be even higher (Fig. 3). Global CPOC discharge would further increase if boreal, subtropical, and tropical mountain forests were considered. We are aware that these estimates are based on extrapolation from a very small continental area and absolute flux has large uncertainty. Nevertheless, the magnitude of the estimate demonstrates the need to better quantify CPOC transfer rates in mountain rivers and track its conveyance through large river systems.

Little is known about the onward fate and routing of CPOC through large rivers. On average $\sim 69\%$ of the CPOM transported by the Erlenbach was water logged, with a density greater than water (Fig. 2B). If this observation applies to other temperate streams where channel morphology can promote transient storage of CPOM,

sampling with drift nets may have missed large fractions of CPOM traveling near the stream bed. Perhaps more important, water-logged CPOM may have a different fate in fluvial networks than if it were to float. During transport in steep channels, water-logged CPOM may be ground by gravel bedload, reducing its size. The size reduction of CPOM by bedload grinding is poorly understood, but the observed magnitude of the CPOC flux means it could be an important in-stream source of $\text{FPOC}_{\text{biosphere}}$ (Hilton et al., 2012).

Furthermore, a high density of CPOM may promote its burial potential in sedimentary basins. If water-logged CPOM is delivered to depositional environments as part of the bedload it is more likely to rapidly accumulate in sedimentary deposits. Observations of large terrestrial organic debris in deep-sea turbidites in Indonesia (Saller et al., 2006), woody clasts and plant debris in modern deep-sea sediments offshore Taiwan (Kao et al., 2014), and mountain rivers draining the west coast of the United States (Leithold and Hope, 1999) all suggest that CPOC can be delivered to deep-marine settings. We substantiate these arguments by estimating the contribution of CPOC to TOC in exhumed turbidite sequences in the Apennines, Italy (see the Data Repository). Despite estimated transport distances of as much as 300 km offshore, CPOC was buried and preserved for 14 m.y. and represents $\sim 10\%$ of the TOC. Water-logged woody debris can be delivered by mountain rivers as CPOC (Fig. 2B), and its presence may enhance the efficiency of carbon burial and associated atmospheric CO_2 sequestration by erosion of mountain belts (Kao et al., 2014; Galy et al., 2015).

CONCLUSIONS

CPOC is the dominant form of OC discharge by the Erlenbach over decadal time

scales, increasing the carbon loss from the biosphere by ~250% over DOC and FPOC^{biosphere}. The majority of CPOC may be transported in water-logged CPOM as part of the bedload. Our observations provide new impetus to study the production, transfer, and routing of CPOC from mountain headwaters, and subsequently through large river systems to fully assess the net impact of erosion on the global carbon cycle (Battin et al., 2008; Hilton et al., 2012, 2015; Galy et al., 2015). Due to anthropogenic CO₂ emissions and global warming, extreme precipitation events may become more frequent (Rajczak et al., 2013), causing an increased number of extreme floods. CPOC transport exhibits a much stronger dependency on water discharge than FPOC and DOC transport (Fig. 1), and could therefore become more important for carbon budgets of mountain streams in the coming decades. This may have implications for forest management, food availability in stream ecosystems, and carbon mobilization by erosion of the terrestrial biosphere.

ACKNOWLEDGMENTS

We thank the current and former members of the mountain hydrology team for help in the field and in the laboratory. M. Sieber prepared the samples for chemical analyses. F. Hagedorn shared his DOC data of the Erlenbach. A. Galy and N. Hovius assisted with fieldwork in the Apennines and subsequent chemical analyses. Discussions with A. Badoux, K. Bunte, V. Galy, N. Hovius, J. Kirchner, K. Krause, and P. Schleppi are acknowledged. We thank E. Leithold, P. Raymond, and an anonymous reviewer for their comments, which improved the work. This study was supported by Schweizerischer Nationalfonds (SNF) grant 200021_124634/1, Engineering and Physical Sciences Research Council (UK) grants EP/P502365/1 and EP/P504120/1, and the Swiss Federal Research Institute for Forest, Snow and Landscape Research.

REFERENCES CITED

- Battin, T.J., Kaplan, L.A., Findlay, S., Hopkinson, C.S., Marti, E., Packman, A.I., Newbold, J.D., and Sabater, F., 2008, Biophysical controls on organic carbon fluxes in fluvial networks: *Nature Geoscience*, v. 1, p. 95–100, doi:10.1038/ngeo101.
- Berner, R.A., 1982, Burial of organic carbon and pyrite sulfur in the modern ocean: Its geochemical and environmental significance: *American Journal of Science*, v. 282, p. 451–473, doi:10.2475/ajs.282.4.451.
- Bianchi, T.S., Geller, J.J., and Allison, M.A., 2007, Hydrodynamic sorting and transport of terrestrially-derived organic carbon in sediments of the Mississippi and Atchafalaya Rivers: *Estuarine, Coastal and Shelf Science*, v. 73, p. 211–222, doi:10.1016/j.ecss.2007.01.004.
- Bunte, K., Swingle, K.W., and Abt, S.R., 2007, Guidelines for using bedload traps in coarse-bedded mountain streams: Construction, installation, operation and sample processing: U.S. Forest Service Rocky Mountain Research Station, General Technical Report RMRS-GTR-191, 91 p., http://www.fs.fed.us/rm/pubs/rmrs_gtr191.pdf.
- Fisher, S.G., and Likens, G.E., 1973, Energy flow in Bear Brook, New Hampshire: An integrative approach to stream ecosystem metabolism: *Ecological Monographs*, v. 43, p. 421–439, doi:10.2307/1942301.
- Galy, V., Peucker-Ehrenbrink, B., and Eglinton, T., 2015, Global carbon export from the terrestrial biosphere controlled by erosion: *Nature*, v. 521, p. 204–207, doi:10.1038/nature14400.
- Goñi, M.A., Hatten, J.A., Wheatcroft, R.A., and Borgeld, J.C., 2013, Particulate organic matter export by two contrasting small mountainous rivers from the Pacific Northwest, U.S.A.: *Journal of Geophysical Research*, v. 118, p. 112–134, doi:10.1002/jgrg.20024.
- Hagedorn, F., Schleppi, P., Waldner, P., and Flühler, H., 2000, Export of dissolved organic carbon and nitrogen from Gleysol dominated catchments—The significance of water flow paths: *Biogeochemistry*, v. 50, p. 138–161, doi:10.1023/A:1006398105953.
- Hilton, R.G., Galy, A., Hovius, N., Kao, S.-J., Horng, M.-J., and Chen, H., 2012, Climatic and geomorphic controls on the erosion of terrestrial biomass from subtropical mountain forest: *Global Biogeochemical Cycles*, v. 26, GB3014, doi:10.1029/2012GB004314.
- Hilton, R.G., Galy, V., Gaillardet, J., Dellinger, M., Bryant, C., O'Regan, M., Gröcke, D.R., Coxall, H., Bouchez, J., and Calmels, D., 2015, Erosion of organic carbon in the Arctic as a geological carbon dioxide sink: *Nature*, v. 524, p. 84–87, doi:10.1038/nature14653.
- Kao, S.-J., et al., 2014, Preservation of terrestrial organic carbon in marine sediments offshore Taiwan: Mountain building and atmospheric carbon dioxide sequestration: *Earth Surface Dynamics*, v. 2, p. 127–139, doi:10.5194/esurf-2-127-2014.
- Kramer, N., and Wohl, E., 2014, Estimating fluvial wood discharge using time-lapse photography with varying sampling intervals: *Earth Surface Processes and Landforms*, v. 39, p. 844–852, doi:10.1002/esp.3540.
- Leithold, E.L., and Hope, R.S., 1999, Deposition and modification of a flood layer on the northern California shelf: Lessons from and about the fate of terrestrial particulate organic carbon: *Marine Geology*, v. 154, p. 183–195, doi:10.1016/S0025-3227(98)00112-1.
- Rajczak, J., Pall, P., and Schär, C., 2013, Projections of extreme precipitation events in regional climate simulations for Europe and the Alpine region: *Journal of Geophysical Research*, v. 118, p. 1–17, doi:10.1002/jgrd.50297.
- Rickenmann, D., Turowski, J.M., Fritschi, B., Klaiber, A., and Ludwig, A., 2012, Bedload transport measurements at the Erlenbach stream with geophones and automated basket samplers: *Earth Surface Processes and Landforms*, v. 37, p. 1000–1011, doi:10.1002/esp.3225.
- Saller, A., Lin, R., and Dunham, J., 2006, Leaves in turbidite sands: The main source of oil and gas in the deep-water Kutei Basin, Indonesia: *American Association of Petroleum Geologists Bulletin*, v. 90, p. 1585–1608, doi:10.1306/04110605127.
- Sands, R., 2005, *Forestry in a global context*: Wallingford, Connecticut, CABI Publishing, 272 p.
- Schleppi, P., Muller, N., Edwards, P.J., and Bucher, J.B., 1999, Three years of increased nitrogen deposition do not affect the vegetation of a montane forest ecosystem: *Phyton*, v. 39, p. 197–204.
- Seo, J.I., Nakamura, F., Nakano, D., Ichiyanagi, H., and Chun, K.W., 2008, Factors controlling the fluvial export of large woody debris, and its contribution to organic carbon budgets at watershed scales: *Water Resources Research*, v. 44, W04428, doi:10.1029/2007WR006453.
- Smith, J.C., Hovius, N., Galy, A., Tye, A.M., Turowski, J.M., and Schleppi, P., 2013, Runoff-driven export of particulate organic carbon from soil in temperate forested uplands: *Earth and Planetary Science Letters*, v. 365, p. 198–208, doi:10.1016/j.epsl.2013.01.027.
- Stallard, R.F., 1998, Terrestrial sedimentation and the carbon cycle: Coupling weathering and erosion to carbon burial: *Global Biogeochemical Cycles*, v. 12, p. 231–257, doi:10.1029/98GB00741.
- Turowski, J.M., Yager, E.M., Badoux, A., Rickenmann, D., and Molnar, P., 2009, The impact of exceptional events on erosion, bedload transport and channel stability in a step-pool channel: *Earth Surface Processes and Landforms*, v. 34, p. 1661–1673, doi:10.1002/esp.1855.
- Turowski, J.M., Badoux, A., Bunte, K., Rickli, C., Federspiel, N., and Jochner, M., 2013, The mass distribution of coarse particulate organic matter exported from an alpine headwater stream: *Earth Surface Dynamics*, v. 1, p. 1–11, doi:10.5194/esurf-1-1-2013.
- West, A.J., Lin, C.-W., Lin, T.-C., Hilton, R.G., Liu, S.-H., Chang, C.-T., Lin, K.-C., Galy, A., Sparkes, R.B., and Hovius, N., 2011, Mobilization and transport of coarse woody debris to the oceans triggered by an extreme tropical storm: *Limnology and Oceanography*, v. 56, p. 77–85, doi:10.4319/lo.2011.56.1.0077.
- Wohl, E., 2013, Floodplains and wood: *Earth-Science Reviews*, v. 123, p. 194–212, doi:10.1016/j.earscirev.2013.04.009.
- Wohl, E., and Ogden, F.L., 2013, Organic carbon export in the form of wood during an extreme tropical storm, Upper Rio Chagres, Panama: *Earth Surface Processes and Landforms*, v. 38, p. 1407–1416, doi:10.1002/esp.3389.

Manuscript received 23 July 2015

Revised manuscript received 29 October 2015

Manuscript accepted 29 October 2015

Printed in USA