

Mendelian Randomization Analysis of n-6 Polyunsaturated Fatty Acid Levels and Pancreatic Cancer Risk



Dalia H. Ghoneim¹, Jingjing Zhu¹, Wei Zheng², Jirong Long², Harvey J. Murff³, Fei Ye⁴, Veronica Wendy Setiawan⁵, Lynne R. Wilkens¹, Nikhil K. Khankari², Philip Haycock⁶, Samuel O. Antwi⁷, Yaohua Yang², Alan A. Arslan⁸, Laura E. Beane Freeman⁹, Paige M. Bracci¹⁰, Federico Canzian¹¹, Mengmeng Du¹², Steven Gallinger¹³, Graham G. Giles^{14,15,16}, Phyllis J. Goodman¹⁷, Charles Kooperberg¹⁸, Loïc Le Marchand¹⁹, Rachel E. Neale¹⁹, Ghislaine Scelo²⁰, Kala Visvanathan^{21,22}, Emily White²³, Demetrius Albanes⁹, Pilar Amiano²⁴, Gabriella Andreotti⁹, Ana Babic²⁵, William R. Bamlet²⁶, Sonja I. Berndt⁹, Lauren K. Brais²⁵, Paul Brennan²⁰, Bas Bueno-de-Mesquita^{27,28,29,30}, Julie E. Buring^{31,32}, Peter T. Campbell³³, Kari G. Rabe²⁶, Stephen J. Chanock⁹, Priya Duggal³⁴, Charles S. Fuchs^{35,36,37}, J. Michael Gaziano^{32,38}, Michael G. Goggins³⁹, Thilo Hackert⁴⁰, Manal M. Hassan⁴¹, Kathy J. Helzlsouer⁴², Elizabeth A. Holly¹⁰, Robert N. Hoover⁹, Verena Katske⁴³, Robert C. Kurtz⁴⁴, I-Min Lee^{31,32}, Núria Malats⁴⁵, Roger L. Milne^{14,15,16}, Neil Murphy⁴⁶, Ann L. Oberg²⁶, Miquel Porta⁴⁷, Nathaniel Rothman⁹, Howard D. Sesso^{31,32}, Debra T. Silverman⁹, Ian M. Thompson Jr⁴⁸, Jean Wactawski-Wende⁴⁹, Xiaoliang Wang²³, Nicolas Wentzensen⁹, Herbert Yu¹, Anne Zeleniuch-Jacquotte⁵⁰, Kai Yu⁹, Brian M. Wolpin²⁵, Eric J. Jacobs⁵¹, Eric J. Duell⁵², Harvey A. Risch⁵³, Gloria M. Petersen²⁶, Laufey T. Amundadottir⁵⁴, Peter Kraft⁵⁵, Alison P. Klein^{21,22,39}, Rachel Z. Stolzenberg-Solomon⁹, Xiao-Ou Shu², and Lang Wu¹

ABSTRACT

Background: Whether circulating polyunsaturated fatty acid (PUFA) levels are associated with pancreatic cancer risk is uncertain. Mendelian randomization (MR) represents a study design using genetic instruments to better characterize the relationship between exposure and outcome.

Methods: We utilized data from genome-wide association studies within the Pancreatic Cancer Cohort Consortium and Pancreatic Cancer Case-Control Consortium, involving approximately 9,269 cases and 12,530 controls of European descent, to evaluate associations between pancreatic cancer risk and genetically predicted plasma n-6 PUFA levels. Conventional MR analyses were performed using individual-level and summary-level data.

Results: Using genetic instruments, we did not find evidence of associations between genetically predicted plasma n-6 PUFA levels

and pancreatic cancer risk [estimates per one SD increase in each PUFA-specific weighted genetic score using summary statistics: linoleic acid odds ratio (OR) = 1.00, 95% confidence interval (CI) = 0.98–1.02; arachidonic acid OR = 1.00, 95% CI = 0.99–1.01; and dihomo-gamma-linolenic acid OR = 0.95, 95% CI = 0.87–1.02]. The OR estimates remained virtually unchanged after adjustment for covariates, using individual-level data or summary statistics, or stratification by age and sex.

Conclusions: Our results suggest that variations of genetically determined plasma n-6 PUFA levels are not associated with pancreatic cancer risk.

Impact: These results suggest that modifying n-6 PUFA levels through food sources or supplementation may not influence risk of pancreatic cancer.

¹Division of Cancer Epidemiology, Population Sciences in the Pacific Program, University of Hawaii Cancer Center, University of Hawaii at Manoa, Honolulu, Hawaii. ²Division of Epidemiology, Department of Medicine, Vanderbilt Epidemiology Center, Vanderbilt-Ingram Cancer Center, Vanderbilt University Medical Center, Nashville, Tennessee. ³Division of General Internal Medicine, Department of Medicine, Vanderbilt University, Nashville, Tennessee. ⁴Department of Biostatistics, Vanderbilt University Medical Center, Nashville, Tennessee. ⁵Department of Preventive Medicine, Keck School of Medicine, University of Southern California, Los Angeles, California. ⁶MRC Integrative Epidemiology Unit, University of Bristol, Bristol, England, United Kingdom. ⁷Department of Health Sciences Research, Mayo Clinic, Jacksonville, Florida. ⁸Departments of Obstetrics and Gynecology, Population Health and Environmental Medicine, NYU Perlmutter Comprehensive Cancer Center, New York, New York. ⁹Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Bethesda, Maryland. ¹⁰Department of Epidemiology and Biostatistics, University of California, San Francisco, San Francisco, California.

¹¹Genomic Epidemiology Group, German Cancer Research Center (DKFZ), Heidelberg, Germany. ¹²Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, New York, New York. ¹³Lunenfeld-Tanenbaum Research Institute, Sinai Health System and University of Toronto, Toronto, Ontario, Canada. ¹⁴Division of Cancer Epidemiology, Cancer Council Victoria, Melbourne, Victoria, Australia. ¹⁵Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, The University of Melbourne, Melbourne, Victoria, Australia. ¹⁶Precision Medicine, School of Clinical Sciences at Monash Health, Monash University, Clayton, Victoria, Australia. ¹⁷SWOG Statistical Center, Fred Hutchinson Cancer Research Center, Seattle, Washington. ¹⁸Division of Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle, Washington. ¹⁹Department of Population Health, QIMR Berghofer Medical Research Institute, Brisbane, Queensland, Australia. ²⁰International Agency for Research on Cancer, Lyon, France. ²¹Department of Epidemiology, Johns Hopkins School of Public Health, Baltimore, Maryland. ²²Department of Oncology, Sidney Kimmel Comprehensive Cancer Center,

Introduction

Pancreatic cancer remains one of the deadliest cancers (1). Polyunsaturated fatty acids (PUFA), linked to the inflammatory process, may influence pancreatic cancer development (2). However, evidence from epidemiologic studies is inconsistent (3). For example, associations with n-3 PUFA intake were inverse, positive, or null, and associations with n-6 PUFA intake were positive or null across different studies. Conventional epidemiologic study designs may suffer from methodologic limitations, such as reverse causation, selection bias, and uncontrolled confounding (4). We, therefore, conducted a Mendelian randomization (MR) analysis using genetic variants as instrumental variables. Higher proportions of variance of n-6 PUFA levels were explained by variants compared with n-3 PUFA levels (5, 6). We thus focused on n-6 PUFA in our analysis.

Materials and Methods

Instrumental variables

We identified SNPs associated with plasma or red blood cell (RBC) levels of n-6 PUFAs [linoleic acid, arachidonic acid, adrenic acid, gamma linolenic acid (GLA), and dihomo-gamma-linolenic acid (DGLA)] from the genome-wide association studies (GWAS) catalog and from published literature (up to November 2018; ref. 6). We selected SNPs associated at $P < 5 \times 10^{-8}$ that were independent from each other ($r^2 < 0.1$). For correlated SNPs, the SNP with a lower P value was selected unless an independent association was reported, in which case both were selected. We used estimates of association with plasma PUFA levels for our analyses. For SNPs initially reported to be associated with RBC PUFA levels, we checked their associations with plasma levels in the GWAS conducted by the Cohorts for Heart and Aging Research in Genomic Epidemiology Consortium (CHARGE;

ref. 6); if estimates were significant ($P < 0.05$) we included them in our analyses.

Genetic association datasets for pancreatic cancer risk

For evaluation of associations with pancreatic cancer risk, we used data from four GWASs conducted in the Pancreatic Cancer Cohort Consortium (PanScan) and the Pancreatic Cancer Case-Control Consortium (PanC4; ref. 7). Detailed information on quality control and imputation has been provided elsewhere (7). Data from approximately 9,269 cases and 12,530 controls of European ancestry were used. Only variants with imputation quality of $r^2 \geq 0.3$ were retained.

MR analysis

We performed separate MR analyses for each type of PUFA. On the basis of power estimation (<https://shiny.cnsgenomics.com/mRnd/>), the minimal detectable ORs per SD of genetically predicted PUFA levels at 80% power and alpha of 0.05 ranged from 1.08 to 1.13 for linoleic acid, 1.06 to 1.21 for arachidonic acid, 1.13 to 1.15 for adrenic acid, 1.17 to 1.27 for GLA, and 1.08 to 1.11 for DGLA. We created a weighted genetic score (wGRS) to represent the genetically estimated PUFA level using information from published GWASs of PUFA plasma levels and data from PanScan/PanC4 GWASs. For each subject, a wGRS was created as the weighted sum of the number of association alleles at each locus multiplied by the point estimate for the association with plasma PUFA level:

$$wGRS = \sum_{i=1}^n \beta_i SNP_i$$

where β_i is the regression coefficient of the i th SNP for the PUFA and SNP_i is the dosage of the association alleles (0, 1, 2) of the i th SNP. All association alleles were converted to correspond to increased PUFA

Johns Hopkins School of Medicine, Baltimore, Maryland. ²³Cancer Prevention Program, Fred Hutchinson Cancer Research Center, Seattle, Washington. ²⁴Ministry of Health of the Basque Government, Public Health Division of Gipuzkoa, Biodonostia Health Research Institute, Donostia-San Sebastian; CIBER Epidemiología y Salud Pública, Madrid, Spain. ²⁵Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, Massachusetts. ²⁶Department of Health Sciences Research, Mayo Clinic College of Medicine, Rochester, Minnesota. ²⁷Department for Determinants of Chronic Diseases (DCD), National Institute for Public Health and the Environment (RIVM), Bilthoven, the Netherlands. ²⁸Department of Gastroenterology and Hepatology, University Medical Centre, Utrecht, the Netherlands. ²⁹Department of Epidemiology and Biostatistics, The School of Public Health, Imperial College London, London, United Kingdom. ³⁰Department of Social and Preventive Medicine, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia. ³¹Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts. ³²Division of Preventive Medicine, Brigham and Women's Hospital, Boston, Massachusetts. ³³Behavioral and Epidemiology Research Group, American Cancer Society, Atlanta, Georgia. ³⁴Department of Epidemiology, Johns Hopkins School of Public Health, Baltimore, Maryland. ³⁵Yale Cancer Center, New Haven, Connecticut. ³⁶Department of Medicine, Yale School of Medicine, New Haven, Connecticut. ³⁷Smilow Cancer Hospital, New Haven, Connecticut. ³⁸Boston Veteran Affairs Healthcare System, Boston, Massachusetts. ³⁹Department of Pathology, Sol Goldman Pancreatic Cancer Research Center, Johns Hopkins School of Medicine, Baltimore, Maryland. ⁴⁰Department of General, Visceral and Transplantation Surgery, University of Heidelberg, Heidelberg, Germany. ⁴¹Department of Gastrointestinal Medical Oncology, The University of Texas MD Anderson Cancer Center, Houston, Texas. ⁴²Epidemiology and Genomics Research Program, Division of Cancer Control and Population Science, National Cancer Institute, National Institutes of Health, Bethesda, Maryland. ⁴³Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany. ⁴⁴Gastroenterology, Hepatology, and Nutrition Service, Memorial Sloan Kettering Cancer Center, New York, New York. ⁴⁵Genetic and Molecular Epidemiology Group, Spanish National

Cancer Research Center, Madrid, Spain. ⁴⁶Section of Nutrition and Metabolism, International Agency for Research on Cancer, Lyon, France. ⁴⁷Hospital del Mar Institute of Medical Research (IMIM), Universitat Autònoma de Barcelona, Barcelona, Spain. ⁴⁸CHRISTUS Santa Rosa Hospital - Medical Center, San Antonio, Texas. ⁴⁹Department of Epidemiology and Environmental Health, University of Buffalo, Buffalo, New York. ⁵⁰Departments of Population Health and Environmental Medicine, NYU Perlmutter Comprehensive Cancer Center, New York, New York. ⁵¹Epidemiology Research Program, American Cancer Society, Atlanta, Georgia. ⁵²Unit of Nutrition and Cancer, Cancer Epidemiology Research Program, Bellvitge Biomedical Research Institute (IDIBELL), Catalan Institute of Oncology (ICO), Barcelona, Spain. ⁵³Department of Chronic Disease Epidemiology, Yale School of Public Health, New Haven, Connecticut. ⁵⁴Laboratory of Translational Genomics, Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Bethesda, Maryland. ⁵⁵Department of Epidemiology, Harvard School of Public Health, Boston, Massachusetts.

Note: D.H. Ghoneim and J. Zhu contributed equally as co-first authors of this article.

Corresponding Authors: Lang Wu, University of Hawaii Cancer Center, 701 Ilalo Street, Building B, Room 520, Honolulu, HI 96813. Phone: 808-564-5965; Fax: 808-586-2982; E-mail: lwu@cc.hawaii.edu; and Xiao-Ou Shu, Division of Epidemiology, Department of Medicine, Vanderbilt Epidemiology Center, Vanderbilt-Ingram Cancer Center, Vanderbilt University Medical Center, 2525 West End Avenue, Suite 600, Nashville, TN 37232. Phone: 615-936-0713; E-mail: xiao-ou.shu@vanderbilt.edu

Cancer Epidemiol Biomarkers Prev 2020;29:2735-9

doi: 10.1158/1055-9965.EPI-20-0651

©2020 American Association for Cancer Research.

levels. We used logistic regression models to assess the associations between wGRS and pancreatic cancer risk. Besides unadjusted analyses, we performed analyses adjusting for age, sex, and the top principal components. We stratified the data by age (<50 years, 50–70 years, and ≥70 years) and sex. We also used the summary statistics of the PanScan/PanC4 GWASs to estimate the MR associations using the fixed effects inverse variance-weighted approach (4).

Results

The instruments used for each of the n-6 PUFA are included in **Table 1**. There was no evidence of association (at $P < 0.01$) between any of the wGRS and common pancreatic cancer risk factors. We did not detect any statistically significant associations between genetically predicted plasma n-6 PUFA levels and pancreatic cancer risk (**Table 2**). The estimates for adrenic acid and GLA had wide confidence intervals, consistent with the estimated lower power for these. The associations remained virtually unchanged regardless of covariate adjustment, analyzing individual-level versus summary statistics data, or within strata of age or sex.

Discussion

We did not observe significant associations between genetically predicted n-6 PUFA levels and pancreatic cancer risk in the PanScan/PanC4 subjects. As the proportion of variance of n-6 PUFA that can be

explained by the summed association magnitudes of these GWAS-identified loci is relatively high, there is reasonable statistical power to detect any meaningful associations. PUFA has been reported to be associated with colorectal cancer risk on the basis of MR analysis (8). Most dietary sources of n-6 PUFA are consumed infrequently. A limitation of this study is that there is no information for the genetic variants associated with total n-6 PUFA levels in previous literature, and thus, we could not determine the association between total n-6 PUFA levels and pancreatic cancer risk using genetic instruments. Alternative designs of a direct assessment of dietary sources and measurement of PUFA levels in blood at repeat timepoints can better characterize the relationship between PUFA and pancreatic cancer risk. Further studies are also needed to investigate the potential relationships in subjects of other populations.

Disclosure of Potential Conflicts of Interest

P. Haycock reports grants from Cancer Research UK during the conduct of the study. M. Du reports grants from NCI (P30CA008748), Geoffrey Beene Foundation, Arnold and Arlene Goldstein Family Foundation, and Society of Memorial Sloan Kettering Cancer Center during the conduct of the study. G.G. Giles reports grants from National Health and Medical Research Council (paid to institution, Cancer Council Victoria) during the conduct of the study. L. Le Marchand reports grants from NCI during the conduct of the study. R.E. Neale reports grants from National Health and Medical Research Council during the conduct of the study. J.E. Buring reports grants from NIH during the conduct of the study. C.S. Fuchs reports personal fees from Agios, Amylin Pharmaceuticals, Bain Capital, CytomX Therapeutics, Daiichi Sankyo, Eli Lilly, Entrinsic Health, EvolveImmune Therapeutics, Genentech, Merck, Taiho, and Unum Therapeutics outside the

Table 1. Genetic instruments for plasma phospholipid levels of n-6 PUFAs (% of total fatty acids) that were genome-wide significant ($P < 5 \times 10^{-8}$) in previous GWASs.

Chr	SNP	GRCh37/hg19 position	Allele ^a	EAF	β	SE	P	% VE ^b per allele	% VE per IV ^c	F-statistic per IV ^d
Linoleic acid (18:2n6)										
10	rs10740118	65101207	G/C	0.56	0.2484	0.0431	8.08×10^{-9}	0.2–0.7	9.4–25.1	452–1461
11	rs174547	61570783	C/T	0.32	1.4737	0.0417	4.98×10^{-274}	7.6–18.1		
11	rs2727270	61603237	T/C	0.44	0.69	0.07	2.60×10^{-21}	0.5–2.4		
16	rs16966952	15135943	A/G	0.31	0.3512	0.0439	1.23×10^{-15}	0.5–2.5		
16	rs2280018	15150833	A/C	0.38	0.38	0.05	3.60×10^{-14}	0.6–1.4		
Arachidonic acid (20:4n6)										
11	rs174547	61570783	T/C	0.68	1.6909	0.0253	3.30×10^{-971}	3.7–37.6	4.1–44	311–5708
11	rs102275	61557803	T/C	0.68	2.49	0.1	6.60×10^{-147}	0.3–5.8		
16	rs16966952	15135943	G/A	0.69	0.1989	0.0314	2.43×10^{-10}	0.1–0.6		
Adrenic acid (22:4n6)										
11	rs174547	61570783	T/C	0.67	0.0483	0.0019	6.26×10^{-140}	7.8–10.9	7.8–10.9	1844–2667
GLA (18:3n6)										
11	rs174547	61570783	T/C	0.67	0.0156	0.0009	2.29×10^{-72}	2.2–4.6	2.5–6.4	186–497
16	rs16966952	15135943	G/A	0.69	0.0061	0.0009	5.05×10^{-11}	0.3–1.8		
11	rs10899123 ^e	75501207	C/G	0.91	0.0055	0.0014	9.97×10^{-5}	NA		
DGLA (20:3n6)										
11	rs174547	61570783	C/T	0.33	0.3550	0.0136	2.63×10^{-151}	8.7–11.1	13.5–26.3	850–1944
11	rs968567	61595564	T/C	0.16	0.29	0.02	1.30×10^{-42}	1.4–7.9		
16	rs16966952	15135943	G/A	0.69	0.2204	0.013	7.55×10^{-65}	2.0–4.5		
16	rs2280018	15150833	C/A	0.61	0.16	0.02	4.50×10^{-25}	1.4–2.8		

Abbreviations: Chr, chromosome; EAF, effect allele frequency; IV, instrumental variable; VE, variation explained.

^aThe first listed allele represents the effect allele associated with an increased level of corresponding PUFA; the second allele represents the alternative allele.

^b% VE = $[2 \times \beta^2 \times \text{EAF} \times (1 - \text{EAF}) / \text{var}(\text{PUFA})] \times 100$, unless indicated in article, such as in Guan and colleagues (6).

^c% VE per IV = sum of the %VE per allele for each SNP included in the IV.

^dF-statistic is a measure of the strength of the genetic instrument and is calculated as follows: $[R^2 \times (n - 1 - k)] / [(1 - R^2) \times k]$, where R^2 = % VE, n = sample size, k = total number of instrumental variables.

^eGenetic variant, rs10899123, showed an association at $5 \times 10^{-8} < P < 0.05$ in the CHARGE studies (Guan and colleagues, 2014; ref. 6), although it showed a GWAS significant association in the study by Hu and colleagues (2016; ref. 9). It was included in the genetic instrument in sensitivity analyses while it did not significantly change the association. The analyses excluding it in the instrument are reported in **Table 2**.

Table 2. Associations between 1 SD increase in PUFA-specific wGRSs and pancreatic cancer risk in PanScan and PanC4 studies^a.

Subgroup	Cases/controls	Linoleic acid		Arachidonic acid		DGLA	
		OR (95% CI)	P	OR (95% CI)	P	OR (95% CI)	P
Overall^b	9,269/12,530	0.99 (0.97–1.01)	0.30	1.01 (1.00–1.02)	0.37	0.94 (0.87–1.01)	0.10
Overall^c	9,206/12,525	1.00 (0.98–1.02)	0.89	1.00 (0.99–1.01)	0.53	0.94 (0.87–1.02)	0.13
Data source^c							
PanScan1	1,746/1,812	1.03 (0.97–1.08)	0.31	0.99 (0.97–1.01)	0.35	0.97 (0.81–1.16)	0.72
PanScan2	1,768/1,841	0.99 (0.93–1.04)	0.59	1.01 (0.99–1.03)	0.47	0.97 (0.82–1.15)	0.73
PanScan3	1,528/5,080	0.99 (0.93–1.05)	0.66	1.01 (0.98–1.03)	0.56	0.91 (0.76–1.10)	0.34
PanC4	4,164/3,792	1.00 (0.96–1.03)	0.91	1.01 (0.99–1.02)	0.53	0.94 (0.83–1.06)	0.30
Overall^d	9,040/12,496	1.00 (0.97–1.03)	0.95	1.00 (0.99–1.02)	0.73	0.95 (0.87–1.03)	0.21
Age^e							
>70	3,494/3,385	1.02 (0.98–1.06)	0.42	1.00 (0.98–1.01)	0.68	1.02 (0.90–1.17)	0.73
50–70	3,917/6,916	0.99 (0.96–1.03)	0.74	1.01 (0.99–1.02)	0.55	0.91 (0.81–1.02)	0.11
≤50	1,795/2,224	0.98 (0.93–1.03)	0.42	1.01 (0.99–1.04)	0.32	0.90 (0.75–1.07)	0.24
Sex^f							
Male	4,985/7,801	1.00 (0.97–1.03)	0.94	1.00 (0.99–1.02)	0.66	0.97 (0.87–1.08)	0.55
Female	4,221/4,225	1.00 (0.97–1.04)	0.99	1.00 (0.99–1.02)	0.71	0.99 (0.95–1.04)	0.77

Abbreviation: CI, confidence interval.

^aResults for adrenic acid and GLA not shown; their associations were not significant, with relatively wide CIs.

^bORs and 95% CIs estimated using individual-level data without adjustment, and represent 1 SD increase in each PUFA-specific wGRS.

^cORs and 95% CIs estimated using individual-level data with adjustment of age (under 50, 50–60, 60–70, 70–80, and above 80), sex, and 10 or seven principal components for PanScan and PanC4 data, respectively, and represent 1 SD increase in each PUFA-specific wGRS.

^dORs and 95% CIs estimated using summary statistics data.

^eORs and 95% CIs estimated using individual-level data with adjustment of age, sex, and 10 or seven principal components for PanScan and PanC4 data, respectively, and represent 1 SD increase in each PUFA-specific wGRS.

^fORs and 95% CIs estimated using individual-level data with adjustment of age (under 50, 50–60, 60–70, 70–80, and above 80) and 10 or seven principal components for PanScan and PanC4 data, respectively, and represent 1 SD increase in each PUFA-specific wGRS.

submitted work; serves as a director for CytomX Therapeutics and owns unexercised stock options for CytomX Therapeutics and Entrinsic Health; is a cofounder of EvolveImmune Therapeutics and has equity in this private company; and has provided expert testimony for Amlyn Pharmaceuticals and Eli Lilly. I.-M. Lee reports grants from NIH during the conduct of the study. R.L. Milne reports grants from National Health and Medical Research Council during the conduct of the study. A.L. Oberg reports grants from NCI (P50 CA102701) during the conduct of the study. I.M. Thompson Jr reports grants from NCI, NIH (several grants in support of conduct and administration of SELECT and PCPT studies) during the conduct of the study. J. Wactawski-Wende reports grants from NIH/National Heart, Lung, and Blood Institute (funding for WHI) during the conduct of the study. A. Zeleniuch-Jacqotte reports grants from NIH/NCI during the conduct of the study. A.P. Klein reports grants from NCI during the conduct of the study. X.-O. Shu reports grants and personal fees from NCI (grant review) during the conduct of the study. L. Wu reports grants from NCI during the conduct of the study. No potential conflicts of interest were disclosed by the other authors.

Disclaimer

The authors assume full responsibility for analyses and interpretation of these data.

Authors' Contributions

D.H. Ghoneim: Formal analysis, writing–review and editing. **J. Zhu:** Formal analysis, investigation, writing–original draft. **W. Zheng:** Methodology, writing–review and editing. **J. Long:** Writing–review and editing. **H.J. Murff:** Writing–review and editing. **F. Ye:** Writing–review and editing. **V.W. Setiawan:** Writing–review and editing. **L.R. Wilkens:** Writing–review and editing. **N.K. Khankari:** Resources, writing–review and editing. **P. Haycock:** Methodology, writing–review and editing. **S.O. Antwi:** Writing–review and editing. **Y. Yang:** Writing–review and editing. **A.A. Arslan:** Resources, data curation. **L.E. Beane Freeman:** Resources, data curation. **P.M. Bracci:** Resources, data curation. **F. Canzian:** Resources, data curation. **M. Du:** Resources, data curation. **S. Gallinger:** Resources, data curation. **G.G. Giles:** Resources, data curation. **P.J. Goodman:** Resources, data curation. **C. Kooperberg:** Resources, data curation. **L. Le Marchand:** Resources, data curation, writing–review and editing. **R.E. Neale:** Resources, data curation, writing–review and editing.

G. Scelo: Resources, data curation. **K. Viswanathan:** Resources, data curation. **E. White:** Resources, data curation. **D. Albanes:** Resources, data curation. **P. Amiano:** Resources, data curation. **G. Andreotti:** Resources, data curation. **A. Babic:** Resources, data curation. **W.R. Bamlet:** Data curation. **S.I. Berndt:** Resources, data curation. **L.K. Brais:** Resources, data curation. **P. Brennan:** Resources, data curation. **B. Bueno-de-Mesquita:** Resources, data curation. **J.E. Buring:** Resources, data curation. **P.T. Campbell:** Resources, data curation. **K.G. Rabe:** Data curation. **S.J. Chanock:** Resources, data curation. **P. Duggal:** Resources, data curation. **C.S. Fuchs:** Resources, data curation. **J.M. Gaziano:** Resources, data curation. **M.G. Goggins:** Resources, data curation. **T. Hackert:** Resources, data curation. **M.M. Hassan:** Resources, data curation. **K.J. Helzlsouer:** Resources, data curation. **E.A. Holly:** Resources, data curation. **R.N. Hoover:** Resources, data curation. **V. Katzev:** Resources, data curation. **R.C. Kurtz:** Resources, data curation. **I.-M. Lee:** Resources, data curation. **N. Malats:** Resources, data curation. **R.L. Milne:** Resources, data curation, writing–review and editing. **N. Murphy:** Resources, data curation. **A.L. Oberg:** Data curation. **M. Porta:** Resources, data curation. **N. Rothman:** Resources, data curation. **H.D. Sesso:** Resources, data curation. **D.T. Silverman:** Resources, data curation. **I.M. Thompson Jr:** Resources, data curation. **J. Wactawski-Wende:** Resources, data curation. **X. Wang:** Data curation. **N. Wentzensen:** Resources, data curation. **H. Yu:** Resources, writing–review and editing. **A. Zeleniuch-Jacqotte:** Resources, data curation. **K. Yu:** Data curation. **B.M. Wolpin:** Resources, data curation. **E.J. Jacobs:** Resources, data curation. **E.J. Duell:** Resources, data curation. **H.A. Risch:** Resources, data curation, writing–review and editing. **G.M. Petersen:** Resources, data curation. **L.T. Amundadottir:** Data curation. **P. Kraft:** Resources, data curation. **A.P. Klein:** Resources, data curation. **R.Z. Stolzenberg-Solomon:** Resources, data curation. **X.-O. Shu:** Conceptualization, resources, supervision, investigation, writing–review and editing. **L. Wu:** Conceptualization, resources, data curation, formal analysis, supervision, funding acquisition, investigation, methodology, writing–original draft, project administration.

Acknowledgments

The authors are indebted to the research team and participants of the PanScan and PanC4 consortia participating studies for their contributions to this study. This study was supported by NCI grants K99 CA218892 and R00 CA218892. The Multiethnic Cohort was supported by grant U01 CA164973. The Women's

Health Initiative program was funded by the National Heart, Lung, and Blood Institute, NIH, and U.S. Department of Health and Human Services through contracts HHSN268201600018C, HHSN268201600001C, HHSN268201600002C, HHSN268201600003C, and HHSN268201600004C. The Connecticut Pancreas Cancer Study was supported, in part, by NCI-NIH grant 5R01CA098870 (to H.A. Risch). The cooperation of 30 Connecticut hospitals, including Stamford Hospital, in allowing patient access, is gratefully acknowledged. This study was approved by the State of Connecticut Department of Public Health Human Investigation Committee. Certain data used in this study were obtained from the Connecticut Tumor Registry in the Connecticut Department of Public Health. A detailed list

of acknowledgments for other PanScan/PanC4 participating studies is included elsewhere (Klein and colleagues; ref. 7).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received May 5, 2020; revised June 21, 2020; accepted September 18, 2020; published first September 23, 2020.

References

- Gong Z, Holly EA, Bracci PM. Survival in population-based pancreatic cancer patients: San Francisco Bay area, 1995–1999. *Am J Epidemiol* 2011;174:1373–81.
- Funahashi H, Satake M, Hasan S, Sawai H, Newman RA, Reber HA, et al. Opposing effects of n-6 and n-3 polyunsaturated fatty acids on pancreatic cancer growth. *Pancreas* 2008;36:353–62.
- Qin B, Xun P, He K. Fish or long-chain (n-3) PUFA intake is not associated with pancreatic cancer risk in a meta-analysis and systematic review. *J Nutr* 2012;142:1067–73.
- Burgess S, Butterworth A, Thompson SG. Mendelian randomization analysis with multiple genetic variants using summarized data. *Genet Epidemiol* 2013;37:658–65.
- Lemaitre RN, Tanaka T, Tang W, Manichaikul A, Foy M, Kabagambe EK, et al. Genetic loci associated with plasma phospholipid n-3 fatty acids: a meta-analysis of genome-wide association studies from the CHARGE Consortium. *PLoS Genet* 2011;7:e1002193.
- Guan W, Steffen BT, Lemaitre RN, Wu JHY, Tanaka T, Manichaikul A, et al. Genome-wide association study of plasma N6 polyunsaturated fatty acids within the Cohorts for Heart and Aging Research in Genomic Epidemiology Consortium. *Circ Cardiovasc Genet* 2014;7:321–31.
- Klein AP, Wolpin BM, Risch HA, Stolzenberg-Solomon RZ, Mocchi E, Zhang M, et al. Genome-wide meta-analysis identifies five new susceptibility loci for pancreatic cancer. *Nat Commun* 2018;9:556.
- Khankari NK, Banbury BL, Borges MC, Haycock P, Albanes D, Arndt V, et al. Mendelian randomization of circulating polyunsaturated fatty acids and colorectal cancer risk. *Cancer Epidemiol Biomarkers Prev* 2020;29:860–70.
- Hu Y, Li H, Lu L, Manichaikul A, Zhu J, Chen YD, et al. Genome-wide meta-analyses identify novel loci associated with n-3 and n-6 polyunsaturated fatty acid levels in Chinese and European-ancestry populations. *Hum Mol Genet* 2016;25:1215–24.