

dietary intervention, even though glyce- mic control improved (HbA_{1c} $8.1 \pm 1.9\%$ to $7.2 \pm 1.3\%$, $P < 0.01$), no significant change was noticed in QTc (415.7 ± 19 to 415.1 ± 16.3 ms, $P = NS$) or QTd (25 ± 7.5 to 23.5 ± 6.7 ms, $P = NS$). Individual changes in QTc and QTd were not significantly correlated with corresponding initial levels or change in HbA_{1c} . The subset of patients who had prolonged QTc or microalbuminuria at baseline did not show statistically significant differences postintervention. There was no correlation between HbA_{1c} or fasting blood glucose and QTc or QTd before or after intervention. However, a significant correlation was observed between QTc and QTd before ($r = 0.590$, $P < 0.05$) and after ($r = 0.405$, $P < 0.05$) intervention.

It is surprising that despite the conservative criterion for the diagnosis of increased QTd (many previous studies have considered QTd >80 ms as abnormally prolonged), no patient was found to have prolonged QTd. This contrasts with other studies that have reported a high prevalence of QT abnormalities in diabetes (5).

In conclusion, this is the first prospective interventional study in a well-defined cohort of newly diagnosed type 2 diabetic patients to demonstrate that dietary treatment does not influence QT abnormalities in the short term. This study also shows that the prevalence of abnormal QTc and QTd in type 2 diabetes may not be as high as previously thought.

RAJEEV KUMAR, MRCP¹
MILES FISHER, MD¹
RHINNANON WHITAKER, BSC²
PETER W. MACFARLANE, FRCP³

From the ¹Department of Diabetes, Glasgow Royal Infirmary, Glasgow, Scotland, U.K.; the ²Institute of Medical and Social Care Research, University of Wales, Bangor, U.K.; and the ³Division of Cardiovascular and Medical Sciences, Section of Cardiology, Royal Infirmary, University of Glasgow, Glasgow, Scotland, U.K.

Address correspondence to Dr. Rajeev Kumar, Department of Diabetes and Endocrinology, Queen Elizabeth II Hospital, Welwyn Garden City, AL7 4HQ, U.K. E-mail: rajeevrs@hotmail.com.

© 2004 by the American Diabetes Association.

Acknowledgments—We are thankful to E. Cameron (diabetes specialist nurse), Bernie Quinn (dietitian), Jean Watts and Kathryn Macclaren (ECG technicians), and Julie Kennedy (ECG coder) for all their assistance.

References

1. Veglio M, Sivieri R, Chinaglia A, Scaglione L, Cavallo-Perin P, for the Neuropathy Study Group of the Italian Society of the Study of Diabetes, Piemonte Affiliate: QT interval prolongation and mortality in type 1 diabetic patients: a 5-year cohort prospective study. *Diabetes Care* 23: 1381–1383, 2000
2. Naas AA, Davidson NC, Thompson C, Cummings F, Ogston SA, Jung RT, Newton RW, Struthers AD: QT and QTc dispersion are accurate predictors of cardiac death in newly diagnosed non-insulin-dependent diabetes: a cohort study. *BMJ* 316:745–746, 1998
3. Willems JL, Arnaud P, van Bommel JH, Degani R, Macfarlane PW, Zywiec C, for the CSE Working Party: Common standards for quantitative electrocardiography: goals and main results. *Meth Inform Med* 29:263–271, 1990
4. Macfarlane PW, Lawrie TV: The normal electrocardiogram and vectorcardiogram. In *Comprehensive Electrocardiology*. Macfarlane PW, Lawrie TV, Eds. Oxford, U.K., Pergamon, 1989, p 407–457
5. Veglio M, Bruno G, Borra M, Macchia G, Barger G, D'Errico N, Pagano GF, Cavallo-Perin P: Prevalence of increased QT interval duration and dispersion in type 2 diabetic patients and its relationship with coronary heart disease: a population-based cohort. *J Intern Med* 25:317–324, 2002

Impact of Metformin on Glucose Metabolism in Nondiabetic, Obese African Americans

A placebo-controlled, 24-month randomized study

Pharmacological therapy using anti-diabetic drugs can delay the onset of type 2 diabetes in individuals with impaired glucose tolerance (IGT) (1–3). These drugs lower blood glucose, but each drug class impacts glucose dysregulation by a different mechanism (4). It is unclear whether simply lowering mean blood glucose can reduce the risk of type 2 diabetes or whether it is the drug effect on particular targets of glucose dysregulation or other unknown drug effects that

allow for the decreased incidence of type 2 diabetes in IGT.

Metformin decreases the incidence of type 2 diabetes in IGT (3). But whether metformin treatment earlier in the disease process might further delay progression to IGT/type 2 diabetes is unknown. We performed a double-blind placebo-controlled study of metformin in first-degree relatives of African Americans with type 2 diabetes for 24 months. The aim of the present study is to examine the effect of metformin on glucose metabolism in normal glucose tolerant (NGT) African Americans at risk for type 2 diabetes.

The study included 126 NGT African Americans. The MET group ($n = 45$) received 500 mg/day metformin and the PLA group ($n = 81$) received placebo for 24 months. Yearly oral glucose tolerance tests and frequently sampled intravenous glucose tolerance tests were performed.

Baseline clinical characteristics were similar between groups. There were no sex differences. The MET group demonstrated a reduction in weight (-1.4 ± 1 kg) and BMI (-0.53 ± 0.4 kg/m²). Weight (1.4 ± 0.8 kg, $P < 0.02$) and BMI (0.5 ± 0.34 kg/m², $P < 0.01$) increased in PLA. Following metformin therapy, acute first- and second-phase glucose, insulin, and C-peptide area under the curve did not change. However, hepatic insulin extraction (HIE) was increased 21% in the MET group vs. 5% in the PLA group.

As increased hepatic glucose production and decreased HIE is commonly seen as a component of glucose dysregulation in type 2 diabetes and other high-risk populations (5–7), improvement in HIE could potentially lead to overall improvement in glucose metabolism. In this study, the MET group was associated with modest weight reduction and enhanced HIE. The reason for the improved HIE is unclear but may reflect a mechanistic change at the level of the liver in response to metformin. Our findings of increased HIE could partly be responsible for the hepatic effect on glucose regulation previously reported (4). This finding is of great interest, demonstrating a “resetting” of the hepatic response to insulin resistance and a return to a more physiologic glucose regulation with even small doses of metformin in nondiabetic at-risk individuals.

Although other glucose-lowering drugs have been found to both improve (8) and worsen (9) HIE, to the best of our knowledge, the current findings repre-

sent a unique response not previously reported with metformin usage in an NGT population. Whether this restoration of HIE translates to a reduced occurrence of IGT/type 2 diabetes in our at-risk African-American population is not known. The drug was well tolerated for the 2-year duration without significant adverse events. Further studies are needed to determine the optimal dose of metformin for maximum benefit on early abnormalities in glucose dysregulation while minimizing side effects. Based on our current data, the treatment of at-risk African Americans with metformin is safe and may be beneficial in preventing conversion to IGT/type 2 diabetes.

DARA SCHUSTER, MD
TRUDY GAILLARD, RN
SCOTT RHINESMITH, MD
DIANE HABASH, PHD
KUAME OSEI, MD

From the Division of Endocrinology, Diabetes and Metabolism, Department of Internal Medicine, Ohio State University College of Medicine and Public Health, Columbus, Ohio.

Address correspondence to Dara Schuster, MD, 491 McCampbell Hall, 1581 Dodd Dr., Columbus, OH 43210. E-mail: schuster.26@osu.edu.

© 2004 by the American Diabetes Association.

References

- Chiaesson J-L, Josse RG, Gomis R, Hanefeld M, Karasik A, Laakso M: Acarbose for prevention of type 2 diabetes mellitus: the STOP-NIDDM randomized trial. *Lancet* 359:2072-2077, 2002
- Buchanan TA: Pancreatic B-cell defects in gestational diabetes: implications for the pathogenesis and prevention of type 2 diabetes. *J Clin Endocrinol Metab* 86:989-993, 2001
- Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, Nathan DM, for the Diabetes Prevention Program Research Group: Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 346:393-403, 2002
- Bailey CJ: Biguanides and NIDDM. *Diabetes Care* 15:755-772, 1992
- Osei K, Schuster DP: Ethnic differences in secretion, sensitivity and hepatic extraction of insulin in black and white Americans. *Diabet Med* 11:755-762, 1994
- Boyko EJ, Keane EM, Marshall JA, Hamman RF: Higher insulin and C-peptide concentrations in Hispanic population at high risk for NIDDM: San Luis Valley Diabetes Study. *Diabetes* 40:509-515, 1991
- Morin-Papunen LC, Vauhkonen I, Koivunen RM, Ruokonen A, Tapanainen JS: Insulin sensitivity, insulin secretion and metabolic and hormonal parameters in healthy women and women with polycystic ovarian syndrome. *Hum Reprod* 15:1266-1274, 2000
- Miyazaki Y, Matsuda M, DeFronzo RA: Dose-response effect of pioglitazone on insulin sensitivity and insulin secretion in type 2 diabetes. *Diabetes Care* 25:517-523, 2002
- Osei K, Rinesmith S, Gaillard T, Schuster DP: Metabolic effects of chronic glipizide gastrointestinal therapeutic system on serum glucose, insulin secretion, insulin sensitivity and hepatic insulin extraction in glucose-tolerant, first-degree relatives of African American patients with type 2 diabetes: new insights on mechanisms of action. *Metabolism* 52:565-572, 2003

Elevated C-Reactive Protein Levels Do Not Correspond to Autoimmunity in Type 1 Diabetes

A large body of data now supports that high normal or elevated C-reactive protein (CRP) levels are a marker of ongoing low-grade chronic inflammatory process and may predict a higher risk for cardiovascular disease. Type 1 diabetes is a T-cell-mediated disease (1), and autoantibodies (2) are considered to be the surrogate markers for the autoimmune process and have been used for disease prediction (3). However, little is known about the relationship between diabetes autoimmunity and the acute-phase response. We studied the relationship of islet cell autoimmunity in type 1 diabetes as reflected by the presence of GAD65 and ICA512 autoantibodies and CRP titers. We analyzed the following five groups of individuals: pre-diabetic ($n = 10$), new-onset ($n = 29$), long-term type 1 diabetic ($n = 16$), and long-term type 2 diabetic ($n = 55$) patients and healthy subjects ($n = 50$). CRP levels were significantly higher in long-term type 1 diabetic patients (0.11 mg/dl [range 0.014-1.39]) than in either new-onset type 1 diabetic patients (0.061 mg/dl [0.012-1.12], $P = 0.04$) or healthy control subjects (0.06 mg/dl [0.007-1.33], $P = 0.02$). Long-term type 2 diabetic patients have higher CRP levels (0.24 mg/dl [0.03-4.41]) than healthy control subjects ($P = 0.04$). We

could not find any correlation between CRP levels and any combination or titer of autoantibodies in pre-diabetic, new-onset, and long-term type 1 diabetic patients.

The lack of elevated CRP in the pre-diabetic and recently diagnosed patients with type 1 diabetes is consistent with the notion that chronic localized autoimmune inflammation does not result in appreciable changes in the CRP levels, thus making it unsuitable as a marker of type 1 diabetic autoimmunity. The difference in CRP levels in long-term and new-onset type 1 diabetic patients may be related in part to the vanish of the intrinsic physiologically secreted insulin once extending the honeymoon period in type 1 diabetes. Insulin has been shown to have anti-inflammatory properties, including downregulation of acute-phase protein production of the liver (4); thus, it may lead to lower CRP levels.

The underlying mechanisms behind elevated CRP levels in long-term type 1 and type 2 diabetic patients might be different, as elevated glucose levels as well as insulin resistance are contributing factors to the inflammatory process. Prolonged elevated blood glucose levels are implicated in chronic inflammatory changes in the tissues in both type 1 and type 2 diabetic patients and reflected by the elevated CRP values. However, in type 2 diabetes, insulin resistance is also likely to be a major factor (5), as elevated CRP values have been found in near-normoglycemic pre-diabetic type 2 diabetic and metabolic syndrome patients.

ANDRAS TRESZL, MD
LASZLO SZEREDAY, MD
ALESSANDRO DORIA, MD
GEORGE L. KING, MD
TIHAMER ORBAN, MD

From the Joslin Diabetes Center, Harvard Medical School, Boston Massachusetts.

Address correspondence to Tihamer Orban, MD, Section of Immunology and Immunogenetics, Joslin Diabetes Center, Harvard Medical School, One Joslin Place, Boston, MA 02215. E-mail: torban@joslin.harvard.edu.

© 2004 by the American Diabetes Association.

References

- Martin S, Wolf-Eichbaum D, Duinkerken G, Scherbaum WA, Kolb H, Noordzij JG, Roep BO: Development of type 1 diabetes despite severe hereditary B-lymphocyte deficiency. *N Engl J Med* 345:1036-1040,

- 2001
- Atkinson MA, Eisenbarth GS: Type 1 diabetes: new perspectives on disease pathogenesis and treatment. *Lancet* 358:221–229, 2001
 - Liu E, Eisenbarth GS: Type 1A diabetes mellitus-associated autoimmunity. *Endocrinol Metab Clin North Am* 31:391–410, 2002 [review vii–viii]
 - Dandona P, Aljada A, Mohanty P: The anti-inflammatory and potential anti-atherogenic effect of insulin: a new paradigm. *Diabetologia* 45:924–930, 2002
 - Haffner SM: Insulin resistance, inflammation, and the prediabetic state. *Am J Cardiol* 92:18J–26J, 2003

COMMENTS AND RESPONSES

Orlistat Augments Postprandial Increases in Glucagon-Like Peptide-1 in Obese Type 2 Diabetic Patients

Response to Damci et al.

Damci et al. (1) suggest that the lipase inhibitor, orlistat, stimulates the postprandial secretion of glucagon-like peptide-1 (GLP-1) after meals containing fat and speculate that this may lead to reductions in both postprandial glycemia and energy intake. The authors observed (in a cohort of 29 type 2 diabetic patients) that after ingestion of a 600 kcal breakfast (comprising 38% fat, 50% carbohydrate, and 12% protein), the increases from baseline in plasma GLP-1 and serum C-peptide were slightly greater and that of serum glucose slightly less, as measured in a single blood sample taken 60 min after commencement of the meal. The authors have apparently failed to appreciate the following: 1) While fat is a potent stimulant of GLP-1 secretion, the latter appears to be dependent on the digestion of fat to fatty acids (2,3), e.g., in healthy subjects, the stimulation of GLP-1 by intraduodenal triglyceride is abolished by concomitant administration of orlistat

(2). 2) As the regulation of gastric emptying of fat is also dependent on lipolysis (4,5), orlistat may accelerate gastric emptying of meals containing fat (4). If the meal also contains carbohydrate, orlistat has the capacity to exacerbate overall postprandial glycemic excursions in type 2 diabetic patients by this mechanism (3). Even relatively minor variations in gastric emptying of carbohydrate are now known to potentially have a major effect on postprandial glycemia and insulin responses in both healthy subjects and type 2 diabetic patients (6,7). 3) In healthy subjects, acute lipase inhibition with orlistat attenuates rather than increases the inhibitory effects of fat on subsequent energy intake (2,8).

The effects of orlistat on glycemic, insulin, and incretin responses to a meal, including the time course of these effects, are likely to be critically dependent on both the macronutrient composition and energy content of a meal, perhaps particularly the ratio of fat to carbohydrate. It should also be recognized that in patients with type 2 diabetes, gastric emptying is frequently abnormal and may influence these responses (7). In relation to the study by Damci et al. (1), we would offer an alternative interpretation, i.e., after administration of orlistat, the carbohydrate, fat, and protein components of their meal (the type of meal is not described) initially emptied from the stomach more rapidly, and this was reflected in a relative increase in C-peptide and GLP-1 at 60 min and a slight reduction in plasma glucose at that time. The stimulation of GLP-1 would reflect the greater small intestinal carbohydrate load during this time as well as the presence of fatty acids that escaped lipase inhibition. We would anticipate that the overall GLP-1 response to the meal would be less and initial postprandial glycemia worse. It would require more frequent blood sampling over a longer period of time to fully interpret the action of orlistat on incretin and insulin secretion and to draw conclusions about GLP-1 as a mediator of the effects of orlistat on body weight in obesity.

MICHAEL HOROWITZ, MB, BS, PHD¹
CHRISTINE FEINLE-BISSET, PHD¹
KAREN JONES, PHD¹
MICHAEL NAUCK, MD²

From the ¹Department of Medicine, University of Adelaide, Royal Adelaide Hospital, Adelaide, Aus-

tralia; and ²Diabeteszentrum, Bad Lauterberg im Harz, Germany.

Address correspondence to Professor Michael Horowitz, Department of Medicine, Royal Adelaide Hospital, North Terrace, Adelaide, South Australia, Australia, 5000. E-mail: michael.horowitz@adelaide.edu.au.

© 2004 by the American Diabetes Association.

References

- Damci T, Yalin S, Balci H, Oser Z, Korugan U, Ozyazar M, Illkova H: Orlistat augments postprandial increases in glucagon-like peptide-1 in obese type 2 diabetic patients. *Diabetes Care* 27:1077–1080, 2004
- Feinle C, O'Donovan D, Doran S, Andrews JM, Wishart J, Chapman I, Horowitz M: Effects of fat digestion on appetite, APD motility, and gut hormones in response to duodenal fat infusion in humans. *Am J Physiol Gastrointest Liver Physiol* 284:G798–G807, 2003
- Pilichiewicz A, O'Donovan D, Feinle C, Lei Y, Wishart JM, Bryant L, Meyer JH, Horowitz M, Jones KL: Effect of lipase inhibition on gastric emptying of, and the glycemic incretin responses to, an oil/ aqueous drink in type 2 diabetes mellitus. *J Clin Endocrinol Metab* 88:3829–3834, 2003
- Schwizer W, Asal K, Kreiss C, Mettraux C, Berovicka J, Remy B, Guzelhan C, Hartmann D, Fried M: Role of lipase in the regulation of upper gastrointestinal function in humans. *Am J Physiol* 273:G612–G620, 1995
- Carney BI, Jones KL, Horowitz M, Sun WM, Penagini R, Meyer JH: Gastric emptying of oil and aqueous meal components in pancreatic insufficiency: effects of posture and on appetite. *Am J Physiol* 268:G925–G932, 1995
- O'Donovan DG, Doran S, Feinle-Bisset C, Jones KL, Meyer JH, Wishart JM, Morris HA, Horowitz M: Effect of variations in small intestinal glucose delivery on plasma glucose, insulin and incretin hormones in healthy subjects and type 2 diabetes. *J Clin Endocrinol Metab* 89:3431–3435, 2004
- Rayner CK, Samsom M, Jones KL, Horowitz M: Relationships of upper gastrointestinal motor and sensory function with glycaemic control. *Diabetes Care* 24:371–381, 2001
- O'Donovan D, Feinle-Bisset C, Wishart J, Horowitz M: Lipase inhibition attenuates the acute inhibitory effects of oral fat on food intake in healthy subjects. *Br J Nutr* 90:849–852, 2003

Orlistat Augments Postprandial Increases in Glucagon-Like Peptide-1 in Obese Type 2 Diabetic Patients

Response to Horowitz et al.

We thank Horowitz et al. (1) for their comments on our article (2) on the effect of orlistat on postprandial glucose, insulin, and incretin levels. They raise criticisms regarding the increase in postprandial GLP-1 levels, asserting that it is a result of fat digestion rather than the fat content of the meal. They also state that to draw more definitive conclusions, more frequent blood sampling over a longer period of time is required rather than a single sample at 60 min after the mixed meal. Ours was an

observatory pilot study that tested the effect of this drug on postprandial incretin levels. Putative mechanisms of action will be addressed in further studies in which we will take these criticisms into account. Horowitz et al. also state that orlistat would be expected to worsen postprandial hyperglycemia since this drug accelerates gastric emptying. Apart from our study, orlistat was previously shown to ameliorate postprandial blood glucose levels in long-term studies (3).

TANER DAMCI, MD¹

SERAP YALIN, MD¹

HURIYE BALCI, PHD²

ZEYNEP OSAR, MD¹

USTUN KORUGAN, MD¹

MUCAHIT OZYAZAR, MD¹

HASAN ILKOVA, MD²

From the ¹Department of Internal Medicine, Division of Endocrinology, Diabetes and Metabolism, Istanbul University Cerrahpasa Medical School, Istanbul, Turkey; and the ²Central Laboratories, Istanbul University Cerrahpasa Medical School, Istanbul, Turkey.

Address correspondence to Taner Damci, MD,

Atakoy 4, Kisim O-67 D:7, 34750 Atakoy, Istanbul, Turkey. E-mail: tdamci@superonline.com.

T.D. is an advisory board member for Aventis and has received honoraria from Roche, Aventis, and Pfizer. H.I. is an advisory board member for Aventis and has received honoraria from Roche, Novo Nordisk, Pfizer, and GlaxoSmithKline.

© 2004 by the American Diabetes Association.

References

1. Horowitz M, Feinle-Bisset C, Jones K, Nauck M: Orlistat augments postprandial increases in glucagon-like peptide-1 in obese type 2 diabetic patients (Letter). *Diabetes Care* 27:2770, 2004
2. Damci T, Yalin S, Balci H, Osar Z, Korugan U, Ozyazar M, Ilkova H: Orlistat augments postprandial increases in glucagon-like peptide 1 in obese type 2 diabetic patients. *Diabetes Care* 27:1077–1080, 2004
3. Holander PA, Elbein SC, Hirsch IB, Kelley D, McGill J, Taylor T: Role of orlistat in the treatment of obese patients with type 2 diabetes: a 1-year randomized double-blind study. *Diabetes Care* 21:1288–1294, 1998