

Neurosensory effects of chronic exposure to arsenic via drinking water in Inner Mongolia: I. signs, symptoms and pinprick testing

Yanhong Li, Yajuan Xia, Linlin He, Zhixiong Ning, Kegong Wu, Baixiao Zhao, X. Chris Le, Richard Kwok, Michael Schmitt, Timothy Wade, Judy Mumford and David Otto

ABSTRACT

This study was designed to assess the effects of exposure to arsenic in drinking water on neurosensory function. A questionnaire including neurological signs and symptoms and a brief neurological exam consisting of pinprick testing of the arms and legs and knee-jerk test were administered to 321 residents of the Bamen region of Inner Mongolia, China. Arsenic in water was measured by hydride generation atomic fluorescence. Participants were divided into three exposure groups—low (non-detectable–20), medium (100–300) and high (400–700 $\mu\text{g}/\text{l}$) arsenic. Significant group differences were observed in pinprick scores for all four limbs. Results indicate that arsenic alters pinprick (pain) thresholds at well-water concentrations as low as 400 $\mu\text{g}/\text{l}$, well below the 1000 $\mu\text{g}/\text{l}$ threshold for neurological effect specified by NRC (1999). Regression models suggest that a 50% increase in pinprick score is associated with a 71–159 ppb increase in arsenic concentration.

Key words | arsenic, drinking water, pain, pinprick, symptoms

Yanhong Li
Yajuan Xia
Kegong Wu
Inner Mongolia Center For Endemic Disease
Control and Research, Huhhot, China

Linlin He
Zhixiong Ning
Bamen Anti-Epidemic Station, Lin He, Inner
Mongolia, China

Baixiao Zhao
Lin He Anti-Epidemic Station, Lin He, Inner
Mongolia, China

X. Chris Le
University of Alberta, Edmonton,
Alberta, Canada

Richard Kwok
RTI International, Research Triangle Park, NC. USA

Michael Schmitt
Timothy Wade
Judy Mumford
David Otto (corresponding author)
U.S. Environmental Protection Agency,
Research Triangle Park, NC. USA
Tel: +1 (919) 966-6226
Fax: +1 (919) 966-6367
E-mail: otto.david@epa.gov

INTRODUCTION

Arsenic occurs naturally in soil and ground water in many parts of the world from the weathering of bedrock or other materials in the crust of the earth (Thornton & Farago 1997). Alluvial and organic soils contain more arsenic than sandy soils (Kabata-Pendias & Pendias 1984). The present study was carried out in the Bayingnormen (Bamen) region of Inner Mongolia. Bamen is a major agricultural area—the melon and sunflower capital of China—located between ancient mountain ranges (Langsan and Da Qingsan) to the north and the Yellow River to the south. Arsenic has accumulated in the soil of the Bamen region over the millennia from erosion of the mountains and periodic flooding of the Yellow River.

Arsenic concentrations found in the underground lake basin, between the mountains and river, range from nondetectable to 1,800 $\mu\text{g}/\text{l}$ (Ma *et al.* 1999). Arsenicism has been known to occur in this area since 1990.

Sensory axonal peripheral neuropathy is a common sequela of acute arsenic poisoning, reported to occur at doses as low as 40–50 $\mu\text{g}/\text{kg}$ (ATSDR 1998; NRC 1999). However, there is considerable uncertainty about the threshold for neurological effects of chronic arsenic exposure in drinking water. Based on a review of data from Taiwan, Argentina and Chile by Hotta (1989) and other available findings, the National Research Council (1999) concluded that there is

doi: 10.2166/wh.2005.022

no consistent evidence of peripheral neuropathy in humans exposed to arsenic in drinking water at levels below 1000 $\mu\text{g/l}$. Similarly, Kreiss *et al.* (1983) failed to find any evidence of arsenic-induced neuropathy in 147 residents of Ester Dome, Alaska who were drinking well water with arsenic levels up to 15,000 $\mu\text{g/l}$. Clinical neurological examinations and nerve-conduction velocity (NCV) tests were administered under blinded conditions in this careful study. However, the authors concluded that NCV was an insensitive screening measure for subclinical arsenic neuropathy.

Adverse neurological effects have been reported in other studies at arsenic levels below 1000 $\mu\text{g/l}$. Hindmarsh *et al.* (1977) found abnormal electromyograms in 32 of 110 Canadians with well water concentrations of 60–1400 $\mu\text{g/l}$. Basu *et al.* (1996) observed sensory polyneuropathy in 8 West Bengali patients chronically exposed to drinking water arsenic levels of 200–2000 $\mu\text{g/l}$. A variety of neurotoxic effects have also been reported in a study (Ma *et al.* 1995) of Inner Mongolian farmers whose drinking water was contaminated by arsenic with concentrations ranging from nondetectable – 1,800 $\mu\text{g/l}$. These effects included hearing loss, atrophy of the optic nerve, narrowing of the visual field and peripheral neuropathy.

A variety of health problems including skin cancer, peripheral neuropathy, gastroenteritis and cardiovascular disease have been associated with arsenic exposure in Bamen residents (Ma *et al.* 1995). The present study was undertaken to evaluate the effects of exposure to arsenic in drinking water on sensory signs and symptoms assessed by questionnaire and on pain sensation measured by pinprick testing. Tests of visual and tactile function were also administered and are reported elsewhere in this volume (Otto *et al.* 2006 (this issue)).

METHODS

Subjects

321 residents from the Bamen region of Inner Mongolia participated in this study. Three exposure groups were defined *a priori* as low (0–20), medium (100–300) and high (400–700 $\mu\text{g/l}$) arsenic in drinking water. A database containing arsenic concentrations for more than 10,000

wells in areas of chronic arsenicosis was available from local anti-epidemic stations. This database was used to identify areas which would yield a wide range of exposure (0–800 $\mu\text{g/l}$). Water samples were then obtained from 363 households. Participants were recruited from this sample to obtain the three exposure groups. Participants were not informed of water arsenic levels prior to neurosensory testing.

Demographic characteristics of these groups are shown in Table 1. Few of the participants had a high school education and none had been to college. The mean age of participants was 34.7 years. 12 subjects were eliminated from statistical analyses due to missing well water data.

Protection of human subjects

This study was conducted according to the Declaration of Helsinki recommendations for the protection of human subjects (World Medical Association 1989). Informed consent was obtained from all subjects to participate in the study. Test procedures and potential risks were described to subjects in the consent form. The protocol was reviewed and approved by the U.S. Environmental Protection Agency.

Assessment of arsenic

Total arsenic in water samples obtained from the wells of individual families or community water sources was assessed by hydride generation atomic fluorescence (Le & Ma 1998). Distribution of arsenic concentrations in the three groups is shown in Figure 1.

Questionnaire

A questionnaire, designed for this study, was administered to obtain demographic information, health and exposure history of participants, and self-reported neurosensory signs and symptoms. Questions included age, gender, education, smoking and drinking history, pesticide exposure, ratings of neurological function and history of central or peripheral nervous system disorders. Neurological symptom items are shown in Table 2. Education level was classified as none, primary, middle school or high school. Drinking

Table 1 | Distribution of Demographic Data by Arsenic Exposure Group

| | Exposure group (arsenic in drinking water) | | |
|--|--|--------------------|------------------|
| | Low Mean(SD) | Medium Mean(SD) | High Mean(SD) |
| Mean As ($\mu\text{g/l}$) | 12.4 (5.7) | 200.9 (55.6) | 567.3 (70.7) |
| Mean age | 35.8 (12.8) | 35.7 (14.0) | 33.7 (13.9) |
| Sample size | N | N | N |
| | 97 | 109 | 103 |
| | N (%) | N (%) | N (%) |
| Gender (Male) | 49 (51%) | 59 (54%) | 58 (56%) |
| Education | | | |
| None | 14 (15%) | 21 (20%) | 19 (19%) |
| Primary | 38 (41%) | 51 (48%) | 43 (42%) |
| Middle | 35 (38%) | 26 (25%) | 36 (35%) |
| High | 5 (5%) | 8 (8%) | 4 (4%) |
| Smoke | | | |
| Yes | 25 (25%) | 40 (41%) | 32 (33%) |
| Drink Alcohol | | | |
| Yes | 20 (25%) | 18 (17%) | 9 (12%) |
| Exposure to pesticides more than 5 years ago | | | |
| Yes | 50 (56%) | 56 (53%) | 41 (42%) |
| Exposure to pesticides within past 5 years | | | |
| Yes | 55 (62%) | 57 (54%) | 48 (49%) |

history was measured simply as ‘Do you drink alcohol more than twice a week (Y/N)?’ Information was obtained on the type and amount of tobacco used and age when smoking commenced. Pack years were calculated as (years smoked) \times (# cigarettes per day)/20 \times 365. Participants were also questioned whether they had been exposed to pesticides during the past five years (Y/N) or prior to that time (Y/N). If subjects were unable to read, questions were administered by interview. The primary purpose of this questionnaire was to identify possible confounders.

Brief neurological examination

A brief neurological examination consisting of pinprick testing of the left and right upper and lower extremities and the knee-jerk reflex was conducted by physicians with extensive clinical experience. The pinprick test was administered from distal-to-proximal positions of the hands and feet in order to determine the extent of pain detection (ability to feel sensation from a blunt needle). Participants were instructed to close both eyes and to extend the right

hand (or foot) for examination. Testing began at the end of the finger (or toe) and continued toward the elbow (or knee) until the subject said (s)he could feel pain. The procedure was then repeated with the left hand and foot.




Figure 1 | Distribution of arsenic concentrations in well water of participants. Bin width on x-axis is 50 $\mu\text{g/l}$; values below x-axis correspond to midpoint of bin. The distribution reflects gaps between low, medium and high exposure groups.

Table 2 | Number and percent of self-reported neurological signs and symptoms

| | Exposure group (arsenic in drinking water) | | |
|-----------------------------------|--|------------|------------|
| | Low | Medium | High |
| Amnesia | N(%) | N(%) | N(%) |
| Yes | 16 (18%) | 24 (23%) | 28 (29%) |
| Impaired autonomic nervous system | | | |
| Yes | 16 (18%)* | 8 (8%)*† | 28 (29%)† |
| Headache | | | |
| Yes | 24 (27%) | 34 (33%) | 27 (28%) |
| Hearing loss | | | |
| Yes | 14 (16%) | 5 (5%) | 8 (8%) |
| Impaired heat/cold sensation | | | |
| Yes | 18 (20%)* | 4 (4%)*† | 22 (22%)*† |
| Numbness | | | |
| Yes | 22 (25%)* | 14 (14%)*† | 29 (30%)*† |
| Pain | | | |
| Yes | 7 (8%)‡ | 3 (3%)† | 33 (34%)*† |
| Impaired sense of smell | | | |
| Yes | 9 (10%)* | 2 (2%)*† | 12 (12%)*† |
| Impaired vibration sensation | | | |
| Yes | 23 (26%)*‡ | 17 (16%)*† | 42 (43%)*† |

*p < 0.05, low vs. medium, (χ^2 -test, 1 degree of freedom)

†p < 0.05, medium vs. high, (χ^2 -test, 1 degree of freedom)

‡p < 0.05, low vs. high, (χ^2 -test, 1 degree of freedom)

Neither the subject nor the examiner was aware of the well-water arsenic level or exposure group at the time of testing, in other words, the test was administered double-blind. Testing with eyes closed also prevented participants from knowing the location of the needle during examination.

Clinical tests were scored as follows: absence of pinprick sensation in upper extremities (1 = none, 2 = end of finger only, 3 = whole finger, 4 = below wrist, 5 = below elbow); absence of pinprick sensation in lower extremities (1 = none, 2 = end of toe, 3 = whole toe, 4 = below ankle, 5 = below knee); and knee jerk reflex (1 = normal, 2 = low, 3 = high). Testing was conducted in medical facilities located in the villages where participants lived.

Statistical analyses

Statistical analyses were conducted in SAS v. 8.0 (2001, SAS Institute, Inc.) and Stata v. 8.2 (2004, Stata Corporation). Chi square tests were used to explore the association of categorical dependent and independent variables. Mean group differences were tested using analysis of variance

(ANOVA). Because the outcome measure was an ordered categorical variable, multivariate models were also performed with ordered logistic regression (McCullagh 1980). Dummy variables were used for categories of arsenic exposure with low exposure as the referent group and modeling arsenic level as a continuous variable. Variables which were significantly related to arsenic exposure or neurological test results were included in the multivariate analysis. P-values < 0.05 (two-sided) were considered to be statistically significant; and p-values < 0.1 (two-sided) to be of borderline statistical significance.

RESULTS

Demographic data broken down by exposure group are shown in Table 1. No demographic variables were significantly ($p < 0.05$) associated with exposure category by χ^2 tests for association, although there was a tendency for alcohol consumption ($p = 0.12$) and pesticide exposure ($p = 0.11$) to be related to arsenic exposure. Table 2 shows the number and percent of respondents by exposure category for neurological signs and symptoms. Self-reported autonomic nervous system dysfunction, heat sensitivity, numbness, pain, impaired smell, and reduced sensitivity to vibration were significantly associated with arsenic exposure category. Although the highest arsenic category generally had the highest proportion reporting the condition, the medium arsenic category, and not the low arsenic category, generally had the lowest proportion reporting the condition.

Less than 20% of participants had reduced sensitivity on any of the pinprick or knee-jerk measures. Since the distributions of pinprick and knee-jerk ratings were highly skewed, ratings were scored as either normal (= 1) or abnormal (> 1). The χ^2 -test was then used to assess the percentage of participants with abnormal pinprick or knee-jerk ratings for each survey or demographic question. Table 3 depicts the association of pinprick score (score > 1) and questionnaire responses. Symptoms associated with the highest percentage of abnormal pinprick response ($p < .005$) were numbness, pain and vibration sensation. Other items (amnesia and autonomic nervous system dysfunction) were also significantly associated with

Table 3 | Percent with higher scores (> 1) in pinprick and knee-jerk tests by demographic characteristics and self-reported neurological signs and symptoms^a

| | | Pinprick test | | | | | Kneejerk |
|---|--------|---------------|------------------|------------------|------------------|------------------|----------|
| | | N | L. Arm | R. Arm | L. Leg | R. Leg | |
| Gender- | Male | 163 | 16% | 16% | 8% | 6%* | 1% |
| | Female | 143 | 19% | 19% | 11% | 11% | 4% |
| Drink alcohol- | Yes | 46 | 15% | 15% | 9% | 7% | 2% |
| | No | 213 | 17% | 17% | 10% | 9% | 2% |
| Education (middle school- or above) | Yes | 112 | 15% | 14% | 5%* | 4% [†] | 2% |
| | No | 176 | 20% | 20% | 12% | 11% | 3% |
| Smoke cigarettes- | Yes | 93 | 14% | 14% | 7% | 4%* | 4% |
| | No | 197 | 20% | 20% | 11% | 11% | 2% |
| Amnesia- | Yes | 68 | 31% [‡] | 31% [‡] | 16% [†] | 16% [†] | 2% |
| | No | 221 | 14% | 14% | 8% | 8% | 3% |
| Autonomic Nervous System impairment | Yes | 52 | 29% [†] | 33% [‡] | 23% [‡] | 17% [†] | 2% |
| | No | 237 | 16% | 15% | 7% | 7% | 3% |
| Headache- | Yes | 85 | 20% | 21% | 12% | 12% | 4% |
| | No | 204 | 18% | 17% | 9% | 7% | 2% |
| Hearing impairment- | Yes | 27 | 26% | 22% | 19% | 15% | 0% |
| | No | 262 | 18% | 18% | 9% | 8% | 3% |
| Numbness- | Yes | 65 | 35%¶ | 37%¶ | 20%¶ | 19%¶ | 2% |
| | No | 222 | 14% | 13% | 7% | 6% | 3% |
| Pain- | Yes | 43 | 93%¶ | 88%¶ | 49%¶ | 42%¶ | 5% |
| | No | 245 | 5% | 6% | 3% | 3% | 2% |
| Pesticide exposure more than 5 years ago- | Yes | 145 | 17% | 17% | 11% | 8% | 4%* |
| | No | 145 | 19% | 20% | 8% | 9% | 1% |
| Pesticide exposure within 5 years- | Yes | 158 | 17% | 16% | 11% | 8% | 4%* |
| | No | 132 | 20% | 20% | 8% | 9% | 1% |
| Impaired sense of smell | Yes | 22 | 32%* | 27% | 23% [†] | 14% | 0% |
| | No | 267 | 17% | 18% | 9% | 8% | 3% |
| Vibration sensation | Yes | 80 | 34%¶ | 35%¶ | 23%¶ | 19%¶ | 1% |
| | No | 209 | 12% | 12% | 5% | 5% | 3% |

*p < 0.1, [†]p < 0.05, [‡]p < 0.01, ¶p < 0.005 (χ^2 -test, 1 degree of freedom)

^anumbers do not add to 309 due to missing responses

pinprick ratings ($p < .01$ or $.05$). Several demographic variables (smoking, education level and gender) were weakly associated with a few pinprick measures.

Figure 2 illustrates pinprick scores by exposure group for each limb. ANOVA tests indicated significant differences in pinprick scores of both arms and legs, but not in knee-jerk reflex ratings. Pinprick tests of all four limbs indicate a trend for the absence of pinprick sensation to increase in a proximal direction with increasing arsenic concentrations in drinking water. Significant group differ-

ences in pinprick scores were observed between low and high, medium and high, but not between low and medium exposure groups. The largest differences in pinprick score were found between low and high exposure groups.

In view of the categorical nature of the dependent variables, categorical multivariate regression analyses were also performed on pinprick scores. Results are shown in Table 4. Knee-jerk response was unrelated to arsenic and was not included in the regression analyses. Significant increases in pinprick score were found for all pinprick measures




Figure 2 | Comparison of mean pinprick test scores for low, medium and high exposure groups in four limbs tested.

comparing high to low arsenic categories, controlling for age, gender, education, smoking and drinking. Increases in pinprick scores were also noted comparing medium to low categories, but none of the tests were statistically significant. Participants in high arsenic groups had predicted probabilities of reduced response ranging from 0.22–0.33. Although more respondents reported increased pinprick score in the hands, regression coefficients were stronger for leg measures. Coefficients are interpreted as the increase in log odds of a one-unit increase in pinprick score for a change from low-to-medium or low-to-high arsenic categories. Exponentiating

the coefficients (Selvin 1996) provides an estimate of the relative risk of a one unit increase in pinprick score associated with an increase in arsenic category. For example, those in the high arsenic group were more than five times as likely ($\exp[1.70] = 5.47$) to have a left arm pinprick score one unit higher than those in the lowest category. Those in the high arsenic category were almost 20 times more likely to have a left leg pinprick score one unit higher than the lowest arsenic exposure category.

Finally, a multivariate regression model was run to determine the quantity of arsenic in drinking water required to increase pinprick score by 50% or 100%. Results from modeling arsenic as a continuous variable are shown in Table 5. A 50% reduction would be equivalent to an increase in pinprick score from 2.0 to 3.0; a 100% increase in score would be equivalent to an increase from 1 to 2 or 2 to 4. Regression coefficients were again stronger in leg compared to arm measures. A 50% increase in pinprick score was associated with a 71–159 ppb increase in arsenic concentration and a 100% increase was associated with a 121–171 ppb increase.

DISCUSSION

The pinprick test is widely used in clinical neurology to assess the integrity of pain and/or touch sensation in

Table 4 | Categorical multivariate analysis^a: mean predicted probability of elevated pinprick score (>1) and regression coefficients^b

| Extremity | Measure | Arsenic exposure category | | |
|------------------------|--|---------------------------|-------------------|---------------|
| | | Low | Medium | High |
| Left Arm | Coefficient (se) | Referent | 0.35 (0.5) | 1.70 (0.47)** |
| | Predicted Probability of Elevated Pinprick Score | 0.09 | 0.12 | 0.33 |
| Right Arm | Coefficient (se) | Referent | 0.62 (0.48) | 1.51 (0.47)** |
| | Predicted Probability of Elevated Pinprick Score | 0.09 | 0.14 | 0.29 |
| Left Leg | Coefficient (se) | Referent | 1.41 (0.85)* | 2.96 (0.82)** |
| | Predicted Probability of Elevated Pinprick Score | 0.02 | 0.07 | 0.24 |
| Right Leg ^c | Coefficient (se) | Referent | NA ^c | 2.16 (0.51)** |
| | Predicted Probability of Elevated Pinprick Score | | 0.04 ^d | 0.22 |

^aOrdered logistic regression, outcome is pinprick score coded 1 through five, controlling for age, gender, smoking, drinking and education. Dummy variables were used for arsenic exposure category, with low exposure as the referent group

^bCoefficients are interpreted as the increase in log odds of a one-unit increase in pinprick score for a change from low to medium or low to high arsenic categories

^cNo reduced pinprick response events in low arsenic category, low and medium were combined

^dEstimate for combined low and medium arsenic exposure categories

* $p < 0.1$, ** $p < 0.005$

Table 5 | Multivariate Regression Model^a: coefficients for continuous arsenic variables and estimates of increase in arsenic required to increase pinprick score by 50% or 100%

| Pinprick measure | Regression coefficient ^b | Standard error | Predicted increase in arsenic needed to increase pinprick score | |
|------------------|-------------------------------------|----------------|---|---------|
| | | | 50% | 100% |
| Left arm | 0.00328* | .0008 | 123 ppb | 211 ppb |
| Right arm | 0.00255* | .0008 | 159 ppb | 271 ppb |
| Left leg | 0.00528* | .0011 | 77 ppb | 131 ppb |
| Right leg | 0.0057* | .0012 | 71 ppb | 121 ppb |

^aOrdered logistic regression; continuous As measure; pinprick sensitivity scored 1-5

^bCoefficients interpreted as the increase in log odds of a one unit increase in pinprick rating

^c50% reduction would be equivalent to an increase in pinprick scores of 2 to a 3, or a 1 to a 1.5. Similarly, a 100% reduction is equivalent to an increase in pinprick scores of 1 to 2, or 2 to 4

*p < 0.005

somatosensory pathways (Greenberg *et al.* 2002). Peripheral-to-central and side-to-side comparisons provide rapid indications of lesions in contralateral sensory pathways or peripheral neuropathy. The pinprick test was used in the present study since peripheral neuropathy is a well-known concomitant of acute arsenic poisoning.

The results indicate subtle, but significant impairment in the perception of pain in the hands and feet associated with arsenic exposure, consistent with case reports of peripheral neuropathy (Feldman 1999). Furthermore the results demonstrate that sensory effects of chronic exposure to arsenic in drinking water occur well below the 1000 µg/l level specified by NRC (1999) as the threshold for neurological impairment. Reduced pain sensation assessed by the pinprick test was observed at levels as low as 400 µg/l based on group comparisons. Regression analyses suggest effects at even lower concentrations. In particular, logistic regression modeling suggests that a 70–160 µg/l increase in drinking water arsenic results in a 50% increase in pinprick scores. The possibility of a threshold effect was not explored due to the limitations associated with a categorical response variable. Further study is needed to clarify the threshold for arsenic-related impairment of pain sensation.

Self-reported symptoms (numbness, pain, and reduced vibration sensation) were significantly associated with increased pinprick scores, providing face validity for pinprick test results. On the other hand, questionnaire

responses were somewhat less consistent than pinprick measures in relation to arsenic exposure category. That is, the highest proportion of respondents reporting neurological abnormalities occurred in the highest arsenic category, but a surprisingly high proportion of respondents in the lowest category also reported these signs or symptoms.

Similar findings were obtained in measures of vibrotactile sensation (Otto *et al.* 2006 (this issue)). Vibration thresholds were measured in two fingers (2nd and 5th) of both hands. The pattern of effects was the same for pain and vibration tests, that is, significant group differences were found between high and medium, high and low, but not medium and low exposure groups. Spline regression modeling of vibration measures suggests an effect threshold of 150–170 µg/l arsenic in drinking water.

These findings are consistent with a recent study of current perception thresholds (CPT) in asymptomatic Taiwanese living in villages where blackfoot disease (BFD) is endemic (Tseng 2003). BFD results from chronic exposure to arsenic in drinking water. Three nerves, trigeminal, median and peroneal, were assessed using standard CPT test frequencies: 5, 250 and 2000 Hz (Katims *et al.* 1986). Higher thresholds were found at all three frequencies in arsenic-exposed villagers compared to controls. However, the probability of abnormal values was higher in longer (peroneal and median) than shorter (trigeminal) nerves and at lower (5 and 250 Hz) than higher (2000 Hz) frequencies. It should

be noted that Tseng did not assess pain or vibration directly and that no environmental or body burden measures of arsenic were obtained in this study.

Few other epidemiological studies of peripheral neuropathy in arsenic-exposed populations have been undertaken. Kreiss *et al.* (1983) measured nerve conduction velocity (NCV) in Alaskan residents exposed to arsenic in well water at levels up to 15,000 $\mu\text{g}/\text{l}$. No significant association of arsenic concentration and NCV was found in that study. Gerr *et al.* (2000) also failed to find any relationship of arsenic exposure (from dust and soil) and NCV in residents living in the vicinity of a pesticide packaging plant, although strong associations were found with measures of postural balance, vibrotactile thresholds and hand tremor. Results of the latter study suggest that NCV is not as sensitive as other measures in assessing the neurological consequences of arsenic exposure, a possible explanation of the negative result in the Alaskan study.

The pinprick method is simple, but imprecise and subjective. The pinprick test is not calibrated and the method is not standardized, although it is widely used in clinical neurology. Fairly crude categories were used in assessing pinprick score, for instance, normal, end of finger, whole finger, below wrist or below elbow. Related measures such as the vibrothesiometer (Otto *et al.* 2006) (this issue) provide a more quantitative measure of sensory (vibration) threshold. The vibrothesiometer yields a continuous measure derived by the ascending method of limits, a well-established psychophysical procedure. However, both methods of assessing sensory thresholds ultimately reduce to subjective judgments. Despite these limitations, both methods provide evidence that arsenic exposure impairs somatosensory function in humans at levels not previously appreciated.

CONCLUSIONS

NRC (1999) concluded that arsenic in drinking water does not impair neurological function at concentrations below 1,000 $\mu\text{g}/\text{l}$. This study demonstrates that arsenic alters pain thresholds at well-water concentrations at least as low as 400 $\mu\text{g}/\text{l}$. Furthermore the results show that a small increase in drinking water arsenic (70–160 $\mu\text{g}/\text{l}$) is associated with a

50% increase in pinprick score. Although further research is needed to clarify the precise threshold for neurological effects, neurosensory measures appear to be a useful non-cancer endpoint for assessing the health effects of arsenic in drinking water. Finally, this study provides the first evidence that chronic exposure to arsenic in drinking water is associated with elevated pain thresholds.

DISCLAIMER

Although this document has been reviewed in accordance with U.S. Environmental Protection Agency policy and approved for publication, approval does not signify that the contents reflect the views or policies of the agency.

REFERENCES

- Agency for Toxic Substances and Disease Registry (ATSDR) 1998 *Toxicological Profile for Arsenic*. U.S. Department of Health and Human Services, Atlanta, GA.
- Basu, D., Dasgupta, J., Mukherjee, A. & Guha Mazumder, D. 1996 Chronic neuropathy due to arsenic intoxication from geochemical source—A five-year follow-up. *JANEI* **1**, 45–47.
- Feldman, R. 1999 *Occupational and Environmental Neurotoxicology*. Lippincott-Raven, New York, pp. 69–91.
- Gerr, F., Letz, R., Ryan, P. & Green, R. 2000 Neurological effects of environmental exposure to arsenic in dust and soil among humans. *NeuroToxicol.* **21**, 475–487.
- Greenberg, D., Aminoff, M. & Simon, R. 2002 *Clinical Neurology*, 5th Ed. Langer Medical Books/McGraw-Hill, New York.
- Hindmarsh, J., McLetchie, O., Hefferman, L., Hayne, O., Ellenberger, H., McCurdy, R. & Thiebaut, T. 1977 Electromyographic abnormalities in chronic environmental arsenicalism. *J. Anal. Toxicol.* **1**, 270–276.
- Hotta, N. 1989 Clinical aspects of chronic arsenic poisoning due to environmental and occupational pollution around a small refinery [Japanese]. *Nippon Taishitsugaku Zasshi [Jpn J Const Med]* **53**(1/2), 49–70.
- Kabata-Pendias, A. & Pendias, H. 1984 *Trace Elements in Soils and Plants*. CRC Press, Boca Raton, Florida.
- Katims, J., Long, D. & Ng, L. 1986 Transcutaneous electrical nerve stimulation (TNS): frequency and waveform specificity in humans. *Appl. Neurophysiol.* **49**, 86–91.
- Kreiss, K., Zack, M., Landrigan, P., Feldman, R., Niles, C., Chirico-Post, J., Sax, D., Boyd, M. & Cox, D. 1983 Neurologic evaluation of a population exposed to arsenic in Alaskan well water. *Arch. Environ. Hlth.* **38**, 116–121.
- Le, X. & Ma, M. 1998 *Short column liquid chromatography with hydride generation atomic fluorescence detection for speciation of arsenic*. *Anal. Chem.* **70**, 1926–1933.
- Ma, H., Guo, X., Yu, G., Wu, K., Xia, Y., Dang, Y., Li, Y., Zheng, Z., Zhou, H., Wang, F., Li, Z., Li, Z. & Wu, R. 1995 Clinical

- features of arsenicism in an endemic area (Inner Mongolia) with arsenic contamination in drinking water. *J. Chin. Endem. Dis. Spec Supp.*, 17–24.
- Ma, H., Xia, Y., Wu, K., Sun, T. & Mumford, J. 1999 Human exposure to arsenic and health effects in Bayingnormen, Inner Mongolia. In *Arsenic Exposure and Health Effects* (eds. Chappell, W., Abernathy, C. & Calderon, R.), Elsevier, New York, pp. 127–132.
- McCullagh, 1980 Regression models for ordinal data. *J. Royal Stat. Soc., Series B* **42**, 109–142.
- National Research Council 1999 *Arsenic in Drinking Water*. National Academy Press, Washington, DC.
- Otto, D., He, L., Xia, Y., Li, Y., Wu, K., Ning, Z., Shao, B., Hudnell, K., Kwok, R., Mumford, J., Geller, A. & Wade, T. 2006 Neurosensory effects of chronic exposure to arsenic via drinking water in Inner Mongolia: II. Vibrotactile and visual function. *J. Wat. Health* **4**(1), 39–48.
- SAS v 8.0 for Windows* 2001 SAS Institute, Cary, NC.
- Stata/SE 8.2 for Windows* 2004 Stata Corp., College Station, TX.
- Thornton, I. & Farago, M. 1997 The geochemistry of arsenic. In *Arsenic: Exposure and Health Effects* (eds. Abernathy, C., Calderon, R. & Chappell, W.), Chapman & Hall, New York, pp. 1–16.
- Tseng, C. 2003 Abnormal current perception thresholds measured by neurometer among residents in blackfoot disease-hyperendemic villages in Taiwan. *Toxicol. Lett.* **146**, 17–36.
- World Medical Association 1989 *Declaration of Helsinki*. 41st World Medical Assembly, Hong Kong, China.

Available online January 2006