

Clinical and Pathological Characteristics of *KEAP1*- and *NFE2L2*-Mutated Non-Small Cell Lung Carcinoma (NSCLC)



Rieke Frank^{1,2}, Matthias Scheffler^{1,2}, Sabine Merkelbach-Bruse^{2,3}, Michaela A. Ihle^{2,3}, Anna Kron^{1,3}, Michael Rauer^{4,5}, Frank Ueckerth^{2,3}, Katharina König^{2,3}, Sebastian Michels^{1,2}, Rieke Fischer^{1,2}, Anna Eisert^{1,2}, Jana Fassunke^{2,3}, Carina Heydt^{2,3}, Monika Serke^{2,6}, Yon-Dschun Ko^{2,7}, Ulrich Gerigk^{2,8}, Thomas Geist^{2,9}, Britta Kaminsky^{2,10}, Lukas C. Heukamp^{2,3}, Mathieu Clement-Ziza^{4,5}, Reinhard Büttner^{2,3}, and Jürgen Wolf^{1,2}

Abstract

Purpose: *KEAP1* and *NFE2L2* mutations are associated with impaired prognosis in a variety of cancers and with squamous cell carcinoma formation in non-small cell lung cancer (NSCLC). However, little is known about frequency, histology dependence, molecular and clinical presentation as well as response to systemic treatment in NSCLC.

Experimental Design: Tumor tissue of 1,391 patients with NSCLC was analyzed using next-generation sequencing (NGS). Clinical and pathologic characteristics, survival, and treatment outcome of patients with *KEAP1* or *NFE2L2* mutations were assessed.

Results: *KEAP1* mutations occurred with a frequency of 11.3% ($n = 157$) and *NFE2L2* mutations with a frequency of 3.5% ($n = 49$) in NSCLC patients. In the vast majority of patients, both mutations did not occur simultaneously. *KEAP1* mutations were found mainly in adenocarcinoma (AD; 72%), while *NFE2L2* mutations were more common in squamous cell

carcinoma (LSCC; 59%). *KEAP1* mutations were spread over the whole protein, whereas *NFE2L2* mutations were clustered in specific hotspot regions. In over 80% of the patients both mutations co-occurred with other cancer-related mutations, among them also targetable aberrations like activating *EGFR* mutations or *MET* amplification. Both patient groups showed different patterns of metastases, stage distribution and performance state. No patient with *KEAP1* mutation had a response on systemic treatment in first-, second-, or third-line setting. Of *NFE2L2*-mutated patients, none responded to second- or third-line therapy.

Conclusions: *KEAP1*- and *NFE2L2*-mutated NSCLC patients represent a highly heterogeneous patient cohort. Both are associated with different histologies and usually are found together with other cancer-related, partly targetable, genetic aberrations. In addition, both markers seem to be predictive for chemotherapy resistance. *Clin Cancer Res*; 24(13): 3087–96. ©2018 AACR.

¹Lung Cancer Group Cologne, Department I of Internal Medicine, University Hospital of Cologne, Cologne, Germany. ²Network Genomic Medicine (Lung Cancer), Cologne, Germany. ³Institute of Pathology, University Hospital of Cologne, Cologne, Germany. ⁴Center for Molecular Medicine Cologne (CMMC), University of Cologne, Cologne, Germany. ⁵Cologne Excellence Cluster for Cellular Stress Responses in Aging-Associated Diseases (CECAD) Research Center and Systems Biology of Ageing Cologne, University of Cologne, Cologne, Germany. ⁶Department for Pulmonology and Thoracic Oncology, Lung Clinic Hemer, Hemer, Germany. ⁷Johanniter Hospital, Evangelical Clinics of Bonn, Bonn, Germany. ⁸Marien-Hospital Bonn, GFO Clinics Bonn, Bonn, Germany. ⁹Center Practice for Lung and Bronchial Diseases, Düsseldorf, Germany. ¹⁰Clinic for Pulmonology and Allergology, Bethanien Hospital Solingen, Solingen, Germany.

Note: Supplementary data for this article are available at Clinical Cancer Research Online (<http://clincancerres.aacrjournals.org/>).

R. Frank and M. Scheffler contributed equally to this article.

R. Büttner and J. Wolf contributed equally to this article.

Corresponding Author: Jürgen Wolf, Lung Cancer Group Cologne, Department I for Internal Medicine, Center for Integrated Oncology, University Hospital of Cologne, Kerpener Str. 62, 50937 Cologne, Germany. Phone: 49-221-478-89050; Fax: 49-221-478-89051; E-mail: juergen.wolf@uk-koeln.de

doi: 10.1158/1078-0432.CCR-17-3416

©2018 American Association for Cancer Research.

Introduction

Systemic therapies targeting oncogenic aberrations in non-small cell lung cancer (NSCLC) have dramatically improved the outcome of genetically defined subgroups. Exemplarily, NSCLC patients with activating *EGFR* mutations, *ALK* or *ROS1* rearrangements benefit from tyrosine kinase inhibitor (TKI) treatment in terms of response and survival (1–6). It thus is a critical need to further identify and characterize genomic aberrations in NSCLC, which could either act as therapeutic targets themselves or modify response to targeted treatment.

The *KEAP1*-NRF2 (protein encoded by the *NFE2L2* gene) pathway plays a critical role in oxidative stress response by triggering antioxidant and anti-inflammatory effects (7). In healthy tissue *KEAP1* counteracts NRF2 by leading to its degradation (7–11). Being exposed to oxidative stress, *KEAP1* is inactivated and no longer able to bind and control NRF2, which is subsequently stabilized and translocates into the nucleus (7, 9, 12). There, *KEAP1* promotes transcription of genes encoding detoxifying enzymes and antioxidative stress proteins (13–15).

Mutations in the *KEAP1*/NRF2-pathway are known to be involved in malignant transformation in various cancer types (16–24). Somatic loss-of-function mutations of *KEAP1* lead to

Translational Relevance

In the present study, we show that *KEAP1* and *NFE2L2* mutations represent a heterogeneous NSCLC subgroup. Despite preclinical models showing a close interaction of these mutations in transformation, they occur nearly mutually exclusive in NSCLC and are associated with different histologies. Their frequent co-occurrence with other cancer-related mutations in line with the clinical heterogeneity argues against a role as "driver" mutations and stimulates further experiments investigating a modifier role, particular in tumors carrying already established drivers. In addition, our results suggest using NGS-based molecular multiplex diagnostics in clinical research to cover not only already established driver mutations but also potentially modifying co-occurring mutations. Finally, this work provides further evidence for a role of both *KEAP1* and *NFE2L2* mutations in chemoresistance.

an increase of NRF2 in the nucleus (16, 19, 25). Somatic gain-of-function mutations of *NFE2L2* are found near or within pivotal binding motifs (17) and interrupt binding of NRF2 to KEAP1 dimers (26). This leads to an increase of (i) intracellular NRF2, (ii) the synthesis of antioxidant and detoxification enzymes, and (iii) the production of drug efflux pumps in cancer cells (16, 19, 21, 26).

KEAP1 or *NFE2L2* mutations promote cell proliferation in tumors and may also participate in causing resistance to chemotherapy (19, 21, 27). Downregulation of *NFE2L2* or overexpression of *KEAP1* both triggered chemotherapy sensitivity (21, 26–29). In a squamous-cell carcinoma lung cancer (LSCC) mouse model, *KEAP1* mutations were associated with carcinoma formation and tumor aggressiveness, and both *KEAP1* and *NFE2L2* mutations promoted resistance against radiotherapy (RT; ref. 30).

It has recently been demonstrated that NSCLC patients with *KEAP1* mutation in addition to an activating *KRAS* mutation have a worse prognosis compared with *KRAS*-mutated patients without *KEAP1* mutation (31). The co-occurrence of *KEAP1* mutations generally describes a biologically unique subtype of *KRAS*-mutated NSCLCs (32). In adenocarcinoma of the lung (AD), patients with a high mutational load and benefit from anti-PD1 treatment also showed a high prevalence of *KEAP1* mutations (33). Furthermore, *NFE2L2* mutations were found to be associated with high PD-L1 expression in LSCC (34). It thus suggests a link between these mutations and a possible benefit from immune-checkpoint inhibition.

So far, however, little is known about the clinical presentation of NSCLC patients harboring mutations in either *KEAP1*, *NFE2L2*, or both as well as on their impact on systemic NSCLC treatment. In the present study, we describe and analyze the clinical and genetic characteristics of NSCLC patients with *KEAP1* or *NFE2L2* mutations and compare both groups with each other. We further evaluated the survival of these patients and their responses on systemic treatments.

Materials and Methods

Patients

Patient data was analyzed consecutively within the Network Genomic Medicine (NGM) Lung Cancer. NGM is a German

health care provider network where next-generation sequencing (NGS)-based molecular diagnostics of lung cancer is performed centrally for about 280 hospitals and private-practices-based oncologists (www.ngm-cancer.com). Incoming formalin-fixed paraffin-embedded (FFPE) lung cancer samples were analyzed from 2011 to 2013 and from May 2015 to August 2015 at the Institute of Pathology, University Hospital of Cologne (Cologne, Germany). Screening procedures and data assessment were performed in accordance with local standards. Data assessment was approved by the responsible ethics committee (ref. number 10-242), and all patients consented for data analysis.

The cohorts consisted of both an establishment cohort and a validation cohort for the implementation of NGS in routine lung cancer diagnostics (35). Patients of both cohorts were not pre-selected regarding smoking history, age, stage, or sex.

Samples and immunohistochemistry

Histopathologic diagnostics was performed centrally per local standard operating procedures. The histopathologic differentiation between AD and SQCC was based on immunohistochemical staining [CK5/6, CK7, p40, and thyroid transcription factor 1 (TTF1)] as previously published (36).

Next-generation sequencing (NGS) and FISH diagnostics

NGS was performed using a MiSeq benchtop sequencer as described previously (Illumina; ref. 35; Supplementary Table S1). We used an in-house algorithm to call for genomic variants of the targeted sequences (37). The variants were then stored in a FileMaker (Filemaker GmbH, Germany) database for further analyses. After reporting, we used COSMIC (<http://cancer.sanger.ac.uk/cosmic>), OncoKB (<http://oncokb.org>), and CancerHotspots (<http://cancerhotspots.org>) databases for further annotation. *In silico* evaluation on the impact of the detected mutations was made with PolyPhen-2 (<http://genetics.bwh.harvard.edu/pph2>).

In a subset of patients, FISH were carried out to determine rearrangements and/or amplifications (Supplementary Table S6). *MET* and *FGFR1* amplifications were tested and classified as reported (38, 39). *FGFR1* was analyzed routinely in all LSCC patients, whereas *MET* amplification testing by FISH was routinely introduced for AD in June 2014.

Clinical characteristics

We assessed age, sex, tumor stage (UICC), TNM status, smoking status (qualitatively following CDC reporting; ref. 40, and in pack-years), exposure to other known pulmonary carcinogens, histology, metastases and treatment modalities and their outcome, if available. Outcome was provided by the network partners and followed Response Evaluation Criteria in Solid Tumors V1.0 (RECIST), proposing that stable disease (SD) and responses [partial response (PR), complete response (CR)] need to be confirmed by a CT scan not earlier than 4 weeks after the initial response assessment. If restaging was not performed or available, we asked for investigators' assessments. Staging and restaging procedures were performed in accordance with local standards from each partner, usually including CT scans of the involved areas and MRI scans for brain metastases in all patients.

Statistics and software

Qualitative variables (e.g., sex or smoking status) were summarized by count and percentage, quantitative variables by mean

and standard deviation or median and range. Distribution of time-to-event was described by the Kaplan–Meier estimator. Follow-up was assessed using "reversed" Kaplan–Meier statistics. Association between categorical variables was assessed using χ^2 tests or Fisher exact tests, if appropriate. Statistical analyses were performed using SPSS v23.0 (IBM Corp.). Plots were generated using GraphPad Prism 6 (GraphPad Software Inc.). For the "lollipop charts," we used cBioPortal for Cancer Genomics' MutationMapper (http://www.cbioportal.org/mutation_mapper.jsp).

Results

KEAP1 mutations

Tumor tissue of 1,391 NSCLC patients was analyzed using NGS (Fig. 1; Supplementary Table S1). *KEAP1* mutations were detected in 157 patients (11.3%). Up to 134 different mutations were identified (Supplementary Table S2). They are spread over the whole *KEAP1* length (Fig. 3A). The most common mutations were Q620 deletions ($n = 5$), R320L ($n = 4$), R320W ($n = 4$), and V369A ($n = 3$). No hotspot mutations could be identified (Fig. 3A).

NFE2L2 mutations

Forty-nine patients (3.5%) had 30 different *NFE2L2* mutations (Supplementary Table S3). The most common mutations were W24R ($n = 5$), R34Q ($n = 5$), W24C ($n = 4$), E79K ($n = 4$), and R34G ($n = 3$). Clusters of mutations were observed in region E79 ($n = 5$), R34 ($n = 11$), and W24 ($n = 9$; Fig. 3A). R34 and E79 have been described as hotspot regions in human cancer, whereas W24 has not (41). These hotspot regions are near or within the ETGE and DLG motifs, which are the *KEAP1* binding domains of *NFE2L2* (ref. 17; Fig. 3A).




Figure 1.

Analytical flow sheet of the study. Euler diagram showing the proportions of mutated patients compared with the total cohort.

Clinical characteristics of patients with *KEAP1* and *NFE2L2* mutations

Detailed clinical characteristics of the cohort are provided in Table 1. A complete listing of the control group is provided in Supplementary Table S8. In both patients with *KEAP1* mutation (*KEAP1* group) and in patients with *NFE2L2* mutation (*NFE2L2* group), more males were affected. The same was true for the control group (Fig. 2A). There was no gender bias in this dataset ($P = 0.51$). The median age at diagnosis was 63.7 years (range, 40–84) for the *KEAP1* group and 66.6 years (range, 30–81) for the *NFE2L2* group.

Smoking status was assessable in 112 patients with *KEAP1* mutation and 35 patients with *NFE2L2* mutation. Both groups were characterized by strong smoking exposure, with only 6 never smokers in the *KEAP1* group and three in the *NFE2L2* group. In contrast, the control group consisted of 19.4% never-smokers (Fig. 2B). Median quantity of pack-years was 35 for *KEAP1* (range, 0–100) and 40 for *NFE2L2* (range, 0–120). Three patients in the *NFE2L2* group had a history of asbestos exposure and one had been exposed to quartz dust, while there was not a comparable exposure documented within the *KEAP1* group.

Patients with *KEAP1* mutation had predominantly AD (72.2%), comparable to the control group with 73.8% AD. In contrast, LSCC was the most frequent histology subtype with 59.2% in the *NFE2L2* group, followed by AD (32.7%; Fig. 2C, $P < 0.01$). In both groups, most patients had metastasized disease (stage IV) at first diagnosis, significantly more frequent in the *KEAP1* group (71.0% vs. 51.2%, $P = 0.03$).

Eastern Cooperative Oncology Group's Performance state (ECOG) was documented in 105 patients in the *KEAP1* group and 26 in the *NFE2L2* group. While ECOG 1 and ECOG 2 had a similar distribution pattern in both groups, there were differences for ECOG 0 (21.9% vs. 26.9%), ECOG 3 (7.6% vs. 3.8%), and ECOG 4 (1.9% vs. 0%). The difference in ECOG 3 plus 4 in both groups was not significant ($P = 0.46$). In the control group, nearly half of the patients presented with ECOG 0 (Fig. 2D).

In the *KEAP1* group, a high frequency of distant metastases affecting bones, brain, liver, skin, and spleen could be seen in stage IV patients, whereas patients in the *NFE2L2* mutation more frequently suffered from local spread (i.e., lung, pleura, mediastinal lymph nodes; Fig. 2E and F).

There were 4 patients with both a *KEAP1* mutation and a *NFE2L2* mutation. All four patients had a history of smoking and three patients had LSCC, while only 1 patient had AD. There were no further similarities regarding clinical characteristics and co-occurring aberrations between those patients. We performed a Fisher exact test, which revealed a z -pooled $P > 2.2 \times 10^{-16}$ for mutually exclusivity.

Co-occurring aberrations

We found additional genomic aberrations in 87.3% of the *KEAP1* group and 83.7% of the *NFE2L2* group (Fig. 3B; Supplementary Tables S4 and S5). *TP53* mutations were the most common co-occurring mutations in both groups (44.9%, *KEAP1*; 40.8%, *NFE2L2*). *EGFR* mutations also showed a similar distribution pattern occurring in 6.3% (*KEAP1*) and 6.1% (*NFE2L2*). In both patient groups *MET* amplifications occurred at high frequency (18.3%, *KEAP1*; 26.3%, *NFE2L2*; Supplementary Table S6). Fitting to the correspondent histological distributions, we detected differences regarding the co-occurrence of *KRAS*

Table 1. Comparison of clinical characteristics of patients with *KEAP1* mutations versus patients with *NFE2L2* mutations

Characteristics	Patients (%) <i>KEAP1</i>	Patients (%) <i>NFE2L2</i>	Patients (%) Control
Number of patients	157 (100)	49 (100)	826 (100)
Age at diagnosis			
Median	63.7	66.6	67.0
Range	40–84	30–81	23–93
Sex			
Male	95 (60.5)	27 (55.1)	471 (57.0)
Female	62 (39.5)	22 (44.9)	355 (43.0)
ECOG			
0	23 (14.6)	7 (14.3)	293 (35.4)
1	55 (35.0)	14 (28.6)	211 (25.5)
2	17 (10.8)	4 (8.2)	54 (6.5)
3	8 (5.1)	1 (2.0)	39 (4.7)
4	2 (1.3)	0 (0.0)	3 (0.3)
5	0 (0.0)	0 (0.0)	3 (0.3)
NA	52 (33.1)	23 (46.9)	233 (28.2)
Smoking history			
Never smoker	6 (3.8)	3 (6.1)	75 (9.1)
Former smoker	37 (23.6)	8 (16.3)	153 (18.5)
Current smoker	69 (43.9)	24 (49.0)	159 (19.2)
N/A	45 (28.7)	14 (28.6)	439 (53.1)
Pack years			
Median (range)	86	28	40 (0–120)
Histology			
Squamous cell carcinoma	158	49	826
Adenocarcinoma	28 (17.7)	29 (59.2)	140 (16.9)
Adenosquamous carcinoma	114 (72.2)	16 (32.7)	610 (73.8)
Others/NA	5 (3.2)	1 (2.0)	4 (0.5)
Others/NA	11 (7.0)	3 (6.1)	72 (8.7)
Stage at diagnosis			
I	158	49	826
II	10 (6.3)	8 (16.3)	20 (2.4)
III	6 (3.8)	4 (8.2)	15 (1.8)
IV	24 (15.2)	9 (18.4)	104 (12.6)
N/A	98 (62.0)	22 (44.9)	451 (54.6)
N/A	20 (12.7)	6 (12.2)	236 (28.6)
T stage (TNM)			
T1	158	49	
T2	20 (12.7)	5 (10.2)	
T3	34 (21.5)	11 (22.4)	
T4	22 (13.9)	8 (16.3)	
T4	43 (27.2)	17 (34.7)	
N/A	39 (24.7)	8 (16.3)	
N stage (TNM)			
N0	158	49	
N1	28 (17.7)	12 (24.5)	
N2	16 (10.1)	6 (12.2)	
N3	41 (25.9)	9 (18.4)	
N3	33 (20.9)	13 (26.5)	
N/A	40 (25.3)	9 (18.4)	
Metastases localization (stage IV), thereof			
Bones	98	22	
Brain	27 (27.6)	2 (9.1)	
Lung	19 (19.4)	2 (9.1)	
Pleura	20 (20.4)	9 (40.9)	
Distant lymph nodes	12 (12.2)	3 (13.6)	
Adrenal gland	9 (9.2)	3 (13.6)	
Liver	7 (7.1)	4 (18.2)	
Skin	11 (11.2)	1 (4.5)	
Spleen	3 (3.1)	0 (0.0)	
Stomach	1 (1.0)	0 (0.0)	
Pancreas	2 (2.0)	0 (0.0)	
Pancreas	0 (0.0)	1 (4.5)	

($P = 0.03$), *PTEN* (n.s.), *PIK3CA* (n.s.) mutations and *FGFR1* amplifications (Supplementary Table S6; Fig. 3B). Out of the reported mutations, we identified only one *TP53* polymorphism (P72R) in both groups, six proposed *MET* polymorphisms (T1010I), and one *EGFR* P848L. The differences in mutational profile is strongly supported by data on gene expression differences and mutational data (42, 43).

Survival

Median follow-up for stage IV patients was only 1.8 months (95% CI, 0.0–3.7 months) for the *KEAP1* group ($n = 80$) and 5.3 months (95% CI, 2.1–8.5 months) for the *NFE2L2* group ($n = 21$). Thus, we abstained from further survival analyses. The median OS of these patients with *KEAP1* mutation was 19.1 months (95% CI, 1.8–36.3 months) and 14.0 months




Figure 2. Clinical characteristics of patients with *KEAP1* and *NFE2L2* mutations. **A**, Gender distribution. **B**, Smoking history. **C**, Histologic subtypes. **D**, Eastern Cooperative Oncology Group's Performance state (ECOG) at diagnosis. **E** and **F**, Distribution of metastatic sites in stage IV patients. Body and organ pictures taken from the Library of Science & Medical Illustrations under the CC BY-NC-SA 4.0 license.

Downloaded from <http://aacrjournals.org/clinccancerres/article-pdf/24/13/3087/2044366/3087.pdf> by guest on 14 December 2024

(95% CI, 5.6–22.3 months) for patients with *NFE2L2* mutation (Fig. 3C).

Response assessment

For 30 patients with *KEAP1* mutation, details about systemic therapy were assessed. Twenty-seven patients had additional aberrations (Supplementary Table S7). These patients received between one and three lines of chemotherapy or TKIs as systemic therapy. Most patients (28) did not respond to first-line chemotherapy [progressive disease (PD)]. Both two patients with

SD had an activating *EGFR* mutation and one of them received erlotinib and bevacizumab, the other had SD under carboplatin/paclitaxel, representing the only non-PD *KEAP1* patient treated with conventional chemotherapy. Due to the short duration of first-line treatment, 17 patients (56.7%) received systemic first-line therapy. Outcome analysis was available for 8 patients, which all developed PD regardless of therapy regimen. In third-line, 3 patients received therapy. Two of them had a PD but the outcome of the other patient is not known (Supplementary Table S7). There was no responder in none of the therapy lines.




Figure 3. Results of sequencing analyses and survival analyses. **A**, Distribution of *KEAP1* mutations and *NFE2L2* mutations among the genomic position. **B**, Frequency of co-occurring mutations in percentage (only mutations with $\geq 5\%$ in one of the subgroups). **C**, Overall survival (Kaplan-Meier curve) in stage IV patients with *KEAP1* mutations and *NFE2L2* mutations ($n = 101$).

Eight patients with *NFE2L2* mutation receiving systemic therapy were evaluable for outcome analysis. Only one patient did not present an additional aberration. Four patients developed PD

during first-line therapy, leading to a significantly better outcome regarding non-PD first-line as compared with the *KEAP1* group ($P = 0.01$, Fig. 4B). Two patients had a SD as best outcome




Figure 4. Treatment outcome. **A**, Responses to systemic therapy according to the therapy line. **B**, PD vs. non-PD in first-line systemic therapy. **C**, Regimen used in first-line therapy, second-line therapy, and third-line therapy. **D**, Progressive disease (PD) versus non-progressive disease (non-PD) of chemotherapy-based and targeted therapy-based therapies regardless of therapy line.

and two patients had a partial response (PR). Of these 4 patients with benefit from therapy, 2 patients received erlotinib and bevacizumab and 1 of these 2 patients had an activating *EGFR* mutation and responded. The third patient harbored an additional *STK11* mutation and received carboplatin and etoposide resulting in a PR, representing the only responder to a platinum-based regimen in the cohort. The fourth patient did not harbor a detectable comutation and received

cisplatin, pemetrexed, and bevacizumab, developing a SD. Two patients received both second- and third-line treatments without response (Fig. 4A).

Out of the 38 patients with an outcome analysis 16 patients harbored an additional *TP53* mutation. Twelve out of 32 patients (37.5%) with a PD during first-line therapy had a *TP53* mutation. In contrast, also 4 out of 6 patients (66.6%) with a non-PD during first-line therapy had *TP53* mutations. We therefore conclude that

chemoresistance in our cohort is not biased by the co-occurrence of *TP53* mutations.

Comparing treatment with regard to the therapy used regardless of therapy line, 62 regimens could be identified in 38 patients, whereof 50 were chemotherapy treatments and 10 erlotinib-based treatments (\pm bevacizumab). Two lines could not be identified retrospectively. In each group, we found one PR and two SD, respectively (Fig. 4D).

Discussion

To the best of our knowledge, this is the largest cohort of NSCLC patients with *KEAP1* and *NFE2L2* mutations analyzed so far regarding clinical and pathologic characteristics. Both mutations are discussed to play a pivotal role in cancer formation and maintenance. We detected *KEAP1* mutations with a frequency of 11.3% and *NFE2L2* mutations with a frequency of 3.5% among a European all-comer NSCLC cohort.

Surprisingly, even though both types of mutations affect the same pathway and show a close functional interaction, they only exceptionally co-occur within the same tumor. Thus, only 4 patients in our cohort harbored a *KEAP1* and a *NFE2L2* mutation simultaneously. Moreover, both mutations are associated with different NSCLC histologies, as well as their known expression patterns (43). While most *KEAP1* mutations occurred in patients with AD, patients with *NFE2L2* mutations mostly presented with LSCC. On a molecular level, *KEAP1* mutations were not found in specific hotspot regions, but heterogeneously spread over the whole protein, whereas *NFE2L2* mutations were found in specific hotspot regions, in line with previous reports (17, 41). Regarding clinical presentation, *KEAP1* patients had a significantly higher frequency of stage IV at primary diagnosis than *NFE2L2* patients. Further, although not significantly, the *KEAP1* group presented in worse performance state and tended to present with a higher frequency of hematogenous spread.

In a small subset of our cohort in which reproducible response on systemic therapies was assessable, response on different lines and regimens was remarkably poor, especially in the *KEAP1* group. None of the patients in both groups responded to second-line or third-line therapy. The co-occurrence of *TP53* mutation did not bias the outcome of these patients. Given the broad spectrum of chemotherapy agents used, these data suggest that these mutations might contribute to chemotherapy resistance, as proposed in preclinical lung cancer models and in other tumor types (19, 21, 26–29). However, in view of the small number of patients and the short follow-up, further data are needed to confirm this finding.

For both mutations, the majority of patients had additional aberrations. In our cohort, in some contrast to recent work (26), even the co-occurrence of targetable *EGFR* mutations was detected in both groups and clinical benefit for patients treated with an *EGFR*-TKI was seen. Further, nearly half of the *NFE2L2* patients had amplifications of either *MET* (26.3%) or *FGFR1* (20.6%). *MET* amplifications were also frequent in *KEAP1*-mutated patients (18.3%), which, in addition, harbor *KRAS* mutations in nearly half of the cases. It remains to be established whether and how co-occurring *NFE2L2* or *KEAP1* mutations affect efficacy of specific targeted treatments against already established targets like activating *EGFR* mutations or against targets under clinical investigation like amplifications

of *MET* and *FGFR1* or *KRAS* mutations and whether they themselves might represent targets for specific therapies. To date, there are efforts reported to overcome the proposed chemotherapy-resistance triggered by the *KEAP1/NRF2* pathway using brusatol as an *NRF2* inhibitor (44). This potential therapeutic option might be considered when reevaluating treatment approaches for *KEAP1/NFE2L2*-mutated NSCLC. Nevertheless, it still remains unclear how the mutations in *KEAP1* and *NFE2L2* interact with *NRF2* inhibition.

We are aware of the main limitations of this study: Even though we conducted parts of this study prospectively, robust therapeutic data are retrospective and available only for a subset of patients. Further, due to the short follow-up period, we could not obtain robust survival data that would be needed to link genomic aberrations with a prognostic impact. The number of patients reported here with the analyzed mutations is also small due to the prevalences of both mutations in our cohort. Nevertheless, the prevalences were in line with data from the TCGA cohorts LUAD and LUSC (17, 45). Further, with the diagnostic setup we used here, it is not possible to make assumptions about *KEAP1* allelic losses, which have been shown to occur frequently in lung cancer (16).

In summary, our data provide a comprehensive description of *NFE2L2* and *KEAP1* mutations regarding frequency and clinical as well as molecular characteristics. The observed heterogeneity of NSCLC harboring these mutations as well as the co-occurrence of further cancer-related mutations in the majority of patients harboring *NFE2L2* or *KEAP1* strongly argue against a function as driver mutations and rather imply a role as modifiers. Our observations also suggest that both mutations may play a role in chemoresistance and advocate for the development of therapeutic approaches different than chemotherapy for these patients. In particular, the interaction of mutations in *KEAP1* and *NFE2L2* with co-occurring targetable genetic aberrations needs to be better understood and underlines the need for NGS-based molecular multiplex diagnostics to cover co-occurring mutations at least in the setting of clinical research.

Disclosure of Potential Conflicts of Interest

R. Fischer is a consultant/advisory board member for and reports receiving speakers bureau honoraria from Lung Cancer. J. Fassunke reports receiving speakers bureau honoraria from AstraZeneca. C. Heydt reports receiving speakers bureau honoraria from AstraZeneca. M. Serke reports receiving speakers bureau honoraria from Bristol, Myers, Squibb; Roche; AstraZeneca; Boehringer; and Celgene, and is a consultant/advisory board member for Bristol, Myers, Squibb; MSD; Roche; Boehringer, and Abbvie. No potential conflicts of interest were disclosed by the other authors.

Authors' Contributions

Conception and design: R. Frank, M. Scheffler, M. Serke, R. Büttner, J. Wolf
Development of methodology: M. Scheffler, K. König, C. Heydt, M. Serke, R. Büttner, J. Wolf

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): R. Frank, M. Scheffler, S. Merkelbach-Bruse, M. Ihle, K. König, R. Fischer, J. Fassunke, C. Heydt, M. Serke, Y.-D. Ko, T. Geist, B. Kaminsky, L.C. Heukamp, R. Büttner, J. Wolf

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): R. Frank, M. Scheffler, S. Merkelbach-Bruse, M. Ihle, A. Kron, M. Rauer, F. Ueckerth, C. Heydt, M. Serke, M. Clement-Ziza, R. Büttner, J. Wolf

Writing, review, and/or revision of the manuscript: R. Frank, M. Scheffler, S. Merkelbach-Bruse, M. Ihle, A. Kron, M. Rauer, S. Michels, R. Fischer, A. Eisert,

J. Fassunke, C. Heydt, M. Serke, U. Gerigk, T. Geist, B. Kaminsky, L.C. Heukamp, M. Clement-Ziza, R. Büttner, J. Wolf
Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): M. Scheffler, A. Eisert, J. Fassunke, U. Gerigk, J. Wolf
Study supervision: M. Scheffler, M. Clement-Ziza, J. Wolf

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received November 15, 2017; revised February 22, 2018; accepted March 27, 2018; published first April 3, 2018.

References

- Mok TS, Wu YL, Thongprasert S, Yang CH, Chu DT, Saijo N, et al. Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma. *N Engl J Med* 2009;361:947–57.
- Shaw AT, Kim DW, Nakagawa K, Seto T, Crino L, Ahn MJ, et al. Crizotinib versus chemotherapy in advanced ALK-positive lung cancer. *N Engl J Med* 2013;368:2385–94.
- Shaw AT, Ou SH, Bang YJ, Camidge DR, Solomon BJ, Salgia R, et al. Crizotinib in ROS1-rearranged non-small-cell lung cancer. *N Engl J Med* 2014;371:1963–71.
- Scheffler M, Schultheis A, Teixido C, Michels S, Morales-Espinosa D, Viteri S, et al. ROS1 rearrangements in lung adenocarcinoma: prognostic impact, therapeutic options and genetic variability. *Oncotarget* 2015;6:10577–85.
- Clinical Lung Cancer Genome P, Network Genomic M. A genomics-based classification of human lung tumors. *Sci Transl Med* 2013;5:209ra153.
- Barlesi F, Mazieres J, Merlio JP, Debieuvre D, Mosser J, Lena H, et al. Routine molecular profiling of patients with advanced non-small-cell lung cancer: results of a 1-year nationwide programme of the French Cooperative Thoracic Intergroup (IFCT). *Lancet* 2016;387:1415–26.
- Itoh K, Wakabayashi N, Katoh Y, Ishii T, Igarashi K, Engel JD, et al. Keap1 represses nuclear activation of antioxidant responsive elements by Nrf2 through binding to the amino-terminal Neh2 domain. *Genes Dev* 1999;13:76–86.
- Zhang DD, Lo SC, Cross JV, Templeton DJ, Hannink M. Keap1 is a redox-regulated substrate adaptor protein for a Cul3-dependent ubiquitin ligase complex. *Mol Cell Biol* 2004;24:10941–53.
- Zhang DD, Hannink M. Distinct cysteine residues in Keap1 are required for Keap1-dependent ubiquitination of Nrf2 and for stabilization of Nrf2 by chemopreventive agents and oxidative stress. *Mol Cell Biol* 2003;23:8137–51.
- Kobayashi A, Kang MI, Okawa H, Ohtsuiji M, Zenke Y, Chiba T, et al. Oxidative stress sensor Keap1 functions as an adaptor for Cul3-based E3 ligase to regulate proteasomal degradation of Nrf2. *Mol Cell Biol* 2004;24:7130–9.
- McMahon M, Itoh K, Yamamoto M, Hayes JD. Keap1-dependent proteasomal degradation of transcription factor Nrf2 contributes to the negative regulation of antioxidant response element-driven gene expression. *J Biol Chem* 2003;278:21592–600.
- Dhakshinamoorthy S, Saiswal AK. Functional characterization and role of INrf2 in antioxidant response element-mediated expression and antioxidant induction of NAD(P)H:quinone oxidoreductase1 gene. *Oncogene* 2001;20:3906–17.
- Itoh K, Chiba T, Takahashi S, Ishii T, Igarashi K, Katoh Y, et al. An Nrf2/small Maf heterodimer mediates the induction of phase II detoxifying enzyme genes through antioxidant response elements. *Biochem Biophys Res Commun* 1997;236:313–22.
- Kwak MK, Itoh K, Yamamoto M, Sutter TR, Kensler TW. Role of transcription factor Nrf2 in the induction of hepatic phase 2 and antioxidative enzymes *in vivo* by the cancer chemoprotective agent, 3H-1, 2-dimethiole-3-thione. *Mol Med* 2001;7:135–45.
- Ishii T, Itoh K, Takahashi S, Sato H, Yanagawa T, Katoh Y, et al. Transcription factor Nrf2 coordinately regulates a group of oxidative stress-inducible genes in macrophages. *J Biol Chem* 2000;275:16023–9.
- Singh A, Misra V, Thimmulappa RK, Lee H, Ames S, Hoque MO, et al. Dysfunctional KEAP1-NRF2 interaction in non-small-cell lung cancer. *PLoS Med* 2006;3:e420.
- Cancer Genome Atlas Research Network. Comprehensive genomic characterization of squamous cell lung cancers. *Nature* 2012;489:519–25.
- Kandoth C, McLellan MD, Vandin F, Ye K, Niu B, Lu C, et al. Mutational landscape and significance across 12 major cancer types. *Nature* 2013;502:333–9.
- Ohta T, Iijima K, Miyamoto M, Nakahara I, Tanaka H, Ohtsuiji M, et al. Loss of Keap1 function activates Nrf2 and provides advantages for lung cancer cell growth. *Cancer Res* 2008;68:1303–9.
- Ooi A, Dykema K, Ansari A, Petillo D, Snider J, Kahnoski R, et al. CUL3 and NRF2 mutations confer an NRF2 activation phenotype in a sporadic form of papillary renal cell carcinoma. *Cancer Res* 2013;73:2044–51.
- Zhang P, Singh A, Yegnasubramanian S, Esopi D, Kombairaju P, Bodas M, et al. Loss of Kelch-like ECH-associated protein 1 function in prostate cancer cells causes chemoresistance and radioresistance and promotes tumor growth. *Mol Cancer Ther* 2010;9:336–46.
- Shibata T, Kokubu A, Gotoh M, Ojima H, Ohta T, Yamamoto M, et al. Genetic alteration of Keap1 confers constitutive Nrf2 activation and resistance to chemotherapy in gallbladder cancer. *Gastroenterology* 2008;135:1358–68. 68.e1–4.
- Nioi P, Nguyen T. A mutation of Keap1 found in breast cancer impairs its ability to repress Nrf2 activity. *Biochem Biophys Res Commun* 2007;362:816–21.
- Kim YR, Oh JE, Kim MS, Kang MR, Park SW, Han JY, et al. Oncogenic NRF2 mutations in squamous cell carcinomas of oesophagus and skin. *J Pathol* 2010;220:446–51.
- Li QK, Singh A, Biswal S, Askin F, Gabrielson E. KEAP1 gene mutations and NRF2 activation are common in pulmonary papillary adenocarcinoma. *J Hum Genet* 2011;56:230–4.
- Shibata T, Ohta T, Tong KI, Kokubu A, Odogawa R, Tsuta K, et al. Cancer related mutations in NRF2 impair its recognition by Keap1-Cul3 E3 ligase and promote malignancy. *Proc Natl Acad Sci U S A* 2008;105:13568–73.
- Homma S, Ishii Y, Morishima Y, Yamadori T, Matsuno Y, Haraguchi N, et al. Nrf2 enhances cell proliferation and resistance to anticancer drugs in human lung cancer. *Clin Cancer Res* 2009;15:3423–32.
- Wang XJ, Sun Z, Villeneuve NF, Zhang S, Zhao F, Li Y, et al. Nrf2 enhances resistance of cancer cells to chemotherapeutic drugs, the dark side of Nrf2. *Carcinogenesis* 2008;29:1235–43.
- Singh A, Boldin-Adamsky S, Thimmulappa RK, Rath SK, Ashush H, Coulter J, et al. RNAi-mediated silencing of nuclear factor erythroid-2-related factor 2 gene expression in non-small cell lung cancer inhibits tumor growth and increases efficacy of chemotherapy. *Cancer Res* 2008;68:7975–84.
- Jeong Y, Hoang NT, Lovejoy A, Stehr H, Newman AM, Gentles AJ, et al. Role of KEAP1/NRF2 and TP53 mutations in lung squamous cell carcinoma development and radiation resistance. *Cancer Discov* 2017;7:86–101.
- Riely GJ, Kim HR, Yu HA, Berger MF, Solit DB, Kris MG, et al. Association of outcomes and co-occurring genomic alterations in patients with KRAS-mutant non-small cell lung cancer. *ASCO Annual Meeting*; 2016: J Clin Oncol 34, 2016 (suppl; abstr 9019).
- Skoulidis F, Byers LA, Diao L, Papadimitrakopoulou VA, Tong P, Izzo J, et al. Co-occurring genomic alterations define major subsets of KRAS-mutant lung adenocarcinoma with distinct biology, immune profiles, and therapeutic vulnerabilities. *Cancer Discov* 2015;5:860–77.
- Rizvi NA, Hellmann MD, Snyder A, Kvistborg P, Makarov V, Havel JJ, et al. Cancer immunology. Mutational landscape determines sensitivity to PD-1 blockade in non-small cell lung cancer. *Science* 2015;348:124–8.
- Scheel AH, Ansen S, Schultheis AM, Scheffler M, Fischer RN, Michels S, et al. PD-L1 expression in non-small cell lung cancer: correlations with genetic alterations. *Oncoimmunology* 2016;5:e1131379.
- Konig K, Peifer M, Fassunke J, Ihle MA, Kunstlinger H, Heydt C, et al. Implementation of amplicon parallel sequencing leads to improvement of diagnosis and therapy of lung cancer patients. *J Thorac Oncol* 2015;10:1049–57.
- Travis WD, Brambilla E, Nicholson AG, Yatabe Y, Austin JH, Beasley MB, et al. The 2015 World Health Organization classification of lung tumors: impact of genetic, clinical and radiologic advances since the 2004 classification. *J Thorac Oncol* 2015;10:1243–60.

37. Peifer M, Fernandez-Cuesta L, Sos ML, George J, Seidel D, Kasper LH, et al. Integrative genome analyses identify key somatic driver mutations of small-cell lung cancer. *Nat Genet* 2012;44:1104–10.
38. Schildhaus HU, Heukamp LC, Merkelbach-Bruse S, Riesner K, Schmitz K, Binot E, et al. Definition of a fluorescence in-situ hybridization score identifies high- and low-level FGFR1 amplification types in squamous cell lung cancer. *Mod Pathol* 2012;25:1473–80.
39. Schildhaus HU, Schultheis AM, Ruschoff J, Binot E, Merkelbach-Bruse S, Fassunke J, et al. MET amplification status in therapy-naive adeno- and squamous cell carcinomas of the lung. *Clin Cancer Res* 2015; 21:907–15.
40. Centers for Disease C, Prevention. Cigarette smoking among adults and trends in smoking cessation—United States, 2008. *MMWR Morb Mortal Wkly Rep* 2009;58:1227–32.
41. Chang MT, Asthana S, Gao SP, Lee BH, Chapman JS, Kandoth C, et al. Identifying recurrent mutations in cancer reveals widespread lineage diversity and mutational specificity. *Nat Biotechnol* 2016;34:155–63.
42. Campbell JD, Alexandrov A, Kim J, Wala J, Berger AH, Pedamallu CS, et al. Distinct patterns of somatic genome alterations in lung adenocarcinomas and squamous cell carcinomas. *Nat Genet* 2016;48:607–16.
43. Solis LM, Behrens C, Dong W, Suraokar M, Ozburn NC, Moran CA, et al. Nrf2 and Keap1 abnormalities in non-small cell lung carcinoma and association with clinicopathologic features. *Clin Cancer Res* 2010;16:3743–53.
44. Harder B, Tian W, La Clair JJ, Tan AC, Ooi A, Chapman E, et al. Brusatol overcomes chemoresistance through inhibition of protein translation. *Mol Carcinog* 2017;56:1493–500.
45. Cancer Genome Atlas Research Network. Comprehensive molecular profiling of lung adenocarcinoma. *Nature* 2014;511:543–50.