

Bim-targeted cancer therapy: A link between drug action and underlying molecular changes

Toru Akiyama,^{1,2} Crispin R. Dass,¹
and Peter F.M. Choong^{1,3}

¹Departments of Orthopaedics and Surgery, University of Melbourne, St. Vincent's Health, Melbourne, Australia; ²Department of Orthopaedic Surgery, Faculty of Medicine, The University of Tokyo, Tokyo, Japan; and ³Bone and Soft Tissue Sarcoma Service, Peter MacCallum Cancer Centre, Melbourne, Australia

Abstract

In the past few years, the pro-apoptotic molecule Bim has attracted increasing attention as a plausible target for tumor therapy. A variety of normal and pathological systems regulated by Bim, dependent on cell type, apoptotic stimulation, and chemotherapeutic agents, have been documented. Bim promotes anoikis of many tumor cells, such as lung cancer, breast cancer, osteosarcoma, and melanoma. Various chemotherapeutic agents use Bim as a mediating executioner of cell death. Hence, Bim suppression supports metastasis and chemoresistance. Imatinib, gefitinib, bortezomib, and Bim protein itself are spotlighted as current and future Bim-targeting therapeutic agents. The potential benefits of Bim-targeted therapies are selectivity of treatment for tumor cells and reduction in tumor-associated phenomena such as chemoresistance and metastasis. Thus, Bim-targeting therapies may provide more effective and unique tumor management modalities in future. This review article discusses all these issues. [Mol Cancer Ther 2009;8(12):3173–80]

Introduction

The important role of "apoptosis" in tumorigenesis and tumor treatment has been the subject of various reviews (1). The completion of tumorigenesis requires a multistep process whereby disturbing physiologic and genetically mediated programs mandates unceasing proliferation and growth (2). Thus, it is essential that the acquisition of these

characters protects cells from being channeled into the normal genetic and programmed cell death system—apoptosis. In addition, metastatic ability and chemotherapy resistance require the abrogation of apoptosis (1, 3, 4). Experimental evidence shows that a single oncogene can be critical to tumorigenesis or that tumors become addicted to the oncogene for their tumorigenesis (2). The latter is an "oncogene addiction" hypothesis. Naturally, these addicted genes pose as suitable targets for tumor control. Bim is one such possible candidate of an addicted oncogene.

This review focuses on the normal Bim regulatory system at first. Subsequently, it discusses the role of Bim suppression in tumorigenesis, focusing on the areas of tumor metastasis and chemotherapy. Some current chemotherapeutic agents, especially molecular-targeting agents, use Bim as an executioner. These agents could be classified as "primitive" Bim-targeting agents. The diversity of Bim-regulatory systems depends on cell type, stimulus, and pathological events, such as metastasis and chemoresistance.

Thus, proper control of Bim expression provides tumor-selective therapeutic effects. This is a major possible advantage for Bim-targeted therapies, which use not only Bim induced by agents but also the Bim protein itself. Modified Bim protein is also a possible candidate for targeted oncological therapy in the future. Although one of the disadvantages of Bim-targeted therapy is broad and strong apoptosis-inducing ability, this in turn can be exploited for therapeutical outcomes. Finally, the review concludes with a discussion on the advantages and disadvantages of Bim-targeted therapies.

The Role of Bim as a Guardian of Tissue Homeostasis

In response to apoptotic signals, various enzymes are activated in a pathway-specific manner and the classical caspase activation chain reaction is set in motion (5). Mammals have mainly two distinct apoptosis signaling pathways, the death receptor pathway and the mitochondrial pathway. In the mitochondrial pathway, the B-cell lymphoma-2 (Bcl-2)-family of proteins have a crucial role. The Bcl-2 family comprises three subfamilies, namely, an anti-apoptotic family, pro-apoptotic multidomain family, and pro-apoptotic BH3-only protein family. The Bcl-2-homology domain 3 only (BH3-only) proteins share only the short BH3 domain with members of the BCL-2 family. BH3-only proteins are strictly regulated through both transcription and post-transcription mechanisms (6). They are essential for

Received 4/30/09; revised 9/21/09; accepted 10/10/09; published OnlineFirst 11/24/09.

Requests for reprints: Crispin R. Dass, St. Vincent's Hospital Melbourne, L3-Daly Wing, 35 Victoria Pde., Fitzroy, Melbourne, VIC 3065 Australia. Phone: 61-3-9288-3990; Fax: 61-3-9416-3610. E-mail: crispin.dass@svhm.org.au

Copyright © 2009 American Association for Cancer Research.

doi:10.1158/1535-7163.MCT-09-0685




Figure 1. A mechanistic scheme of how Bim activation can act to promote apoptosis. Cytokine withdrawal, chemotherapeutic agents, and cell detachment trigger apoptosis by upregulation of Bim. Bim upregulation leads to disruption of the outer mitochondrial membrane, resulting in release of cytochrome c. Released cytochrome c promotes apoptosome-mediated caspase-9 activation. The apoptosome is composed of Apaf1 (apoptotic-protease-activating factor 1) and released cytochrome c. Once activated, caspase-9 cleaves executioner caspase-3, -6, and -7 and leads to apoptosis in the stimulated cell.

initiation of various physiological apoptotic situations, including developmentally programmed cell death and stress-induced apoptosis (7). So far, at least eight BH3-only proteins have been discovered in mice and humans. Bim is one of these BH3-only proteins. Bim upregulation triggers cytochrome c release from mitochondria, which consequently induces a chain reaction that entails the formation of the apoptosome and the activation of its effector, caspase-9. Thus, Bim upregulation induces apoptosis (Fig. 1).

Bim was independently identified by two groups as a Bcl-2- or Mcl1-binding protein, by screening of a λ -phage expression library (from a T-cell lymphoma) and by screening of a yeast two-hybrid library (from ovarian tissue), respectively. Alternative splicing generates at least three Bim isoforms, including Bim_S, Bim_L, and Bim_{EL}, which differ in their proapoptotic activity. Bim_{EL} is composed of 196 amino acids, and contains not only the BH3 domain but also two ubiquitination sites, three ERK phosphorylation sites, one JNK phosphorylation site, and one transmembrane domain (Fig. 2). *In vitro*, Bim is essential for apoptosis of various cell types, including lymphocytes, osteoclasts, osteoblasts, mast cells, epithelial cells, endothelial cells, and neurons (6–8). Bim is a critical regulator for the development and normal responses of various immune systems. Bim deficiency causes abnormal

accumulation of lymphoid and myeloid cells. The phenotype of Bim^{-/-} mice is fatal systemic lupus erythematosus-like autoimmune disease. In the area of tissue development and homeostasis, Bim plays a key role in skeletal homeostasis, spermatogenesis, and mammary gland formation (9–11).

The Bim Regulatory System

The pro-apoptotic activity of Bim is controlled by transcriptional and post-transcriptional systems. The forkhead-like transcription factor FOXO3a (forkhead box O3a) and tumor suppressor gene *Runt3* (runt-related protein 3) are key transcriptional regulators of Bim. Cytokine withdrawal or other apoptotic stimuli cause the upregulation of Bim mRNA level through activation of FOXO3a in primary osteoblast, hepatocyte, neurons, and paclitaxel-treated cancer cells (7, 8, 12, 13).

Paclitaxel is used for node-positive early-stage breast cancer, metastatic breast cancer, and advanced ovarian cancer, interacting with cellular microtubules (13, 14). Paclitaxel upregulates the Bim EL expression level via increasing FOXO3a expression. MCF-7, a paclitaxel-sensitive breast cancer cell line, expresses high basal levels of FOXO3a. Paclitaxel treatment increases Bim protein dramatically without affecting the levels of other Bcl-2 family members. Thus, only the

Bim expression level correlated with apoptosis induction. However this was not observed in MDA-MB231 cells, which expressed low levels of FOXO3a and Bim. Therefore, paclitaxel sensitivity is controlled by FOXO3a and Bim expression levels (13).

Paclitaxel sensitivity correlates with non-small cell lung cancer (NSCLC) cell lines as well (15). Brain-derived neurotrophic factor (BDNF) is a biomarker of poor prognosis of neuroblastoma patients. BDNF provides chemoresistance to neuroblastoma via activation of MAPK and PI3K pathways. Interestingly, Bim was involved in paclitaxel but not etoposide- or cisplatin-induced neuroblastoma apoptosis. Paclitaxel-induced cell death via Bim expression is abrogated by BDNF. BDNF activates the PI3K/AKT pathway and suppresses FOXO3a activity (16). Thus, paclitaxel is a Bim-targeting agent. Other drugs are discussed below.

FOXO3a activity is suppressed through AKT-mediated phosphorylation. The deficiency of the tumor suppressor gene *RUNX3* evokes gastric mucosa hyperplasia, and around one half of human gastric cancer cells do not significantly express *RUNX3* (17). *RUNX3* is responsible for transforming growth factor- β (TGF- β)-induced gastric epithelial cells apoptosis. *RUNX3* upregulates Bim transcription with FOXO3a association and cells undergo TGF- β -induced apoptosis (18). TGF- β induces apoptosis signals not only through upregulation of Bim mRNA but also through the post-transcriptional pathway. It upregulates Bim through its inhibition of phosphorylation by extracellular signal-regulated kinase (ERK). ERK-mediated phosphorylation targets Bim for ubiquitylation and induces proteasomal degradation (7, 19). TGF- β transcriptionally induces the mitogen-activated protein kinase (MAPK) phosphatase (MKP2) through

SMAD3. MKP2 abrogates ERK activity and ERK-induced Bim ubiquitination. Thus, MKP2 induced by TGF- β upregulates Bim protein levels (20).

Growth factor-mediated stimulation can also activate ERK and downregulate Bim in a post-translational manner. On the contrary, JUN amino-terminal kinase (JNK)-mediated phosphorylation causes decreased binding of Bim to the anti-apoptotic protein Bcl2 and activates Bim as a result (19). In the T-cell acute lymphoblastic leukemia (T-ALL) cell line Sup-T, JNK-mediated phosphorylation of Bim promotes proteasomal Bim degradation (21). However, the role of JNK for Bim is still unclear.

Bim and Tumorigenesis

The breakdown of tissue homeostasis leads to various pathological situations including tumor formation. Thus, Bim plays important roles for tumorigenesis and tumor treatment (Fig. 3). Baby mouse kidney epithelial (BMK) cells transformed by E1A and dominant negative p53 (p53DD) form tumors in nude mice. The formation and growth of the tumors in nude mice are supported by Bim deficiency (22). These data suggest that Bim is possibly a key regulator of epithelial tumors as well. However, the absence of Bim plays a more important role for the formation of tumor metastasis and the acquisition of resistance to chemotherapy.

The Role of Bim in Metastasis

The acquisition of "anchorage-independence" is an indispensable step for tumor metastasis. Cells are normally dependent on anchorage and undergo apoptosis after their




Figure 2. The structural domains of BimEL. The schematic structure of BimEL is illustrated. Two sites of ubiquitin binding are shown at Lys-3 and Lys-108. Three sites of ERK phosphorylation are Ser-55, Ser-65, and Ser-73. One site of JNK phosphorylation is Thr-112. The BH3 domain, the dynein light chain 1 (DLC1) binding motif, and transmembrane domain are depicted.




Figure 3. Flow chart of Bim regulation in tumorigenesis and cancer treatment. Bim expression level controls the fate of the tumor cell. There are several transcriptional and post-transcriptional regulation points for Bim. The two main direct regulation systems are the ERK and AKT pathways. ERK phosphorylates Bim and promotes Bim degradation post-transcriptionally. AKT suppresses Bim transcription factor FOXO3a. Tumor cells mainly use ERK-proteasome axis in order to escape from apoptosis induced by Bim.

loss of attachment with their neighboring cells or their extracellular matrix. Therefore, apoptosis induced by cell detachment or “anoikis” is a crucial barrier against metastases. Anoikis stimuli can activate both the death receptor and mitochondrial pathways. Anoikis stimuli-activating mitochondrial pathways recruit or suppress several molecules, such as Bim, Bmf, Omi/HtrA2, Bcl-XL, and Mcl-1. Bim plays a key role in the anoikis of a variety of tumor cells, such as breast cancer, lung cancer, osteosarcoma, fibrosarcoma, and melanoma (23–25). These tumor cells have to bypass or abrogate Bim-mediated cell death one way or another in order to metastasize.

The cell detachment signal arises from the cell surface. Thus, cell surface molecules play important roles in anoikis. The cell surface glycoprotein mesothelin can activate a Bim suppression system via ERK activation in the human breast cancer cell line MDA-MB231 as well (24). A particularly interesting new cysteine-histidine-rich protein, (PINCH)-1, a cytoplasmic component of cell-extracellular matrix adhesions, can activate not only ERK-mediated Bim suppression but also the suppression of Bim transcriptionally (26).

In fact, several reports propose the relationship between various cell surface molecules and Bim. However, epidermal growth factor receptor (EGFR) is one of the most important cell surface molecules. In the human breast cancer cell line MCF-10A, Bim functions as a key regulator of anoikis

downstream of the EGFR-ERK pathway. Cell detachment attenuates the integrin signal. The integrin signal is indispensable for the maintenance of EGFR expression. Loss of integrin suppresses EGFR expression and inhibits ERK signaling. As a result, EGFR suppression by cell detachment upregulates Bim expression. Downregulation of Bim with RNA interference (RNAi) inhibits MCF-10A anoikis. Over-expressed EGFR maintains ERK activation after cell detachment and blocks Bim expression and anoikis (27).

Cell detachment-induced reactive oxygen species (ROS) are candidates upstream of the EGFR-ERK-Bim suppression system. Extracellular matrix-cell contact increases intracellular ROS through integrin. ROS oxidizes and activates tyrosine kinase Src. The active form of Src stimulates signaling downstream of EGFR including both AKT and ERK in a ligand-independent manner, culminating in the downregulation of Bim (28). Acting downstream to cell surface signals, both ERK and AKT play pivotal roles. The breast cancer cell line MDA-MB231 and HBC4 cells evade anoikis by the suppression of Bim with constitutive activation of the MEK-ERK pathway. MEK inhibitors sensitize these cell lines to anoikis by blocking proteasomal degradation of Bim_{EL} (29). In mutant MCF-10A cells expressing ΔRaf-ER, Raf-ERK signaling is activated by 4-OHT stimulation. Mutant MCF-10A cells show Bim downregulation and anoikis resistance with 4-OHT stimulation (30).

The 14-3-3 family is a critical regulator of various cellular responses such as proliferation, differentiation, cycling, apoptosis, and tumorigenesis. 14-3-3 ζ is a member of 14-3-3 family and forms a positive feedback loop with AKT. 14-3-3 ζ is highly expressed in NSCLC tissue and involved in anoikis resistance of lung cancer cells. 14-3-3 ζ knockdown causes the restitution of anchorage-dependence by Bim and Bad upregulation and Mcl-1 downregulation. As a result, lung cancer cells lose resistance to anoikis. Bim, but not Bad, can induce Bax activation and anoikis. Bim downregulation with RNAi abrogates restitution of anchorage-dependence by 14-3-3 ζ knockdown (31). B-Raf and N-Ras, commonly activated in human cutaneous melanoma, provide melanoma cells with resistance to anoikis. The constitutively active mutants of N-Ras and B-Raf downregulate Bim expression levels in melanocytes or melanoma cell lines via ERK activation. Bim suppression by the constitutively active mutants of N-Ras or by RNAi inhibits melanocyte anoikis (32, 33). Tumor metastasis is one of the critical determinants of prognosis. These data suggest that the Bim expression level is inversely related to prognosis. Indeed, reduced Bim expression is significantly correlated with poor prognosis of human cutaneous malignant melanoma (34). Currently, an increasing number of *in vitro* studies suggest that the ERK-Bim axis plays a pivotal role in tumor metastasis. Further analyses will reveal the role played by Bim in metastasis in patients.

The Role of Bim in Chemotherapy

Bim plays important roles not only in tumor metastasis but also in chemotherapy (Table 1). Bcr/Abl is a fusion oncogene formed by the fragment of Bcr and c-Abl, the product of the reciprocal t (q34; q22; refs. 9, 22) chromosomal translocation, the so-called Philadelphia (Ph1) chromosome. It is essential for leukemogenesis in chronic myelogenous leukemia (CML) and Ph1-positive acute lymphoblastic leukemia (ALL). Thus, Bcr/Abl was a good target candidate for novel CML or Ph1-positive ALL. Imatinib has dramatically improved clinical outcome for treatment of Bcr/Abl-positive leukemia, especially for early chronic phase CML patients. Bcr/Abl is an uncontrolled chimeric tyrosine kinase and can promote multimodal oncogenic function, including anti-apoptotic effects (35). Bim plays a pivotal role in the ability of imatinib to induce apoptosis. Imatinib induces the mito-

chondrial pathway apoptosis in Bcr/Abl-expressing cells. Imatinib increases Bim and Bmf transcription and activates Bim and Bad post-transcriptionally. Bim knockdown by RNAi abrogates imatinib-induced apoptosis in Bcr/Abl positive human leukemia cell lines (35). These Bim accumulations are attributed to inhibition of transcriptionally FoxO3a and post-transcriptional proteasomal degradation (36, 37). In K562 leukemia cells, Bim^{EL} phosphorylation and proteasome degradation is promoted by the ERK signal (36) or by beta-1 integrin attachment signal without ERK activation (38). In BaF3 Bcr/Abl-expressing cell lines, dephosphorylation of FoxO3a by imatinib promotes Bim transcription (37).

Dasatinib and nilotinib have been developed for imatinib-resistant leukemia, both of which can upregulate Bim (39, 40). The ERK pathway is important for Bim regulation by dasatinib as well. The MEK inhibitor PD184352 sensitizes the leukemia cell line K562 to dasatinib activity (41). Data derived from the use of the mouse DA1-3b BCR-ABL+ leukemic cell line suggest that Bim downregulation contributes to imatinib and dasatinib resistance as well. The imatinib- and dasatinib-resistant cell line secretes interleukin 3 (IL3) and is able to confer chemoresistance to nonmutated cells via a paracrine manner. IL3 activates both the MEK-ERK1/2 and the JAK2-STAT5 pathways. Both pathways can downregulate Bim, and JAK2 inhibition exhibits stronger Bim suppression. Thus, Bim, at least partially, controls the antitumor effects of imatinib and other related agents.

EGFR plays a critical role not only in anchorage-independence but also in tumorigenic features, such as uncontrollable cellular proliferation. Abnormal EGFR signaling is found in carcinomas of the lung, breast, and colon, making it a good target for chemotherapeutic agents. An EGFR-targeting agent, gefitinib, has been successful in the treatment of certain cancers, especially NSCLCs (42). Gefitinib evaluation for breast cancer is now at the phase II trial stage. Gefitinib causes rapid Bim accumulation in NSCLC cell lines through the blockade of the MEK/ERK pathway downstream of EGFR (42).

Erlotinib is another EGFR-targeting drug to treat lung cancer and shows survival improvement of NSCLC patients. Erlotinib upregulates Bim expression in lung cancer cell lines *in vitro* and in mouse xenograft models (43). Bim siRNA protects cells against apoptosis induced by both gefitinib and erlotinib. EGFR mutation is one of the adverse prognostic factors in gefitinib-treated patients

Table 1. Current Bim-targeted drugs

Agent	Tumor	Target gene or protein	Bim Regulation Signal
Paclitaxel Imatinib, dasatinib, and nilotinib	Breast cancer CML, Ph1(+) ALL	Microtubules Bcr/Abl	FOXO3a Proteasome degradation promoted by ERK signal, Integrin signal
Gefitinib, erlotinib	Breast, colon cancer, and NSCLC	EGFR	Proteasome degradation promoted by ERK signal
Bortezomib	Multiple myeloma	26S proteasome	Proteasome degradation

NOTE: Except for paclitaxel, all drugs target proteasome degradation terminally. Tumor cells may suppress Bim mainly through proteasome degradation.

with advanced lung adenocarcinoma. T790M mutation of EGFR abrogates Bim induction and subsequent apoptosis induced by gefitinib (44). Thus, these reports suggest that Bim is a terminal target of gefitinib and erlotinib.

Bortezomib (Velcade, PS341) is a 20S proteasome inhibitor used for treatment of multiple myeloma, and is currently in clinical trials enrolling patients suffering from various types of solid tumors (45). As mentioned above, Bim phosphorylation by ERK promotes Bim degradation through the proteasome-ubiquitination system. Therefore, bortezomib can be classified as a Bim-targeted tumor therapy agent. Although bortezomib activates Bim in order to terminate multiple myeloma, the activation system is not a direct inhibition of proteasome degradation of Bim as expected. Bortezomib upregulates the BH3 domain-only protein Noxa, not Bim. Accumulated Noxa dissociates Bim from Mcl-1, a member of the anti-apoptotic Bcl2 family. Dissociated Bim can activate the Bax/Bak complex and channel cells into apoptosis (46). The function of bortezomib as a Bim-targeting agent is basically as a sensitizer of other agents rather than as a direct tumor killer.

Bortezomib sensitizes prostate cancer cells to apoptosis through the extrinsic pathway by the death receptor ligand TRAIL. Downstream of TRAIL, Bim and Bik activation by bortezomib is important for apoptosis. Downregulation of Bim and Bik by RNAi impairs TRAIL-induced apoptosis. Interestingly, the TRAIL-induced apoptosis of breast cancer cell lines is not amplified by bortezomib (47). In the human prostate cancer cell lines PC3 and LNCaP-Pro5, apoptosis induced by bortezomib is not affected by Bim deficiency (48). However, bortezomib can abrogate paclitaxel-resistance induced by H-Ras. H-Ras promotes Bim phosphorylation and proteasome degradation of Bim upregulated by paclitaxel. Transformed BMK cells are sensitive to paclitaxel *in vitro* and *in vivo*, although Bim deficiency cancels this sensitivity. Bortezomib can abrogate H-Ras-promoted paclitaxel resistance (22). Thus, bortezomib can select cell types and enhance chemotherapeutic agents through Bim activation.

Future Directions in Developing Bim-Targeted Tumor Treatments

As a future direction, first we discuss the prospect of Bim protein transduction. Various tumors are immortalized and transformed with overexpression of anti-apoptotic Bcl-2 family proteins. Recently, plenty of BH3 mimetics have been developed and some of them show successful outcomes both *in vitro* and in clinical trials. One of the most potent BH3 mimetics so far is ABT-737. ABT-737 shows tumor suppression efficacy against small-cell lung carcinoma and various types of lymphoma and leukemia. However, both Mcl-1 and A1 overexpression cause ABT-737 resistance because the affinity of ABT-737 to Mcl-1 is weak. Mcl-1 is expressed in various tumors, so more work has to be done to increase coverage.

Bim can abrogate the functions of all of the anti-apoptotic Bcl-2 family proteins including Mcl-1 and A1 (49). Interestingly, in some tumors, Bim expression is inversely correlated to Mcl-1 expression (25, 31). Thus, Bim protein injection into cells

is a potent candidate of Bim-targeted tumor treatment, especially for tumors resistant to BH3 mimetics. Actually, it is reported that the TAT-Bim fusion protein can induce apoptosis of several cell lines, lymphoma, melanoma, and pancreatic cancer. TAT motif is a peptide transduction domain derived from the HIV-1. Thus, the modified Bim protein is a potent candidate of antitumor agent, although rigorous *in vivo* testing in clinically relevant models of neoplasia is required and forms the necessary next step in development in this area.

Bim downregulation is crucial not only for chemotherapy resistance but also for metastasis. The effects of some Bim-targeted agents can be restricted to the restitution of anchorage-dependence or chemotherapy sensitivity. This restriction can reduce side effects and provide possible new cocktail regimens of chemotherapeutic agents. On the other hand, the strong pro-apoptotic ability of Bim can abrogate all of the anti-apoptotic Bcl-2 molecules. This ability can make Bim-targeted agents act only as strong cytotoxins without tumor selectivity, and is a significant disadvantage.

Nevertheless, tumor-selective kill is a distinct possibility with Bim-targeted therapies. The diversity of Bim regulatory systems provides researchers tumor-selective Bim-targeted agents. Accordingly, correction of abnormal Bim-regulation systems abrogates tumorigenesis, metastatic ability, and chemoresistance. As previously explained above, Bim downregulation is mainly promoted by ERK phosphorylation and subsequent proteasomal degradation, although Bim is rigorously regulated by various transcriptional and post-transcriptional systems in normal mammalian cells. Thus, a restitution of the ubiquitination-proteasome Bim degradation system is the key to furnishing tumor selectivity. Furthermore, proteasome inhibitors such as bortezomib cannot target all Bim-downregulated tumors. This phenomenon suggests that the Bim proteasome degradation system is a cell-specific target. However, the search for proper targets in the ubiquitin-proteasome system is relatively young, and much more work needs to be done to furnish better options for therapy.

The specificity of ubiquitination-proteasome degradation is determined by distinct E3 ligase complexes. E3 ligase recognizes specific proteins and signals (Fig. 4). The proteasome complex is a common and ubiquitous system, although E3 ligase-mediated degradation of Bim is unique to Bim. Thus, if a Bim E3 ligase inhibitor would be developed, the drug could yield quite unique and tumor-selective effects. However, the intricate mechanisms of E3 ligase activity on Bim has not been revealed completely, so this notion has to be treated with caution.

Previously, involvement of c-Cbl in Bim ubiquitin-proteasome degradation was suggested, although recently adverse data have been reported (9, 50). The most potent candidate is an ElonginB/C-Cullin2-CIS E3 ligase complex. The receptor for activated C-kinase (RACK)-1 promotes the formation and activation of this E3 ligase complex and contributes to paclitaxel resistance of MDA-MB468 and MCF7 breast cancer cell lines *in vitro* and *in vivo* (51). This report suggests that RACK1 contributes to the paclitaxel resistance ability through the promotion of Bim degradation. Further analysis is required to reveal more precisely whether the Bim E3




Figure 4. Scheme of Bim ubiquitination. The process of ubiquitination and proteasome degradation is an ATP-dependent chain reaction. Ubiquitin is first conjugated to E1 and the ubiquitin moiety is then transferred to E2. Finally, ubiquitin is transferred to Bim via E3. Polyubiquitin-conjugated Bim is degraded by 26S proteasome. Proteasome degradation regenerates free ubiquitin for reuse through disassembling ubiquitin chains from Bim. Ub, ubiquitin; E1, ubiquitin-activating enzyme; E2, ubiquitin-conjugating enzymes; E3, ubiquitin ligases.

ligase mechanism, RACK1, and an ElonginB/C-Cullin2-CIS E3 ligase complex are potential targets for the development of tumor-selective Bim-targeted agents.

Conclusions

Physiologically, the pro-apoptotic BH3-only protein Bim plays a key role in induction of apoptosis of various type cells and development of some organs. Recently, the knowledge pertaining to the importance of Bim in cancer is increasing. Current reports suggest that Bim downregulation is important for tumorigenesis, especially for metastatic ability. The ERK-mediated proteasomal degradation system is frequently used for Bim downregulation in order for cancer cells to obtain metastatic ability. Chemotherapeutic agents, such as imatinib, gefitinib, and bortezomib activate Bim in order to kill tumor cells. These agents are, as it were, primitive Bim-targeted agents. Bim downregulation contributes to chemoresistance as well. A direct Bim protein transduction system is a potent candidate of tumor treatment because BH3 mimetics show successful tumor treatment data. Furthermore, Bim protein transduction may possibly overcome tumor resistance to BH3 mimetics.

A possible disadvantage of the application of drugs targeting Bim is the potential for a strong and broad range apoptosis induction. However, this point can be easily turned into an advantage, if Bim selectivity could be engineered into the candidate molecule. The potent advantage of Bim-targeting therapy is tumor cell selectivity because of the diversity of Bim controlling systems within cells. Furthermore, targeting of the Bim ubiquitination-proteasome degradation system, which is frequently used by tumor cells as an escape route from chemotherapeutic agents or anoikis, can provide tumor selectivity. Bim suppression is important not only for chemotherapy resistance but also for metastasis. The effects of some Bim-targeted agents are more a restitution of anchorage-dependence or chemotherapy sensitivity than as direct tumor killers. This potent feature may provide a new combination recipe for chemotherapeutic agents clinically. The area of Bim-targeted cancer treatment is really in an early phase of development. Further critical analysis of the Bim regulatory system is required in order to genuinely test the boundaries for such a not-so-novel concept.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

References

1. Adams JM, Cory S. The Bcl-2 apoptotic switch in cancer development and therapy. *Oncogene* 2007;26:1324–37.
2. Felsher DW. Oncogene addiction versus oncogene amnesia: perhaps more than just a bad habit? *Cancer Res* 2008;68:3081–6, discussion 6.
3. Pedraza-Farina LG. Mechanisms of oncogenic cooperation in cancer initiation and metastasis. *Yale J Biol Med* 2006;79:95–103.
4. Mehlen P, Puisieux A. Metastasis: a question of life or death. *Nat Rev Cancer* 2006;6:449–58.
5. Hengartner MO. The biochemistry of apoptosis. *Nature* 2000;407:770–6.
6. Youle RJ, Strasser A. The BCL-2 protein family: opposing activities that mediate cell death. *Nat Rev Mol Cell Biol* 2008;9:47–59.
7. Strasser A. The role of BH3-only proteins in the immune system. *Nat Rev Immunol* 2005;5:189–200.
8. Kawamura N, Kugiyama Y, Oshima Y, et al. Akt1 in osteoblasts and osteoclasts controls bone remodeling. *PLoS One* 2007;2:e1058.
9. Akiyama T, Bouillet P, Miyazaki T, et al. Regulation of osteoclast apoptosis by ubiquitylation of proapoptotic BH3-only Bcl-2 family member Bim. *EMBO J* 2003;22:6653–64.
10. Mailloux AA, Overholtzer M, Schmelzle T, Bouillet P, Strasser A, Brugge JS. BIM regulates apoptosis during mammary ductal morphogenesis, and its absence reveals alternative cell death mechanisms. *Dev Cell* 2007;12:221–34.
11. Coultas L, Bouillet P, Loveland KL, et al. Concomitant loss of proapoptotic BH3-only Bcl-2 antagonists Bik and Bim arrests spermatogenesis. *EMBO J* 2005;24:3963–73.
12. Barreyro FJ, Kobayashi S, Bronk SF, Werneburg NW, Malhi H, Gores GJ. Transcriptional regulation of Bim by FoxO3A mediates hepatocyte lipopapoptosis. *J Biol Chem* 2007;282:27141–54.
13. Sunters A, Fernandez de Mattos S, Stahl M, et al. FoxO3a transcriptional regulation of Bim controls apoptosis in paclitaxel-treated breast cancer cell lines. *J Biol Chem* 2003;278:49795–805.
14. Olsen SR. Taxanes and COX-2 inhibitors: from molecular pathways to clinical practice. *Biomed Pharmacother* 2005;59[Suppl 2]:S306–10.
15. Li R, Moudgil T, Ross HJ, Hu HM. Apoptosis of non-small-cell lung cancer cell lines after paclitaxel treatment involves the BH3-only proapoptotic protein Bim. *Cell Death Differ* 2005;12:292–303.
16. Li Z, Zhang J, Liu Z, Woo CW, Thiele CJ. Downregulation of Bim by brain-derived neurotrophic factor activation of TrkB protects neuroblastoma cells from paclitaxel but not etoposide or cisplatin-induced cell death. *Cell Death Differ* 2007;14:318–26.
17. Li QL, Ito K, Sakakura C, et al. Causal relationship between the loss of RUNX3 expression and gastric cancer. *Cell* 2002;109:113–24.
18. Yano T, Ito K, Fukamachi H, et al. The RUNX3 tumor suppressor up-regulates Bim in gastric epithelial cells undergoing transforming growth factor β -induced apoptosis. *Mol Cell Biol* 2006;26:4474–88.
19. Hubner A, Barrett T, Flavell RA, Davis RJ. Multisite phosphorylation regulates Bim stability and apoptotic activity. *Mol Cell* 2008;30:415–25.
20. Ramesh S, Qi XJ, Wilder GM, et al. TGF β -mediated BIM expression and apoptosis are regulated through SMAD3-dependent expression of the MAPK phosphatase MKP2. *EMBO Rep* 2008;9:990–7.
21. Leung KT, Li KK, Sun SS, Chan PK, Ooi VE, Chiu LC. Activation of the JNK pathway promotes phosphorylation and degradation of BimEL—a novel mechanism of chemoresistance in T-cell acute lymphoblastic leukemia. *Carcinogenesis* 2008;29:544–51.
22. Tan TT, Degenhardt K, Nelson DA, et al. Key roles of BIM-driven apoptosis in epithelial tumors and rational chemotherapy. *Cancer Cell* 2005;7:227–38.
23. Simpson CD, Anyiwe K, Schimmer AD. Anoikis resistance and tumor metastasis. *Cancer Lett* 2008;272:177–85.
24. Uehara N, Matsuoka Y, Tsubura A. Mesothelin promotes anchorage-independent growth and prevents anoikis via extracellular signal-regulated kinase signaling pathway in human breast cancer cells. *Mol Cancer Res* 2008;6:186–93.
25. Woods NT, Yamaguchi H, Lee FY, Bhalla KN, Wang HG. Anoikis, initiated by Mcl-1 degradation and Bim induction, is deregulated during oncogenesis. *Cancer Res* 2007;67:10744–52.
26. Chen K, Tu Y, Zhang Y, Blair HC, Zhang L, Wu C. PINCH-1 regulates the ERK-Bim pathway and contributes to apoptosis resistance in cancer cells. *J Biol Chem* 2008;283:2508–17.
27. Reginato MJ, Mills KR, Paulus JK, et al. Integrins and EGFR coordinately regulate the pro-apoptotic protein Bim to prevent anoikis. *Nat Cell Biol* 2003;5:733–40.
28. Giannoni E, Buricchi F, Grimaldi G, et al. Redox regulation of anoikis: reactive oxygen species as essential mediators of cell survival. *Cell Death Differ* 2008;15:867–78.
29. Fukazawa H, Noguchi K, Masumi A, Murakami Y, Uehara Y. BimEL is an important determinant for induction of anoikis sensitivity by mitogen-activated protein/extracellular signal-regulated kinase inhibitors. *Mol Cancer Ther* 2004;3:1281–8.
30. Marani M, Hancock D, Lopes R, Tenev T, Downward J, Lemoine NR. Role of Bim in the survival pathway induced by Raf in epithelial cells. *Oncogene* 2004;23:2431–41.
31. Li Z, Zhao J, Du Y, et al. Down-regulation of 14–3–3 ζ suppresses anchorage-independent growth of lung cancer cells through anoikis activation. *Proc Natl Acad Sci U S A* 2008;105:162–7.
32. Boisvert-Adamo K, Aplin AE. Mutant B-RAF mediates resistance to anoikis via Bad and Bim. *Oncogene* 2008;27:3301–12.
33. Goldstein NB, Johannes WU, Gadeliya AV, et al. Active N-Ras and B-Raf Inhibit anoikis by downregulating bim expression in melanocytic cells. *J Invest Dermatol* 2009;129:432–7.
34. Dai DL, Wang Y, Liu M, Martinka M, Li G. Bim expression is reduced in human cutaneous melanomas. *J Invest Dermatol* 2008;128:403–7.
35. Kuroda J, Puthalakath H, Cragg MS, et al. Bim and Bad mediate imatinib-induced killing of Bcr/Abl+ leukemic cells, and resistance due to their loss is overcome by a BH3 mimetic. *Proc Natl Acad Sci U S A* 2006;103:14907–12.
36. Luciano F, Jacquel A, Colosetti P, et al. Phosphorylation of Bim-EL by Erk1/2 on serine 69 promotes its degradation via the proteasome pathway and regulates its proapoptotic function. *Oncogene* 2003;22:6785–93.
37. Essafi A, Fernandez de Mattos S, Hassen YA, et al. Direct transcriptional regulation of Bim by FoxO3a mediates STI571-induced apoptosis in Bcr-Abl-expressing cells. *Oncogene* 2005;24:2317–29.
38. Hazlehurst LA, Argilagos RF, Dalton WS. Beta1 integrin mediated adhesion increases Bim protein degradation and contributes to drug resistance in leukaemia cells. *Br J Haematol* 2007;136:269–75.
39. Liu J, Joha S, Idziorek T, et al. BCR-ABL mutants spread resistance to non-mutated cells through a paracrine mechanism. *Leukemia* 2008;22:791–9.
40. Belloc F, Moreau-Gaudry F, Uhalde M, et al. Imatinib and nilotinib induce apoptosis of chronic myeloid leukemia cells through a Bim-dependant pathway modulated by cytokines. *Cancer Biol Ther* 2007;6:912–9.
41. Nguyen TK, Rahmani M, Harada H, Dent P, Grant S. MEK1/2 inhibitors sensitize Bcr/Abl+ human leukemia cells to the dual Abl/Src inhibitor BMS-354/825. *Blood* 2007;109:4006–15.
42. Cragg MS, Kuroda J, Puthalakath H, Huang DC, Strasser A. Gefitinib-induced killing of NSCLC cell lines expressing mutant EGFR requires BIM and can be enhanced by BH3 mimetics. *PLoS Med* 2007;4:1681–89; discussion 90.
43. Gong Y, Somwar R, Politi K, et al. Induction of BIM is essential for apoptosis triggered by EGFR kinase inhibitors in mutant EGFR-dependent lung adenocarcinomas. *PLoS Med* 2007;4:e294.
44. Costa DB, Halmos B, Kumar A, et al. BIM mediates EGFR tyrosine kinase inhibitor-induced apoptosis in lung cancers with oncogenic EGFR mutations. *PLoS Med* 2007;4:1669–79; discussion 80.
45. Fennell DA, Chacko A, Mutti L. BCL-2 family regulation by the 20S proteasome inhibitor bortezomib. *Oncogene* 2008;27:1189–97.
46. Gomez-Bougie P, Wulleme-Toumi S, Menoret E, et al. Noxa up-regulation and Mcl-1 cleavage are associated to apoptosis induction by bortezomib in multiple myeloma. *Cancer Res* 2007;67:5418–24.
47. Nikrad M, Johnson T, Puthalath H, Coultas L, Adams J, Kraft AS. The proteasome inhibitor bortezomib sensitizes cells to killing by death receptor ligand TRAIL via BH3-only proteins Bik and Bim. *Mol Cancer Ther* 2005;4:443–9.
48. Canfield SE, Zhu K, Williams SA, McConkey DJ. Bortezomib inhibits docetaxel-induced apoptosis via a p21-dependent mechanism in human prostate cancer cells. *Mol Cancer Ther* 2006;5:2043–50.
49. Zhang L, Ming L, Yu J. BH3 mimetics to improve cancer therapy; mechanisms and examples. *Drug Resist Updat* 2007;10:207–17.
50. Wiggins CM, Band H, Cook SJ. c-Cbl is not required for ERK1/2-dependent degradation of BimEL. *Cell Signal* 2007;19:2605–11.
51. Zhang W, Cheng GZ, Gong J, et al. RACK1 and CIS mediate the degradation of BimEL in cancer cells. *J Biol Chem* 2008;283:16416–26.