



BMI and Coronary Heart Disease Risk Among Low-Income and Underinsured Diabetic Patients

Nan Li,^{1,2} Peter T. Katzmarzyk,¹
 Ronald Horswell,¹ Yonggang Zhang,¹
 Weiqin Li,^{1,2} Wenhui Zhao,¹ Yujie Wang,¹
 Jolene Johnson,³ and Gang Hu¹

Diabetes Care 2014;37:3204–3212 | DOI: 10.2337/dc14-1091

OBJECTIVE

The association between obesity and coronary heart disease (CHD) risk remains debatable, and no studies have assessed this association among diabetic patients. The aim of our study was to investigate the association between BMI and CHD risk among patients with type 2 diabetes.

RESEARCH DESIGN AND METHODS

The sample included 30,434 diabetic patients (10,955 men and 19,479 women) 30–95 years of age without a history of CHD or stroke in the Louisiana State University Hospital-Based Longitudinal Study.

RESULTS

During a mean follow-up period of 7.3 years, 7,414 subjects developed CHD. The multivariable-adjusted hazard ratios for CHD across levels of BMI at baseline (18.5–24.9, 25–29.9, 30–34.9, 35–39.9, and ≥ 40 kg/m²) were 1.00, 1.14 (95% CI 1.00–1.29), 1.27 (1.12–1.45), 1.54 (1.34–1.78), and 1.42 (1.23–1.64) ($P_{\text{trend}} < 0.001$) in men and 1.00, 0.95 (0.85–1.07), 0.95 (0.84–1.06), 1.06 (0.94–1.20), and 1.09 (1.00–1.22) ($P_{\text{trend}} < 0.001$) in women, respectively. When we used an updated mean or last visit value of BMI, the positive association between BMI and CHD risk did not change in men. However, the positive association of BMI with CHD changed to a U-shaped association in women when we used the last visit value of BMI.

CONCLUSIONS

Our study suggests that there is a positive association between BMI at baseline and during follow-up with the risk of CHD among patients with type 2 diabetes. We indicate a U-shaped association between BMI at the last visit and the risk of CHD among women with type 2 diabetes.

Obesity and diabetes are two important public health problems in the U.S. (1). Two in three adults in the U.S. are currently classified as overweight or obese (BMI ≥ 25 kg/m²), and one-third of them are frankly obese (BMI ≥ 30 kg/m²) (1). The estimated number of adults with diabetes in the U.S. is 26.1 million in 2005–2010 or $\sim 12\%$ of the population (2). Among U.S. diabetic patients, the prevalence of overweight or obesity has increased to $\geq 80\%$ (3). Obesity is associated with increased risks of several cardiometabolic diseases, including hypertension (4), diabetes (5), coronary heart disease (CHD) (6,7), heart failure (8), and stroke (9).

Cardiovascular diseases, especially CHD, are the leading causes of death worldwide. In recent years, several prospective studies have assessed the association

¹Pennington Biomedical Research Center, Baton Rouge, LA

²Tianjin Women's and Children's Health Center, Tianjin, China

³Louisiana State University Health, Baton Rouge, LA

Corresponding author: Gang Hu, gang.hu@pbrc.edu.

Received 30 April 2014 and accepted 25 August 2014.

© 2014 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered.

between obesity and the risks of total and CVD mortality among diabetic patients, and the results are inconsistent. To date, many studies have reported positive associations (10,11), inverse associations (12–14), U-shaped associations (15–17), or no associations (18) between BMI and mortality among patients with diabetes. All of these studies were focused on the association between BMI and CVD mortality; however, no studies assessed the association between BMI and the risk of incident CHD among diabetic patients. In this study, we examined the association between BMI and the risk of CHD among patients with type 2 diabetes in the Louisiana State University Hospital-Based Longitudinal Study (LSUHLS).

RESEARCH DESIGN AND METHODS

Study Population

Between 1997 and 2012, the LSU Health Care Services Division (LSUHCS) operated seven public hospitals and affiliated clinics in Louisiana providing quality medical care to the residents of Louisiana regardless of their income or insurance coverage (19–26). Overall, LSUHCS facilities have served ~1.6 million patients (35% of the Louisiana population) since 1997. Administrative, anthropometric, laboratory, clinical diagnosis, and medication data collected at these facilities are available in electronic form for both inpatients and outpatients from 1997. Using these data, we have established the LSUHLS (19). A cohort of diabetic patients was established by using the ICD-9 (code 250) between 1 January 1999 and 31 December 2009. Both inpatients and outpatients were included, and all patients were under primary care. Confirmation of diabetes diagnoses was made by applying the American Diabetes Association criteria: a fasting plasma glucose level ≥ 126 mg/dL; 2-h glucose level ≥ 200 mg/dL after a 75-g 2-h oral glucose tolerance test; and one or more classic symptoms plus a random plasma glucose level ≥ 200 mg/dL (27,28). The first record of diabetes diagnosis was used to establish the baseline for each patient in the present analyses due to the design of the cohort study. Before diagnosis with diabetes, these patients have used the LSU system for an average 5.0 years. We have validated the diabetes diagnosis in LSUHCS hospitals. The

agreement of diabetes diagnosis was 97%: 20,919 of a sample of 21,566 hospital discharge diagnoses based on ICD codes also had physician-confirmed diabetes by using the American Diabetes Association diabetes diagnosis criteria (27).

The current study included 30,434 newly diagnosed patients (10,955 men and 19,479 women) with type 2 diabetes who were 30–95 years of age without a history of CHD and stroke at the time of diabetes diagnosis and with complete repeated data on all risk factor variables. We only included African Americans and whites because the patient numbers of Hispanics, Asians, and Native Americans are very small in the LSUHCS hospitals. Patients were excluded if they were underweight (BMI < 18.5 kg/m²) because of limited statistical power for this group. Compared with diabetic patients excluded from the present analyses due to missing data, those included in the analyses were younger (51.0 vs. 57.6 years old), had a higher frequency of African Americans (58.6 vs. 45.4%), and less males (37.2 vs. 47.1%). The study and analysis plan was approved by the Pennington Biomedical Research Center and LSU Health Sciences Center Institutional Review Boards, LSU System. We did not obtain informed consent from participants involved in our study because we used anonymized data compiled from electronic medical records.

Baseline Measurements

The patient's characteristics, including age at diabetes diagnosis, sex, race/ethnicity, family income, smoking status, types of health insurance, BMI, blood pressure, total cholesterol, HDL cholesterol, LDL cholesterol, triglycerides, glycosylated hemoglobin (HbA_{1c}), estimated glomerular filtration rate (eGFR), and medication (antihypertensive drug, cholesterol-lowering drug, and antidiabetes drug) within a half year before the diabetes diagnosis (baseline), during follow-up after the diabetes diagnosis (follow-up), and the last visit were extracted from the computerized hospitalization records. Height and weight were measured without shoes and with light clothing according to a standardized protocol. BMI was calculated by dividing weight in kilograms by the square of

height in meters. Values of BMI, blood pressure, HbA_{1c}, LDL cholesterol, and eGFR over time were measured firstly at baseline and secondly as an updated mean of annual measurements, calculated for each participant from baseline to each year of follow-up. For example, at 1 year, the updated mean is the average of the baseline and 1-year values, and at 3 years, it is the average of baseline, 1-year, 2-year, and 3-year values. In the case of an event during follow-up, the period for estimating updated mean values was from baseline to the year before this event occurred (29). The average number of BMI measurements during the follow-up period was 15.0.

Prospective Follow-up

Follow-up information was obtained from the LSUHLS inpatient and outpatient database by using the unique number assigned to every patient who visits the LSUHCS hospitals. The diagnosis of CHD was the primary end point of interest of the study and defined according to the following ICD-9: CHD (ICD-codes 410–414). Follow-up of each cohort member continued until the date of the diagnosis of CHD, the date of the last visit if the subject stopped use of LSUHCS hospitals, or the date of death, determined from linking to the Louisiana Office of Public Health Vital Records Registry, or 31 May 2012 (23,26).

Statistical Analyses

The association between BMI and the risk of CHD was analyzed by using Cox proportional hazards models. BMI was evaluated in the following two ways: 1) as five weight categories (18.5–24.9 [reference group], 25–29.9, 30–34.9, 35–39.9, and ≥ 40 kg/m²) and 2) as a continuous variable. The trend over different categories of BMI was tested in models with the median of each category as a continuous variable. All analyses were adjusted for age (continuous variable) and race (African American and white) (model 1) and further for smoking (never, past, and current), income (continuous variable), and types of insurance (free, self-pay, Medicaid, Medicare, and commercial) (model 2), and additionally for systolic blood pressure (continuous variable), HbA_{1c} (continuous variable), LDL cholesterol

(continuous variable), HDL cholesterol (continuous variable), triglycerides (continuous variable), eGFR (≥ 90 , 60–89, 30–59, 15–29, and < 15 mL/min/1.73 m²), use of antihypertensive drugs (no use, ACE inhibitor, angiotensin II receptor blockers, β -blockers, calcium channel blocker, diuretics, and other antihypertensive drugs), use of diabetes medications (no use, oral hypoglycemic agents, and insulin), and use of cholesterol-lowering agents (no use, statins, and other cholesterol-lowering agents) (model 3). We stratified the samples by sex because there was a significant interaction between sex and BMI on the risk of CHD. Since the interactions between race and BMI on the risk of CHD were not statistically significant, data for white and African Americans were combined in some analyses. Statistical significance was considered to be $P < 0.05$. All statistical analyses were performed with PASW

for Windows, version 20.0 (IBM SPSS Inc., Chicago, IL).

RESULTS

General characteristics of the study population at baseline are presented in Table 1.

During a mean follow-up period of 7.3 years, 7,414 subjects (2,926 men and 4,488 women) developed CHD. Patients who developed CHD during follow-up were older and used more glucose-lowering, lipid-lowering, and antihypertensive medication compared with those who did not develop CHD.

The multivariable-adjusted (age, race, smoking, income, and types of insurance) (model 2) hazard ratios (HRs) for CHD at different levels of BMI at baseline (18.5–24.9 [reference group], 25–29.9, 30–34.9, 35–39.9, and ≥ 40 kg/m²) were 1.00, 1.14 (95% CI 1.00–1.29), 1.27 (1.12–1.45), 1.54 (1.34–

1.78), and 1.42 (1.23–1.64) ($P_{\text{trend}} < 0.001$) in men and 1.00, 0.95 (0.85–1.07), 0.95 (0.84–1.06), 1.06 (0.94–1.20), and 1.09 (1.00–1.22) ($P_{\text{trend}} < 0.001$) in women, respectively (Table 2). After further adjustment for other confounding factors (systolic blood pressure, HbA_{1c}, LDL cholesterol, HDL cholesterol, triglycerides, eGFR, use of antihypertensive drugs, use of diabetes medications, and use of cholesterol-lowering agents), this association remained significant among men ($P_{\text{trend}} < 0.001$) and women ($P_{\text{trend}} = 0.006$).

When BMI was examined as a continuous variable, the multivariable-adjusted (model 2) HRs of CHD for each one-unit increase in BMI at baseline were 1.015 (95% CI 1.011–1.020) in men and 1.004 (95% CI 1.001–1.008) in women (Table 2). There was a significant interaction between sex and BMI on CHD risk ($\chi^2 = 9.86$; 1df, $P < 0.005$), which indicated that this positive

Table 1—Baseline characteristics of patients with type 2 diabetes by the outcome during follow-up^a

	Men			Women		
	No CHD	CHD	<i>P</i> value	No CHD	CHD	<i>P</i> value
Number of participants	8,029	2,926		14,991	4,488	
Age, mean (SE) (years)	50.2 (0.11)	54.1 (0.18)	<0.001	51.1 (0.08)	53.9 (0.15)	<0.001
Income, mean (SE) (\$/family)	18,935 (319)	22,207 (521)	<0.001	18,850 (203)	20,273 (371)	<0.001
BMI at baseline, mean (SE) (kg/m ²)	32.5 (0.09)	33.3 (0.14)	<0.001	35.6 (0.07)	35.7 (0.12)	0.36
BMI during follow-up, mean (SE) (kg/m ²)	32.3 (0.08)	33.2 (0.13)	<0.001	35.5 (0.07)	35.7 (0.12)	0.18
BMI at last visit, mean (SE) (kg/m ²)	32.2 (0.09)	33.0 (0.14)	<0.001	35.3 (0.07)	35.4 (0.13)	0.42
Race, <i>N</i> (%)			<0.001			<0.001
African American	4,731 (58.9)	1,369 (46.8)		9,168 (61.2)	2,380 (53.0)	
White	3,298 (41.1)	1,557 (53.2)		5,823 (38.8)	2,108 (47.0)	
HbA _{1c} , mean [% (mmol/mol)]	8.02 (64)	7.99 (64)	0.59	7.51 (59)	7.71 (61)	<0.001
HDL cholesterol, mean (SE) (mg/dL)	39.5 (0.1)	38.8 (0.2)	0.006	46.0 (0.1)	45.0 (0.2)	<0.001
LDL cholesterol, mean (SE) (mg/dL)	109 (0.5)	105 (0.8)	<0.001	116 (0.3)	113 (0.6)	<0.001
Triglycerides, mean (SE) (mg/dL)	151 (1.0)	160 (1.7)	<0.001	140 (0.6)	149 (1.2)	<0.001
GFR, <i>N</i> (%) (mL/min/1.73 m ²)			<0.001			<0.001
≥ 90	4,261 (53.1)	1,215 (41.5)		7,256 (48.4)	1,824 (40.7)	
60–89	2,977 (37.1)	1,208 (41.3)		6,027 (40.2)	1,820 (40.5)	
30–59	656 (8.2)	430 (14.7)		1,539 (10.3)	752 (16.8)	
15–29	84 (1.0)	52 (1.8)		113 (0.7)	73 (1.6)	
< 15	51 (0.6)	21 (0.7)		56 (0.4)	19 (0.4)	
Current smoker, <i>N</i> (%)	2,930 (36.5)	987 (33.7)	0.014	3,808 (25.4)	1,178 (26.3)	0.46
Types of insurance, <i>N</i> (%)			<0.001			<0.001
Free	5,999 (74.7)	1,969 (67.3)		12,559 (83.8)	3,399 (75.7)	
Self-pay	698 (8.7)	141 (4.8)		601 (4.0)	125 (2.8)	
Medicaid	368 (4.6)	150 (5.1)		692 (4.6)	325 (7.3)	
Medicare	732 (9.1)	591 (20.2)		844 (5.6)	561 (12.5)	
Commercial	232 (2.9)	75 (2.6)		295 (2.0)	78 (1.7)	
Uses of medications, <i>N</i> %						
Glucose-lowering medication	5,196 (64.7)	1,989 (68.0)	<0.001	9,730 (64.9)	3,085 (68.7)	<0.001
Lipid-lowering medication	3,852 (48.0)	1,879 (64.2)	<0.001	8,188 (54.6)	2,994 (66.7)	<0.001
Antihypertensive medication	5,495 (68.4)	2,183 (74.6)	<0.001	11,017 (73.5)	3,484 (77.6)	<0.001

^aData represent means or percentages. All data except age are adjusted for age and race.

Table 2—HRs of CHD according to different levels of BMI at baseline, during follow-up, and at last visit among patients with type 2 diabetes

	BMI (kg/m ²)					P for trend	Each 1-kg/m ² increase
	<25.0	25.0–29.9	30–34.9	35–39.9	≥40		
Baseline							
Men	1,561	3,008	2,948	1,789	1,649		
Number of cases	350	774	811	540	451		
Person-years	11,085	21,192	19,990	11,382	10,889		
Adjustment	HR (95% CI)						
Model 1 ^a	1.00	1.12 (0.98–1.27)	1.24 (1.09–1.41)	1.50 (1.31–1.72)	1.39 (1.20–1.61)	<0.001	1.015 (1.010–1.020)
Model 2 ^b	1.00	1.14 (1.00–1.29)	1.27 (1.12–1.45)	1.54 (1.34–1.78)	1.42 (1.23–1.64)	<0.001	1.015 (1.011–1.020)
Model 3 ^c	1.00	1.16 (1.00–1.33)	1.24 (1.07–1.43)	1.47 (1.26–1.72)	1.45 (1.24–1.70)	<0.001	1.015 (1.009–1.020)
Women	1,681	3,873	4,719	3,968	5,238		
Number of cases	420	904	1,054	938	1,172		
Person-years	12,664	30,032	36,548	30,102	39,538		
Adjustment	HR (95% CI)						
Model 1 ^a	1.00	0.93 (0.83–1.05)	0.92 (0.82–1.03)	1.03 (0.91–1.16)	1.05 (0.94–1.18)	0.010	1.004 (1.000–1.007)
Model 2 ^b	1.00	0.95 (0.85–1.07)	0.95 (0.84–1.06)	1.06 (0.94–1.20)	1.09 (1.00–1.22)	<0.001	1.004 (1.001–1.008)
Model 3 ^c	1.00	0.93 (0.82–1.06)	0.92 (0.82–1.05)	1.02 (0.90–1.16)	1.07 (0.95–1.22)	0.006	1.005 (1.001–1.009)
Follow-up							
Men	1,494	3,108	3,028	1,797	1,528		
Number of cases	335	814	813	529	435		
Person-years	10,691	21,876	20,542	11,629	9,806		
Adjustment	HR (95% CI)						
Model 1 ^a	1.00	1.14 (1.00–1.30)	1.22 (1.07–1.39)	1.49 (1.30–1.72)	1.50 (1.29–1.74)	<0.001	1.017 (1.012–1.023)
Model 2 ^b	1.00	1.16 (1.02–1.32)	1.26 (1.11–1.44)	1.53 (1.33–1.77)	1.55 (1.33–1.80)	<0.001	1.018 (1.013–1.023)
Model 3 ^c	1.00	1.18 (1.02–1.36)	1.24 (1.07–1.44)	1.45 (1.24–1.69)	1.55 (1.31–1.82)	<0.001	1.017 (1.011–1.023)
Women	1,579	3,897	4,874	3,965	5,164		
Number of cases	388	905	1,086	948	1,161		
Person-years	12,098	30,034	37,749	29,862	39,139		
Adjustment	HR (95% CI)						
Model 1 ^a	1.00	0.97 (0.86–1.10)	0.95 (0.84–1.07)	1.10 (0.98–1.25)	1.10 (0.98–1.24)	<0.001	1.004 (1.001–1.008)
Model 2 ^b	1.00	0.99 (0.88–1.12)	0.98 (0.87–1.11)	1.14 (1.01–1.29)	1.14 (1.01–1.29)	<0.001	1.006 (1.002–1.009)
Model 3 ^c	1.00	1.00 (0.88–1.15)	1.00 (0.88–1.14)	1.12 (0.98–1.28)	1.12 (0.99–1.28)	0.003	1.005 (1.000–1.009)
Last visit							
Men	1,715	3,064	2,892	1,713	1,571		
Number of cases	406	794	782	494	450		
Person-years	12,241	21,431	19,682	11,108	10,076		
Adjustment	HR (95% CI)						
Model 1 ^a	1.00	1.12 (0.99–1.27)	1.20 (1.06–1.36)	1.44 (1.25–1.64)	1.48 (1.28–1.70)	<0.001	1.016 (1.011–1.020)
Model 2 ^b	1.00	1.15 (1.01–1.30)	1.25 (1.10–1.41)	1.49 (1.30–1.70)	1.53 (1.32–1.76)	<0.001	1.016 (1.011–1.021)
Model 3 ^c	1.00	1.11 (0.97–1.26)	1.19 (1.04–1.37)	1.33 (1.15–1.54)	1.46 (1.25–1.70)	<0.001	1.015 (1.009–1.020)
Women	1,891	3,989	4,708	3,837	5,054		
Number of cases	502	907	1,039	913	1,127		
Person-years	14,259	30,874	36,346	28,962	38,442		
Adjustment	HR (95% CI)						
Model 1 ^a	1.00	0.87 (0.78–0.97)	0.87 (0.78–0.97)	1.01 (0.90–1.13)	1.00 (0.90–1.12)	0.022	1.004 (1.000–1.007)
Model 2 ^b	1.00	0.89 (0.80–1.00)	0.90 (0.81–1.00)	1.05 (0.93–1.17)	1.04 (0.93–1.17)	0.003	1.004 (1.001–1.008)
Model 3 ^c	1.00	0.90 (0.80–1.02)	0.91 (0.81–1.03)	1.03 (0.92–1.17)	1.00 (0.89–1.13)	0.085	1.003 (0.999–1.006)

^aAdjusted for age and race. ^bAdjusted for age, race, types of insurance, income, and smoking. ^cAdjusted for age, race, types of insurance, income, smoking, systolic blood pressure, LDL cholesterol, HDL cholesterol, triglycerides, HbA_{1c}, eGFR, use of antihypertensive drugs (none, ACE inhibitor, angiotensin II receptor blockers, β-blockers, calcium channel blocker, diuretics, other antihypertensive drugs, and any two or more of above treatments), glucose-lowering agents (none, oral hypoglycemic agents, and insulin), and cholesterol-lowering agents (none, statins, and other cholesterol-lowering agents).

association was stronger in men than in women.

When we did an additional analysis by using an updated mean value of BMI, we

found the same positive association between BMI and CHD risk among both men ($P_{\text{trend}} < 0.001$) and women ($P_{\text{trend}} < 0.001$) (Table 2). When we did another

additional analysis by using the last visit value of BMI, we found a positive association between BMI and CHD risk among men ($P_{\text{trend}} < 0.001$) and a

U-shaped association between BMI and CHD risk among women ($P_{\text{trend}} = 0.003$) (Table 2). Women who were overweight and had class I obesity (BMI 25–34.9 kg/m²) at last visit had a lower risk of CHD compared with normal-weight women (BMI <25 kg/m²).

After excluding subjects who were diagnosed with CHD during the first 2 years of follow-up ($n = 3,207$), the multivariable-adjusted HRs (model 2) of CHD for each one-unit increase in BMI at baseline, during follow-up, and at the last visit were 1.014 (95% CI 1.010–1.019), 1.017 (1.012–1.022), and 1.015 (1.009–1.019) in men and 1.005 (1.001–1.009), 1.006 (1.002–1.010), and 1.005 (1.000–1.008) in women (data not shown), respectively.

In stratified analyses, the multivariable-adjusted positive association between BMI and CHD risk was present among

men with different smoking status (Tables 3 and 4). When stratified by age, race, and use of antidiabetic drugs, this positive association of BMI at baseline, during follow-up, and at the last visit with CHD risk was still present among men in all subgroups and among women in some of the subgroups (Tables 3 and 4).

CONCLUSIONS

Our study found a positive association of BMI at baseline and during follow-up with the risk of CHD among both men and women with type 2 diabetes, and this association was stronger among men than among women. In addition, we found that a positive association between BMI and the risk of CHD was present in both African Americans and whites with type 2 diabetes and in nonsmokers and smokers.

The positive association did not change among men but changed to a U-shaped association among women with type 2 diabetes when we assessed BMI of the last visit with CHD risk.

Only a few prospective studies have evaluated the association between obesity and total and CVD mortality among diabetic patients, and the results are controversial including inverse associations (12–14), positive associations (10,11), U-shaped associations (15–17), or no association (18). The current study was the first, to our knowledge, to assess the association between BMI and the risk of incident CHD among diabetic patients. The results of our study indicated a positive association between BMI and the risk of CHD among patients with type 2 diabetes. We found this positive association of CHD risk by

Table 3—HRs (95% CIs) of CHD according to different levels of BMI at baseline among various subpopulations^a

	BMI (kg/m ²)					P for trend	P for interaction
	<25.0	25.0–29.9	30–34.9	35–39.9	≥40		
Men							
Age groups (years)							>0.50
<50	1.00	1.07 (0.86–1.33)	1.19 (0.96–1.48)	1.47 (1.19–1.85)	1.44 (1.15–1.80)	<0.001	
50–59	1.00	1.28 (1.01–1.60)	1.48 (1.17–0.85)	1.57 (1.23–2.00)	1.56 (1.22–2.01)	<0.001	
≥60	1.00	1.01 (0.8–1.27)	1.07 (0.84–1.35)	1.42 (1.10–1.85)	0.83 (0.59–1.15)	0.38	
Race							>0.10
African American	1.00	1.18 (0.99–1.40)	1.26 (1.05–1.51)	1.54 (1.26–1.88)	1.53 (1.24–1.89)	<0.001	
White	1.00	1.08 (0.89–1.32)	1.27 (1.05–1.54)	1.54 (1.26–1.88)	1.35 (1.10–1.66)	<0.001	
Smoking status							<0.05
Never	1.00	1.17 (0.95–1.44)	1.35 (1.10–1.67)	1.68 (1.35–2.09)	1.70 (1.36–2.12)	<0.001	
Ever or current	1.00	1.33 (1.09–1.62)	1.41 (1.15–1.73)	1.61 (1.28–2.03)	1.38 (1.08–1.76)	<0.001	
Glucose-lowering medication							>0.25
No use	1.00	1.56 (1.19–2.06)	1.40 (1.05–1.88)	1.66 (1.20–2.29)	1.85 (1.33–2.56)	0.001	
Oral hypoglycemic agents	1.00	1.25 (0.95–1.65)	1.46 (1.11–1.92)	1.66 (1.23–2.23)	1.57 (1.15–2.14)	<0.001	
Metformin	1.00	1.08 (0.87–1.34)	1.31 (1.06–1.62)	1.56 (1.24–1.95)	1.47 (1.16–1.85)	<0.001	
Sulfonylureas	1.00	1.04 (0.83–1.32)	1.24 (0.98–1.56)	1.47 (1.15–1.88)	1.32 (1.02–1.71)	<0.001	
Other oral agents	1.00	1.13 (0.78–1.66)	1.35 (0.93–1.95)	1.52 (1.03–2.22)	1.76 (1.20–2.59)	<0.001	
Insulin	1.00	1.09 (0.88–1.34)	1.29 (1.05–1.58)	1.56 (1.26–1.94)	1.49 (1.19–1.86)	<0.001	
Women							
Age groups (years)							>0.05
<50	1.00	1.10 (0.89–1.37)	1.04 (0.84–1.29)	1.09 (0.88–1.35)	1.06 (0.86–1.30)	0.70	
50–59	1.00	0.89 (0.73–1.08)	0.88 (0.73–1.07)	1.02 (0.84–1.24)	1.06 (0.87–1.28)	0.048	
≥60	1.00	0.89 (0.72–1.09)	0.89 (0.73–1.09)	0.99 (0.79–1.23)	1.06 (0.85–1.33)	0.46	
Race							>0.20
African American	1.00	0.90 (0.77–1.05)	0.85 (0.72–0.99)	1.06 (0.90–1.24)	1.07 (0.92–1.26)	0.005	
White	1.00	1.03 (0.86–1.23)	1.09 (0.91–1.30)	1.08 (0.90–1.29)	1.13 (0.95–1.35)	0.17	
Smoking status							>0.75
Never	1.00	0.92 (0.77–1.08)	0.96 (0.82–1.13)	1.06 (0.90–1.26)	1.11 (0.94–1.31)	0.005	
Ever or current	1.00	1.05 (0.85–1.29)	1.04 (0.85–1.28)	1.11 (0.89–1.37)	1.10 (0.89–1.37)	0.21	
Glucose-lowering medication							>0.05
No use	1.00	1.05 (0.82–1.34)	1.08 (0.84–1.38)	1.08 (0.83–1.40)	1.30 (1.01–1.67)	0.029	
Oral hypoglycemic agents	1.00	0.94 (0.75–1.17)	0.87 (0.70–1.08)	0.93 (0.74–1.16)	0.89 (0.71–1.12)	0.43	
Metformin	1.00	1.04 (0.86–1.26)	1.04 (0.86–1.26)	1.16 (0.95–1.40)	1.12 (0.93–1.36)	0.073	
Sulfonylureas	1.00	0.85 (0.69–1.05)	0.92 (0.75–1.12)	0.99 (0.80–1.22)	0.99 (0.81–1.22)	0.17	
Other oral agents	1.00	0.79 (0.60–1.03)	0.77 (0.59–0.99)	0.83 (0.64–1.09)	0.83 (0.64–1.07)	0.80	
Insulin	1.00	0.90 (0.74–1.10)	0.97 (0.80–1.18)	1.12 (0.92–1.36)	1.14 (0.94–1.38)	0.001	

^aAdjusted for age, race, types of insurance, income, and smoking, other than the variable for stratification.

Table 4—HRs (95% CIs) of CHD according to different levels of BMI during follow-up and at last visit among various subpopulations^a

	BMI (kg/m ²)					P for trend	P for interaction
	<25.0	25.0–29.9	30–34.9	35–39.9	≥40		
Follow-up							
Men							
Age groups (years)							
<50	1.00	1.07 (0.86–1.34)	1.21 (0.97–1.51)	1.43 (1.13–1.81)	1.58 (1.25–2.00)	<0.001	>0.25
50–59	1.00	1.21 (0.90–1.52)	1.36 (1.08–1.71)	1.59 (1.25–2.02)	1.52 (1.18–1.95)	<0.001	
≥60	1.00	1.12 (0.89–1.41)	1.10 (0.86–1.39)	1.34 (1.02–1.77)	1.07 (0.76–1.49)	0.22	
Race							
African American	1.00	1.08 (0.91–1.29)	1.14 (0.95–1.36)	1.50 (1.23–1.82)	1.62 (1.31–2.01)	<0.001	>0.10
White	1.00	1.25 (1.03–1.53)	1.41 (1.16–1.72)	1.62 (1.31–1.99)	1.56 (1.26–1.94)	<0.001	
Smoking status							
Never	1.00	1.23 (0.99–1.53)	1.40 (1.12–1.73)	1.73 (1.38–2.17)	1.93 (1.53–2.43)	<0.001	<0.05
Ever or current	1.00	1.27 (1.04–1.55)	1.34 (1.09–1.64)	1.61 (1.29–2.03)	1.36 (1.05–1.76)	<0.001	
Glucose-lowering medication							
No use	1.00	1.55 (1.18–2.05)	1.46 (1.08–1.96)	1.71 (1.24–2.37)	1.86 (1.32–2.62)	0.001	>0.25
Oral hypoglycemic agents	1.00	1.11 (0.85–1.45)	1.33 (1.02–1.73)	1.44 (1.07–1.93)	1.60 (1.18–2.18)	<0.001	
Metformin	1.00	1.18 (0.95–1.48)	1.36 (1.09–1.70)	1.62 (1.28–2.05)	1.64 (1.29–2.10)	<0.001	
Sulfonylureas	1.00	0.98 (0.77–1.24)	1.13 (0.90–1.43)	1.37 (1.07–1.75)	1.35 (1.04–1.76)	<0.001	
Other oral agents	1.00	1.18 (0.78–1.78)	1.41 (0.94–2.09)	1.56 (1.03–2.35)	1.79 (1.18–2.71)	<0.001	
Insulin	1.00	1.18 (0.95–1.46)	1.32 (1.06–1.64)	1.64 (1.31–2.05)	1.63 (1.29–2.07)	<0.001	
Women							
Age groups (years)							
<50	1.00	1.11 (0.87–1.40)	1.16 (0.92–1.45)	1.21 (0.97–1.52)	1.14 (0.92–1.42)	0.18	>0.25
50–59	1.00	1.01 (0.82–1.24)	0.93 (0.76–1.14)	1.11 (0.90–1.36)	1.15 (0.94–1.40)	0.030	
≥60	1.00	0.87 (0.71–1.07)	0.84 (0.69–1.03)	1.04 (0.84–1.29)	1.01 (0.81–1.27)	0.42	
Race							
African American	1.00	0.94 (0.80–1.10)	0.97 (0.83–1.14)	1.11 (0.95–1.31)	1.15 (0.98–1.36)	<0.001	>0.25
White	1.00	1.06 (0.89–1.28)	1.01 (0.84–1.20)	1.19 (1.00–1.43)	1.15 (0.96–1.38)	0.044	
Smoking status							
Never	1.00	0.99 (0.83–1.18)	1.01 (0.86–1.20)	1.15 (0.96–1.36)	1.19 (1.01–1.42)	<0.001	>0.90
Ever or current	1.00	1.14 (0.92–1.41)	1.15 (0.93–1.42)	1.31 (1.06–1.64)	1.20 (0.96–1.49)	0.036	
Glucose-lowering medication							
No use	1.00	1.08 (0.84–1.40)	1.06 (0.82–1.37)	1.33 (1.02–1.73)	1.38 (1.06–1.79)	0.002	>0.10
Oral hypoglycemic agents	1.00	0.96 (0.77–1.20)	0.94 (0.76–1.17)	0.94 (0.75–1.18)	0.90 (0.72–1.13)	0.35	
Metformin	1.00	1.11 (0.91–1.35)	1.13 (0.93–1.37)	1.21 (0.99–1.47)	1.21 (0.99–1.47)	0.035	
Sulfonylureas	1.00	0.99 (0.79–1.24)	1.01 (0.82–1.25)	1.08 (0.87–1.35)	1.08 (0.87–1.35)	0.17	
Other oral agents	1.00	0.96 (0.71–1.31)	0.82 (0.61–1.11)	0.88 (0.64–1.20)	0.92 (0.68–1.25)	0.76	
Insulin	1.00	1.05 (0.85–1.31)	1.10 (0.89–1.35)	1.26 (1.02–1.56)	1.29 (1.05–1.59)	<0.001	
Last visit							
Men							
Age groups (years)							
<50	1.00	1.03 (0.83–1.28)	1.14 (0.92–1.41)	1.45 (1.16–1.81)	1.50 (1.20–1.87)	<0.001	>0.25
50–59	1.00	1.25 (1.01–1.55)	1.37 (1.10–1.69)	1.56 (1.24–1.96)	1.50 (1.19–1.91)	<0.001	
≥60	1.00	1.10 (0.89–1.36)	1.13 (0.90–1.41)	1.24 (0.95–1.63)	1.17 (0.86–1.60)	0.09	

Continued on p. 3210

Table 4—Continued

	BMI (kg/m ²)					P for trend	P for interaction
	<25.0	25.0–29.9	30–34.9	35–39.9	≥40		
Race							
African American	1.00	1.06 (0.89–1.25)	1.15 (0.97–1.36)	1.42 (1.17–1.72)	1.61 (1.32–1.97)	<0.001	>0.10
White	1.00	1.26 (1.04–1.51)	1.36 (1.13–1.64)	1.58 (1.30–1.93)	1.53 (1.25–1.87)	<0.001	
Smoking status							
Never	1.00	1.25 (1.02–1.52)	1.44 (1.19–1.76)	1.66 (1.35–2.05)	1.95 (1.58–2.41)	<0.001	>0.05
Ever or current	1.00	1.15 (0.95–1.38)	1.19 (0.98–1.44)	1.50 (1.20–1.88)	1.25 (0.98–1.59)	0.003	
Glucose-lowering medication							
No use	1.00	1.54 (1.19–2.00)	1.41 (1.06–1.86)	1.69 (1.25–2.30)	1.65 (1.18–2.32)	0.004	>0.10
Oral hypoglycemic agents	1.00	1.06 (0.83–1.36)	1.32 (1.03–1.69)	1.42 (1.08–1.88)	1.61 (1.21–2.14)	<0.001	
Metformin	1.00	0.99 (0.82–1.21)	1.23 (1.01–1.49)	1.40 (1.13–1.73)	1.48 (1.19–1.83)	<0.001	
Sulfonylureas	1.00	0.97 (0.78–1.21)	1.19 (0.96–1.47)	1.40 (1.11–1.78)	1.34 (1.06–1.71)	<0.001	
Other oral agents	1.00	0.97 (0.66–1.41)	1.26 (0.88–1.81)	1.38 (0.94–2.02)	1.46 (1.00–2.14)	<0.001	
Insulin	1.00	1.04 (0.85–1.26)	1.22 (1.00–1.48)	1.38 (1.12–1.70)	1.47 (1.19–1.82)	<0.001	
Women							
Age groups (years)							
<50	1.00	0.93 (0.75–1.14)	0.97 (0.79–1.19)	1.04 (0.85–1.28)	0.96 (0.79–1.16)	0.75	>0.25
50–59	1.00	0.86 (0.72–1.04)	0.86 (0.72–1.03)	0.99 (0.83–1.20)	1.05 (0.87–1.25)	0.055	
≥60	1.00	0.87 (0.72–1.05)	0.80 (0.66–0.97)	0.99 (0.81–1.22)	1.02 (0.82–1.27)	0.66	
Race							
African American	1.00	0.83 (0.72–0.97)	0.88 (0.76–1.02)	1.02 (0.88–1.19)	1.04 (0.90–1.21)	0.001	>0.25
White	1.00	0.97 (0.82–1.15)	0.93 (0.79–1.09)	1.08 (0.91–1.28)	1.06 (0.90–1.25)	0.14	
Smoking status							
Never	1.00	0.88 (0.75–1.03)	0.92 (0.79–1.07)	1.05 (0.90–1.23)	1.07 (0.92–1.25)	0.005	>0.90
Ever or current	1.00	1.03 (0.84–1.26)	1.06 (0.87–1.29)	1.18 (0.96–1.45)	1.08 (0.88–1.33)	0.26	
Glucose-lowering medication							
No use	1.00	0.93 (0.73–1.19)	1.15 (0.91–1.45)	1.13 (0.88–1.45)	1.23 (0.96–1.58)	0.016	0.262
Oral hypoglycemic agents	1.00	0.94 (0.77–1.14)	0.87 (0.71–1.06)	0.95 (0.77–1.17)	0.85 (0.69–1.05)	0.23	
Metformin	1.00	1.02 (0.86–1.22)	0.97 (0.82–1.16)	1.14 (0.96–1.36)	1.08 (0.91–1.29)	0.11	
Sulfonylureas	1.00	0.91 (0.74–1.11)	0.88 (0.72–1.07)	1.05 (0.86–1.28)	0.97 (0.79–1.18)	0.39	
Other oral agents	1.00	0.83 (0.63–1.08)	0.74 (0.57–0.96)	0.81 (0.62–1.06)	0.80 (0.62–1.04)	0.40	
Insulin	1.00	0.85 (0.70–1.03)	0.88 (0.73–1.05)	1.07 (0.89–1.30)	1.03 (0.86–1.24)	0.013	

^aAdjusted for age, race, types of insurance, income, and smoking, other than the variable for stratification.

BMI at baseline and during follow-up. In addition, this positive association was present in different race, antidiabetes medication, and smoking groups. It is noteworthy that there was a U-shaped association between BMI at the last visit and the risk of CHD among women with type 2 diabetes in the current study. Our study found that diabetic women who were overweight and had class I obesity (BMI 25–34.9 kg/m²) at the last visit had a lower risk of CHD compared with normal-weight women (BMI <25 kg/m²).

It is well known that women with diabetes have a greater or equal relative risk of CHD than men with diabetes (30,31). The current study found a significant positive association of BMI and CHD risk among both men and women with type 2 diabetes, and this association is stronger among men than among women. The finding from our study is noteworthy for us to prevent CHD among patients with type 2 diabetes. In addition, more studies are needed to confirm the different effect size of BMI with CHD risk among men and women with type 2 diabetes.

It has been suggested that three potential methodological concerns should be considered when assessing the associations between obesity and health outcomes (32). The most serious concern is reverse causation associated with CHD and death risk. People with a history of CVD and several other chronic diseases frequently lose weight, and thus, people with a lower weight might increase the estimated risk of death. A recent analysis pooling five longitudinal studies has found that patients who have normal weight at the time of diabetes diagnosis have a higher mortality risk than those who are overweight or obese (12). They suggest that diabetic individuals with metabolically obese normal-weight may reflect underlying illness that predisposes to mortality (33). Despite having a normal BMI, these diabetic individuals have hyperinsulinemia, insulin resistance, and dyslipidemia, and all of these factors predispose individuals to death (33). In the current study, we excluded patients with a history of CHD and stroke at time of diabetes diagnosis, which can minimize the influence of reverse causation. Moreover, we performed another sensitivity analysis by excluding the subjects who were diagnosed with CHD during the first 2 years of follow-up ($n =$

3,207), and the positive association of BMI at baseline and during follow-up with CHD risk was still present. The second major concern is that confounding factors may distort the association between body weight and CHD. Smoking is a particularly important factor because smokers tend to weigh less and have much higher CHD risk than nonsmokers. In the current study, smoking status was considered as a confounding factor in the multivariable model, and the positive association between BMI and the risk of CHD was found in both never-smokers and smokers. The third methodological concern in some analyses between weight and CHD risk is that the physiologic effects of excess fatness, such as hypertension, diabetes, and dyslipidemia, were controlled for statistically, thus artificially removing some of the effects of being overweight. Obesity has been found as a strong risk factor for hypertension (4), high levels of HbA_{1c} (5), and high serum cholesterol among diabetic patients (34) and has also been the key or important component of the metabolic syndrome (35). All of these factors are associated with an increased risk of CHD (35–37) and considered as mediating factors for the physiologic effects of obesity on the CHD risk. In the current study, the adjustment for systolic blood pressure, LDL cholesterol, HDL cholesterol, triglycerides, HbA_{1c}, eGFR, and treatment attenuated the association between BMI and CHD risk, but BMI as a continuous variable remained a statistically significant predictor of CHD in the multivariable model.

There are several strengths in our study, including the large sample size, high proportion of African Americans, and the use of administrative databases to avoid differential recall biases. We have used baseline BMI levels, updated mean values of BMI during follow-up, and the last visit value of BMI in the analyses, which can avoid potential bias from a single baseline measurement. In addition, participants in this study used the same public health care system that minimizes the influence of accessibility to health care, particularly in comparing men and women. One limitation of our study is that our analysis was not performed on a representative sample of the population, which limits the generalizability of this study; however, LSUHCS hospitals are public

hospitals and cover >1.6 million patients, most of whom are low-income persons in Louisiana. The results of the current study will have wide applicability for the population with low income and without health insurance in the U.S. Another limitation of our study is that we did not have data on other obesity indicators, such as waist, hip, and thigh circumferences, and did not assess abdominal height, although these adiposity predictors have been shown to be associated with CVD risk (6,38,39). Third, while body weight was measured at each clinic visit, clinically measured BMI might not be as accurate as BMI measured in carefully conducted laboratory studies (40). Fourth, even though our analyses adjusted for an extensive set of confounding factors, residual confounding due to the measurement error in the assessment of confounding factors, unmeasured factors such as heart rates, physical activity, education, and dietary factors, cannot be excluded.

In summary, we found a positive association between BMI at baseline and during follow-up with the risk of CHD among men and women with type 2 diabetes, and this association was stronger among men than among women. We also found a positive association between BMI at the last visit and the risk of CHD among men with type 2 diabetes and a U-shaped association between BMI at the last visit and the risk of CHD among women with type 2 diabetes.

Funding. This work was supported by LSU's Improving Clinical Outcomes Network, the Louisiana Clinical Data Research Network, and the National Institute of General Medical Sciences of the National Institutes of Health, which funds the Louisiana Clinical & Translational Science Center.

Duality of Interest. No potential conflicts of interest relevant to this article were reported.

Author Contributions. N.L. and G.H. wrote the manuscript and researched data. P.T.K., R.H., and J.J. reviewed and edited the manuscript. Y.Z., W.L., W.Z., and Y.W. researched data. All authors read and approved the final manuscript. G.H. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

References

1. Flegal KM, Carroll MD, Ogden CL, Curtin LR. Prevalence and trends in obesity among US adults, 1999–2008. *JAMA* 2010;303:235–241

2. Cheng YJ, Imperatore G, Geiss LS, et al. Secular changes in the age-specific prevalence of diabetes among U.S. adults: 1988-2010. *Diabetes Care* 2013;36:2690-2696
3. Kanaya AM, Adler N, Moffet HH, et al. Heterogeneity of diabetes outcomes among Asians and Pacific Islanders in the US: the Diabetes Study of Northern California (DISTANCE). *Diabetes Care* 2011;34:930-937
4. Hu G, Barengo NC, Tuomilehto J, Lakka TA, Nissinen A, Jousilahti P. Relationship of physical activity and body mass index to the risk of hypertension: a prospective study in Finland. *Hypertension* 2004;43:25-30
5. Hu G, Lindström J, Valle TT, et al. Physical activity, body mass index, and risk of type 2 diabetes in patients with normal or impaired glucose regulation. *Arch Intern Med* 2004;164:892-896
6. Klein S, Allison DB, Heymsfield SB, et al.; Association for Weight Management and Obesity Prevention; NAASO, The Obesity Society; American Society for Nutrition; American Diabetes Association. Waist circumference and cardiometabolic risk: a consensus statement from Shaping America's Health: Association for Weight Management and Obesity Prevention; NAASO, The Obesity Society; the American Society for Nutrition; and the American Diabetes Association. *Am J Clin Nutr* 2007;85:1197-1202
7. Hu G, Tuomilehto J, Silventoinen K, Barengo N, Jousilahti P. Joint effects of physical activity, body mass index, waist circumference and waist-to-hip ratio with the risk of cardiovascular disease among middle-aged Finnish men and women. *Eur Heart J* 2004;25:2212-2219
8. Hu G, Jousilahti P, Antikainen R, Katzmarzyk PT, Tuomilehto J. Joint effects of physical activity, body mass index, waist circumference, and waist-to-hip ratio on the risk of heart failure. *Circulation* 2010;121:237-244
9. Hu G, Tuomilehto J, Silventoinen K, Sarti C, Männistö S, Jousilahti P. Body mass index, waist circumference, and waist-hip ratio on the risk of total and type-specific stroke. *Arch Intern Med* 2007;167:1420-1427
10. Hu G, Jousilahti P, Barengo NC, Qiao Q, Lakka TA, Tuomilehto J. Physical activity, cardiovascular risk factors, and mortality among Finnish adults with diabetes. *Diabetes Care* 2005;28:799-805
11. Katzmarzyk PT, Hu G, Cefalu WT, Mire E, Bouchard C. The importance of waist circumference and BMI for mortality risk in diabetic adults. *Diabetes Care* 2013;36:3128-3130
12. Carnethon MR, De Chavez PJ, Biggs ML, et al. Association of weight status with mortality in adults with incident diabetes. *JAMA* 2012;308:581-590
13. Kokkinos P, Myers J, Faselis C, Doulas M, Kheirbek R, Nylen E. BMI-mortality paradox and fitness in African American and Caucasian men with type 2 diabetes. *Diabetes Care* 2012;35:1021-1027
14. McEwen LN, Kim C, Karter AJ, et al. Risk factors for mortality among patients with diabetes: the Translating Research Into Action for Diabetes (TRIAD) Study. *Diabetes Care* 2007;30:1736-1741
15. So WY, Yang X, Ma RC, et al. Risk factors in V-shaped risk associations with all-cause mortality in type 2 diabetes-The Hong Kong Diabetes Registry. *Diabetes Metab Res Rev* 2008;24:238-246
16. Khalangot M, Tronko M, Kravchenko V, Kulchinska J, Hu G. Body mass index and the risk of total and cardiovascular mortality among patients with type 2 diabetes: a large prospective study in Ukraine. *Heart* 2009;95:454-460
17. Tobias DK, Pan A, Jackson CL, et al. Body-mass index and mortality among adults with incident type 2 diabetes. *N Engl J Med* 2014;370:233-244
18. McEwen LN, Karter AJ, Waitzfelder BE, et al. Predictors of mortality over 8 years in type 2 diabetic patients: Translating Research Into Action for Diabetes (TRIAD). *Diabetes Care* 2012;35:1301-1309
19. Li W, Wang Y, Chen L, et al. Increasing prevalence of diabetes in middle or low income residents in Louisiana from 2000 to 2009. *Diabetes Res Clin Pract* 2011;94:262-268
20. Hu G, Horswell R, Wang Y, et al. Body mass index and the risk of dementia among Louisiana low income diabetic patients. *PLoS ONE* 2012;7:e44537
21. Wang Y, Katzmarzyk PT, Horswell R, et al. Racial disparities in diabetic complications in an underinsured population. *J Clin Endocrinol Metab* 2012;97:4446-4453
22. Zhao W, Katzmarzyk PT, Horswell R, et al. Blood pressure and stroke risk among diabetic patients. *J Clin Endocrinol Metab* 2013;98:3653-3662
23. Zhao W, Katzmarzyk PT, Horswell R, et al. Aggressive blood pressure control increases coronary heart disease risk among diabetic patients. *Diabetes Care* 2013;36:3287-3296
24. Zhao W, Katzmarzyk PT, Horswell R, et al. HbA1c and lower-extremity amputation risk in low-income patients with diabetes. *Diabetes Care* 2013;36:3591-3598
25. Zhao W, Katzmarzyk PT, Horswell R, Wang Y, Johnson J, Hu G. HbA1c and heart failure risk among diabetic patients. *J Clin Endocrinol Metab* 2014;99:E263-E267
26. Zhao W, Katzmarzyk PT, Horswell R, Wang Y, Johnson J, Hu G. HbA1c and coronary heart disease risk among diabetic patients. *Diabetes Care* 2014;37:428-435
27. Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care* 1997;20:1183-1197
28. Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care* 2003;26(Suppl. 1):S5-S20
29. Adler AI, Stratton IM, Neil HA, et al. Association of systolic blood pressure with macrovascular and microvascular complications of type 2 diabetes (UKPDS 36): prospective observational study. *BMJ* 2000;321:412-419
30. Hu G; DECODE Study Group. Gender difference in all-cause and cardiovascular mortality related to hyperglycaemia and newly-diagnosed diabetes. *Diabetologia* 2003;46:608-617
31. Kalyani RR, Lazo M, Ouyang P, et al. Sex differences in diabetes and risk of incident coronary artery disease in healthy young and middle-aged adults. *Diabetes Care* 2014;37:830-838
32. Willett WC, Dietz WH, Colditz GA. Guidelines for healthy weight. *N Engl J Med* 1999;341:427-434
33. Conus F, Rabasa-Lhoret R, Péronnet F. Characteristics of metabolically obese normal-weight (MONW) subjects. *Appl Physiol Nutr Metab* 2007;32:4-12
34. Gostynski M, Gutzwiller F, Kuulasmaa K, et al.; WHO MONICA Project. Analysis of the relationship between total cholesterol, age, body mass index among males and females in the WHO MONICA Project. *Int J Obes Relat Metab Disord* 2004;28:1082-1090
35. Grundy SM, Cleeman JI, Daniels SR, et al.; American Heart Association; National Heart, Lung, and Blood Institute. Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation* 2005;112:2735-2752
36. Hu G, Sarti C, Jousilahti P, et al. The impact of history of hypertension and type 2 diabetes at baseline on the incidence of stroke and stroke mortality. *Stroke* 2005;36:2538-2543
37. Hu G, Qiao Q, Tuomilehto J, Balkau B, Borch-Johnsen K, Pyörälä K; DECODE Study Group. Prevalence of the metabolic syndrome and its relation to all-cause and cardiovascular mortality in nondiabetic European men and women. *Arch Intern Med* 2004;164:1066-1076
38. Gustat J, Elkasabany A, Srinivasan S, Berenson GS. Relation of abdominal height to cardiovascular risk factors in young adults: the Bogalusa Heart Study. *Am J Epidemiol* 2000;151:885-891
39. Heitmann BL, Frederiksen P. High circumference and risk of heart disease and premature death: prospective cohort study. *BMJ* 2009;339:b3292
40. Bhaskaran K, Forbes HJ, Douglas I, Leon DA, Smeeth L. Representativeness and optimal use of body mass index (BMI) in the UK Clinical Practice Research Datalink (CPRD). *BMJ Open* 2013;3:e003389