

# Oral Semaglutide

Vinh Bui and Joshua J. Neumiller

## Introduction

The number of antihyperglycemic medications approved for the treatment of type 2 diabetes (T2D) has increased dramatically in recent years (1). Since the initial approval of exenatide (Byetta) by the U.S. Food and Drug Administration (FDA) in 2005 (2), agents in the glucagon-like peptide 1 (GLP-1) receptor agonist class of medications have quickly gained popularity as a treatment option for the management of hyperglycemia in T2D (3). The increase in use of GLP-1 receptor agonists is due, at least in part, to their demonstrated clinical utility in improving glycemic control and inducing weight loss in people with T2D (4).

One practical barrier to GLP-1 receptor agonist use for some individuals is the need to administer these agents via subcutaneous injection. For patients unwilling or unable to initiate a GLP-1 receptor agonist for this reason, an alternative may be available in the future—namely, an oral formulation of semaglutide that is currently in phase 3 investigation for the treatment of T2D (5). This article provides a brief discussion of once-daily oral semaglutide and the future role it may play in the management of T2D.

## Indications

Since 2005, seven GLP-1 receptor agonists have received FDA approval as adjuncts to diet and exercise to improve glycemic control in adults with T2D (3). If approved by the

FDA, oral semaglutide would likely carry the same indication as other GLP-1 receptor agonist agents currently available in the United States. A once-weekly injectable formulation of semaglutide is also under investigation for the treatment of obesity (3), but it is not yet known whether the manufacturer plans to pursue an obesity indication for the oral formulation in the future.

## Mechanisms of Action

Activation of GLP-1 receptors has been shown to improve glycemic control through several mechanisms, including 1) increased glucose-dependent insulin secretion from pancreatic  $\beta$ -cells, 2) decreased glucagon secretion from pancreatic  $\alpha$ -cells, and 3) delaying gastric emptying (6). GLP-1 receptor activation in the central nervous system is additionally involved in satiety signaling, thus contributing to weight loss (6). Similar to the clinical benefits expected from currently available GLP-1 receptor agonists, clinical evidence with oral semaglutide use to date demonstrates improvements in glycemic control and weight (7).

Oral semaglutide is co-formulated within a tablet with the absorption enhancer sodium *N*-(8-[2-hydroxybenzoyl] amino) caprylate, or “SNAC” for short (7). SNAC helps facilitate semaglutide absorption in the stomach by increasing the local pH, which in turn leads to increased drug solubility and protects against proteolytic degradation (7).

Washington State University College of Pharmacy, Spokane, WA

Corresponding author: Joshua J. Neumiller, [jneumiller@wsu.edu](mailto:jneumiller@wsu.edu)

<https://doi.org/10.2337/cd18-0034>

©2018 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0> for details.

**Potential Advantages**

If approved by the FDA, the primary advantage of oral semaglutide will be its oral route of administration. As the first orally available agent in its class (if approved), it will provide patients and clinicians with a noninjectable GLP-1 receptor agonist option. In addition, emerging clinical data suggest that this agent may have advantages over currently available therapies in terms of glycemic efficacy and weight loss.

Results from the PIONEER 1 trial were recently reported at the American Diabetes Association’s (ADA’s) 78th Scientific Sessions (8). PIONEER 1 was a placebo-controlled phase 3 trial in people with T2D uncontrolled with diet and exercise. Participants were randomized to receive one of three doses of semaglutide (3, 7, or 14 mg once daily) or placebo. Participants were adults (≥18 years of age) with an A1C of 7.0–9.5%. After 26 weeks of treatment, participants receiving the 14-mg dose realized a mean decrease in A1C of 1.5% and a mean weight loss of 4.3 kg (8).

Headline results from several other trials have been reported by the manufacturer, showing superior improvements in A1C with oral semaglutide compared to empagliflozin (9) and sitagliptin (10) in the PIONEER 2 and PIONEER 3 studies, respectively. Headline results from the PIONEER 4 study additionally reported that the 14-mg dose of oral semaglutide achieved noninferiority for A1C reduction compared to 1.8 mg liraglutide, with superior weight reduction compared to liraglutide after 26 weeks of treatment (11).

**Potential Disadvantages**

The potential disadvantages of oral semaglutide therapy are similar to those of other agents in the GLP-1 receptor agonist class. In the largest trial completed to date with oral semaglutide, the most common adverse events associated with its use were gastrointestinal in nature (e.g.,

nausea and vomiting) (7). This was similarly seen in the same trial with the once-weekly injectable semaglutide formulation (7). The association of oral semaglutide with other key adverse events common to the GLP-1 receptor agonist class, such as acute kidney injury and acute pancreatitis, will be investigated during phase 3 trials. Ongoing and future studies of oral semaglutide will continue to elucidate the tolerability and safety considerations of this agent.

**Cost**

The future cost of oral semaglutide, should it receive FDA approval, is unknown.

**Commentary**

The GLP-1 receptor agonist class of medications has emerged as a noteworthy advancement in the management of T2D. Guidelines from major organizations such as the ADA and the American Association of Clinical Endocrinologists/American College of Endocrinology highlight GLP-1 receptor agonists as treatment options for many patients with T2D (3,12). Although diversity does exist within the GLP-1 receptor agonist class in terms of administration frequency (twice-daily, once-daily, or once-weekly administration) and pen device functionality, all currently available agents are injectables. If approved, oral semaglutide would provide a viable choice within the class for patients who are unable or unwilling to initiate injectable therapy. Given the glycemic and weight benefits that can be realized with the use of GLP-1 receptor agonist therapy, an orally administered product would be a welcome addition to the T2D treatment toolkit.

**Bottom Line**

Clearly, oral semaglutide has many hurdles ahead before it attains FDA approval. That said, evidence published to date is promising. Over the past several years, the GLP-1 receptor agonist class has become a bit crowded, with one product voluntarily discontinued

by the manufacturer due to a lack of market uptake (13). If approved, this oral GLP-1 receptor agonist would be a truly unique agent within the class. Emerging phase 3 data suggest that treatment with this product results in significant A1C reduction and weight loss. Only time will tell if the commercial availability of this product will become a reality.

**Duality of Interest**

No potential conflicts of interest relevant to this article were reported.

**Author Contributions**

Both authors researched data and wrote the manuscript. J.J.N. is the guarantor of this work and, as such, had full access to all of the references cited and takes responsibility for the accuracy of content.

**References**

- White JR Jr. A brief history of the development of diabetes medications. *Diabetes Spectr* 2014;27:82–86
- Exenatide [Byetta] prescribing information. Wilmington, Del., AstraZeneca Pharmaceuticals, February 2015 revision. Available from [www.azpicentral.com/byetta/pi\\_byetta.pdf#page=1](http://www.azpicentral.com/byetta/pi_byetta.pdf#page=1). Accessed 28 April 2018
- American Diabetes Association. 8. Pharmacologic approaches to glycemic treatment: *Standards of Medical Care in Diabetes—2018*. *Diabetes Care* 2018;41(Suppl. 1):S73–S85
- Neumiller JJ. Incretin-based therapies. *Med Clin North Am* 2015;99:107–129
- Novo Nordisk. R&D pipeline. Available from [www.novonordisk.com/rnd/rd-pipeline.html](http://www.novonordisk.com/rnd/rd-pipeline.html). Accessed 28 April 2018
- Neumiller JJ. Incretin pharmacology: a review of the incretin effect and current incretin-based therapies. *Cardiovasc Hematol Agents Med Chem* 2012;10:276–288
- Davies M, Pieber TR, Hartoft-Nielsen ML, et al. Effect of oral semaglutide compared with placebo and subcutaneous semaglutide on glycemic control in patients with type 2 diabetes: a randomized clinical trial. *JAMA* 2017;318:1460–1470
- Aroda VR, Rosenstock J, Terauchi Y, et al. Effect and safety of oral semaglutide monotherapy in type 2 diabetes: PIONEER 1 trial (Poster 2-LB). Presented at the American Diabetes Association’s 78th Scientific Sessions, Orlando, Fla., 22–26 June 2018
- Novo Nordisk. Oral semaglutide shows superior improvement in HbA1c vs. empagliflozin in the PIONEER 2 trial.

Downloaded from <http://diabetesjournals.org/clinical/article-pdf/36/4/327/500644/327.pdf> by guest on 28 May 2022

Available from [www.novonordisk.com/media/news-details.2195846.html](http://www.novonordisk.com/media/news-details.2195846.html). Accessed 4 July 2018

10. Novo Nordisk. Oral semaglutide shows superior reductions in HbA1c and weight compared to sitagliptin in the long-term safety and efficacy trial, PIONEER 3. Available from [www.novonordisk.com/media/news-details.2201779.html](http://www.novonordisk.com/media/news-details.2201779.html). Accessed 4 July 2018

11. Novo Nordisk. Oral semaglutide shows statistically significant greater reductions in HbA1c and weight compared to Victoza and sitagliptin in the PIONEER 4 and 7 trials. Available from [www.novonordisk.com/media/news-details.2200408.html](http://www.novonordisk.com/media/news-details.2200408.html). Accessed 4 July 2018

12. Garber AJ, Abrahamson MJ, Barzilay JI, et al. Consensus statement by the American Association of Clinical

Endocrinologists and American College of Endocrinology on the comprehensive type 2 diabetes management algorithm—2018 executive summary. *Endocr Pract* 2018;24:91–120

13. GlaxoSmithKline. Tanzeum (albiglutide) discontinuation: Q&A. Available from [www.tanzeum.com/pdfs/consumer-faq.pdf](http://www.tanzeum.com/pdfs/consumer-faq.pdf). Accessed 29 April 2018