

3 Jones, I. R., and Edwards, D. H., "An Experimental Study of the Forces Generated by the Collapse of Transient Cavities in Water," *Journal of Fluid Mechanics*, Vol. 7, Part 4, Apr. 1960, pp. 596-609.

4 Naudé, C. F., and Ellis, A. T., "On the Mechanism of Cavitation Damage by Nonhemispherical Cavities Collapsing in Contact with a Solid Boundary," *ASME Journal of Basic Engineering*, Vol. 83, No. 4, Dec. 1961, pp. 648-656.

5 Shutler, N. D., and Mesler, R. B., "A Photographic Study of the Dynamics and Damage Capabilities of Bubbles Collapsing Near Solid Boundaries," *ASME Journal of Basic Engineering*, Vol. 87, No. 3, Sept. 1965, pp. 648-656.

6 Ellis, A. T., "On Jets and Shock Waves from Cavitation," *Proceedings of Sixth Symposium on Naval Hydrodynamics*, Washington, 1966, pp. 137-161.

7 Brunton, J. H., Wilson, M. P. W., and Camus, J. J., "The High-Speed Photography of Liquid Flow in Collapsing Cavities and Impinging Liquid Drops," *Proceedings of the 8th International Congress on High-Speed Photography*, Stockholm, 1968, pp. 467-470.

8 Lauterborn, W., "High-Speed Photography of Laser-Induced Breakdown in Liquids" *Applied Physics Letters*, Vol. 21, No. 1, July 1972, pp. 27-29.

9 Kling, C. L., and Hammit, F. G., "A Photographic Study of Spark-Induced Cavitation Bubble Collapse," *ASME Journal of Basic Engineering*, Vol. 94, No. 4, Dec. 1972, pp. 825-833.

10 Smith, R. H., and Mesler, R. B., "A Photographic Study of the Effect of an Air Bubble on the Growth and Collapse of a Vapor Bubble Near a Surface," *ASME Journal of Basic Engineering*, Vol. 94, No. 4, Dec. 1972, pp. 933-942.

11 Lauterborn, W., and Bolle, H., "Experimental Investigations of Cavitation Bubble Collapse in Neighbourhood of a Solid Boundary," *Journal of Fluid Mechanics*, Vol. 72, Part 2, Nov. 1975, pp. 391-399.

12 Chahine, G. L., "Interaction between an Oscillating Bubble and a Free Surface," *ASME JOURNAL OF FLUIDS ENGINEERING*, Vol. 99, No. 4, Dec. 1977, pp. 709-716.

13 Fujikawa, S., and Akamatsu, T., "Experimental Investigations of Cavitation Bubble Collapse by a Water Shock Tube," *Bulletin of the JSME*, Vol. 21, No. 152, Feb. 1978, pp. 223-230.

14 Fujikawa, S., and Akamatsu, T., "Effects of the Non-Equilibrium Condensation of Vapour on the Pressure Wave Produced by the Collapse of a Bubble in a Liquid," *Journal of Fluid Mechanics*, Vol. 97, Part 3, Apr. 1980, pp. 481-512.

15 Shima, A., Takayama, K., Tomita, Y., and Miura, N., "An Experimental Study on Effects of a Solid Wall on the Motion of Bubble and Shock Waves in Bubble Collapse," *Acustica*, Vol. 48, No. 5, 1981, pp. 293-301.

16 Rayleigh, Lord., "On the Pressure Developed in a Liquid During the Collapse of a Spherical Cavity," *Philosophical Magazine*, Vol. 34, 1917, pp. 94-98.

17 Rattray, M., Jr., "Perturbation Effects in Cavitation Bubble Dynamics," California Institute of Technology, Ph.D. thesis, 1951.

18 Trilling, L., "The Collapse and Rebound of a Gas Bubble," *Journal of Applied Physics*, Vol. 23, No. 1, Jan. 1952, pp. 14-17.

19 Shima, A., "The Behavior of a Spherical Bubble in the Vicinity of a Solid Wall," *ASME Journal of Basic Engineering*, Vol. 90, No. 1, Mar. 1968, pp. 75-89.

20 Tulin, M. P., "On the Creation of Ultra-Jets," L. I. Sedov 60th Anniversary Volume, Problems of Hydrodynamics and Continuum Mechanics, Moscow, 1969, pp. 725-747.

21 Plesset, M. S., and Chapman, R. B., "Collapse of an Initially Spherical Vapour Cavity in the Neighbourhood of a Solid Boundary," *Journal of Fluid Mechanics*, Vol. 47, Part 2, May 1971, pp. 283-290.

22 Mitchell, T. M., and Hammit, F. G., "Asymmetric Cavitation Bubble Collapse," *ASME Journal of Basic Engineering*, Vol. 95, No. 1, Mar. 1973, pp. 29-37.

23 Shima, A., and Tomita, Y., "On the Impulse Pressure Accompanying Spherical Bubble Collapse in Liquids," Reports of the Institute of High Speed Mechanics, Tohoku University, Vol. 31, No. 281, 1975, pp. 97-135.

24 Tomita, Y., and Shima, A., "On the Behavior of a Spherical Bubble and the Impulse Pressure in a Viscous Compressible Liquid," *Bulletin of the JSME*, Vol. 20, No. 149, Nov. 1977, pp. 1453-1460.

25 Shima, A., and Nakajima, K., "The Collapse of a Non-Hemispherical Bubble Attached to a Solid Wall," *Journal of Fluid Mechanics*, Vol. 80, Part 2, Apr. 1977, pp. 369-391.

26 Tomita, Y., and Shima, A., "The Effect of Heat Transfer on the Behavior of a Bubble and the Impulse Pressure in a Viscous Compressible Liquid," *ZAMM*, Bd. 59, Heft 7, Juli 1979, S. 297-306.

27 Shima, A., and Sato, Y., "The Collapse of a Bubble Attached to a Solid Wall," *Ingenieur-Archiv*, Bd. 48, Heft 2, 1979, S. 85-95.

28 Sato, Y., and Shima, A., "The Collapse of an Initially Spherical Bubble Near a Solid Wall," Reports of the Institute of High Speed Mechanics, Tohoku University, Vol. 42, No. 325, 1980, pp. 1-24.

29 Fujikawa, S., and Akamatsu, T., "On the Mechanisms of Cavitation Bubble Collapse," *Proceedings of the 10th IAHR Symposium*, Tokyo, Vol. 1, 1980, pp. 91-102.

30 Shima, A., and Tomita, Y., "The Behavior of a Spherical Bubble Near a Solid Wall in a Compressible Liquid," *Ingenieur-Archiv*, Bd. 51, Heft 3/4, 1981, S. 243-255.

31 Bowden, F. P., and Brunton, J. H., "The Deformation of Solids by

Liquid Impact at Supersonic Speeds," *Proceedings of the Royal Society of London*, Series A, Vol. 263, No. 1315, Oct. 1961, pp. 433-450.

32 Hancox, N. L., and Brunton, J. H., "The Physics of Impact and Deformation: Multiple Impact, The Erosion of Solids by the Repeated Impact of Liquid Drops," *Philosophical Transactions of the Royal Society of London*, Series A, Vol. 260, 1966, pp. 121-139.

33 Singer, B. G., and Harvey, S. J., "Cavitation Damage Studies Using Plasticine," *International Journal of Mechanical Science*, Vol. 21, 1979, pp. 409-416.

DISCUSSION

A. Prosperetti¹

The authors are to be congratulated for reporting in this paper results of interest for their bearing on the mechanism of cavitation damage and bubble dynamics in general.

The shock wave which in a real flow situation is irradiated by collapsing bubbles is obtained here by means of a spark discharge and subsequent growth of a vapor bubble. This results in a good simulation of the real process if the parameters of the shock wave (notably its width and strength) are comparable in the two cases. Maybe the authors can comment on this point referring to their collapse shock results shown e.g. in Fig. 10.

I was also interested in the bubble-bubble interaction results shown in Fig. 5 in view of some numerical work based on potential flow which we did on the problem some time ago (L. Guerri, G. Lucca, and A. Prosperetti, "A Numerical Method for the Dynamics of Nonspherical Cavitation Bubbles," in *Proceedings of the Second International Colloquium on Drops and Bubbles*, Monterey, November 19-21, 1981, D. H. Le Croisette Ed., Jet Propulsion Laboratory Publication 82-7, pp. 175-181). The figure reproduced here shows some results for the case of axisymmetric collapse of one spherical and one hemispherical bubble attached to a solid wall (or, by reflection in the horizontal plane, of three equally spaced bubbles). Successive shapes of the bubbles are shown by thick lines, while the thin lines indicate particle paths. The situation is different from that of the paper because the bubbles begin to collapse at the same time. However these results show that the hemispherical bubble is impeded in its collapse by the neighbouring one and takes on an elongated shape qualitatively similar to that of the vapor bubble in the experiments (Fig. 5(b), frames 4-6). The numerical results also show that the time to collapse is affected by the presence of other bubbles, a factor which is not taken into account in the calculations of the paper. I would suggest that the discrepancy between calculated and measured collapse times for large R_e which can be seen in Fig. 7 is caused by the flow field of the spark-generated bubble. When the bubbles come close together (which, for the conditions of the experiment, implies large radii and greater collapse times) the interaction is not only due to the pressure wave but also to the actual flow field. Maybe some other effects depending on R_e reported in the paper are due to this factor.

Finally I would like to make a remark, and ask the authors' opinion, on the difficult question of shock waves versus microjets as agents of cavitation damage. The unambiguous identification of a shock wave irradiated by the bubble upon rebound and a failure to identify pressure pulses caused by the microjet does not necessarily demonstrate that cavitation damage is not to be imputed to the latter. If the action of the microjets is to cause very minute fractures of the surface of the solid, they could have such an effect without producing

¹Istituto di Fisica, 20133 Milano, Italy.




Fig. 12 Simultaneous axisymmetric collapse of one hemispherical and one spherical bubble in an inviscid incompressible liquid. The figures show successive bubble shapes (thick lines) and particle paths (thin lines). The initial distance between the bubble centers is, from left to right, 2.5, 2.2, and 2.1 initial radii. The shapes shown are for: A, $t = 0$; B, $t = 0.950$; C, $t = 1.063$; D, $t = 1.105$ (left and right), $t = 1.103$ (center); E, $t = 1.130$; F, $t = 1.135$. Distances are made dimensionless with respect to the initial radius R_0 and times with respect to $R_0 (\rho / (p_\infty - p_i))^{1/2}$, where ρ is the liquid density and p_∞ and p_i denote the ambient and internal pressure taken as constants.

notable pressures. It is possible that shock waves become effective in causing weight loss only after the surface has been exposed to "working" by the microjets? In this case both mechanisms would be important and, once more, the truth would be in the middle.

Authors' Closure

The authors wish to thank Professor Prosperetti for his useful discussions.

In the present experiment, shock waves with strength $p_s <$ about 5 MPa and pulse width $w_s \approx 5 \mu s$ are used in order to study the shock wave-bubble interaction. These values were determined by referring to the experimental study on the ultrasonic cavitation [34] in which the shock amplitudes of about 5 MPa were measured at the position of 1 mm apart from the center of collapse by means of optical methods. By the way, parameters of the shock wave produced in Fig. 10 are measured as $p_{G,\max} = 10.4$ MPa and $w = 2.6 \mu s$. The peak pressure $p_{g,\max}$ will reduce to the value of about 3 MPa at $L = 5$ mm along a $1/L$ decrease curve.

The authors are also thankful to Professor Prosperetti for his suggestion concerning the bubble-bubble interaction, and believe that the numerical results obtained by him and co-worker provide valuable addition to this paper. As a gas bubble increases in size, the interaction of it with a spark bubble becomes remarkable. This relation is shown in Fig. 13, where $\bar{R}_{v,0}$ indicates an averaged curve of five trials. It is clearly seen that when R_e increases, the discrepancy of H_v from $\bar{R}_{v,0}$ becomes larger. This means that a gas bubble will be affected by a spark bubble with increasing R_e . In general, there exists Bjerknes force between pulsating bubbles at a




Fig. 13 Interaction between a gas bubble and a spark bubble

distance $2d$ apart. Two bubbles pulsating in phase attract each other, and bubbles pulsating 180 deg out of phase repel each other. These are mathematically equivalent to the motion of a bubble at a distance d from a solid wall for the former and from a free surface for the latter, respectively. The periods of collapse will be lengthened near a solid wall and shortened near a free surface. In our experiment the motions of two bubbles are 180 deg out of phase in the initial stage of their motions. As a result, the calculated collapse times would be overestimated than measured collapse times because the interaction of a gas bubble with a spark bubble is not taken

into account in our calculations, and thus the discrepancy between them becomes larger with increasing R_e .

Now the authors would make a remark on the mechanism of impact pressure generation. We do believe that three types of damage forces are possible, and basically agree with the discussor's opinion. They are resulted from a shock wave, from a liquid jet and from both a shock wave and a liquid jet. The induced impact wall pressure modes are fairly corresponding to the bubble collapse modes depending on L/R_{\max} , where L is the distance between the electrodes and the

solid wall and R_{\max} a maximum bubble radius (see reference [35] for details).

Additional References

34 Hinsch, K., and Brinkmeyer, E., "Investigation of Very Short Cavitation Shock Waves by Coherent Optical Methods," *Proceedings of the Society of Photo-optical Instrumentation Engineers*, Vol. 97, 1977, pp. 166-171.

35 Shima, A., Takayama, K., Tomita, Y. and N. Ohsawa, "Mechanism of Impact Pressure Generation from Spark-Generated Bubble Collapse Near a Wall," *AIAA Journal*, Vol. 21, No. 1, Jan. 1983, pp. 55-59.