

# Breastfeeding and Mammographic Breast Density: A Cross-sectional Study

Kayla R. Getz<sup>1</sup>, Babatunde Adedokun<sup>2</sup>, Shuai Xu<sup>1</sup>, and Adetunji T. Toriola<sup>1</sup>



## ABSTRACT

Breastfeeding is inversely associated with breast cancer risk but the associations of breastfeeding with mammographic breast density (MBD) are not clear. We investigated the association between breastfeeding and volumetric measures of MBD [volumetric percent density (VPD), dense volume (DV), and non-dense volume (NDV)] and evaluated whether it differs by race, menopausal status, and body mass index (BMI). The study population was comprised of 964 women (67% non-Hispanic White, 29% non-Hispanic Black) who had screening mammography at Washington University School of Medicine, St. Louis, MO. VPD, DV and NDV were  $\log_{10}$  transformed. We performed multivariable linear regression models adjusted for age, BMI, family history of breast cancer, race, and age at menarche among all participants and exclusively in parous women. Mean age was 50.7 years. VPD was 12% lower among women who breastfed 0–6 months, [ $10^{\beta} = 0.88$ , 95% confidence interval (CI); 0.79–0.98] compared with nulliparous women. Breastfeeding was not associated with VPD among women who breastfed >7 months. Breastfeeding was inversely associated

with DV [parous never breastfed:  $10^{\beta} = 0.93$ ; 95% CI (0.83–1.04), breastfed 0–6 months:  $10^{\beta} = 0.91$ , 95% CI (0.79–1.05), breastfed 7–12 months:  $10^{\beta} = 0.94$ ; 95% CI (0.81–1.10), breastfed >12 months:  $10^{\beta} = 0.87$ , 95% CI (0.78–0.98),  $P_{\text{trend}} = 0.03$ ]. BMI modified the association between breastfeeding and VPD. Women who breastfed for 0–6 months and had a BMI < 25 kg/m<sup>2</sup> had lower VPD compared with nulliparous women, but among women with a BMI  $\geq$  25 kg/m<sup>2</sup> there was no association ( $P_{\text{interaction}} = 0.04$ ). In this diverse study population, the association of breastfeeding with VPD appears to be modified by BMI, but not by race or menopausal status. Future research exploring the associations of breastfeeding with other mammographic features are needed.

**Prevention Relevance:** Breastfeeding for up to 6 months may be associated with lower VPD among women with a BMI < 25 kg/m<sup>2</sup>. The potential role of MBD in mediating the associations of breastfeeding with breast cancer risk in a select group of women deserves further evaluation.

See related *Spotlight*, p. 309

## Introduction

Mammographic breast density (MBD) reflects the proportion of epithelial and stromal tissues relative to adipose tissue in the breast (1). There is a positive association between breast density and risk of developing breast cancer. Women with  $\geq$ 75% density experience an approximately 5-fold increased risk of developing breast cancer compared with women with <5% density (2). In addition, maintaining dense breasts or developing more dense breast tissue over time is positively associated with the risk of developing breast cancer (3).

Breastfeeding is inversely associated with breast cancer risk (4, 5). Reproductive factors such as higher parity and older age at first birth are inversely and positively associated

with MBD, respectively, but studies have reported conflicting associations between breastfeeding and MBD (6–14). The inconsistent findings may arise from how breastfeeding and MBD were measured and differences in study populations (6–8, 10–14). To the best of our knowledge, there are no studies on the association between breastfeeding and volumetric measures of MBD, as previous studies used either categorical or area-based measures.

Existing studies also suggest differences in the association between breastfeeding and MBD by menopausal status. The Nurses' Health Study observed positive associations between breastfeeding duration with dense and non-dense area among premenopausal women but no associations among postmenopausal women (6). There was a positive association between breastfeeding duration and percent density among premenopausal women in a study performed among women in the Mexican Teacher's Cohort, but no relationship was observed among postmenopausal women (13). Various demographic factors are related to breastfeeding behaviors, including race (15–17). Differences in breastfeeding initiation and duration exist between non-Hispanic White and non-Hispanic Black women, with non-Hispanic White women more likely to breastfeed compared with non-Hispanic Black women (16, 17). Not only are breastfeeding practices and duration different across race, but race is also associated with MBD

<sup>1</sup>Division of Public Health Sciences, Department of Surgery, Washington University School of Medicine, St. Louis, Missouri. <sup>2</sup>Center for Clinical Cancer Genetics and Global Health, Department of Medicine, University of Chicago, Chicago, Illinois.

**Corresponding Author:** Adetunji T. Toriola, Washington University in St. Louis School of Medicine, 660 S. Euclid Ave, St. Louis, MO 63110. Phone: 314-286-2668; E-mail: a.toriola@wustl.edu

Cancer Prev Res 2023;16:353–62

doi: 10.1158/1940-6207.CAPR-22-0482

©2023 American Association for Cancer Research

(18–20). Moore and colleagues recently reported that non-Hispanic Black women were more likely to have extremely dense breasts but less likely to have heterogeneously dense breasts compared with non-Hispanic White women. However, they did not investigate breastfeeding in their study (19). Although there has been an increase in breastfeeding initiation and duration among non-Hispanic Black women over time, it is important to investigate the association between breastfeeding and MBD by race because the prevalence of breastfeeding remains lower in non-Hispanic Black women across the United States compared with non-Hispanic White women (17, 21). Breastfeeding is inversely associated with risk of estrogen receptor–negative (ER<sup>-</sup>) breast cancer and non-Hispanic Black women have a higher risk of and ER<sup>-</sup> breast cancer (22). If breastfeeding is associated with MBD in non-Hispanic Black women, it could provide an insight into how breastfeeding duration influences breast parenchyma and may provide supporting evidence to conduct large cohort studies to assess the association between breastfeeding duration and breast cancer development in this population.

The objective of this study is to investigate the association between breastfeeding and volumetric measures of MBD in a racially diverse study population and to determine whether menopausal status, race or body mass index (BMI) modify this association.

## Materials and Methods

### Study population and design

This study population consists of women who received their annual screening mammogram at the Joanne Knight Breast Health Center (BHC), at Siteman Cancer Center at Washington University School of Medicine, St. Louis, MO. Women who had their annual screening mammogram at the BHC were approached and screened for participation in the study. Eligibility criteria included: (i) age 35 to 64 years, (ii) ability to comply with all required study procedures and schedule; (iii) not having serious medical condition that would prevent the participant from returning for annual mammogram in 12 months, and (iv) not being pregnant. Exclusion criteria included: (i) history of any cancer, including breast cancer; (ii) history of breast augmentation, reduction, or implants; and (iii) history of selective estrogen receptor modulators during the previous 6 months.

Participants completed a detailed questionnaire on demographic characteristics, reproductive factors, medication use, family history of breast cancer, alcohol use, tobacco use and adiposity at various ages. Further, participants' height and weight were measured using a stadiometer and OMRON Full Body Sensor Body Composition Monitor and Scale Model HBF-514FC, respectively. Approval for the study was granted by the Institutional Review Board at Washington University School of Medicine, St. Louis, MO. All study participants provided written informed consent and the study was conducted in accordance with the Declaration of Helsinki.

### Breastfeeding

Breastfeeding duration was derived by combining the number of months the women reported breastfeeding for their first, second, and last child. The original breastfeeding duration variable was categorized into four groups (“0–3 months,” “3–6 months,” “7–12 months,” and “>12 months”). To calculate cumulative duration of breastfeeding for all children, this variable combined breastfeeding duration across children and assumed the mother breastfed for the maximum number of months within each category for recoding purposes. Women who were nulliparous and those who had children but never breastfed were assigned to separate categories. The categories of breastfeeding duration analyzed were: “nulliparous,” “parous never breastfed,” “breastfed for 0–6 months,” “breastfed for 7–12 months,” and “breastfed for >12 months.” Of the 906 parous women in the study, data on breastfeeding duration were available for 779 women, while 127 parous women were missing breastfeeding duration.

### MBD assessment

Volpara version 1.5 (Matakina Technology Ltd) was used to evaluate volumetric measures of MBD—volumetric percent density (VPD), dense volume (DV), and non-dense volume (NDV) of the breast (23, 24). Volpara assesses measures of DV by averaging the mediolateral oblique and cranial-caudal views of the left and right breasts. DV is the volume of fibroglandular tissue in the breast (cm<sup>3</sup>). NDV is calculated as the difference between total breast volume (cm<sup>3</sup>) and DV. VPD is the ratio of the volume of fibroglandular tissue (i.e., DV) to the total breast volume, expressed as a percentage.

### Statistical analysis

We calculated the participants' characteristics using percentages for categorical variables as well as means and SDs for continuous variables. We evaluated these characteristics across the different breastfeeding duration categories and performed  $\chi^2$  and ANOVA tests to assess univariate associations. We performed complete case analyses of the associations between breastfeeding duration and MBD on 964 participants (excluding 127 parous women who were missing breastfeeding duration, 32 women who were missing MBD and additional 18 women who were missing values for confounding variables; flowchart of analytic sample presented in Supplementary Fig. S1) and performed log<sub>10</sub> transformations on VPD, DV, and NDV due to the skewed distribution.

We built multivariable-adjusted linear regression models to assess the associations between breastfeeding duration and MBD adjusting for age (continuous), BMI (continuous), family history of breast cancer in a first-degree relative (no, yes), race (non-Hispanic White, non-Hispanic Black, other), and age at menarche (continuous). Models were also adjusted for alcohol use as well as use of oral contraceptives and hormone replacement therapy among premenopausal and postmenopausal women, respectively. We additionally adjusted for body shape at age 10 which is based on the Stunkard scale/the figure rating scale. We recoded the original scale from nine categories to four

(1 = body shapes 1 and 2, 2 = body shapes 3 and 4, 3 = body shape 5, and 4 = body shapes 6–9; ref. 25). Use of alcohol, oral contraceptives, hormone replacement therapy and body shape at age 10 were not included in the final models because they did not change the point estimates by >10%. Coefficients and 95% confidence intervals (CI) were backtransformed for easier interpretation. Tests for trend were performed by treating breastfeeding duration as an ordinal variable. To determine whether menopausal status, race and BMI were effect modifiers on the association between breastfeeding duration and MBD, we added interaction terms to the multivariable linear regression models and conducted stratified analyses.

In addition, we investigated the association between breastfeeding and MBD among parous women only. Excluding nulliparous women from the analyses may help to isolate the impact of breastfeeding, which may be influenced by parity. Time since last birth was considered as a confounding variable in these analyses but was not included in the final models because it did not influence the point estimates by >10%. We

also performed additional sensitivity analyses to address women who had missing values for breastfeeding duration. Inverse probability weights were calculated to mimic the population where no participants had breastfeeding duration missing. Weights were added to the models and participants with breastfeeding duration missing were excluded.

All statistical tests were two sided, and *P* values < 0.05 were considered statistically significant. All analyses were performed using SAS version 9.4 (SAS Institute Inc., Statistical Analysis System, RRID:SCR\_008567).

#### Data availability

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

## Results

### Descriptive statistics

The complete case analysis included 964 women with a mean age of 50.7 years (Table 1). Approximately 67% of the women

**Table 1.** Characteristics of 964 women recruited during annual screening mammography at the Joanne Knight BHC, Washington University School of Medicine, St. Louis, MO.<sup>a</sup>

	Total ( <i>N</i> = 964) <i>N</i> (%) or mean (SD)	Nulliparous ( <i>N</i> = 228) <i>N</i> (%) or mean (SD)	Parous never breastfed ( <i>N</i> = 289) <i>N</i> (%) or mean (SD)	0–6 months ( <i>N</i> = 115) <i>N</i> (%) or mean (SD)	7–12 months ( <i>N</i> = 91) <i>N</i> (%) or mean (SD)	>12 months ( <i>N</i> = 241) <i>N</i> (%) or mean (SD)	<i>P</i>
<b>Age, year</b>	50.7 (7.1)	48.8 (7.1)	52.2 (6.7)	52.1 (7.0)	53.3 (6.9)	49.2 (6.8)	<0.0001
<b>Age at menarche, year</b>	12.7 (1.6)	12.6 (1.5)	12.7 (1.7)	12.8 (1.6)	12.6 (1.6)	12.9 (1.6)	0.31
<b>Age at first birth, year<sup>b</sup></b>	25.6 (6.2)	—	23.3 (6.1)	27.3 (6.4)	26.0 (5.8)	27.3 (5.4)	<0.0001
<b>Body mass index, kg/m<sup>2</sup></b>	30.6 (7.8)	31.2 (8.8)	32.4 (7.9)	30.0 (6.1)	29.6 (7.1)	28.5 (7.2)	<0.0001
<b>Race</b>							
Non-Hispanic White	645 (66.9)	176 (77.2)	140 (48.4)	73 (63.5)	66 (72.5)	190 (78.8)	
Non-Hispanic Black	279 (28.9)	43 (18.9)	144 (49.8)	38 (33.0)	20 (22.0)	34 (14.1)	<0.0001
Other race	40 (4.2)	9 (4.0)	5 (1.7)	4 (3.5)	5 (5.5)	17 (7.1)	
<b>Menopause status</b>							
Premenopausal	613 (63.6)	167 (73.3)	157 (54.3)	63 (54.8)	46 (50.6)	180 (74.7)	<0.0001
Postmenopausal	351 (36.4)	61 (26.8)	132 (45.7)	52 (45.2)	45 (49.5)	61 (25.3)	
<b>Family history of breast cancer</b>							
Yes	228 (23.7)	65 (28.5)	52 (18.0)	27 (23.5)	30 (33.0)	54 (22.4)	0.01
No	736 (76.4)	163 (71.5)	237 (82.0)	88 (76.5)	61 (67.0)	187 (77.6)	
<b>Parity number</b>							
Nulliparous	228 (23.7)	228 (100)	—	—	—	—	
1–2 children	480 (49.8)	—	210 (72.7)	90 (78.3)	63 (69.2)	117 (48.6)	<0.0001 <sup>c</sup>
≥3 children	256 (26.6)	—	79 (27.3)	25 (21.7)	28 (30.8)	124 (51.5)	
<b>Current alcohol use<sup>d</sup></b>							
Yes	627 (65.2)	152 (66.7)	155 (53.8)	79 (69.3)	64 (70.3)	177 (73.4)	<0.0001
No	335 (34.8)	76 (33.3)	133 (46.2)	35 (30.7)	27 (29.7)	64 (26.6)	
<b>Mammographic density<sup>e</sup></b>							
Volumetric percent density, %	8.5 (6.4)	9.4 (7.3)	6.8 (4.9)	7.6 (4.9)	9.0 (6.5)	9.9 (7.3)	<0.0001
Dense volume, cm <sup>3</sup>	96.1 (83.9)	104.2 (92.3)	103.6 (95.0)	87.4 (56.0)	104.1 (111.0)	80.5 (53.2)	0.02
Non-dense volume, cm <sup>3</sup>	1450.4 (1423.7)	1442.5 (1502.8)	1777.6 (1666.7)	1416.5 (1319.7)	1318.7(1105.0)	1131.5 (1067.8)	<0.0001
<b>log<sub>10</sub>-Mammographic density</b>							
Volumetric percent density, %	0.8 (0.3)	0.9 (0.3)	0.8 (0.2)	0.8 (0.2)	0.9 (0.3)	0.9 (0.3)	<0.0001
Dense volume, cm <sup>3</sup>	1.9 (0.3)	1.9 (0.3)	1.9 (0.3)	1.9 (0.2)	1.9 (0.3)	1.8 (0.2)	0.02
Non-dense volume, cm <sup>3</sup>	3.0 (0.4)	3.0 (0.4)	3.1 (0.4)	3.0 (0.3)	3.0 (0.3)	2.9 (0.4)	<0.0001

Abbreviations: *N*, number; %, column percentage; SD, standard deviation.

<sup>a</sup>Presented as *N* (%) for categorical variables or mean (SD) for continuous variables.

<sup>b</sup>Four parous women were missing age at first birth.

<sup>c</sup> $\chi^2$  test across parous never breastfed, 0–6 months, 7–12 months, and >12 months groups.

<sup>d</sup>Two participants are missing alcohol use information.

<sup>e</sup>Mammographic density indicators are not normally distributed. Kruskal–Wallis tests were conducted instead of ANOVA. These indicators were log<sub>10</sub> transformed to follow approximately normal distributions.

were non-Hispanic White, and 29% were non-Hispanic Black. The average BMI was 30.6 kg/m<sup>2</sup>. Approximately 30% of the women in the study were parous but never breastfed, followed by women who breastfed for >12 months (25.0%), nulliparous women (23.7%), women who breastfed for 0–6 months (11.9%), and women who breastfed for 7–12 months (9.4%). Nulliparous women were younger and had a younger age at menarche than women who reported breastfeeding. Parous women who never breastfed had the youngest age at first birth (23.3 years) and the highest BMI (32.4 kg/m<sup>2</sup>) compared with the other women. Women who breastfed for >12 months were likely to have the most children and also had the highest VPD (9.9%) but lowest DV (80.5 cm<sup>3</sup>) and NDV (1,131.5 cm<sup>3</sup>).

### Multivariable analysis

#### All study participants

VPD was 12% lower among women who were parous but never breastfed, [ $10^{\beta} = 0.88$ , 95% CI (0.81–0.96)] and women who breastfed 0–6 months, [ $10^{\beta} = 0.88$ , 95% CI (0.79–0.98)] compared with nulliparous women (Table 2). Breastfeeding was not associated with VPD among women who breastfed 7–12 months or >12 months ( $P_{\text{trend}} = 0.92$ ). There was an inverse association between breastfeeding and DV [parous never breastfed:  $10^{\beta} = 0.93$ , 95% CI (0.83–1.04), breastfed 0–6 months:  $10^{\beta} = 0.91$ , 95% CI (0.79–1.05), breastfed 7–12 months:  $10^{\beta} = 0.94$ , 95% CI (0.81–1.10), and breastfed >12 months:  $10^{\beta} = 0.87$ , 95% CI (0.78–0.98)  $P_{\text{trend}} = 0.03$ ] (Table 2). There were no associations between breastfeeding and NDV.

#### Parous women only

In the analyses limited to parous women, there was a positive association between breastfeeding for 7–12 months and VPD [ $10^{\beta} = 1.12$ , 95% CI (1.00–1.25)]. There was no association

between breastfeeding and DV, but an inverse trend was observed between breastfeeding duration and NDV [breastfed 0–6 months:  $10^{\beta} = 1.01$ , 95% CI (0.89–1.16), breastfed 7–12 months:  $10^{\beta} = 0.91$ , 95% CI (0.78–1.06), and breastfed >12 months:  $10^{\beta} = 0.90$ , 95% CI (0.80–1.01)  $P_{\text{trend}} = 0.04$ ] (Table 2).

### Interaction analyses

#### All study participants

BMI modified the association between breastfeeding duration and VPD. Among women with BMI < 25 kg/m<sup>2</sup>, women who breastfed for 0–6 months had lower VPD compared with nulliparous women, but among women with a BMI  $\geq 25$  kg/m<sup>2</sup> there was no association ( $P_{\text{interaction}} = 0.04$ ; Table 3). Menopausal status and race did not modify the associations of breastfeeding with VPD, DV, or NDV (Table 3).

#### Parous women only

Similar findings were observed in the analyses limited to parous women; BMI, menopausal status, and race did not modify the association between breastfeeding and any measure of MBD (Table 4).

#### Missing data

The results from the analysis that utilized inverse probability weights to deal with missing data were consistent with the findings from the complete case analysis (Supplementary Table S1) hence; the results from the complete case analysis are not likely to be biased because of missing data.

## Discussion

In the analyses across all study participants, we observed that women who breastfed for 0–6 months had lower VPD

**Table 2.** Associations of breastfeeding duration with MBD measures.<sup>a</sup>

		Volumetric percent density	Dense volume (cm <sup>3</sup> )	Non-dense volume (cm <sup>3</sup> )
	N	$10^{\beta}$ (95% CI)	$10^{\beta}$ (95% CI)	$10^{\beta}$ (95% CI)
All women	964			
Nulliparous	228	Reference	Reference	Reference
Parous never breastfed	289	<b>0.88 (0.81–0.96)</b>	0.93 (0.83–1.04)	1.07 (0.95–1.20)
0–6 months	115	<b>0.88 (0.79–0.98)</b>	0.91 (0.79–1.05)	1.07 (0.93–1.24)
7–12 months	91	1.00 (0.89–1.13)	0.94 (0.81–1.10)	0.96 (0.82–1.11)
>12 months	241	0.95 (0.87–1.04)	<b>0.87 (0.78–0.98)</b>	0.94 (0.84–1.05)
$P_{\text{trend}}$		0.92	<b>0.03</b>	0.10
Parous women	736			
Parous never breastfed	289	Reference	Reference	Reference
0–6 months	115	0.99 (0.89–1.09)	0.98 (0.86–1.12)	1.01 (0.89–1.16)
7–12 months	91	<b>1.12 (1.00–1.25)</b>	1.02 (0.88–1.18)	0.91 (0.78–1.06)
>12 months	241	1.06 (0.98–1.16)	0.94 (0.84–1.06)	0.90 (0.80–1.01)
$P_{\text{trend}}$		0.07	0.38	<b>0.04</b>

Note: Bold indicates statistical significance ( $P < 0.05$ ).

Abbreviations: CI, confidence intervals; MBD, mammographic breast density.

<sup>a</sup>Models were adjusted for age (continuous), age at menarche (continuous), BMI (continuous), race (non-Hispanic White, non-Hispanic Black, other), and family history of breast cancer (yes, no). Mammographic breast density measures were log<sub>10</sub> transformed because of the skewed distribution. Coefficients were backtransformed for easy interpretation.

**Table 3.** Associations of breastfeeding duration with MBD measures by menopause status, race and BMI.<sup>a</sup>

	N	Volumetric percent density		Dense volume (cm <sup>3</sup> )		Non-dense volume (cm <sup>3</sup> )	
		10 <sup>B</sup> (95% CI)	P <sub>trend</sub>	10 <sup>B</sup> (95% CI)	P <sub>trend</sub>	10 <sup>B</sup> (95% CI)	P <sub>trend</sub>
<i>Menopausal status</i>							
Premenopausal <sup>b</sup>							
Nulliparous	167	Reference	0.75	Reference	0.06	Reference	0.19
Parous never breastfed	157	<b>0.86 (0.77–0.96)</b>		0.96 (0.86–1.07)		1.11 (0.99–1.24)	
0–6 months	63	0.90 (0.78–1.04)		1.05 (0.92–1.21)		<b>1.20 (1.04–1.38)</b>	
7–12 months	46	1.00 (0.85–1.18)		0.95 (0.81–1.11)		0.95 (0.81–1.12)	
>12 months	180	0.94 (0.85–1.04)		<b>0.90 (0.82–1.00)</b>		0.97 (0.87–1.07)	
Postmenopausal <sup>b</sup>							
Nulliparous	61	Reference	0.68	Reference	0.30	Reference	0.28
Parous never breastfed	132	0.96 (0.84–1.09)		0.86 (0.68–1.10)		0.95 (0.75–1.21)	
0–6 months	52	0.91 (0.78–1.07)		<b>0.74 (0.55–1.00)</b>		0.85 (0.64–1.14)	
7–12 months	45	1.08 (0.91–1.27)		0.90 (0.66–1.22)		0.85 (0.63–1.15)	
>12 months	61	0.98 (0.84–1.14)		0.83 (0.62–1.10)		0.89 (0.67–1.17)	
P <sub>interaction</sub>		0.68		0.11		0.21	
<i>Race</i>							
Non-Hispanic White <sup>c</sup>							
Nulliparous	176	Reference	0.59	Reference	<b>0.04</b>	Reference	0.25
Parous never breastfed	140	0.93 (0.84–1.04)		0.93 (0.81–1.07)		1.01 (0.87–1.16)	
0–6 months	73	<b>0.87 (0.76–1.00)</b>		0.90 (0.76–1.07)		1.07 (0.90–1.28)	
7–12 months	66	1.02 (0.88–1.17)		0.93 (0.78–1.12)		0.93 (0.77–1.12)	
>12 months	190	0.95 (0.86–1.05)		<b>0.87 (0.76–0.99)</b>		0.94 (0.82–1.07)	
Non-Hispanic Black <sup>c</sup>							
Nulliparous	43	Reference	0.55	Reference	0.97	Reference	0.60
Parous never breastfed	144	0.87 (0.76–1.01)		0.98 (0.79–1.20)		1.14 (0.93–1.40)	
0–6 months	38	0.97 (0.81–1.16)		0.98 (0.75–1.27)		1.02 (0.79–1.32)	
7–12 months	20	0.96 (0.77–1.20)		1.08 (0.79–1.48)		1.13 (0.83–1.55)	
>12 months	34	0.97 (0.80–1.17)		0.95 (0.73–1.25)		0.98 (0.75–1.28)	
P <sub>interaction</sub>		0.48		0.96		0.67	
<i>BMI</i>							
BMI < 25 <sup>d</sup>							
Nulliparous	59	Reference	0.81	Reference	0.33	Reference	0.52
Parous never breastfed	49	0.81 (0.65–1.01)		<b>0.71 (0.56–0.92)</b>		0.92 (0.69–1.22)	
0–6 months	28	<b>0.75 (0.58–0.96)</b>		0.86 (0.65–1.15)		1.22 (0.88–1.68)	
7–12 months	30	1.00 (0.78–1.27)		1.00 (0.76–1.33)		1.02 (0.74–1.40)	
>12 months	96	0.91 (0.77–1.09)		0.82 (0.67–1.00)		0.91 (0.73–1.15)	
BMI ≥ 25 <sup>d</sup>							
Nulliparous	169	Reference	0.64	Reference	0.05	Reference	0.07
Parous never breastfed	240	0.94 (0.85–1.04)		0.98 (0.87–1.11)		1.06 (0.93–1.20)	
0–6 months	87	1.03 (0.91–1.17)		0.92 (0.79–1.08)		0.91 (0.77–1.07)	
7–12 months	61	1.06 (0.92–1.22)		0.93 (0.78–1.11)		0.89 (0.74–1.07)	
>12 months	145	0.98 (0.88–1.09)		0.89 (0.78–1.01)		0.93 (0.81–1.07)	
P <sub>interaction</sub>		<b>0.04</b>		0.13		0.20	

Note: Bold indicates statistical significance ( $P < 0.05$ ), some CI may not appear significant due to rounding.

Abbreviations: BMI, body mass index; CI, confidence intervals; MBD, mammographic breast density.

<sup>a</sup>Mammographic breast density measures were log<sub>10</sub> transformed because of the skewed distribution. Coefficients were backtransformed.

<sup>b</sup>Models were adjusted for age (continuous), age at menarche (continuous), BMI (continuous), race (non-Hispanic White, non-Hispanic Black, other), and family history of breast cancer (yes, no).

<sup>c</sup>Models were adjusted for age (continuous), age at menarche (continuous), BMI (continuous), and family history of breast cancer (yes, no). Other races were excluded from the interaction analysis because of the small sample size.

<sup>d</sup>Models were adjusted for age (continuous), age at menarche (continuous), race (non-Hispanic White, non-Hispanic Black, other), and family history of breast cancer (yes, no).

compared with nulliparous women, which appears to be driven by women with BMI < 25 kg/m<sup>2</sup>. Breastfeeding was also inversely associated with DV in the analyses involving all study participants and with NDV in analyses limited to parous women.

Our findings with VPD are consistent with the current literature (6, 10–12). Tehranifar and colleagues observed sim-

ilar associations between breastfeeding duration and percent density when assessing digitized films utilizing Cumulus software to calculate area measures of MBD (10). They investigated this association in a comparable study population including non-Hispanic Black and White women in their analyses (10). Although their results are similar to ours, our sample size was larger,  $N = 964$  compared with  $N = 191$  in their study, which

**Table 4.** Associations of breastfeeding duration with MBD measures by menopause status, race, and BMI among parous women.<sup>a</sup>

	N	Volumetric percent density		Dense volume (cm <sup>3</sup> )		Non-dense volume (cm <sup>3</sup> )	
		10 <sup>B</sup> (95% CI)	P <sub>trend</sub>	10 <sup>B</sup> (95% CI)	P <sub>trend</sub>	10 <sup>B</sup> (95% CI)	P <sub>trend</sub>
<i>Menopausal status</i>							
Premenopausal <sup>b</sup>							
Parous never breastfed	157	Reference	0.16	Reference	0.23	Reference	<b>0.02</b>
0-6 months	63	1.04 (0.90-1.19)		1.10 (0.96-1.27)		1.09 (0.94-1.25)	
7-12 months	46	1.15 (0.98-1.35)		1.00 (0.85-1.17)		0.88 (0.75-1.03)	
>12 months	180	1.07 (0.96-1.20)		0.95 (0.85-1.06)		<b>0.89 (0.80-1.00)</b>	
Postmenopausal <sup>b</sup>							
Parous never breastfed	132	Reference	0.49	Reference	0.98	Reference	0.68
0-6 months	52	0.94 (0.82-1.08)		0.86 (0.67-1.11)		0.92 (0.71-1.18)	
7-12 months	45	1.10 (0.95-1.28)		1.04 (0.80-1.37)		0.92 (0.70-1.21)	
>12 months	61	1.02 (0.89-1.16)		0.97 (0.75-1.24)		0.96 (0.74-1.23)	
P <sub>interaction</sub>		0.52		0.20		0.31	
<i>Race</i>							
Non-Hispanic White <sup>c</sup>							
Parous never breastfed	140	Reference	0.50	Reference	0.45	Reference	0.31
0-6 months	73	0.93 (0.81-1.07)		0.97 (0.82-1.16)		1.07 (0.90-1.29)	
7-12 months	66	1.09 (0.94-1.26)		1.02 (0.85-1.23)		0.94 (0.78-1.14)	
>12 months	190	1.01 (0.91-1.13)		0.94 (0.82-1.08)		0.95 (0.82-1.10)	
Non-Hispanic Black <sup>c</sup>							
Parous never breastfed	144	Reference	<b>0.04</b>	Reference	0.87	Reference	0.18
0-6 months	38	1.10 (0.96-1.27)		1.00 (0.81-1.23)		0.89 (0.72-1.11)	
7-12 months	20	1.12 (0.93-1.33)		1.11 (0.84-1.46)		0.99 (0.75-1.30)	
>12 months	34	1.14 (0.99-1.33)		0.99 (0.79-1.24)		0.85 (0.67-1.06)	
P <sub>interaction</sub>		0.42		0.91		0.70	
<i>BMI</i>							
BMI < 25 <sup>d</sup>							
Parous never breastfed	49	Reference	0.12	Reference	0.34	Reference	0.56
0-6 months	28	0.91 (0.70-1.17)		1.23 (0.92-1.65)		1.38 (0.99-1.91)	
7-12 months	30	1.20 (0.92-1.56)		<b>1.44 (1.07-1.94)</b>		1.17 (0.84-1.64)	
>12 months	96	1.12 (0.92-1.38)		1.15 (0.91-1.44)		1.00 (0.77-1.29)	
BMI ≥ 25 <sup>d</sup>							
Parous never breastfed	240	Reference	0.18	Reference	0.13	Reference	<b>0.04</b>
0-6 months	87	1.10 (0.98-1.23)		0.93 (0.81-1.08)		0.86 (0.73-1.01)	
7-12 months	61	1.13 (0.99-1.29)		0.94 (0.80-1.12)		0.84 (0.70-1.01)	
>12 months	145	1.06 (0.96-1.17)		0.90 (0.79-1.03)		0.87 (0.76-1.00)	
P <sub>interaction</sub>		0.09		0.12		0.10	

Note: Bold indicates statistical significance ( $P < 0.05$ ), some CI may not appear significant due to rounding.

Abbreviations: BMI, body mass index; CI, confidence intervals; MBD, mammographic breast density.

<sup>a</sup>Mammographic breast density measures were log<sub>10</sub> transformed because of the skewed distribution. Coefficients were backtransformed.

<sup>b</sup>Models were adjusted for age (continuous), age at menarche (continuous), BMI (continuous), race (non-Hispanic White, non-Hispanic Black, other), and family history of breast cancer (yes, no).

<sup>c</sup>Models were adjusted for age (continuous), age at menarche (continuous), BMI (continuous), and family history of breast cancer (yes, no). Other races were excluded from the interaction analysis because of the small sample size.

<sup>d</sup>Models were adjusted for age (continuous), age at menarche (continuous), race (non-Hispanic White, non-Hispanic Black, other), and family history of breast cancer (yes, no).

allowed us to stratify by race. Similar to the mostly null results between breastfeeding and VPD in our analyses limited to parous women, Yaghjian and colleagues also found no association between breastfeeding duration and percent density among parous premenopausal and postmenopausal women in the Nurses' Health Study, which also used digitized mammograms and Cumulus software to measure percent density based on breast area (6). Another study in Korea observed no significant associations between breastfeeding and percent density within twins (12). All of the aforementioned studies measured percent density using digitized mammograms and calculated area-based measures of MBD while we, to the best of

our knowledge, are the first study to use volumetric measures of MBD calculated using Volpara.

We observed an inverse association between breastfeeding duration and DV such that women who breastfed for >12 months had lower DV than nulliparous women who had never breastfed. Similar to our study, a study, by Li and colleagues, found an inverse association between breastfeeding and dense area among Chinese women, but a positive association among Australian women (14). Other studies that investigated the association between breastfeeding duration and dense area observed positive associations (6, 12, 14). Yaghjian and colleagues found a positive association between

breastfeeding duration and dense area particularly among premenopausal women (6). Because this study was conducted among women in the Nurses' Health Study, it is possible that study subjects are demographically and socioeconomically different from our study population.

Breastfeeding duration may be influenced by various factors including educational background and socioeconomic status (15, 26). The relationship between breastfeeding duration and dense area/volume may not be transportable across the study populations due to these potential demographic and socioeconomic differences. Other studies that observed a positive association between breastfeeding duration and dense area were conducted among women who lived in countries other than the United States (12, 14). Breastfeeding behaviors differ by not only race but also country (27). Non-Hispanic Black women breastfeed for shorter durations of time compared with women of other race/ethnicities in the United States (17, 21). Our study also showed that approximately half of parous women who never breastfed were non-Hispanic Black, but there was no difference in the association between breastfeeding duration and DV by race in this study. Also MBD may be different by ethnicity. Because the association between breastfeeding duration and MBD is not consistent across race and geographical location, it is possible breastfeeding may be influencing dense area/volume differently across races other than non-Hispanic Black versus non-Hispanic White.

We did not observe any associations between breastfeeding duration and NDV in analyses involving all study participants, but we observed an inverse trend between breastfeeding duration and NDV in the analyses limited to parous women. After stratifying by menopausal status, there is no association between breastfeeding duration and NDV among postmenopausal parous women, which is similar to findings in the Mexican Teacher's Cohort and Nurses' Health Study (6, 13). Our study found an inverse association between breastfeeding and NDV in premenopausal parous women, which differs from the nonsignificant findings in the Mexican teacher's cohort and positive association observed in the Nurses' Health Study (6, 13). Observations of an inverse association between breastfeeding duration and NDV could potentially be explained by changes in adiposity from breastfeeding, but this relationship is not clearly established (28). Future research is necessary to understand why these differences across the studies are present.

Women who are parous and breastfeed experience structural, cellular, and DNA changes in the breast that may prevent the development of certain subtypes of breast cancer (4, 29–31). Breastfeeding is protective against breast cancer beyond the impact of parity and mechanisms such as reduction in lifetime ovulatory cycles, which reduces exposure to estrogen may explain this effect (5, 32). A meta-analysis investigating the association between breastfeeding and breast cancer by subtype found the strongest inverse association between breastfeeding and ER<sup>-</sup> breast cancer (31, 33). Breastfeeding duration is often higher in low- and middle-income countries compared with

high-income countries but the inverse relationship with ER<sup>-</sup> cancer remains consistent (34–36). Certain changes in the breast tissue appear to benefit women who breastfed for long durations (>12 months), hence breastfeeding has only been consistently associated with a reduction in breast cancer risk among women who breastfeed for long durations (4). Given that MBD is a strong risk factor for breast cancer and the inconsistencies on the associations of breastfeeding with MBD, especially percent density, it is possible that other features of breast parenchyma may mediate the association between breastfeeding and breast cancer development.

### Strengths/limitations

There are several strengths to this study including a relatively large sample size with information on reproductive and demographic variables that were assessed as confounders and effect modifiers. An additional strength is that this study has a modest sample size of non-Hispanic Black women, which provided the opportunity to investigate the association between breastfeeding and MBD measures among non-Hispanic Black women and non-Hispanic White women separately. Also, to our knowledge, this study is the first to explore the association between breastfeeding duration and volumetric measures of MBD. Previous studies have mainly utilized area-based and qualitative measures of MBD. Volumetric and area measures of percent density have been found to be highly correlated ( $r = 0.93$ ), but the correlation between DV and dense area is not as strong ( $r = 0.55$ ; ref. 37). Area-based measures may be impacted by compression and density may appear different visually (38).

There are limitations of this study including potential misclassification of the exposure. Breastfeeding duration was categorized into approximately 3- to 6-month time periods for the first, second, and last children. Categories were combined across children to achieve the final breastfeeding duration for each woman. This may lead to a loss of information and potential misclassification of the exposure if women breastfed the minimum duration of each breastfeeding category per child. It may also mitigate the presence of a linear relationship. Because breastfeeding appears to have a threshold effect with breast cancer, benefitting women who breastfeed for very long durations (>12 months) the most, categorizing breastfeeding duration should not impact the findings in this study (4). In addition, breastfeeding duration data was available only for the first, second, and last children. Participants who had more than 3 children may be misclassified into lower breastfeeding duration categories if they breastfed the children born between the second and last for a longer duration. However, the majority of study participants had three children or less and the unavailability of data for other children is unlikely to have any impact on the results given the consistently null findings. Also, we are unable to differentiate between women who breastfed exclusively compared with those who did not. Future studies with information on exclusive breastfeeding practices can evaluate this. It is possible the full extent of the association between

breastfeeding and MBD was not apparent in our study due to unmeasured confounding or inability to adjust for certain variables because of potential collinearity, especially between parity and breastfeeding duration even after performing analyses on parous women separately. Because we investigated the association between breastfeeding duration and several measures of MBD and assessed this association across potential effect modifiers it is possible, due to the multiple comparisons, significant associations observed were due to chance. Finally, although this study had a relatively large sample of non-Hispanic Black and non-Hispanic White women, there was a limited number of women from other races/ethnic groups, hence, study results may not be generalizable to them.

In conclusion, we observed an inverse association between breastfeeding and VPD for women who breastfeed for 0–6 months, but not women who breastfeed for longer periods. Breastfeeding may protect against breast cancer development by acting through other features and patterns within the breast parenchyma that are not captured in volumetric measures of MBD. Future studies looking at the associations of breastfeeding with breast radiomics and microscopic features are needed.

### Authors' Disclosures

K.R. Getz reports grants from NIH/NCI during the conduct of the study. No disclosures were reported by the other authors.

### References

- Boyd NF, Guo H, Martin LJ, Sun L, Stone J, Fishell E, et al. Mammographic density and the risk and detection of breast cancer. *N Engl J Med* 2007;356:227–36.
- McCormack VA, dos Santos Silva I. Breast density and parenchymal patterns as markers of breast cancer risk: a meta-analysis. *Cancer Epidemiol Biomarkers Prev* 2006;15:1159–69.
- Kerlikowske K, Ichikawa L, Miglioretti DL, Buist DS, Vacek PM, Smith-Bindman R, et al. Longitudinal measurement of clinical mammographic breast density to improve estimation of breast cancer risk. *J Natl Cancer Inst* 2007;99:386–95.
- Faupel-Badger JM, Arcaro KF, Balkam JJ, Eliassen AH, Hassiotou F, Lebrilla CB, et al. Postpartum remodeling, lactation, and breast cancer risk: summary of a National Cancer Institute-sponsored workshop. *J Natl Cancer Inst* 2013;105:166–74.
- Collaborative Group on Hormonal Factors in Breast Cancer. Breast cancer and breastfeeding: collaborative reanalysis of individual data from 47 epidemiological studies in 30 countries, including 50302 women with breast cancer and 96973 women without the disease. *Lancet* 2002;360:187–95.
- Yaghjian L, Colditz GA, Rosner B, Bertrand KA, Tamimi RM. Reproductive factors related to childbearing and mammographic breast density. *Breast Cancer Res Treat* 2016;158:351–9.
- Lope V, Perez-Gomez B, Sanchez-Contador C, Santamarina MC, Moreo P, Vidal C, et al. Obstetric history and mammographic density: a population-based cross-sectional study in Spain (DDM-Spain). *Breast Cancer Res Treat* 2012;132:1137–46.
- Riza E, dos Santos Silva I, De Stavola B, Perry N, Karadedou-Zafiriadou E, Linos D, et al. Correlates of high-density mammographic parenchymal patterns by menopausal status in a rural population in Northern Greece. *Eur J Cancer* 2005;41:590–600.
- Titus-Ernstoff L, Tosteson AN, Kasales C, Weiss J, Goodrich M, Hatch EE, et al. Breast cancer risk factors in relation to breast density (United States). *Cancer Causes Control* 2006;17:1281–90.
- Tehrani P, Reynolds D, Flom J, Fulton L, Liao Y, Kudadjie-Gyamfi E, et al. Reproductive and menstrual factors and mammographic density in African American, Caribbean, and White women. *Cancer Causes Control* 2011;22:599–610.
- Butler LM, Gold EB, Greendale GA, Crandall CJ, Modugno F, Oestreicher N, et al. Menstrual and reproductive factors in relation to mammographic density: the study of women's health across the nation (SWAN). *Breast Cancer Res Treat* 2008;112:165–74.
- Sung J, Song YM, Stone J, Lee K, Lee D. Reproductive factors associated with mammographic density: a Korean co-twin control study. *Breast Cancer Res Treat* 2011;128:567–72.
- Rice MS, Bertrand KA, Lajous M, Tamimi RM, Torres G, Lopez-Ridaura R, et al. Reproductive and lifestyle risk factors and mammographic density in Mexican women. *Ann Epidemiol* 2015;25:868–73.
- Li T, Li J, Heard R, Gandomkar Z, Ren J, Dai M, et al. Understanding mammographic breast density profile in China: a Sino-Australian comparative study of breast density using real-world data from cancer screening programs. *Asia Pac J Clin Oncol* 2022;18:696–705.
- Kitsantas P, Gaffney KF, Kornides ML. Prepregnancy body mass index, socioeconomic status, race/ethnicity and breastfeeding practices. *J Perinat Med* 2011;40:77–83.
- McKinney CO, Hahn-Holbrook J, Chase-Lansdale PL, Ramey SL, Krohn J, Reed-Vance M, et al. Racial and ethnic differences in breastfeeding. *Pediatrics* 2016;138:e20152388.
- Li R, Perrine CG, Anstey EH, Chen J, MacGowan CA, Elam-Evans LD. Breastfeeding trends by race/ethnicity among US children born from 2009 to 2015. *JAMA Pediatr* 2019;173:e193319.

### Authors' Contributions

K.R. Getz: Formal analysis, writing—original draft, writing—review and editing. B. Adedokun: Writing—original draft. S. Xu: Formal analysis, writing—original draft, writing—review and editing. A.T. Toriola: Conceptualization, supervision, funding acquisition, writing—original draft, writing—review and editing.

### Acknowledgments

This work was supported by funding from NIH/NCI (R21CA216515, R01CA246592, to A.T. Toriola), and K.R. Getz was supported by T32CA190194. The content is solely the responsibility of the authors and does not represent the official view of the NIH.

We would like to thank all of the women who participated in this study.

The publication costs of this article were defrayed in part by the payment of publication fees. Therefore, and solely to indicate this fact, this article is hereby marked “advertisement” in accordance with 18 USC section 1734.

### Note

Supplementary data for this article are available at Cancer Prevention Research Online (<http://cancerprevres.aacrjournals.org/>).

Received November 1, 2022; revised January 23, 2023; accepted March 14, 2023; published first March 17, 2023.



18. Heller SL, Hudson S, Wilkinson LS. Breast density across a regional screening population: effects of age, ethnicity and deprivation. *Br J Radiol* 2015;88:20150242.
19. Moore JX, Han Y, Appleton C, Colditz G, Toriola AT. Determinants of mammographic breast density by race among a large screening population. *JNCI Cancer Spectr* 2020;4:pkaa010.
20. del Carmen MG, Halpern EF, Kopans DB, Moy B, Moore RH, Goss PE, et al. Mammographic breast density and race. *AJR Am J Roentgenol* 2007;188:1147–50.
21. Anstey EH, Chen J, Elam-Evans LD, Perrine CG. Racial and geographic differences in breastfeeding - United States, 2011–2015. *MMWR Morb Mortal Wkly Rep* 2017;66:723–7.
22. Anstey EH, Shoemaker ML, Barrera CM, O'Neil ME, Verma AB, Holman DM. Breastfeeding and breast cancer risk reduction: implications for black mothers. *Am J Prev Med* 2017;53:S40–S6.
23. Ellison-Loschmann L, McKenzie F, Highnam R, Cave A, Walker J, Jeffreys M. Age and ethnic differences in volumetric breast density in New Zealand women: a cross-sectional study. *PLoS One* 2013;8:e70217.
24. Lee HN, Sohn Y-M, Han KH. Comparison of mammographic density estimation by volpara software with radiologists' visual assessment: analysis of clinical–radiologic factors affecting discrepancy between them. *Acta Radiol* 2015;56:1061–8.
25. Han Y, Berkey CS, Herman CR, Appleton CM, Alimujiang A, Colditz GA, et al. Adiposity change over the life course and mammographic breast density in postmenopausal women. *Cancer Prev Res* 2020;13:475–82.
26. Heck KE, Braveman P, Cubbin C, Chavez GF, Kiely JL. Socioeconomic status and breastfeeding initiation among California mothers. *Public Health Rep* 2006;121:51–9.
27. Chung SH, Kim HR, Choi YS, Bae CW. Trends of breastfeeding rate in Korea (1994–2012): comparison with OECD and other countries. *J Korean Med Sci* 2013;28:1573–80.
28. Lambrinou CP, Karaglani E, Manios Y. Breastfeeding and postpartum weight loss. *Curr Opin Clin Nutr Metab Care* 2019;22:413–7.
29. Yang L, Jacobsen KH. A systematic review of the association between breastfeeding and breast cancer. *J Womens Health* 2008;17:1635–45.
30. Russo J, Moral R, Balogh GA, Mailo D, Russo IH. The protective role of pregnancy in breast cancer. *Breast Cancer Res* 2005;7:131–42.
31. Ambrosone CB, Higgins MJ. Relationships between breast feeding and breast cancer subtypes: lessons learned from studies in humans and in mice. *Cancer Res* 2020;80:4871–7.
32. Britt K, Ashworth A, Smalley M. Pregnancy and the risk of breast cancer. *Endocr Relat Cancer* 2007;14:907–33.
33. Islami F, Liu Y, Jemal A, Zhou J, Weiderpass E, Colditz G, et al. Breastfeeding and breast cancer risk by receptor status—a systematic review and meta-analysis. *Ann Oncol* 2015;26:2398–407.
34. Joko-Fru WY, Jedy-Agba E, Korir A, Ogunbiyi O, Dzamala CP, Chokunonga E, et al. The evolving epidemic of breast cancer in sub-Saharan Africa: results from the African cancer registry network. *Int J Cancer* 2020;147:2131–41.
35. Figueroa JD, Davis Lynn BC, Edusei L, Titiloye N, Adjei E, Clegg-Lamprey JN, et al. Reproductive factors and risk of breast cancer by tumor subtypes among Ghanaian women: a population-based case-control study. *Int J Cancer* 2020;147:1535–47.
36. Dey S, Boffetta P, Mathews A, Brennan P, Soliman A, Mathew A. Risk factors according to estrogen receptor status of breast cancer patients in Trivandrum, South India. *Int J Cancer* 2009;125:1663–70.
37. Brand JS, Czene K, Shepherd JA, Leifland K, Heddson B, Sundbom A, et al. Automated measurement of volumetric mammographic density: a tool for widespread breast cancer risk assessment. *Cancer Epidemiol Biomarkers Prev* 2014;23:1764–72.
38. Alonzo-Proulx O, Mawdsley GE, Patrie JT, Yaffe MJ, Harvey JA. Reliability of automated breast density measurements. *Radiology* 2015;275:366–76.

