

## Detection of assemblages A and B of *Giardia duodenalis* in water and sewage from São Paulo state, Brazil

Licia Natal Fernandes, Patrícia Pereira de Souza, Ronalda Silva de Araújo, Maria Tereza Pepe Razzolini, Rodrigo Martins Soares, Maria Inês Zanolli Sato, Elayse Maria Hachich, Silvana Audrá Cutolo, Glavur Rogério Matté and Maria Helena Matté

### ABSTRACT

*Giardia duodenalis* is a protozoan that parasitizes humans and other mammals and causes giardiasis. Although its isolates have been divided into seven assemblages, named A to G, only A and B have been detected in human faeces. Assemblage A isolates are commonly divided into two genotypes, AI and AII. Even though information about the presence of this protozoan in water and sewage is available in Brazil, it is important to verify the distribution of different assemblages that might be present, which can only be done by genotyping techniques. A total of 24 raw and treated sewage, surface and spring water samples were collected, concentrated and purified. DNA was extracted, and a nested PCR was used to amplify an 890 bp fragment of the *gdh* gene of *G. duodenalis*, which codes for glutamate dehydrogenase. Positive samples were cloned and sequenced. Ten out of 24 (41.6%) samples were confirmed to be positive for *G. duodenalis* by sequencing. Phylogenetic analysis grouped most sequences with *G. duodenalis* genotype AII from GenBank. Only two raw sewage samples presented sequences assigned to assemblage B. In one of these samples genotype AII was also detected. As these assemblages/genotypes are commonly associated to human giardiasis, the contact with these matrices represents risk for public health.

**Key words** | genotyping, *Giardia*, glutamate dehydrogenase, sewage, water

Licia Natal Fernandes (corresponding author)  
 Patrícia Pereira de Souza  
 Ronalda Silva de Araújo  
 Maria Tereza Pepe Razzolini  
 Silvana Audrá Cutolo  
 Glavur Rogério Matté  
 Maria Helena Matté  
 School of Public Health,  
 University of Sao Paulo, Av. Dr. Arnaldo,  
 715 - Cerqueira César - CEP 01246-904/São Paulo (SP),  
 Brazil  
 Tel.: 55-11-30617753  
 Fax: 55-11-3083 3501  
 E-mail: [licianatal@usp.br](mailto:licianatal@usp.br)

Ronaldo Silva de Araújo  
 Hospital da Clínicas, School of Medicine, University  
 of Sao Paulo, Av. Dr. Enéas de Carvalho Aguiar,  
 255 - Cerqueira César - CEP 05403-000/São Paulo (SP),  
 Brazil

Rodrigo Martins Soares  
 School of Veterinary Medicine and Zootechny,  
 University of Sao Paulo,  
 Av. Prof. Dr. Orlando Marques de Paiva,  
 87 - Cidade Universitária - CEP 05508-270/São  
 Paulo (SP),  
 Brazil

Maria Inês Zanolli Sato  
 Elayse Maria Hachich  
 CETESB - Sao Paulo State Company for Sanitation  
 and Technology, Av. Prof. Frederico Hermann Jr.,  
 345 - Alto de Pinheiros - CEP 05459-900/São Paulo (SP),  
 Brazil

Silvana Audrá Cutolo  
 Polytechnic School, University of São Paulo,  
 Av. Prof. Luciano Gualberto,  
 Travessa 3 n°380 - Cidade Universitária - CEP  
 05508-970/ São Paulo (SP),  
 Brazil

### INTRODUCTION

*Giardia duodenalis* (syn. *lamblia* or *intestinalis*) is a flagellated protozoan that parasitizes the small intestine of humans and other mammals, causing giardiasis. Isolates of this species are morphologically similar, but have been

divided into seven assemblages, named A to G, on the basis of genetic heterogeneity. To date only assemblages A and B have been found in humans (Thompson 2004; Monis *et al.* 2009).

doi: 10.2166/wh.2011.098

In developed countries, the prevalence of human giardiasis varies from 2 to 5%. In Brazil, studies show that it ranges from 16 to 26% (Mascarini & Donalísio 2006; Silva *et al.* 2009; Tashima *et al.* 2009).

*G. duodenalis* is a waterborne pathogen. Cysts of the organism can survive for long periods in water and are resistant to some of the processes used by the water industry, such as filtration and disinfection. To protect public health, therefore, it is important to search for *Giardia* cysts in water and sewage samples (Faubert 2000; Slifko *et al.* 2000; Smith *et al.* 2006; Carmena *et al.* 2007).

Only molecular characterization techniques can be used to differentiate *G. duodenalis* assemblages. These techniques have been used to characterize *Giardia* isolates from environmental sources in a number of countries (Cacciò *et al.* 2003; Sulaiman *et al.* 2004; Bertrand & Schwartzbrod 2007; Castro-Hermida *et al.* 2008; Plutzer *et al.* 2008).

In Brazil, this protozoan has been detected in water and sewage samples by immunofluorescence assay (Franco *et al.* 2001; Cantusio Neto & Franco 2004; Hachich *et al.* 2004; Heller *et al.* 2004; Cantusio Neto *et al.* 2006; Nishi *et al.* 2009; Razzolini *et al.* 2010); however, the molecular characterization of *Giardia* cysts from these sources has not yet been published. Thus, the significance of cysts in the environment for public health is not known in this country.

The objective of this study was to detect and characterize the genotypes of *Giardia* sp. in water and sewage samples from São Paulo state, Brazil, discussing the importance of findings for public health.

## MATERIALS AND METHODS

### Water and sewage samples

A total of 24 water and sewage samples were collected from May 2008 to March 2009 in São Paulo state, Brazil (18 in São Paulo Metropolitan Region and 6 in two small municipalities within the state: Piracicaba and São Lourenço da Serra).

Raw (6) and treated (6) sewage samples were collected at six different wastewater treatment plants that operate by stabilization ponds, activated sludge, anaerobic reactors or by a match of two of these systems. Effluent from these plants

is discharged into water bodies and, in some cases, is directed to non-potable industrial uses. Two raw sewage samples were collected at different entrances of a lake that has been used as a water supply.

Surface water grab samples (11) were taken from a reservoir, close to the abstraction point supplying a drinking water treatment plant. The reservoir is also used as a recreational area. A spring water sample (1) was also collected. All samples were processed within 24 hours after collection.

### Sample concentration

Samples (10 L) were concentrated by the modified membrane filtration technique (Araújo *et al.* 2010), which consists of filtration and centrifugation steps. Membranes with pores of 1 µm and diameters of 14 cm (Millipore<sup>®</sup>, Brazil) were employed. Pre-filters were also required for treated sewage filtration.

Raw sewage samples (100 ml) were concentrated by centrifugation. Sewage was distributed into 50 ml conical tubes and centrifuged at  $7428 \times g$  for 15 minutes at 24°C. The sediments were resuspended and pooled into one tube and then washed with ultra-pure water (Milli Q<sup>®</sup>). The suspension was centrifuged ( $7.428 \times g/15'/24^\circ\text{C}$ ) and supernatant was discharged.

### Sample purification

Sediment obtained from the concentration step was purified by fluctuation in zinc sulfate solution (33%) according to Faust *et al.* (1938), with some modifications. Briefly, sediment was washed with ultra-pure water (Milli Q<sup>®</sup>). Supernatant was discharged after centrifugation ( $7428 \times g/15'/24^\circ\text{C}$ ). Ten millilitres of zinc sulfate solution was added. After homogenization, tubes were centrifuged ( $3714 \times g/3'/24^\circ\text{C}$ ). The pellet was removed with a 10 mL pipette. The supernatant left in the tube (2 ml) was washed twice with ultra-pure water (Milli Q<sup>®</sup>). The pellet was resuspended in ultra-pure water (Milli Q<sup>®</sup>).

### DNA extraction and amplification


Genomic DNA was extracted using the phenol-chloroform-isomilic alcohol protocol (Araújo *et al.* 2010), with small

modifications as follows: all concentrated sediment was submitted to DNA extraction and the incubation period at 65°C was reduced to 80 minutes.

The PCR mix consisted of 25 ng of DNA, 10 × buffer, 2 mM of MgCl<sub>2</sub>, 1% of PVP, 300 μM of deoxynucleoside triphosphate mix, 0.5 μM of each primer, 1.25 U of *Tth* DNA polymerase (Biotools<sup>®</sup>, Madrid, Spain) and ultra-pure water (Milli Q<sup>®</sup>) to make up to a final volume of 25 μl. The same PCR mix, but without the PVP, was used for the second amplification. Amplification primers and conditions are described in Table 1.

The nested PCR product was resolved by electrophoresis using a 1.8% agarose gel prepared with 1 × TAE buffer and stained with ethidium bromide. A 100 bp DNA Ladder Plus (Fermentas) was included as a size marker. Gel images were obtained by UV transillumination and captured by a gel documentation system (Epi Chemi II Darkroom and Software Labworks – UVP).

Following successful PCR, the products were purified (Illustratm GFXtm PCR DNA and Gel Band Purification Kit, GE<sup>®</sup> Healthcare, UK). Molecular cloning and sequencing were done according to the methods of Ivanova et al. (2001). Analyses of 825 positions of the *gdh* gene sequences of the clones were compared with their closest relatives in the GenBank database by BLAST searches (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). Further phylogenetic and molecular evolutionary analyses were performed with the MEGA 4.1 program (Tamura et al. 2007).



**Figure 1** | Ethidium bromide stained agarose gel electrophoresis of the 890 bp fragment amplified from *gdh* gene of *Giardia duodenalis*. Lane 1: 100 bp DNA Ladder Plus (Fermentas); Lane 2: negative control; Lane 3: positive control; Lane 4: raw sewage positive sample; Lane 5: treated sewage positive sample; Lane 6: surface water positive sample.

## RESULTS AND DISCUSSION

PCR amplification of the 890 bp *gdh* gene fragment was detected in 14 (58.3%) out of 24 water and sewage samples analysed (Figure 1). Eleven of the positive samples were cloned and sequenced; ten (41.6%) were confirmed to be

**Table 1** | Primers sequence and PCR cycles used for amplification of *Giardia gdh* genes from environmental samples

PCR reaction	Primer ID	Sequence*	PCR conditions							Band size
			ID	Cycles	D	A	E	FE		
1 <sup>st</sup>	GdhR (F)	5'CGAGCGCGAGCCGAAGT ATATCC3'	95°C, 7 min	40	95°C, 2 min	50°C 1 min	72°C 1.5 min	72°C 5 min	1190 bp	
	#579II (R)	5'GATGTTYGCRCCCATCTG RTAGTTC3'								
2 <sup>nd</sup>	GDH-FI (F)	5'AAYGAGGTYATGCGCTT CTGCCA3'	94°C, 2 min	30	94°C, 35 s	56°C 35 s	72°C 1 min		890 bp	
	#579II (R)	5'GATGTTYGCRCCCATCT GRTAGTTC3'								

ID = Initial denaturation, D = Denaturation, A = Annealing, E = Extension, FE = Final Extension and \* = position 659–1485 from the start of the gene.

positive by sequencing (Table 1). One to ten clones of each sample were sequenced, resulting in a total of 71 nucleotide sequences. Some samples had clones with identical sequences, therefore only 64 nucleotide sequences were submitted to GenBank (GenBank accession numbers: GQ503101–GQ503141, GQ503152–GQ503157 and GQ503159–GQ503174). Phylogenetic analyses grouped most of the 64 sequences with *G. duodenalis* genotype AII sequences from GenBank. Only two raw sewage samples contained sequences assigned to assemblage B. In one of

these samples, both genotype AII and assemblage B were detected (Table 2, Figure 2).

The presence of *Giardia* in water and sewage is well documented worldwide (Karaniš *et al.* 2006; Robertson *et al.* 2006; Anceno *et al.* 2007; Bertrand & Schwartzbrod 2007; Castro-Hermida *et al.* 2008; Plutzer *et al.* 2008). In Brazil, the detection of this protozoan has also been documented in some studies (Franco *et al.* 2000; Cantusio Neto & Franco 2004; Hachich *et al.* 2004; Heller *et al.* 2004; Cantusio Neto *et al.* 2006; Nishi *et al.* 2009; Razzolini *et al.* 2010).

**Table 2** | PCR, sequencing and genotyping results of *Giardia* isolates detected in water and sewage samples from São Paulo, Brazil

Sample origin	Sample number	PCR results	Number of sequenced clones	Sequencing results Number of clones successfully sequenced/GenBank accession numbers	
				AII	B
Raw sewage	1	P	5	5/GQ503101–GQ503104 <sup>a</sup>	0
	2	P	10	0	10/GQ503106–GQ503115
	3	P	9	9/GQ503116–503120 <sup>b</sup>	0
	4	P	4	4/GQ503121–GQ503124	0
	5	N	–	–	–
	6	P	7	6/GQ503125–GQ503130	1/GQ503131
Treated sewage	7	N	–	–	–
	8	P	–	–	–
	9	P	–	–	–
	10	P	9	9/GQ503152–GQ503158 <sup>c</sup>	0
	11	P	1	–	–
	12	N	–	–	–
Surface water	13	P	8	1/GQ503105	0
	14	N	–	–	–
	15	N	–	–	–
	16	N	–	–	–
	17	N	–	–	–
	18	N	–	–	–
	19	P	10	10/GQ503132–GQ503141	0
	20	P	10	10/GQ503159–GQ503168	0
	21	P	6	6/GQ503169–GQ503174	0
	22	N	–	–	–
23	N	–	–	–	
Spring water	24	N	–	–	–


P = positive sample

N = negative sample

<sup>a</sup> = 2 identical sequences (GQ503102)

<sup>b</sup> = 5 identical sequences (GQ503116)

<sup>c</sup> = 3 identical sequences (GQ503154)



**Figure 2** | Phylogenetic tree based on the alignment of 64 nucleotide sequences of *gdh* gene fragment. Tree was constructed by Neighbour-Joining method in MEGA 4.1 Beta program using maximum composite likelihood model and bootstrap of 1000 repetitions. Gene Bank reference sequences were: All (EF685688, AY178736, AY178737), AI (AY178735, M846004), B (AY178738, AY178739, AY178749, AY178750, AY178751, AY178752, AY178753, AY178754, AY178755, AY178756), C (U60982, U60983), D (U60986), E (AY178740, AY178741), F (AY178742, AY178743, AY178744, AF069057), G (AY178745, AY178746, AY178747, AY178748, AF069058) and *G. ardeae* (AF069060).

In the present study, 10 out of 24 (41.6%) water and sewage samples analysed were found to be positive for *G. duodenalis*, corroborating previous reports that *G. duodenalis* is the only species of *Giardia* found in water sources.

Data on the genotypes of *Giardia* isolates found in Brazil are scarce and mainly derived from analysis of human and animal stool specimens (Rocha et al. 2003; Souza et al. 2007; Voloão et al. 2007, 2008). Although Paulino (2005) employed the technique of PCR to detect *G. duodenalis* in surface water samples, genotyping of the isolates was not performed.

The present study is the first to genotype *G. duodenalis* from environmental samples in Brazil and to demonstrate that assemblages A (genotype AII) and B of *G. duodenalis* are present in different water sources including surface water and raw and treated sewage. These findings are similar to those

reported worldwide demonstrating the presence of assemblages A and B in environmental samples (Cacciò et al. 2003; Sulaiman et al. 2004; Robertson et al. 2006; Anceno et al. 2007; Bertrand & Schwartzbrod 2007; Castro-Hermida et al. 2008; Plutzer et al. 2008).

As observed in other studies, assemblage A of *G. duodenalis* was more frequently observed than assemblage B, which was detected in only two raw sewage samples (Cacciò et al. 2003; Sulaiman et al. 2004; Bertrand & Schwartzbrod 2007). In a study carried out in São Paulo, in 2007, Souza et al. (2007) demonstrated that genotype AII was observed in 29 out of 32 clinical samples analysed, and assemblage B in only eight samples. This finding indicates that, in the present study, genotype AII observed in sewage samples may represent the genotype circulating in the population.

An unexpected number of Single Nucleotide Polymorphisms (SNPs) was detected in the majority of sequences obtained in this work. Most sequences had from one to five SNPs. None of these SNPs was observed in sequences recovered from GenBank. Besides, most SNPs resulted in non-synonym substitutions. Once sequencing presented high-quality and very well-defined peaks, these substitutions were not considered to be real, but were assumed to have occurred due to nucleotide incorporation problems during PCR, possibly because *Tth* DNA polymerase, a low-fidelity enzyme, was used.

The presence of *G. duodenalis* genotypes associated to human giardiasis in raw sewage indicates that there is risk in the reuse of sub-products of the sewage treatment process as observed in Table 2. Detection of genotype AII and assemblage B in treated sewage reveals that more attention should be given to sewage treatment processes, since effluents of sewage treatment plants can be used for domestic, agricultural or industrial purposes. Furthermore, this matrix can be released in surface waters that are subjected to indirect reuse including drinking water treatment. Sewage sludge may contain  $10^5$  to  $10^6$  *Giardia* cysts/kg<sup>-1</sup> (Straub et al. 1993; Thiriat et al. 1997), and even though this matrix was not analysed in the present work it will be important to characterize *G. duodenalis* isolates recovered from this kind of sample, especially because of the lack of data (Rimhanen-Finne et al. 2001).

The detection of genotype AII in surface waters also represents risk for public health because this matrix is commonly used as recreation areas or as a source of water supply for humans.

## CONCLUSIONS

The presence of genotype AII and assemblage B, commonly associated to human giardiasis, in water and sewage samples from São Paulo state indicates that contact with these matrices is a risk to public health. Therefore, more attention should be given to water and sewage treatment processes and to the uses of treated sewage and its discharges in watersheds, which should be protected against sources of contamination.

Studies of detection, quantification, infectivity and genotyping of *G. duodenalis* cysts from environmental samples should be encouraged in order to provide enough data for the decisions that need to be taken by public health authorities.

## ACKNOWLEDGEMENTS

This work was financially supported by FINEP, Financiadora de Estudos e Projetos, (Project: 0934/07) and CNPq, Conselho Nacional de Desenvolvimento Científico e Tecnológico (process: 134810/2007-4).

## REFERENCES

- Anceno, A. J., Katayama, H., Houpt, E. R., Chavalitshewinkoon-Petmitr, P., Chuluun, B. & Shipin, O. V. 2007 *IMS-free DNA extraction for the PCR-based quantification of *Cryptosporidium parvum* and *Giardia lamblia* in surface and waste water*. *Int. J. Environ. Health Res.* **17**(4), 297–310.
- Araújo, R. S., Carvalho, T. T. R., Matté, G. R., Fernandes, L. N., Balsalobre, L. C. & Matté, M. H. 2010 Modified method for detection of *Cryptosporidium* oocysts using DNA templates extracted from environmental samples. *Revista do Instituto Adolfo Lutz*. In press.
- Bertrand, I. & Schwartzbrod, J. 2007 *Detection and genotyping of *Giardia duodenalis* in wastewater: relation between assemblages and faecal contamination origin*. *Water Res.* **41**(16), 3675–3682.
- Cacciò, S. M., De Giacomo, M., Alicino, F. A. & Pozio, E. 2003 **Giardia* cysts in wastewater treatment plants in Italy*. *Appl. Environ. Microbiol.* **69**(6), 3393–3398.
- Cantusio Neto, R. & Franco, R. M. B. 2004 *Occurrence of *Giardia* cysts and *Cryptosporidium* oocysts in activated sludge samples in Campinas, SP, Brazil (Ocorrência de oocistos de *Cryptosporidium* spp. e cistos de *Giardia* spp. em diferentes pontos do processo de tratamento de água, em Campinas, São Paulo, Brasil)*. *Higiene Alimentar.* **18**, 52–59.
- Cantusio Neto, R., Santos, J. U. & Franco, R. M. B. 2006 *Evaluation of activated sludge treatment and the efficiency of the disinfection of *Giardia* species cysts and *Cryptosporidium* oocysts by UV at a sludge treatment plant in Campinas, south-east Brazil*. *Water Sci. Technol.* **54**(3), 89–94.
- Carmena, D., Aguinalgalde, X., Zigorraga, C., Fernández-Crespo, J. C. & Ocio, J. A. 2007 *Presence of *Giardia* cysts and *Cryptosporidium* oocysts in drinking water supplies in northern Spain*. *J. Appl. Microbiol.* **102**, 619–629.
- Castro-Hermida, J. A., García-Preseido, I., Almeida, A., González-Warleta, M., Da Costa, J. M. C. & Mezo, M. 2008 *Contribution of treated wastewater to the contamination of recreational river areas with *Cryptosporidium* spp. and *Giardia duodenalis**. *Water Res.* **42**, 3528–3538.
- Faubert, G. 2000 *Immune response to *Giardia duodenalis**. *Clin. Microbiol. Rev.* **13**(1), 35–54.
- Faust, E. C., D'Antoni, J. S., Odom, V., Miller, M. J., Peres, C., Sawitz, W., Thomen, L. F., Tobie, J. E. & Walker, J. H. 1938 *A critical study of clinical laboratory technics for the diagnosis of protozoan cysts and helminth eggs in feces I. Preliminary communication*. *Am. J. Trop. Med.* **18**, 169–183.

- Franco, R. M., Rocha-Eberhardt, R. & Cantusio Neto, R. 2001 Occurrence of *Cryptosporidium* oocysts and *Giardia* cysts in raw water from the Atibaia river, Campinas, Brazil. *Rev. Inst. Med. Trop.* **43**(2), 109–111.
- Hachich, E. M., Sato, M. I., Galvani, A. T., Menegon, J. R. & Mucci, J. L. 2004 *Giardia* and *Cryptosporidium* in source waters of São Paulo State, Brazil. *Water Sci. Technol.* **50**(1), 239–245.
- Heller, L., Bastos, R. K. X., Vieira, M. B. C. M., Bevilacqua, P. D., de Brito, L. L. A., Mota, S. M. M., Oliveira, A. A., Machado, P. M., Salvador, D. P. & Cardoso, A. B. 2004 *Cryptosporidium* oocysts and *Giardia* cysts: environmental circulation and health risks (Oocistos de *Cryptosporidium* e cistos de *Giardia*: circulação no ambiente e riscos à saúde humana). *Epidemiologia e Serviços de Saúde* **13**(2), 79–92.
- Ivanova, E. P., Romanenko, L. A., Matté, M. H., Matté, G. R., Lysenko, A. M., Simidu, U., Kita-Tsukamoto, K., Sawabe, T., Vysotskii, M. V., Frolova, G. M., Mikhailov, V., Christen, R. & Colwell, R. R. 2001 Retrieval of the species *Alteromonas tetraodonis* Simidu et al. 1990 as *Pseudoalteromonas tetraodonis* comb. nov. and emendation of description. *Int. J. Syst. Evol. Microbiol.* **51**(3), 1071–1078.
- Karanis, P., Sotiriadou, I., Kartashev, V., Kourenti, C., Tsvetkova, N. & Stojanova, K. 2006 Occurrence of *Giardia* and *Cryptosporidium* in water supplies of Russia and Bulgaria. *Environ. Res.* **102**, 260–271.
- Mascarini, L. M. & Donalísio, M. R. 2006 Giardiasis and cryptosporidiosis in children institutionalized at daycare centers in the State of São Paulo (Giardiase e cryptosporidiose em crianças institucionalizadas em creches no Estado de São Paulo). *Rev. Soc. Bras. Med. Trop.* **39**(6), 577–579.
- Monis, P. T., Caccio, S. M. & Thompson, R. C. A. 2009 Variation in *Giardia*: towards a taxonomic revision of the genus. *Trends Parasitol.* **25**(2), 93–100.
- Nishi, L., Baesso, M. L., Santana, R. G., Fregadolli, P., Falavigna, D. L. & Falavigna-Guilherme, A. L. 2009 Investigation of *Cryptosporidium* spp. and *Giardia* spp. in a public water-treatment system. *Zoonoses Public Health* **56**(5), 221–228.
- Paulino, R. C. 2005 Molecular detection of *Giardia* sp. from faecal and water samples: genomic DNA extraction, PCR and RFLP (Thesis) (Detecção molecular de *Giardia* sp. em amostras fecais e água: extração de DNA genômico, PCR e RFLP). Universidade Federal do Paraná, Curitiba (PR), Brazil.
- Plutzer, J., Karanis, P., Domokos, K., Torokné, A. & Márialigeti, K. 2008 Detection and characterization of *Giardia* and *Cryptosporidium* in Hungarian raw, surface and sewage water samples by IFT, PCR and sequence analysis of the SSU rRNA and GDH genes. *Int. J. Hyg. Environ. Health* **211**(5–6), 524–533.
- Razzolini, M. T., da Silva Santos, T. F. & Bastos, V. K. 2010 Detection of *Giardia* and *Cryptosporidium* cysts/oocysts in watersheds and drinking water sources in Brazil urban areas. *J. Water Health* **8**(2), 399–404.
- Rimhanen-Finne, R., Ronkainen, P. & Hanninen, M.-L. 2001 Simultaneous detection of *Cryptosporidium parvum* and *Giardia* in sewage sludge by IC-PCR. *J. Appl. Microbiol.* **91**, 1030–1035.
- Robertson, L. J., Hermansen, L., Gjerde, B. K., Strand, E., Alvsvag, J. O. & Langeland, N. 2006 Application of genotyping during an extensive outbreak of waterborne giardiasis in Bergen, Norway, during autumn and winter 2004. *Appl. Environ. Microbiol.* **72**(3), 2212–2217.
- Rocha, M. O., Gomes, M. A., Costa, A. O., Furst, C. & Silva, E. F. 2003 Molecular characterization of Brazilian human *Giardia duodenalis* isolates using isoenzyme and random amplified polymorphic DNA analysis. *Diagn. Microbiol. Infect. Dis.* **46**, 273–278.
- Silva, R. R., da Silva, C. A. M., Pereira, C. A. J., Nicolato, R. L. C., Negrão-Corrêa, D., Lamounier, J. A. & Carneiro, M. 2009 Association between nutritional status, environmental and socio-economic factors and *Giardia lamblia* infections among children aged 6–71 months in Brazil. *Trans. R. Soc. Trop. Med. Hyg.* **103**(5), 512–519.
- Slifko, T. R., Smith, H. V. & Rose, J. B. 2000 Emerging parasite zoonoses associated with water and food. *Int. J. Parasitol.* **30**, 1379–1393.
- Smith, H. V., Cacciò, S. M., Tait, A., Mclachlin, J. & Thompson, R. C. A. 2006 Tools for investigating the environmental transmission of *Cryptosporidium* and *Giardia* infection in humans. *Trends Parasitol.* **22**(4), 160–167.
- Souza, S. L. P., Gennari, S. M., Richtzenhain, L. J., Pena, H. F. J., Funada, M. R., Cortez, A., Gregori, F. & Soares, R. M. 2007 Molecular identification of *Giardia duodenalis* isolates from humans, dogs, cats and cattle from the state of São Paulo, Brazil, by sequence analysis of fragment of glutamate dehydrogenase (*gdh*) coding gene. *Vet. Parasitol.* **149**(3–4), 258–264.
- Straub, T. M., Pepper, I. L. & Gerba, C. P. 1993 Hazards from pathogenic microorganisms in land-disposed sewage sludge. In: *Reviews of Environmental Contamination and Toxicology*, Ware, G. W. (ed.), Springer-Verlag, New York, pp. 55–91.
- Sulaiman, I. M., Jiang, J., Singh, A. & Xiao, L. 2004 Distribution of *Giardia duodenalis* genotypes and subgenotypes in raw urban wastewater in Milwaukee, Wisconsin. *Appl. Environ. Microbiol.* **70**, 3776–3780.
- Tamura, K., Dudley, J., Nei, M. & Kumar, S. 2007 MEGA 4: molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol. Biol. Evol.* **24**, 1596–1599.
- Tashima, N. T., Simões, M. J., Leite, C. Q., Fluminhan, A., Nogueira, M. A. & Malaspina, A. C. 2009 Classic and molecular study of *Giardia duodenalis* in children from a daycare center in the region of Presidente Prudente, Sao Paulo, Brazil. *Rev. Inst. Med. Trop. Sao Paulo* **51**(1), 19–24.
- Thiriart, L., Bogot, V. & Schwartzbrod, J. 1997 Evaluation of a procedure for detection of viable *Giardia* cysts in wastewater sludge. *Water Sci. Technol.* **35**, 377–380.
- Thompson, R. C. A. 2004 The zoonotic significance and molecular epidemiology of *Giardia* and giardiasis. *Vet. Parasitol.* **126**, 15–35.
- Volotão, A. C., Costa-Macedo, L. M., Haddad, F. S., Brandão, A., Peralta, J. M. & Fernandes, O. 2007 Genotyping of *Giardia duodenalis* from human and animal samples from Brazil using beta-giardin gene: a phylogenetic analysis. *Acta Trop.* **102**, 10–19.
- Volotão, A. C. C., Souza Júnior, J. C., Grassini, C., Peralta, J. M. & Fernandes, O. 2008 Genotyping of *Giardia duodenalis* from Southern Brown Howler Monkeys (*Alouatta clamitans*) from Brazil. *Vet. Parasitol.* **158**, 133–137.