

Type II Diabetes Mellitus and Polymorphism of Insulin-Receptor Gene in Mexican Americans

SOUFIA H. RABOUDI, BRAXTON D. MITCHELL, MICHAEL P. STERN, CLAYTON W. EIFLER, STEVEN M. HAFFNER, HELEN P. HAZUDA, AND MARSHA L. FRAZIER

Resistance to insulin action is a well-established feature of type II (non-insulin-dependent) diabetes and is believed by many to contribute to the etiology of this condition. We therefore characterized restriction-fragment-length polymorphisms of the insulin-receptor gene with the restriction enzyme *Rsa* I in 242 Mexican Americans and non-Hispanic Whites with type II diabetes and 202 age-, sex-, and ethnicity-matched control subjects who participated in a population-based study in San Antonio. Alleles of 6.7 kilobases (kb) (A allele), 6.2 kb (B allele), and 3.4 kb (C allele) were identified. The C allele was observed in Mexican Americans only, where its frequency among nondiabetic control subjects was 17.7%. Diabetic Mexican Americans were twice as likely as control subjects to be homozygous for the C allele. The crude odds ratio for diabetes in CC homozygotes compared with the other two genotypes was 2.22, although this result was not statistically significant ($\chi^2 = 1.57$, $P = .21$). The Mantel-Haenszel odds ratio, adjusting for age, however, indicated a 4.71-fold increased risk of diabetes among Mexican Americans with the CC genotype compared with Mexican Americans without this genotype ($\chi^2_{M-H} = 5.38$, $P = .020$). The age of onset of diabetes was also slightly younger in CC homozygote cases (45.4 ± 9.2 yr) than in CX or XX cases (47.7 ± 9.0 and 48.6 ± 9.6 yr, respectively), although this difference was not statistically significant ($P = .467$). The C allele may be involved directly in the etiology of type II diabetes, or it may be in linkage disequilibrium with a functionally significant region of the insulin receptor gene. *Diabetes* 38:975–80, 1989

From the Section of Gastrointestinal Oncology and Digestive Diseases, Department of Medical Oncology, Division of Medicine, University of Texas MD Anderson Cancer Center, Houston; the Division of Clinical Epidemiology, Department of Medicine, University of Texas Health Science Center at San Antonio; and the University of Texas Health Science Center at Houston, Texas.

Address correspondence and reprint requests to Michael P. Stern, MD, Division of Clinical Epidemiology, Department of Medicine, University of Texas Health Science Center at San Antonio, 7703 Floyd Curl Drive, San Antonio, TX 78284-7873.

Received for publication 17 November 1988 and accepted in revised form 2 March 1989.

A strong genetic component to the etiology of type II (non-insulin-dependent) diabetes is suggested by the marked racial and ethnic differences in prevalence (1–4), the association between prevalence and genetic admixture in hybrid populations (5–8), and the high degree of concordance (60–90%) for this condition in monozygotic twins (9,10). Despite this evidence, the search for specific susceptibility genes for type II diabetes has thus far been disappointing. The recent rapid advances in identifying and characterizing restriction-fragment-length polymorphisms (RFLPs) of various genes have made it attractive to search for associations between type II diabetes and RFLPs in candidate genes, i.e., genes that code for proteins known to be involved in carbohydrate metabolism. Among such candidate genes are insulin, insulin-receptor, and glucose-transporter genes. Although initial reports seemed to suggest an association between type II diabetes and polymorphism of the insulin gene (11,12), later reports did not bear this out (13,14).

Resistance to insulin action is a well-established feature of type II diabetes and is believed by many to play an important role in the etiology of this condition (15,16). Moreover, insulin resistance has been shown to be familial in Pima Indians (17), which suggests that it may have a genetic basis. Bogardus et al. (18) have recently shown that insulin resistance assessed by the euglycemic insulin-clamp technique is trimodally distributed in Pima Indians, which is consistent with an autosomal, codominant mode of inheritance. We recently reported that the proportion of nondiabetic Mexican Americans who are hyperinsulinemic, an indirect indicator of insulin resistance, increases in a stepwise fashion according to whether they have zero, one, or two diabetic parents (19). Because pedigrees with progressively stronger family histories of diabetes may be expected to contain progressively greater numbers of prediabetic subjects, these data lend further support to the concept that prediabetic subjects are hyperinsulinemic (and, therefore, possibly in-

sulin resistant) and that this defect is at least familial if not inherited.

Although it has been claimed that insulin resistance in type II diabetes is due to a post-insulin-receptor defect, this concept is based primarily on studies of receptor number and binding (20,21). Recent data, however, have clearly established that the intracellular portion of the β -subunit of the insulin receptor exhibits tyrosine kinase activity, which appears to be involved in receptor function (22). Therefore, insulin-receptor function related to postbinding events could contribute to in vivo insulin resistance. Moreover, tyrosine kinase activity has been reported to be reduced in obese type II diabetic subjects compared to obese nondiabetic subjects in hepatic (23) and adipose (24) tissue.

In this article, we examine the association between type II diabetes and RFLPs of the insulin-receptor gene in Mexican Americans, a high-risk population for diabetes (2,5,25).

RESEARCH DESIGN AND METHODS

The study subjects are derived from the San Antonio Heart Study, a population-based study of cardiovascular disease and diabetes in Mexican Americans and non-Hispanic Whites. Detailed descriptions of the study design, sampling procedures, response rates, and field procedures have appeared elsewhere (2,5,6,25,26). The study was approved by the Institutional Review Board of the University of Texas Health Science Center at San Antonio, and informed consent was obtained from all subjects. A total of 3302 Mexican Americans and 1857 non-Hispanic Whites have participated in the San Antonio Heart Study. This report concerns a case-control study of diabetic subjects and matched control subjects nested within the larger parent study (26). The subjects with diabetes thus represent a population-based case series. Diabetes was diagnosed according to the National Diabetes Data Group criteria (27), as modified for epidemiologic studies (2,5). Only type II diabetic subjects are considered in this report (2,5). Age at diagnosis among previously diagnosed cases was ascertained by patient self-report. For subjects who were newly diagnosed at the time of their survey examination, age at diagnosis was taken to be their attained age at the time of the survey. A total of 376 Mexican Americans and 87 non-Hispanic Whites with type II diabetes were identified, of whom 300 (79.8%) Mexican Americans and 69 (79.3%) non-Hispanic Whites participated in a diabetic-complications examination (26). Gene-polymorphism studies were performed on a random subset of these individuals that consisted of 208 Mexican Americans and 34 non-Hispanic Whites. Controls were selected at random from among the nondiabetic members of the study population. The sampling of controls was stratified by sex, ethnicity, and decade of age, with stratum-specific sampling fractions chosen to match the age, sex, and ethnic distribution of the diabetic subjects. Gene-polymorphism studies were performed on a total of 202 control subjects, of whom 186 were Mexican American.

Whole blood specimens were obtained in heparinized containers and were kept frozen in sterile polypropylene tubes until analyzed. DNA was extracted as previously described (28) and digested with the restriction endonuclease *Rsa* I according to the manufacturer's directions (International Biotechnology, New Haven, CT). The DNA (10

$\mu\text{g}/\text{lane}$) was then separated by size on 0.8% agarose gel and blotted onto a nylon membrane by the method of Southern (29). Prehybridization and hybridization conditions have been described previously (30). RFLPs were detected by use of a 1.4 kb *Bgl* I fragment of the insulin receptor cDNA (kindly provided by G.I. Bell; 31). This fragment corresponds with bases 1601–2963 from the 5' end of the insulin-receptor cDNA described by Ullrich et al. (32). For hybridization, DNA was oligolabeled (33) to a specific activity of $1-3 \times 10^9$ cpm/ μg with [^{32}P]dCTP (New England Nuclear, Boston, MA). After washing under stringent conditions (30), the filters were subjected to autoradiography by use of Kodak XAR film with an intensifying screen at 80°C for 19–60 h. Assessment of insulin-receptor gene RFLPs was performed blind i.e., without knowledge of whether the specimen came from a diabetic or a control subject.

Body mass index (BMI), defined as kg/m^2 , was used as an indicator of overall adiposity, and the ratio of the subscapular to the triceps skinfold (centrality index) was used as an index of body-fat distribution. Anthropometric methods have been described previously (2,34,35).

Differences in age, anthropometric indexes, and age at diagnosis (cases only) according to genotype were evaluated by analysis of variance (36). The odds ratio for diabetes according to genotype, adjusted for age, was computed by the Mantel-Haenszel procedure and evaluated for statistical significance with the Mantel-Haenszel χ^2 statistic (37). The 95% confidence interval of the odds ratio was computed by the method of Robins et al. (38).

RESULTS

Hybridization of the 1.4-kb insulin-receptor-cDNA probe to *Rsa* I-digested human genomic DNA revealed three alleles of 6.7, 6.2, and 3.4 kb (Fig. 1). For purposes of this report,




FIG. 1. Hybridization of 1.4-kilobase (kb) probe to *Rsa* I-digested human DNA. Individuals are indicated by number at top of each lane, and sizes of fragments are indicated at right in kb. *Individuals with variant alleles of 7.5 kb (V_1 , sample 01) and 6.4 kb (V_2 , sample 33).

these alleles have been designated A, B, and C, respectively. Two variant alleles of 7.5 and 6.4 kb, designated as V₁ and V₂, respectively, were also observed (Fig. 1). The A and B alleles were observed in Mexican Americans and non-Hispanic Whites, whereas the C allele was observed exclusively in Mexican Americans, among whom its frequency in nondiabetic control subjects was 17.7%, calculated from Table 1 as

$$\frac{22 + 33 + 5(2) + 1}{2 \times 186} = 0.177$$

The two variant alleles were also observed in Mexican Americans only. The remaining analyses are confined to Mexican Americans.

The distribution of genotypes in Mexican-American diabetic and control subjects is presented in Table 1. The genotype frequencies conformed to Hardy-Weinberg equilibrium ($3\chi^2 = 1.32$, $P = .72$). Few differences were observed between diabetic and control subjects except for the CC genotype, which was more than twice as common in diabetic as in control subjects. For this reason, and because the C allele was observed exclusively in Mexican Americans, the genotypes were aggregated into three categories: CC homozygotes (2 C alleles); CX heterozygotes (1 C allele); and XX (no C alleles). The crude odds ratio for diabetes in CC homozygotes versus the other two genotypes was 2.22, although this result was not statistically significant ($\chi^2 = 1.57$, $P = .21$). This crude analysis, however, does not take into account possible differences between diabetic and control subjects in other diabetes-related risk factors.

Table 2 shows the mean age, BMI, and centrality index according to genotype for diabetic and control subjects. There were no significant differences in the anthropometric indexes according to genotype in either group. Mean age, however, was significantly younger in control subjects with the CC genotype than in control subjects with the other two genotypes. (Note that although the overall group of diabetic and control subjects were matched by age, there is no guarantee that this matching will persist within each genotype.) Because type II diabetes usually does not develop until the 5th or 6th decade of life, the CC control subjects are, by virtue of their younger age, at greater risk of becoming diabetic than the CX or XX control subjects. The effect of such

differential misclassification would be to underestimate the effect of the C allele on diabetes risk.

To take into account the greater risk of future diabetes among CC control subjects, we adjusted the odds ratio for age by the Mantel-Haenszel procedure. The results indicate that the odds of diabetes are 4.71 times greater in individuals with the CC genotype than in individuals with the other two genotypes ($P = .020$; Table 3). Note that three of the five CC control subjects were <30 yr of age. Diabetic subjects with the CC genotype also had a younger age of onset (45.4 ± 9.2 yr) than diabetic subjects with the CX or XX genotype (47.7 ± 9.0 and 48.6 ± 9.6 yr, respectively), although this difference was not statistically significant ($P = .467$).

DISCUSSION

With the restriction endonuclease *Rsa* I in a study of 51 unrelated non-Hispanic Whites, Shaw and Bell (31) described a two-allele polymorphism at the insulin-receptor locus with bands of 6.7 and 6.2 kb. The presence of a 3.4 kb RFLP after digestion with *Rsa* I, identical to our C allele, has been previously reported in Pima Indians but not in non-Hispanic Whites or Blacks (39). No association between the C allele and type II diabetes was found in this study, although only 54 Pima subjects were studied, which raises the possibility of a type II error.

Although we have reported an association between the C allele and type II diabetes, some caution is in order, because this association was not statistically significant in the crude analysis and was detected only after adjustment for age. The case for an association is strengthened, however, by the fact that the C allele has thus far been detected only in Pima Indians and Mexican Americans, two high-risk populations for diabetes that share Native American ancestry (25,40). The absence of the C allele in other populations is based on studies involving 58 Blacks (39) and 193 non-Hispanic Whites (50 non-Hispanic Whites in this study, 51 in the study by Shaw and Bell [31], and 92 in the study by Elbein et al. [39]).

The frequency of the C allele in Pima Indians was reported to be 34% (39). The 17.7% frequency of this allele in Mexican Americans is compatible with the 15–45% Native American admixture that we previously reported in this population (5,6). It is also compatible with the lower prevalence of type II diabetes in Mexican Americans compared with Pima Indians (~15 vs. 40%; 25,40). Although these ecological associations argue for an association between the C allele and diabetes, it is also possible that the presence of this allele in Mexican-American diabetic subjects simply reflects their greater degree of Native American genetic admixture relative to nondiabetic Mexican Americans (6). Nevertheless, that this allele occurs in a gene that codes for a protein of obvious relevance to carbohydrate metabolism (the insulin-receptor gene) suggests that it may have functional significance with respect to the etiology of diabetes or may be in linkage disequilibrium with a functionally significant region of the insulin-receptor gene.

If alterations of the insulin-receptor gene confer increased susceptibility to type II diabetes, this might arise as a result of mutations in the structural or regulatory portions of the gene. Further studies on the nucleotide sequence of this

TABLE 1
Distribution of observed genotypes in Mexican-American diabetic and control subjects

Genotype	Diabetic		Control	
	n	%	n	%
AA	21	10.1	21	11.3
AB	61	29.3	57	30.6
AC	30	14.4	22	11.8
BB	47	22.6	46	24.7
BC	33	15.9	33	17.7
CC	12	5.8	5	2.7
BV ₁	1	0.5	0	
BV ₂	0		1	0.5
AV ₂	1	0.5	0	
CV ₂	2	1.0	1	0.5

TABLE 2
Age and anthropometric variables according to genotype in diabetic and control subjects

	Diabetic			Control		
	CC	CX	XX	CC	CX	XX
n	12	65	131	5	56	125
Age (yr)	50.2 ± 9.7	52.1 ± 8.4	52.6 ± 8.6	36.0 ± 13.0	50.5 ± 10.6	51.4 ± 9.0
	P = .726			P = .560		
	P = .352			P = .001		
Body mass index (kg/m ²)	33.2 ± 5.9	31.3 ± 6.7	31.2 ± 6.1	29.7 ± 6.6	28.8 ± 5.5	29.5 ± 5.7
	P = .955			P = .446		
	P = .294			P = .948		
Centrality index	1.54 ± 0.76	1.35 ± 0.61	1.47 ± 0.54	1.22 ± 0.29	1.37 ± 0.47	1.32 ± 0.45
	P = .167			P = .432		
	P = .690			P = .655		

Values are means ± SD.

gene among individuals with C alleles may help clarify this issue. The C allele is apparently the result of an alteration in the nucleotide sequence giving rise to an additional *Rsa* I restriction-endonuclease-recognition site. Because the total nucleotide sequence of the insulin-receptor gene has not yet been established, we cannot determine if this *Rsa* I site falls in an exon or intron. If it falls in an exon, it could give rise to an altered insulin receptor as a result of a missense or nonsense mutation. If the *Rsa* I site of the C allele produces a missense mutation, it would fall in the region of cDNA bases 1928–2478, described by Ullrich et al. (32), because the coding sequences homologous to the C allele are located in this region (39). It is unlikely to be a nonsense mutation

because this would result in the absence of some or all of the β-subunit of the insulin receptor.

Another possibility is that the *Rsa* I restriction-endonuclease-cleavage site falls in an intron but is associated with a mutation in the structural portion of the gene, resulting in the production of an abnormal insulin receptor. An example of an RFLP that is in linkage disequilibrium with a mutation in the structural portion of the gene occurs in sickle cell anemia (41).

The mechanism(s) whereby alterations of the insulin-receptor gene might contribute to the development of type II diabetes is unknown. Recently, a patient with type A insulin resistance was found to be homozygous for a point mutation within the processing site of the insulin proreceptor, which resulted in a conversion from Arg-Lys-Arg-Arg to Arg-Lys-Arg-Ser (42). The resulting insulin receptors are not cleaved to mature α- and β-subunits and are therefore unable to bind insulin efficiently. In another case, a patient with insulin resistance secondary to leprechaunism was found to have inherited a different insulin-receptor mutant from each parent (43). One of the alleles contained a missense mutation, resulting in the replacement of Lys with Glu at amino acid 460 in the α-subunit. The other allele contained a nonsense mutation, resulting in premature termination of the protein after amino acid 671.

Two studies have reported associations between an *Sst* I RFLP of the insulin-receptor gene and type II diabetes (44,45). In one study, a negative association was reported between a 5.8-kb RFLP and type II diabetes in Blacks (44), whereas in the other study (45), a positive association was found in a predominantly White population. Although opposite in direction, both associations are compatible with linkage disequilibrium between an endonuclease-restriction site and a diabetes-susceptibility site on the insulin-receptor gene. Interestingly, the second study included six Hispanics, four of whom were diabetic. However, because *Rsa* I polymorphisms were not reported in this study, it is not known whether any of these subjects had the C allele. Also, in the second study, the four nondiabetic White subjects with the *Sst* I 5.8-kb RFLP (the nominal susceptibility genotype) were considered to be hyperinsulinemic, suggesting that they might be prediabetic. Two pedigree studies have attempted to document possible linkage between *Rsa* I and *Sst* I poly-

TABLE 3
Distribution of diabetic and control subjects by age and genotype

Age (yr)		Genotype			Total
		CC	CX	XX	
25–29	Diabetic	0	0	1	1
	Control	3	4	2	9
30–34*	Diabetic	2	3	5	10
	Control	0	3	5	8
35–39*	Diabetic	0	3	7	10
	Control	0	3	7	10
40–44	Diabetic	1	6	13	20
	Control	0	2	9	11
45–49	Diabetic	2	11	14	27
	Control	1	11	23	35
50–54	Diabetic	4	13	27	44
	Control	1	8	31	40
55–59*	Diabetic	0	13	28	41
	Control	0	14	19	33
60–64*	Diabetic	3	16	36	55
	Control	0	11	29	40
Total	Diabetic	12	65	131	208
	Control	5	56	125	186

Mantel-Haenszel odds ratio and 95% confidence interval for CC homozygotes vs. CX and XX combined = 4.71 (1.19–18.6); $\chi^2_{M-H} = 5.38, P = 0.20$.

*To avoid 0 marginals, the age groups 30–34 and 35–39 were combined, as were the age groups 55–59 and 60–64. Mantel-Haenszel analysis was also performed with 10-yr strata to avoid necessity of combining strata. Results remained statistically significant ($P = .05$), although the odds ratio decreased to 3.08.

morphisms of the insulin-receptor gene and type II diabetes (46,47). However, too few pedigrees were studied to confirm or exclude genetic linkage. Finally, an *Xba* I RFLP of a glucose-transporter gene was reported to be associated with type II diabetes in a study of Europeans and Japanese (48). Unlike our study, however, none of these studies were population based.

In conclusion, the C allele appears to be associated with type II diabetes in Mexican Americans. Moreover, this allele is observed only in Pima Indians and Mexican Americans, two high-risk populations for type II diabetes (25,40), and not in non-Hispanic Whites or Blacks. The explanation for the association with diabetes remains to be elucidated. Further studies on the nucleic-acid sequence of insulin-receptor genes among individuals with C alleles and family studies to determine if the C allele cosegregates with diabetes should help clarify the role of this gene in type II diabetes among Mexican Americans.

ACKNOWLEDGMENTS

This work was supported by Grant R01-HL-24799 from the National Heart, Lung, and Blood Institute.

REFERENCES

- West KM: *Epidemiology of Diabetes and Its Vascular Lesions*. New York, Elsevier, 1978, p. 199–201
- Stern MP, Rosenthal M, Haffner SM, Hazuda HP, Franco LJ: Sex difference in the effects of sociocultural status on diabetes and cardiovascular risk factors in Mexican Americans: the San Antonio Heart Study. *Am J Epidemiol* 120:834–51, 1984
- Zimmet P: Epidemiology of diabetes and its macrovascular complications in Pacific populations: the medical effects of social progress. *Diabetes Care* 2:144–53, 1979
- Jackson WPU: Epidemiology of diabetes in South Africa. *Adv Metab Disord* 9:111–46, 1978
- Gardner LI, Stern MP, Haffner SM, Gaskill SP, Hazuda HP, Relethford JH, Eifler CW: Prevalence of diabetes in Mexican Americans: relationship to percent of gene pool derived from Native American sources. *Diabetes* 33:86–92, 1984
- Chakraborty R, Ferrell RE, Stern MP, Haffner SM, Hazuda HP, Rosenthal M: Relationship of prevalence of non-insulin dependent diabetes mellitus to Amerindian admixture in the Mexican Americans of San Antonio, Texas. *Genet Epidemiol* 3:435–54, 1986
- Serjeantson SW, Owerbach D, Zimmet P, Nerup J, Thoma K: Genetics of diabetes in Nauru: effects of foreign admixture, HLA antigens and the insulin-gene-linked polymorphism. *Diabetologia* 25:13–17, 1983
- Brosseau JD, Eelkema RC, Crawford AC, Abe TA: Diabetes among the three affiliated tribes: correlation with degree of Indian inheritance. *Am J Public Health* 69:1277–78, 1979
- Newman B, Selby JV, King ML, Slemenda C, Fabsitz R, Friedman GD: Concordance for type 2 (non-insulin dependent) diabetes mellitus in male twins. *Diabetologia* 30:763–69, 1987
- Barnett AH, Eff C, Leslie RDG, Pyke DA: Diabetes in identical twins: a study of 200 pairs. *Diabetologia* 20:87–93, 1981
- Rotwein PS, Chirgwin J, Province M, Knowler WC, Pettitt DJ, Cordell B, Goodman HM, Permutt MA: Polymorphism in the 5'-flanking region of the human insulin gene: a genetic marker for non-insulin dependent diabetes. *N Engl J Med* 308:65–71, 1983
- Owerbach D, Nerup J: Restriction fragment length polymorphism of the insulin gene in diabetes mellitus. *Diabetes* 31:275–55, 1982
- Knowler WC, Pettitt DJ, Vasquez B, Rotwein PS, Andreone TL, Permutt MA: Polymorphism in the 5'-flanking region of the human insulin gene: relationships with non-insulin-dependent diabetes mellitus, glucose and insulin concentrations, and diabetes treatment in the Pima Indians. *J Clin Invest* 74:2129–35, 1984
- Elbein S, Rotwein P, Permutt MA, Bell GI, Sanz N, Karam JH: Lack of association of the polymorphic locus in the 5'-flanking region of the human insulin gene and diabetes in American Blacks. *Diabetes* 34:433–39, 1985
- Stern MP: Type II diabetes mellitus: Interface between clinical and epidemiological investigation. *Diabetes Care* 11:119–26, 1988
- DeFronzo RA, Ferrannini E, Koivisto V: New concepts in the pathogenesis and treatment of non-insulin-dependent diabetes mellitus. *Am J Med* 74 (Suppl.):52–81, 1983
- Lillioja S, Mott DM, Zawadzki JK, Young AA, Abbott WGH, Knowler WC, Bennett PH, Moll P, Bogardus C: In vivo insulin action is a familial characteristic in non-diabetic Pima Indians. *Diabetes* 36:1329–35, 1987
- Bogardus C, Lillioja S, Nyomba BL, Freymond D, Zurlo F, Swinburn B, Knowler W, Howard B, Bennett P: Evidence for a single gene, co-dominant mode of inheritance of insulin resistance in Pima Indians (Abstract). *Diabetes* 37 (Suppl. 1):91A, 1988
- Haffner SM, Stern MP, Hazuda HP, Mitchell BD, Patterson JK: Increased insulin concentrations in non-diabetic offspring of diabetic parents. *N Engl J Med* 319:1297–301, 1988
- Pollett RJ: Insulin receptors and action in clinical disorders of carbohydrate tolerance. *Am J Med* 75 (Suppl.):15–22, 1983
- Kolterman OG, Prince MJ, Olefsky JM: Insulin resistance in non-insulin-dependent diabetes mellitus: impact of sulfonylurea agents in vivo and in vitro. *Am J Med* 74 (Suppl.):82–101, 1983
- Rosen OM: After insulin binds. *Science* 237:1452–58, 1987
- Caro JF, Ittoop O, Pories WJ, Meelheim D, Flickinger EG, Thomas F, Jenquin M, Silverman JF, Khazanie PG, Sinha MK: Studies on the mechanism of insulin resistance in the liver from humans with noninsulin-dependent diabetes. *J Clin Invest* 78:249–58, 1986
- Freidenberg GR, Henry RR, Klein HH, Reichart DR, Olefsky JM: Decreased kinase activity of insulin receptors from adipocytes of non-insulin-dependent diabetic subjects. *J Clin Invest* 79:240–50, 1987
- Diehl AK, Stern MP: Special health problems of Mexican Americans: obesity, gallbladder disease, diabetes mellitus, and cardiovascular disease. *Adv Intern Med* 34:79–96, 1989
- Haffner SM, Fong D, Stern MP, Pugh JA, Hazuda HP, Patterson JK, Van Heuven WAJ, Klein R: Diabetic retinopathy in Mexican Americans and non-Hispanic whites. *Diabetes* 37:878–84, 1988
- National Diabetes Data Group: Classification and diagnosis of diabetes mellitus and other categories of glucose intolerance. *Diabetes* 28:1039–57, 1979
- Frazier ML, Ferrell RE, Arem R, Chamakhi S, Field JB: Restriction fragment length polymorphism of the human insulin gene region among type II diabetic Mexican Americans and Tunisians. *Res Commun Chem Pathol Pharmacol* 52:371–85, 1986
- Southern EM: Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J Mol Biol* 98:503–17, 1975
- Amasino RM: Acceleration of nucleic acid hybridization rate by polyethylene glycol. *Anal Biochem* 152:304–30, 1986
- Shaw DJ, Bell GI: *Rsa* I polymorphism at the insulin receptor locus (INSR) on chromosome 19. *Nucleic Acids Res* 13:8661, 1985
- Ullrich A, Bell GI, Chem EY, Herrera R, Petruzzelli LM, Dull TJ, Gray A, Couseus L, Liao YC, Tsubokawa M, Mason A, Seeburg PH, Grunfeld C, Rosen OM, Ramachandran J: Human insulin receptor and its relationship to the tyrosine kinase family of oncogenes. *Nature (Lond)* 313:756–61, 1985
- Feinberg AP, Vogelstein B: A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal Biochem* 132:6–13, 1983
- Haffner SM, Stern MP, Hazuda HP, Rosenthal M, Knapp JA: The role of behavioral variables and fat patterning in explaining ethnic differences in serum lipids and lipoproteins. *Am J Epidemiol* 123:830–39, 1986
- Haffner SM, Stern MP, Hazuda HP, Rosenthal M, Knapp JA, Malina RM: Role of obesity and fat distribution in non-insulin-dependent diabetes mellitus in Mexican Americans and non-Hispanic Whites. *Diabetes Care* 9:153–61, 1986
- Wilkinson L: *SYSTAT: The System for Statistics*. Evanston, IL, SYSTAT, 1988, p. 484–91
- Schlesselman JJ: *Case-Control Studies: Design, Conduct, Analysis*. New York, Oxford Univ. Press, 1982, p. 183–85
- Robins J, Greenland S, Breslow NE: A general estimator for the variance of the Mantel-Haenszel odds ratio. *Am J Epidemiol* 124:719–23, 1986
- Elbein SC, Corsetti L, Ullrich A, Permutt MA: Multiple restriction fragment length polymorphisms at the insulin receptor locus: a highly informative marker for linkage analysis. *Proc Natl Acad Sci USA* 83:5223–27, 1986
- Knowler WC, Pettitt DJ, Savage PJ, Bennett PH: Diabetes incidence in Pima Indians: contributions of obesity and parental diabetes. *Am J Epidemiol* 113:144–56, 1981
- Kan YW, Dozy AM: Polymorphism of DNA sequence adjacent to human beta-globin structural gene: relationship to sickle mutation. *Proc Natl Acad Sci USA* 75:5631–35, 1978
- Yoshimasa Y, Seino S, Whittaker J, Kakehi T, Kosaki A, Kuzuya H, Imura H, Bell GI, Steiner DF: Insulin-resistant diabetes due to a point mutation that prevents insulin proreceptor processing. *Science* 240:784–87, 1988
- Kadowaki T, Bevins CL, Cama A, Ojamaa K, Marcus-Samuels B, Kadowaki H, Beitz L, McKeon C, Taylor SI: Two mutant alleles of the insulin receptor gene in a patient with extreme insulin resistance. *Science* 240:787–90, 1988
- McGill J, Corsetti L, Elbein SC: Restriction fragment length polymorphisms (RFLP) at the insulin receptor (HIR) in a population of diabetic and non-diabetic Blacks (Abstract). *Diabetes* 36 (Suppl. 1):21A, 1987

45. McClain DA, Henry RR, Ullrich A, Olefsky JM: Restriction-fragment-length polymorphism in insulin-receptor gene and insulin resistance in NIDDM. *Diabetes* 37:1071-75, 1988
46. O'Rahilly S, Trembath RC, Patel P, Galton DJ, Turner RC, Wainscoat JS: Linkage analysis of the human insulin receptor gene in type 2 (non-insulin-dependent) diabetic families and a family with maturity onset diabetes of the young. *Diabetologia* 31:792-97, 1988
47. Elbein SC, Borecki I, Corsetti L, Fajans SS, Hansen AT, Nerup J, Province M, Permutt MA: Linkage analysis of the human insulin receptor gene and maturity-onset diabetes of the young. *Diabetologia* 30:641-47, 1987
48. Li SR, Baroni MG, Oelbaum RS, Stock J, Galton DJ: Association of genetic variant of the glucose transporter with non-insulin-dependent diabetes mellitus. *Lancet* 2:368-70, 1988