

The Changing Face of Young-Onset Diabetes: Type 1 Optimism Mellowed by Type 2 Concerns

Until recently the outlook for a youth or young adult diagnosed with diabetes, which was almost universally type 1, was bleak. Indeed, using data from the National Health Interview Survey as recent as from 1984 to 2000, it was estimated that U.S. children diagnosed with diabetes at 10 years of age had a life expectancy approximately 19 years less than seen in the general population (1). However, more recent data from the Pittsburgh Epidemiology of Diabetes Complications (EDC) study suggest those diagnosed with childhood-onset diabetes between 1965 and 1980 have a life expectancy of almost 69 years, which is less than 4 years lower than the comparable U.S. population (2). This good news has been accompanied by the observation from the Finnish Diabetic Nephropathy (FinnDiane) study that virtually all of the excess mortality seen in type 1 diabetes is related to the development of micro- or macroalbuminuria (3). This seminal observation has been confirmed and extended for up to a 20-year period in the EDC population (4).

The improved prognosis, in terms of mortality, has been accompanied by a dramatic reduction (5) or delay (6) in the incidence of end-stage renal disease. Interestingly, the decline in cardiovascular disease (CVD), the leading cause of overall mortality in diabetes, is less marked (5). One cautionary note, however, has to be made concerning the improvement in mortality of patients with type 1 diabetes. In a recent analysis of over 17,000 individuals in Finland, diagnosed between 1970 and 1999, Harjutsalo et al. (7) compared the time trends of mortality for those diagnosed at an age less than 15 years to those diagnosed at an age of 15 through 29 years. Although a very significant fall was seen in mortality over time for the young-onset group, consistent with the Pittsburgh EDC population (who were all diagnosed before the age of 17), mortality for the older-onset group increased over time reflecting an increasing number deaths related to alcohol, drugs, and acute complications (7). This raises the possibility that

type 1 diabetes mortality patterns may differ markedly by age of onset.

The picture becomes more confusing, and disturbing, when one considers the recent increased incidence of apparent type 2 diabetes occurring in youth and young adults (8). One major challenge is that of typology, or our ability to distinguish between type 1 and type 2 diabetes, which is particularly difficult in an overweight or obese young adult. The SEARCH for Diabetes in Youth (SEARCH) study has examined this issue in some depth and described four groups based on the presence or absence of diabetes autoantibodies and of insulin resistance (9). How well this schema would work in the future in terms of predicting outcome remains to be seen but it is likely to be quite relevant as a number of studies have suggested that even in clear type 1 individuals it is those with evidence of insulin resistance or an insulin resistance/type 2 diabetes family background that have increased cardiovascular and renal disease (10–15). The complexity of this issue is further demonstrated by the observation that many classic type 1 diabetic subjects may retain some residual β -cell function for many years after diagnosis (16), which may partly relate to the benign natural history seen in many of the patients from the Joslin 50-Year Medalist Study who have survived 50 years of type 1 diabetes (17).

So what do we know about the prognosis of type 2 diabetes in youth and young adults? A number of studies have suggested that individuals with type 2 diabetes have worse cardiovascular risk factors than similarly aged individuals with type 1 diabetes. Indeed, the SEARCH study has shown more adverse cardiovascular risk profiles, including blood pressure (18) and lipid levels (19), and a higher prevalence of microalbuminuria (20) in youth-onset type 2 diabetes compared with type 1. Up to now, however, there have been few data on mortality or major outcomes of diabetes comparing type 1 and type 2 diabetes where onset occurred in youth or young

adulthood. Hillier and Pedula (21) some years ago suggested that type 2 diabetes with an onset between age 18 and 44 years ran a more aggressive course than cases diagnosed later, particularly in terms of relative impact compared with the age-matched general population. The results of the Treatment Options for Type 2 Diabetes in Adolescents and Youth (TODAY) study support such a conclusion in terms of metabolic deterioration and have been recently reviewed (22).

In this issue, Constantino et al. (23) now provide further data concerning young adult-onset type 2 diabetes. Using a diabetes clinical database, and matching to the Australian National Death Index, these investigators were able to compare clinical and mortality outcomes from 354 patients with type 2 diabetes and 470 with type 1 diabetes.

Strikingly there was a twofold greater mortality in the type 2 cohort predominantly due to an excess of cardiovascular deaths. Although the clinical data were largely collected through routine encounters, a standardized protocol was used and the data quality is thus likely to be generally high. Likewise, the linkage with the Australian National Death Index is validated and mortality ascertainment data are likely to be complete. A significant weakness of the study, however, is the reliance purely on death certificates alone for cause of death, which were only available for 72% of deaths at the time of analysis. A number of studies have demonstrated the pathways and contributors to death are quite complex in diabetes (24) and the study would be greatly enhanced by the investigation and standardized recording of causes of death. Nevertheless, these data are unique and extremely valuable and support the growing concern that type 2 diabetes with a youth/young adult-onset has a particularly high risk of adverse vascular outcomes. Some of the figures from Constantino et al. (23) are quite concerning with prevalence rates of ischemic heart disease reaching as high as 13% at an age of 40 years compared with only 3% in

- type 1 diabetes: nationwide population based cohort study. *BMJ* 2011;343:d5364
8. Liese AD, D'Agostino RB Jr Hamman RF, et al.; SEARCH for Diabetes in Youth Study Group. The burden of diabetes mellitus among US youth: prevalence estimates from the SEARCH for Diabetes in Youth Study. *Pediatrics* 2006;118:1510–1518
 9. Dabelea D, Pihoker C, Talton JW, et al.; SEARCH for Diabetes in Youth Study. Etiological approach to characterization of diabetes type: the SEARCH for Diabetes in Youth Study. *Diabetes Care* 2011;34:1628–1633
 10. Purnell JQ, Dev RK, Steffes MW, et al. Relationship of family history of type 2 diabetes, hypoglycemia, and autoantibodies to weight gain and lipids with intensive and conventional therapy in the Diabetes Control and Complications Trial. *Diabetes* 2003;52:2623–2629
 11. Erbey JR, Kuller LH, Becker DJ, Orchard TJ. The association between a family history of type 2 diabetes and coronary artery disease in a type 1 diabetes population. *Diabetes Care* 1998;21:610–614
 12. Orchard TJ, Olson JC, Erbey JR, et al. Insulin resistance-related factors, but not glycemia, predict coronary artery disease in type 1 diabetes: 10-year follow-up data from the Pittsburgh Epidemiology of Diabetes Complications Study. *Diabetes Care* 2003;26:1374–1379
 13. Soedamah-Muthu SS, Chaturvedi N, Toeller M, et al.; EURODIAB Prospective Complications Study Group. Risk factors for coronary heart disease in type 1 diabetic patients in Europe: the EURODIAB Prospective Complications Study. *Diabetes Care* 2004;27:530–537
 14. Dabelea D, Kinney G, Snell-Bergeon JK, et al.; Coronary Artery Calcification in Type 1 Diabetes Study. Effect of type 1 diabetes on the gender difference in coronary artery calcification: a role for insulin resistance? The Coronary Artery Calcification in Type 1 Diabetes (CACTI) Study. *Diabetes* 2003;52:2833–2839
 15. Orchard TJ, Chang YF, Ferrell RE, Petro N, Ellis DE. Nephropathy in type 1 diabetes: a manifestation of insulin resistance and multiple genetic susceptibilities? Further evidence from the Pittsburgh Epidemiology of Diabetes Complication Study. *Kidney Int* 2002;62:963–970
 16. Palmer JP. C-peptide in the natural history of type 1 diabetes. *Diabetes Metab Res Rev* 2009;25:325–328
 17. Keenan HA, Sun JK, Levine J, et al. Residual insulin production and pancreatic β -cell turnover after 50 years of diabetes: Joslin Medalist Study. *Diabetes* 2010;59:2846–2853
 18. Rodriguez BL, Dabelea D, Liese AD et al., SEARCH Study Group. Prevalence and correlates of elevated blood pressure in youth with diabetes mellitus: the SEARCH for Diabetes in Youth Study. *J Pediatr* 2010;157:245–251.e1.
 19. Kershner AK, Daniels SR, Imperatore G, et al. Lipid abnormalities are prevalent in youth with type 1 and type 2 diabetes: the SEARCH for Diabetes in Youth Study. *J Pediatr* 2006;149:314–319
 20. Maahs DM, Snively BM, Bell RA, et al. Higher prevalence of elevated albumin excretion in youth with type 2 than type 1 diabetes: the SEARCH for Diabetes in Youth study. *Diabetes Care* 2007;30:2593–2598
 21. Hillier TA, Pedula KL. Complications in young adults with early-onset type 2 diabetes: losing the relative protection of youth. *Diabetes Care* 2003;26:2999–3005
 22. Linder BL, Fradkin JE, Rodgers GP. The TODAY Study: an NIH perspective on its implications for research. *Diabetes Care* 2013;36:1775–1776
 23. Constantino MI, Molyneaux L, Limacher-Gisler F, et al. Long-term complications and mortality in young-onset diabetes: type 2 diabetes is more hazardous and lethal than type 1 diabetes. *Diabetes Care* 2013;36:3863–3869
 24. Diabetes Epidemiology Research International Mortality Study Group. Major cross-country differences in risk of dying for people with IDDM. *Diabetes Care* 1991;14:49–54
 25. Collins R, Armitage J, Parish S, Sleight P, Peto R; Heart Protection Study Collaborative Group. MRC/BHF Heart Protection Study of cholesterol-lowering with simvastatin in 5963 people with diabetes: a randomised placebo-controlled trial. *Lancet* 2003;361:2005–2016
 26. Nathan DM, Zinman B, Cleary PA, et al.; Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications (DCCT/EDIC) Research Group. Modern-day clinical course of type 1 diabetes mellitus after 30 years' duration: the Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications and Pittsburgh Epidemiology of Diabetes Complications experience (1983–2005). *Arch Intern Med* 2009;169:1307–1316
 27. Asleh R, Levy AP. In vivo and in vitro studies establishing haptoglobin as a major susceptibility gene for diabetic vascular disease. *Vasc Health Risk Manag* 2005;1:19–28
 28. Costacou T, Ferrell RE, Orchard TJ. Haptoglobin genotype: a determinant of cardiovascular complication risk in type 1 diabetes. *Diabetes* 2008;57:1702–1706
 29. Costacou T, Ferrell RE, Ellis D, Orchard TJ. Haptoglobin genotype and renal function decline in type 1 diabetes. *Diabetes* 2009;58:2904–2909
 30. Blum S, Vardi M, Brown JB, et al. Vitamin E reduces cardiovascular disease in individuals with diabetes mellitus and the haptoglobin 2-2 genotype. *Pharmacogenomics* 2010;11:675–684