

# Diabetic Autonomic Neuropathy in BB Rats and Effect of ARI Treatment on Heart-Rate Variability and Vagus Nerve Structure

WEI-XIAN ZHANG, SUBRATA CHAKRABARTI, DOUGLAS A. GREENE, AND ANDERS A.F. SIMA

**The preventive effect of the aldose reductase inhibitor (ARI) ponalrestat on heart-rate variability and the development of autonomic neuropathy in the vagus nerve was investigated in the spontaneously diabetic BB rat. ARI treatment completely prevented the characteristic decrease in heart-rate variability and axonal atrophy of the vagus nerve for 4 mo in hyperglycemic BB rats. After 6 mo of treatment, the preventive effect on heart-rate variability was partial, and the vagus nerve demonstrated an increase in regenerating myelinated and unmyelinated fibers. These data suggest that autonomic neuropathy involving the vagus nerve is metabolically induced by demonstrating that inhibition of the polyol pathway significantly delays the occurrence of functional and structural autonomic neuropathy despite the presence of hyperglycemia. *Diabetes* 39:613–18, 1990**

**A**utonomic nerve dysfunction is a serious and common complication in diabetes mellitus and is responsible for an increased morbidity and mortality in diabetic subjects (1–3). It leads to vagal denervation and sensorimotor and reflex dysfunction of the cardiovascular, urogenital, and gastrointestinal systems (4–8).

We have previously reported on autonomic nerve dysfunction and neuroanatomic abnormalities in various autonomic nerves in spontaneously diabetic BB rats (9–13). From these studies, it appears that functional abnormalities such as decreased heart-rate variability and impaired micturition-reflex function precede ultrastructurally detectable neuroan-

atomic lesions in the respective autonomic nerves (11,12). This sequence of events suggests an early potentially reversible metabolic defect in autonomic nerves that may initiate and sustain the subsequent development of structural changes similar to the sequence of events occurring in diabetic somatic peripheral nerve (14,15). In the diabetic mixed sensory and motor sciatic nerve, increased ambient glucose concentrations reduce nerve *myo*-inositol by activation of the polyol pathway, which in turn impairs nerve  $\text{Na}^+\text{-K}^+\text{-ATPase}$  via decreased activity of protein kinase C (16,17). The  $\text{Na}^+\text{-K}^+\text{-ATPase}$  defect has been associated with the early slowing of nerve conduction velocity and the occurrence of structural lesions in somatic nerves (14,15,18,19).

To examine whether activated polyol-pathway and subsequent metabolic abnormalities may underlie autonomic nerve dysfunction and neuropathy, diabetic BB rats were treated with the aldose reductase inhibitor (ARI) ponalrestat from the onset of diabetes. They were examined with respect to heart-rate variability and structural lesions of the vagus nerve after 4 and 6 mo of treatment.

## RESEARCH DESIGN AND METHODS

Twenty prediabetic male BB rats and 10 age- and sex-matched non-diabetes-prone BB rats were obtained from the Department of Pathology, University of Massachusetts (Worcester). All animals were maintained in individual air-filtered metabolic cages with ad libitum access to water and rat chow (Wayne Lab Blox F-6, Wayne, Chicago, IL). Body weight, urine volume, ketonuria, and glucosuria (Keto-Diastix, Miles Canada, Etobicoke) were monitored daily, and blood glucose was measured weekly from tail vein blood samples (Glucometer, Ames, Elkhart, IN). Glycosylated hemoglobin was measured from tail vein blood samples every 2nd mo and expressed as percentage of glycosylated hemoglobin (Glyco-Test, Pierce, Rockford, IL). At detection of glucosuria, diabetic rats were started on small daily doses (0.5–3.0 U/day) of protamine zinc insulin (Connaught-Novo, Toronto) designed to maintain an insulin-deficient state with blood glucose levels ranging between 15 and 20 mM. Three

From the Neuropathology Research Laboratory, Department of Pathology, University of Manitoba, Winnipeg, Manitoba, Canada; and the Diabetes Research and Training Center, Medical Center, University of Michigan, Ann Arbor, Michigan.

Address correspondence and reprint requests to Dr. Anders A.F. Sima, Neuropathology Research Laboratory, Department of Pathology, University of Manitoba, 770 Bannatyne Avenue, Winnipeg, Manitoba R3E 0W3, Canada.

Received for publication 28 August 1989 and accepted in revised form 4 January 1990.

weeks after onset of diabetes, animals were divided into two groups: 1) insulin-deficient untreated diabetic and 2) insulin-deficient diabetic fed ponalrestat-supplemented (ICI, Wilmington, DE) rat chow (80 g ponalrestat/100 kg rat chow), equivalent to a daily dose of 25 mg ponalrestat/kg body wt (actual dose for diabetic rats 32.6 mg/kg body wt). Age- and sex-matched non-diabetes-prone BB rats served as controls. Half of the animals in each experimental group were killed 4 mo after onset of diabetes (5 animals/group) and half at 6 mo postdetection. End-point measurements were performed by investigators unaware of animal identity.

Heart-rate variability was assessed by computed statistic  $\bar{R}$  as previously described in detail (11,20). Simultaneous analog movement, ECG, and respiration signals were obtained from a movement monitor and subcutaneous electrodes. The respiration data were amplified and fed to a Wheatstone's bridge apparatus (Barclay Whyte, Winnipeg, Canada), which discriminates differences in transthoracic impedance due to respiration. The ECG signals were fed to an ECG recorder (Cardiofax ECG-6201, Nikon Kohden, Tokyo) for amplification. Movement, ECG, and respiration signals output was converted by an analog-to-digital converter, and input was sent to a microprocessor for circular time wrapping of R-R intervals per respiration cycle. The  $\bar{R}$  is expressed as the length of the mean vector of  $\bar{R}$  points on a unit circle equal to the length of one respiration cycle arbitrarily set to 1 (11,20).  $\bar{R}$  values were collected at 2, 4, and 6 mo of diabetes.

Animals were perfused with 2.5% glutaraldehyde fixative buffered with 0.1 M sodium cacodylate (pH 7.3) under pentobarbital sodium anesthesia (50 mg/kg body wt). The left unifascicular vagus nerve was dissected and immersed for 4 h at 4°C in the same fixative. The specimens were postfixed in 1% osmium tetroxide in 0.1 M sodium cacodylate (pH 7.3) for 2 h at 4°C. After dehydration through ascending concentrations of ethanol, the tissue was embedded in Epon.

Morphometric abnormalities were assessed by fiber size frequency distributions, density, and occupancy and mean fiber size as previously described in detail (10). These data were obtained from semithin cross sections of the vagus nerve embedded in plastic. From each vagus nerve, all myelinated fibers were digitized with the aid of an HP 9872A digitizer interfaced to an HP 9825A desk computer and plotter (Hewlett-Packard, Fort Collins, CO). Myelinated-fiber oc-

cupancy was expressed as percentage of endoneurial area occupied by myelinated fibers.

To assess axonal atrophy, linear regression analyses of the natural logarithm of axonal area and the number of myelin lamellae were examined as previously described (18).

Size frequency distributions of unmyelinated fibers, fiber density, and fiber occupancy were measured from systematic randomly chosen areas of the vagus nerve. These measurements were obtained from electron micrographs with a total magnification of  $\times 24,000$ , and the calculations were performed in the same way as for myelinated fibers (10).

Data are expressed as means  $\pm$  SE. Significance of differences between groups was calculated by analysis of variance (ANOVA) and modified *t* test. Linear regression analysis was performed by the least-squares method.

## RESULTS

Prediabetic rats developed diabetes at a mean age of  $97 \pm 5$  days. Ponalrestat treatment had no preventive effect on the characteristic decrease in body weight gain seen in untreated diabetic BB rats or on blood glucose or glycosylated hemoglobin levels (Table 1).

After 2 mo of insulin-deficient diabetes, the  $\bar{R}$  values were significantly ( $P < 0.02$ ) decreased compared with nondiabetic control values (Table 2). This difference became even more pronounced ( $P < 0.001$ ) after 4 and 6 mo of untreated diabetes. In insulin-deficient diabetic rats, ARI treatment achieved a complete prevention of the defect in heart-rate variability for 4 mo, whereas after 6 mo, there was only a partial preventive effect that amounted to 40% of the deficit (Table 2).

Myelinated fiber size distribution in untreated 4-mo diabetic rats showed a moderate shift toward smaller fibers compared with age-matched nondiabetic controls (Fig. 1A). This shift in the frequency distribution of myelinated fibers was prevented by 4 mo of ponalrestat treatment, so that no significant difference could be demonstrated compared with the fiber size distribution in nondiabetic controls (Fig. 1B). No differences were demonstrated in myelinated fiber density or occupancy between any of the groups at 4 mo.

Six months of ponalrestat treatment failed to prevent the characteristic myelinated fiber atrophy in diabetic rats. Instead, myelinated fiber size was significantly less ( $P < 0.05$ ) in ponalrestat-treated diabetic than untreated diabetic rats

TABLE 1  
Effect of ponalrestat treatment on body weight and hyperglycemia at 4 and 6 mo of diabetes

| Group                        | Body weight (g) | Blood glucose (mM) | Glycosylated hemoglobin (%) |
|------------------------------|-----------------|--------------------|-----------------------------|
| 4 mo                         |                 |                    |                             |
| Control                      | 460 $\pm$ 12    | 5.7 $\pm$ 0.5      | 3.9 $\pm$ 0.4               |
| Diabetic                     | 353 $\pm$ 10    | 17.8 $\pm$ 3.0     | 8.5 $\pm$ 0.4               |
| Ponalrestat-treated diabetic | 363 $\pm$ 17    | 17.3 $\pm$ 1.2     | 8.2 $\pm$ 0.6               |
| 6 mo                         |                 |                    |                             |
| Control                      | 500 $\pm$ 21    | 3.4 $\pm$ 0.6      | 4.4 $\pm$ 0.5               |
| Diabetic                     | 383 $\pm$ 12    | 20.8 $\pm$ 2.1     | 8.4 $\pm$ 0.9               |
| Ponalrestat-treated diabetic | 363 $\pm$ 17    | 24.3 $\pm$ 0.7     | 7.7 $\pm$ 0.5               |

$n = 5$  for each group.  $P < 0.005$  for body weight, blood glucose, and glycosylated hemoglobin by analysis of variance. All other comparisons not significant.

TABLE 2  
 $\bar{R}$  values ( $\times 10^3$ ) in control, untreated diabetic, and ponalrestat-treated diabetic BB rats

| Group                        | Diabetes duration |                |                |
|------------------------------|-------------------|----------------|----------------|
|                              | 2 mo*             | 4 mo†          | 6 mo†          |
| Control                      | 27.6 $\pm$ 2.0    | 24.5 $\pm$ 2.1 | 21.1 $\pm$ 1.3 |
| Diabetic                     | 20.0 $\pm$ 0.9    | 11.8 $\pm$ 0.5 | 11.1 $\pm$ 0.4 |
| Ponalrestat-treated diabetic | 27.1 $\pm$ 2.3    | 26.7 $\pm$ 1.5 | 15.1 $\pm$ 1.0 |

$P < 0.02$  (Control vs Diabetic, 2 mo);  $P < 0.001$  (Control vs Diabetic, 4 mo);  $P < 0.001$  (Control vs Diabetic, 6 mo);  $P < 0.02$  (Diabetic vs Ponalrestat-treated, 2 mo);  $P < 0.001$  (Diabetic vs Ponalrestat-treated, 4 mo);  $P < 0.02$  (Diabetic vs Ponalrestat-treated, 6 mo);  $P < 0.002$  (Control vs Ponalrestat-treated, 6 mo)

For definition of  $\bar{R}$ , see RESEARCH DESIGN AND METHODS and refs. 11 and 20 for review.  $n = 5$  for each group.

\* $P < 0.05$  by analysis of variance (ANOVA).

† $P < 0.005$  by ANOVA. All other comparisons not significant.

(Table 3; Fig. 2). In contrast, ponalrestat treatment appeared to have a small nonsignificant protective effect on fiber number as reflected by fiber density but did not prevent the diminished ( $P < 0.02$ ) fiber occupancy demonstrated in untreated diabetic BB rats (Table 3). Microscopic examination of semithin cross sections of the vagus nerve embedded in plastic revealed no interstitial inflammation, endoneurial

granulomata, or exacerbation of axonal degeneration in ponalrestat-treated rats.

Four months of ARI treatment completely prevented the characteristic axonal atrophy of myelinated fibers reflected by an unchanged axon-myelin ratio compared with nondiabetic controls (Fig. 3, *top*). However, this protective effect was not sustained at 6 mo, when the axon-myelin ratio in ponalrestat-treated diabetic rats did not differ significantly from that in untreated insulin-deficient rats (Fig. 3, *bottom*).

Diabetes untreated for 4 mo resulted in a marked shift in unmyelinated fiber size distribution toward smaller fiber sizes (Fig. 4A). This shift, reflecting axonal atrophy, was markedly but not completely prevented by ponalrestat treatment (Fig. 4B). Unmyelinated fiber density was not affected by 4 mo of diabetes.

After 6 mo of treatment, no effect could be demonstrated on mean unmyelinated fiber size, whereas fiber density was markedly increased compared with both untreated diabetic rats ( $P < 0.002$ ) and nondiabetic controls ( $P < 0.001$ ) (Table 3). Furthermore, ponalrestat treatment had a protective effect ( $P < 0.02$ ) on unmyelinated fiber occupancy compared with untreated diabetic rats (Table 3).




FIG. 1. **A:** myelinated fiber size frequency distributions in 4-mo untreated diabetic (solid bars;  $n = 5$ ) and control (open bars;  $n = 5$ ) rats. Diabetic rats show fiber atrophy with shift to smaller size frequencies. **B:** 4 mo of ponalrestat treatment (hatched bars;  $n = 5$ ) prevented this shift of myelinated fiber size frequency. Open bars, control ( $n = 5$ ). \*\* $P < 0.01$ .

## DISCUSSION

Diabetic autonomic neuropathy affecting the vagus nerve in the spontaneously diabetic BB rat is characterized by progressive axonal atrophy of myelinated and unmyelinated fibers (10), which is preceded and accompanied by vagal dysfunction substantiated by impaired heart-rate variability (11). Our findings in untreated diabetic BB rats confirm the previous findings and suggest a metabolic pathogenesis for vagal dysfunction and neuropathy. Complete prevention by the ARI ponalrestat of decrease in heart-rate variability for 4 mo in hyperglycemic animals supports the notion that hyperglycemia-induced activation of the polyol pathway plays an important inciting role in the development of autonomic neuropathy. However, this beneficial effect was only partial at 6 mo of diabetes, suggesting that inhibition of the polyol pathway in the presence of severe continuous hyperglycemia cannot be sustained by ponalrestat. These findings appear to concur with a beneficial effect of ARI treatment on autonomic nerve dysfunction in diabetic patients (21). On the other hand, it is conceivable that nonenzymatic glycosylation of long-lived axonal proteins (22) unaffected by aldose reductase inhibition may account for incomplete prevention of axonal atrophy in the persistently hyperglycemic

TABLE 3  
Effect of ponalrestat treatment on myelinated and unmyelinated fiber morphometry at 6 mo

| Group                        | Myelinated fiber morphometry         |   |                      | Unmyelinated fiber morphometry       |  |                      |
|------------------------------|--------------------------------------|---|----------------------|--------------------------------------|--|----------------------|
|                              | Mean fiber size ( $\mu\text{m}^2$ )* | Fiber density ( $\text{m}/\text{mm}^2$ )† | Fiber occupancy (%)‡ | Mean fiber size ( $\mu\text{m}^2$ )* | Fiber density ( $\times 1000/\text{mm}^2$ )† | Fiber occupancy (%)‡ |
| Control                      | $7.03 \pm 0.15$                      | $10,525 \pm 979$                          | $7.40 \pm 0.68$      | $0.96 \pm 0.05$                      | $339 \pm 15$                                 | $32.37 \pm 0.91$     |
| Diabetic                     | $5.92 \pm 0.48$                      | $7328 \pm 759$                            | $4.82 \pm 0.57$      | $0.82 \pm 0.04$                      | $358 \pm 10$                                 | $29.37 \pm 0.79$     |
| Ponalrestat-treated diabetic | $4.65 \pm 0.26$                      | $9429 \pm 523$                            | $4.67 \pm 0.43$      | $0.80 \pm 0.03$                      | $445 \pm 20$                                 | $35.91 \pm 2.46$     |

\* $P < 0.05$  by analysis of variance (ANOVA).  
† $P < 0.01$  by ANOVA.  
‡ $P < 0.005$  by ANOVA. All other comparisons not significant.




FIG. 2. Myelinated fiber size distribution in 6-mo ponalrestat-treated diabetic (hatched bars;  $n = 5$ ) versus untreated diabetic (solid bars;  $n = 5$ ) rats. Paradoxical shift to smaller myelinated fiber sizes in treated rats probably represents addition of small regenerated fibers. \* $P < 0.05$ .

ARI-treated rats at 6 mo. Alternatively, some investigators have suggested that axonal atrophy in murine diabetes may be consequent to maturational and/or nutritional factors in the generally leaner diabetic animals (23). However, this is an unlikely explanation for the residual axonal atrophy that occurred at 6 mo of ARI treatment, because at 4 mo, axonal atrophy was completely prevented in ARI-treated rats, being as stunted as in untreated diabetic rats in whom significant axonal atrophy was present.

In the vagus nerve, the myelinated fiber population represents visceral afferents, and most of the unmyelinated fibers are presynaptic parasympathetic fibers (24). The preventive effect of ponalrestat treatment on axonal atrophy of myelinated fibers, as demonstrated by the axon-myelin ratio after 4 mo of treatment, is analogous to the effect of ARI treatment on somatic afferents in diabetic BB rats (25). Because axonal atrophy in diabetic neuropathy is believed to be a consequence of slowed axonal transport secondary to the  $\text{Na}^+/\text{K}^+$ -ATPase defect (26,27), it is expected that correction of the metabolic defect responsible for the impaired  $\text{Na}^+/\text{K}^+$ -ATPase activity would ameliorate the axonal transport defect and axonal atrophy. This construct is consistent with the normalization of axonal transport after ARI treatment reported by Tomlinson et al. (28–30).

However, after 6 mo of ARI treatment of hyperglycemic rats, this beneficial effect on axonal atrophy was not sustained. Instead, mean myelinated fiber size was markedly reduced even compared with untreated diabetic rats, which is most likely accounted for by the addition of small regenerating myelinated fibers reflected in a nonsignificant 29% increase in myelinated fiber density.

A similar effect of ponalrestat treatment was demonstrated on the perhaps functionally more relevant unmyelinated fiber population of the vagus nerve. The significant shift of fibers toward smaller sizes in untreated diabetic rats was largely ameliorated after 4 mo of ponalrestat treatment, indicating a protective effect on unmyelinated fiber atrophy.




FIG. 3. *Top*: axon-myelin ratio at 4 mo shows complete prevention of characteristic axonal atrophy as signified by decreased slope of regression (*b*) in untreated diabetic rats. A, control ( $n = 5$ ):  $a = 0.20 \pm 0.07$ ,  $b = 0.034 \pm 0.004$  ( $P < 0.05$  by analysis of variance [ANOVA]); B, diabetic ( $n = 5$ ):  $a = 0.36 \pm 0.05$ ,  $b = 0.024 \pm 0.001$ ; C, ponalrestat-treated diabetic ( $n = 5$ ):  $a = 0.24 \pm 0.09$ ,  $b = 0.035 \pm 0.004$ . For *b*,  $P < 0.05$ , A vs. B and B vs. C. *Bottom*: preventive effect above was not sustained at 6 mo. A ( $n = 5$ ):  $a = 0.07 \pm 0.09$ ,  $b = 0.038 \pm 0.004$  ( $P < 0.005$  by ANOVA); B ( $n = 5$ ):  $a = 0.42 \pm 0.06$ ,  $b = 0.019 \pm 0.003$ ; C ( $n = 5$ ):  $a = 0.28 \pm 0.10$ ,  $b = 0.002 \pm 0.003$ . For *b*,  $P < 0.002$  A vs. B and  $P < 0.01$  A vs. C.

Similar to the effect on myelinated fibers, the effect on unmyelinated parasympathetic fibers was not sustained at 6 mo of treatment. Instead, mean unmyelinated fiber size was not significantly different from that of untreated rats, whereas both unmyelinated fiber density and occupancy were increased  $>20\%$  compared with untreated diabetic BB rats, suggesting a significant addition of small regenerating fibers.

These findings are similar to those reported in the sural nerve of the BB rat after long-term prevention with an ARI (25) and after intervention therapy with the ARI sorbinil in human diabetic neuropathy (31), demonstrating a substantial regeneration of myelinated fibers. Unlike the data reported by Sharma et al. (32), we were not able to demon-

strate exacerbation of axonal degeneration after combined ponalrestat-insulin treatment in the vagus or the sural nerve (25). These data suggest that the metabolic perturbations in diabetic nerve not only cause nerve degeneration but also impede the normal regenerative capacity of peripheral nerve. It is possible that the same abnormality of the polyol pathway and subsequent *myo*-inositol-related protein kinase C-signaling mechanism suggested for the  $\text{Na}^+\text{-K}^+$ -ATPase defect may have an inhibitory effect on the responsiveness of neurotrophic factors necessary for nerve fiber regeneration (31).

In summary, ponalrestat treatment of the hyperglycemic BB rat has a significant effect on autonomic neuropathy by delaying the characteristic decrease in heart-rate variability and the neuroanatomic abnormalities of the vagus nerve. In addition, the same treatment regimen appears to markedly promote nerve fiber regeneration in the vagus nerve via an unknown mechanism.

#### ACKNOWLEDGMENTS

This study was supported in part by grants from ICI-Americas, Wilmington, Delaware; the Medical Research Council




FIG. 4. *A*: unmyelinated fiber size frequency distribution in diabetic rats at 4 mo is significantly shifted to smaller fiber sizes, indicating unmyelinated fiber atrophy. Open bars, control ( $n = 5$ ); solid bars, untreated diabetic ( $n = 5$ ). *B*: shift in fiber size is largely prevented by ponalrestat treatment. Open bars, control ( $n = 5$ ); hatched bars, ponalrestat-treated diabetic ( $n = 5$ ). \* $P < 0.05$ ; \*\* $P < 0.01$ .

of Canada; and the Canadian Diabetes Association. W.-X.Z. is a visiting scientist from the Medical University of Shanghai, People's Republic of China, supported by ICI-Americas. S.C. is supported by a scholarship from Diabetes Canada.

We are indebted to T.A.J. McEwen for technical assistance and Jackie McKane for preparing the manuscript.

## REFERENCES

- Ellenberg M: Diabetic neuropathy: clinical aspects. *Metabolism* 25:1627-55, 1976
- Dyrberg T, Benn J, Sandahl-Christiansen J, Hilsted J, Nerup J: Prevalence of diabetic autonomic neuropathy. *Diabetologia* 20:190-94, 1981
- Ewing DJ, Campbell IW, Clarke BF: Mortality in diabetic autonomic neuropathy. *Lancet* 1:601-603, 1976
- Pfeifer MA, Weinberg CR, Cook D, Best JD, Reenan A, Halter JB: Differential changes of autonomic nervous system function with age in man. *Am J Med* 75:249-58, 1983
- Pfeifer MA, Weinberg CR, Cook DL, Reenan A, Halter JB, Ensinnck JW, Porte D Jr: Autonomic neural dysfunction in recently diagnosed diabetic subjects. *Diabetes Care* 7:447-53, 1984
- Hosking DJ, Bennett T, Hampton JR: Diabetic autonomic neuropathy. *Diabetes* 27:1043-54, 1978
- Hilsted J: Pathophysiology in diabetic autonomic neuropathy: cardiovascular, hormonal, and metabolic studies. *Diabetes* 31:730-37, 1982
- Sima AAF: Structural and functional characterization of the neuropathy in the spontaneously diabetic BB-Wistar rat. In *Diabetic Neuropathy*. Goto Y, Horiuchi A, Koigure K, Eds. Amsterdam, Elsevier, 1982, p. 39-49
- Yagihashi S, Sima AAF: Diabetic autonomic neuropathy in the BB rat: ultrastructural and morphometric changes in sympathetic nerves. *Diabetes* 34:558-64, 1985
- Yagihashi S, Sima AAF: Diabetic autonomic neuropathy in BB rat: ultrastructural and morphometric changes in parasympathetic nerves. *Diabetes* 35:733-43, 1986
- McEwen TAJ, Sima AAF: Autonomic neuropathy in BB rat: assessment by improved method for measuring heart-rate variability. *Diabetes* 36:251-55, 1987
- Paro M, Prosdociami M, Zhang W-X, Sutherland G, Sima AAF: Autonomic neuropathy in BB rats and alterations in bladder function. *Diabetes* 38:1023-30, 1989
- Yagihashi S, Sima AAF: Neuroaxonal dystrophy in diabetic autonomic neuropathy: classification and topographic distribution in the BB rat. *J Neuropathol Exp Neurol* 45:545-65, 1986
- Sima AAF: Annotation: can the BB-rat help to unravel diabetic neuropathy? *Neuropathol Appl Neurobiol* 11:253-64, 1985
- Greene DA, Yagihashi S, Lattimer SA, Sima AAF: Nerve Na<sup>+</sup>-K<sup>+</sup>-ATPase, conduction and myo-inositol in the insulin deficient BB-rat. *Am J Physiol* 247:E534-39, 1984
- Greene DA, Lattimer SA: Impaired rat sciatic nerve sodium-potassium ATPase in acute streptozotocin diabetes and its correction by dietary myo-inositol supplementation. *J Clin Invest* 72:1058-63, 1983
- Lattimer SA, Sima AAF, Greene DA: In vitro correction of impaired Na<sup>+</sup>-K<sup>+</sup>-ATPase in diabetic nerve by protein kinase C-agonists. *Am J Physiol* 256:E264-69, 1989
- Sima AAF, Lattimer SA, Yagihashi S, Greene DA: Axo-glial dysjunction: a novel structural lesion that accounts for poorly reversible slowing of nerve conduction in the spontaneously diabetic BB-rat. *J Clin Invest* 77:474-84, 1986
- Greene DA, Chakrabarti S, Lattimer SA, Sima AAF: Role of sorbitol accumulation and myo-inositol depletion in paranodal swelling of large myelinated nerve fibers in the insulin deficient spontaneously diabetic Bio-Breeding rat: reversal by insulin replacement, an aldose reductase inhibitor and myo-inositol. *J Clin Invest* 79:1479-85, 1987
- McEwen TAJ, Sima AAF: Microcomputer collection and analysis of RR-interval data in the BB-rat. *Comput Biol Med*. In press
- Green A, Jaspan J, Kavin H, Chung S, Schoenberg H: Influence of long-term aldose reductase inhibitor therapy on autonomic dysfunction of urinary bladder, stomach and cardiovascular system in diabetic patients. *Diabetes Res Clin Pract* 1:67-75, 1987
- Williams SK, Howarth NL, Devenny JJ, Bitensky MW: Structural and functional consequences of increased tubulin glycosylation in diabetes mellitus. *Proc Natl Acad Sci USA* 79:6546-50, 1982
- Sharma AK, Bajada S, Thomas PK: Influence of streptozotocin-induced diabetes on myelinated nerve fiber maturation and body growth in the rat. *Acta Neuropathol* 53:257-65, 1981
- Leslie RA, Gwyn DG, Hopkins DA: The central distribution of the cervical vagus nerve and gastric afferent and efferent projections in the rat. *Brain Res Bull* 8:37-43, 1982
- Sima AAF, Prashar A, McEwen T: Diabetic neuropathy in the BB-rat and the effect of Stail-treatment (Abstract). *Diabetes* 38 (Suppl. 2):93A, 1989
- Medori R, Autilio-Gambetti L, Jenich H, Gambetti P: Changes in axonal size and slow axonal transport are related in experimental diabetic neuropathy. *Neurology* 38:597-601, 1988
- Medori R, Jenich H, Autilio-Gambetti L, Gambetti P: Experimental diabetic neuropathy: similar changes of slow axonal transport and axonal size in different animal models. *J Neurosci* 8:1814-21, 1988
- Tomlinson DR, Holmes PR, Mayer JH: Reversal by treatment with an aldose reductase inhibitor of impaired axonal transport and motor nerve conduction velocity in experimental diabetes. *Neurosci Lett* 31:189-93, 1982
- Tomlinson DR, Moriarty RJ, Mayer JH: Prevention and reversal of defective axonal transport and motor nerve conduction velocity in rats with experimental diabetes by treatment with the aldose reductase inhibitor sorbinil. *Diabetes* 33:470-76, 1984
- Tomlinson DR, Townsend J, Fretten P: Prevention of defective axonal transport in streptozotocin-diabetic rats by treatment with "Stail" (ICI 128436), an aldose reductase inhibitor. *Diabetes* 34:970-72, 1985
- Sima AAF, Brill V, Nathaniel V, McEwen TAJ, Brown MB, Lattimer SA, Greene DA: Regeneration and repair of myelinated fibers in sural nerve biopsy specimens from patients with diabetic neuropathy treated with sorbinil. *N Engl J Med* 319:548-55, 1988
- Boyrul S, Sharma AK, Stribling D, Mirrless D, Peterson RG, Farber MO, Thomas PK: Ultrastructural observations on myelinated fibres in experimental diabetes: effect of the aldose reductase inhibitor ponalrestat given alone or in conjunction with insulin therapy. *J Neurol Sci* 85:131-47, 1988