

# The Cannabinoid Receptors Are Required for Ultraviolet-Induced Inflammation and Skin Cancer Development

Duo Zheng, Ann M. Bode, Qing Zhao, Yong-Yeon Cho, Feng Zhu, Wei-Ya Ma, and Zigang Dong

Hormel Institute, University of Minnesota, Austin, Minnesota

## Abstract

**Solar UV irradiation is an important carcinogen that leads to the development of skin cancer, which is the most common human cancer. However, the receptors that mediate UV-induced skin carcinogenesis have not yet been unequivocally identified. Here we showed that UV irradiation directly activates cannabinoid receptors 1 and 2 (CB1/2). Notably, our data indicated that the absence of the CB1/2 receptors in mice results in a dramatic resistance to UVB-induced inflammation and a marked decrease in UVB-induced skin carcinogenesis. A marked attenuation of UVB-induced activation of mitogen-activated protein kinases and nuclear factor- $\kappa$ B was associated with CB1/2 deficiency. These data provide direct evidence indicating that the CB1/2 receptors play a key role in UV-induced inflammation and skin cancer development.** [Cancer Res 2008;68(10):3992–8]

## Introduction

Skin cancer is the most common type of human cancer and is associated with excessive exposure to UV solar irradiation. Exposure to UV irradiation can activate various oncogenes and inactivate many tumor suppressor genes. The net result is abnormal proliferation of keratinocytes that might harbor DNA damage leading to the onset of skin cancer (1). Exposure to UVB irradiation has been shown to induce activation, aggregation, and internalization of cell-surface receptors for epidermal growth factor (EGF), tumor necrosis factor  $\alpha$  (TNF- $\alpha$ ), and interleukin-1 (IL-1; refs. 2, 3).

Manipulation of the cannabinoid receptors has been useful in the pain management and treatment of osteoporosis, inflammation, and cancer (4), but the mechanisms of these effects are still not fully understood. Two cannabinoid receptors from mammalian tissues have been cloned and characterized (5, 6). Cannabinoid receptor 1 (CB1) is highly expressed in the brain, whereas the CB2 receptor is found mainly in the immune system (5, 6). However, no direct experimental evidence has confirmed that these membrane-bound receptors are required for UV-induced skin carcinogenesis.

CB1 and CB2 are members of a superfamily of seven-transmembrane-spanning (7-TM) receptors, which have a protein structure defined by an array of seven-membrane-spanning helices with intervening intracellular loops and a COOH-terminal domain that can associate with G proteins. A CB1 (7) and CB2 homology model (8) was constructed using the published X-ray crystal structure of bovine rhodopsin (9), a photosensitive G-protein-

coupled receptor (GPCR) found in the retina. Arrestin-1 (or visual arrestin) and arrestin-4 (or cone arrestin) modulate the action of rhodopsin (10, 11) and arrestin-2 and arrestin-3 modulate non-retinal GPCRs (12, 13). Recently, desensitization of the CB1 receptor was reported to be mediated by the interaction of the COOH-terminal residues (419–439) of CB1 with arrestin-2 (14), indicating that CB1 and CB2 might act as photoreceptors in nonretinal tissues. However, a role for CB1 and CB2 in skin cancer has not been reported. The identification of the “UV receptor” associated with skin cancer is still to be elucidated and the CB1/2 receptors are potential candidates.

The function of CB1 and CB2 in cancer is controversial. The expression levels of both CB1 and CB2 are significantly higher in CA-human papillomavirus-10 cells, which are virally transformed cells derived from human prostate adenocarcinoma, and also in other human prostate cancer cell lines, including LNCaP, DU145, PC3, and CWR22Rnu1, compared with human prostate epithelial and PZ-HPV-7 cells that are virally transformed cells derived from normal human prostate tissues (15). Moreover, the CB1 antagonist Rimonabant or SR141716 inhibited human breast cancer cell proliferation and was more effective in highly invasive metastatic MDA-MB-231 cells than in the less invasive T47D or MCF-7 cells (16). In leukemic precursor cells, the CB2 receptor is an oncoprotein that blocks neutrophilic differentiation when overexpressed in myeloid precursor cells (17). However, treatment of LNCaP cells with WIN# 55,212-2, a mixed CB1/CB2 agonist, resulted in inhibition of cell growth and induction of apoptosis (15, 18). Furthermore, cannabinoids have been shown to suppress the growth of several models of tumor xenografts in rats and mice (19). The CB1 and CB2 receptors are expressed in normal skin and skin tumors, and local activation of cannabinoid receptors induced the apoptotic death of tumorigenic epidermal cells, suppressed the proliferation of melanoma cells, and inhibited the growth and angiogenesis of skin tumor xenografts in nude mice (20, 21). However, cannabinoids may inhibit human keratinocyte proliferation through a non-CB1/CB2 mechanism (22). Therefore, although CB1 and CB2 are physiologically linked with cancer, the functional role of these receptors in cancer is not clear.

Here, we used wild-type CB1/2 (*CB1/2<sup>+/+</sup>*) and CB1/2-deficient (*CB1/2<sup>-/-</sup>*) mice and respective mouse embryonic fibroblasts (MEF) to elucidate the function of endogenous CB1 and CB2 in UVB-induced inflammation and skin cancer development.

## Materials and Methods

**Plasmids and chemicals.** The mouse CB1 plasmid (pcDNA3-CB1) was kindly provided by Dr. Beat Lutz and the mouse CB2 plasmid (pcDNA3-CB2) was kindly provided by Dr. Ruud Delwel. Chemical reagents for molecular biology and buffer preparation, including (*R*)-(+)-WIN55212-2, pertussis toxin, Tris, NaCl, and SDS, were purchased from Sigma-Aldrich Corp. The luciferase assay substrate was from Promega. Restriction enzymes were purchased from Roche Diagnostics Corp. and Taq DNA polymerase was from Qiagen, Inc. The DNA ligation kit (version 2.0) was

**Note:** D. Zheng, A.M. Bode, and Q. Zhao contributed equally to this work. The University of Minnesota is an equal opportunity educator and employer.

**Requests for reprints:** Zigang Dong, Hormel Institute, University of Minnesota, 801 16th Avenue Northeast, Austin, MN 55912. Phone: 507-437-9600; Fax: 507-437-9606; E-mail: zgdong@hi.umn.edu.

©2008 American Association for Cancer Research.  
doi:10.1158/0008-5472.CAN-07-6594

**Table 1.** Effect of UV radiation in CB1 or CB2 on ligand binding affinities

Receptor	vs [ <sup>3</sup> H]CP55940		
	UV radiation	Dose (kJ/m <sup>2</sup> )	Control/UV treated K <sub>i</sub> ratio
CB1	UVB	9	1.9:1
CB1	UVA	60	2.1:1
CB1	UVA	120	2.9:1
CB2	UVB	9	1.4:1
CB2	UVA	60	2.6:1

NOTE: Competition binding studies, using [<sup>3</sup>H]CP55940 as a tracer, were conducted with membrane preparations isolated from cells overexpressing CB1 or CB2 (Perkin-Elmer).

purchased from TAKARA Bio, Inc. Cell culture media and other supplements were purchased from Life Technologies, Inc. [<sup>3</sup>H]CP55940 and membrane fractions, which were isolated from HEK293 EBNA cells overexpressing human CB1 or CB2 receptors, were obtained from Perkin-Elmer Corp. The HisG antibody was from Invitrogen Corp. and antibodies against phosphorylated (p-) c-Jun NH<sub>2</sub>-terminal kinases (JNK), total JNKs, p-p38, total p38, p-extracellular signal-regulated kinases (ERK), and total ERKs were from Cell Signaling Technology, Inc. The antibody to detect phosphorylated serine residues was from Abcam, Inc., and the antibody against TNF- $\alpha$  was purchased from Santa Cruz Biotechnology, Inc.

**Mice.** The CB1/2-deficient and wild-type mice were provided by Dr. Andreas Zimmer.

**In vivo mouse studies.** Age- and gender-matched *CB1/2<sup>+/+</sup>* and *CB1/2<sup>-/-</sup>* mice were divided into groups and initiated by topical application of 200 nmol of 7,12-dimethyl benz(a)anthracene (DMBA), and then 2 wk later, 1 group each of *CB1/2<sup>+/+</sup>* and *CB2<sup>-/-</sup>* mice were treated with increasing doses (increase of 1.5 kJ/m<sup>2</sup>/wk) of UVB thrice a week as follows: week 1, 3 min for 1.5 kJ/m<sup>2</sup> (3 doses); week 2, 6 min for 3.0 kJ/m<sup>2</sup> (3 doses); week 3, 9 min for 4.5 kJ/m<sup>2</sup> (3 doses); week 4, 12 min for 6.0 kJ/m<sup>2</sup> (3 doses); week 5, 15 min for 7.5 kJ/m<sup>2</sup> (3 doses); week 6, 18 min for 9.0 kJ/m<sup>2</sup> (3 doses); week 7 through week 34, continued at 9.0 kJ/m<sup>2</sup> (3 doses/wk). Group 1 (*CB1/2<sup>+/+</sup>*; *n* = 30) and group 2 (*CB1/2<sup>-/-</sup>*; *n* = 30) were untreated, and group 3 (*CB1/2<sup>+/+</sup>*; *n* = 30) and group 4 (*CB1/2<sup>-/-</sup>*; *n* = 30) were treated with UVB only as indicated above. Mice were weighed, photographed, and tumors counted and measured once a week beginning when the first measurable tumors (1 mm<sup>3</sup>) were observed.

**Histopathology and immunohistochemistry.** Depilated adult *CB1/2<sup>+/+</sup>* and *CB1/2<sup>-/-</sup>* mice (6–8 wk old, six per group) were irradiated with 4 kJ/m<sup>2</sup> UVB. Dorsal trunk skin was harvested 24 h after irradiation, immediately fixed in 10% neutral buffered formalin, and processed for H&E staining and immunohistochemistry staining with an antibody against TNF- $\alpha$ .

**Binding analysis.** The ready-to-use membrane fractions were purchased and the overexpression of CB1 or CB2 was confirmed by a [<sup>3</sup>H]CP55940 binding assay following the manufacturer's (Perkin-Elmer) suggested protocol. In a competition binding assay, 5  $\mu$ g of membrane fractions were mixed with 0.5 nmol/L [<sup>3</sup>H]CP55940 and challenged with increasing concentrations (0.1–10  $\mu$ mol/L) of unlabeled ligand (*R*)-(+)-WIN55212-2. These fractions were or were not exposed to UVA (60 or 120 kJ/m<sup>2</sup>) or UVB (9 kJ/m<sup>2</sup>) and incubated at 30°C for 90 min. Nonspecific binding was assessed with 10  $\mu$ mol/L unlabeled ligand. Reactions were terminated by filtration through Whatman GF/C paper and washed with ice-cold incubation buffer [50 mmol/L Tris-HCl (pH 7.5), 2.5 mmol/L EGTA, 5 mmol/L MgCl<sub>2</sub>, 1.0 mg/mL bovine serum albumin]. Bound radioactivity was then determined by liquid scintillation counting. Results were analyzed using GraphPad Prism (version 4, GraphPad Software, Inc.).

**Cell culture and transfections.** *CB1/2<sup>+/+</sup>* and *CB1/2<sup>-/-</sup>* MEFs were isolated as previously described for similar cell lines (1). HEK293 cells and *CB1/2<sup>+/+</sup>* and *CB1/2<sup>-/-</sup>* MEFs were cultured at 37°C in a 5% CO<sub>2</sub> incubator in DMEM supplemented with 10% fetal bovine serum (FBS). When cells reached 50% to 60% confluence, transfection of the expression vectors was done with the Lipofectamine reagent (Invitrogen) following the manufacturer's suggested protocol.

**Construction of His-fusion proteins.** The His-fusion proteins for the CB1 and CB2 receptors were constructed with PCR-amplified open reading frame DNA fragments and the pcDNA4/HisMax vector (Invitrogen). The reading frame of each plasmid construct was confirmed by restriction mapping and DNA sequencing.

**Extraction of membrane and cytosolic proteins.** HEK293 cells were harvested, suspended in cell lysis buffer [20 mmol/L Tris-HCl (pH 8.0), 150 mmol/L NaCl], and disrupted by freezing and thawing. The supernatant fraction including cytosolic and nuclear soluble proteins was recovered by centrifugation at 15,000  $\times$  *g* for 10 min at 4°C and the pellet was resuspended with cell suspension buffer [50 mmol/L Tris-HCl (pH 8.0), 150 mmol/L NaCl, 1% Triton X-100]. The cell suspension was stirred on ice for 1 h, centrifuged at 15,000  $\times$  *g* for 10 min at 4°C, and membrane proteins were recovered in the supernatant fraction.

**Immunoprecipitation and Western blotting.** Immunoblotting was done following the instructions from Cell Signaling Technology. In brief, HEK293 cells or *CB1/2<sup>+/+</sup>* or *CB1/2<sup>-/-</sup>* MEFs were starved for 24 h in DMEM supplemented with 0.1% FBS. Cells were then treated with UVB (4 kJ/m<sup>2</sup>) and harvested in cell lysis or suspension buffer. Samples were equalized for protein and analyzed by Western blotting. To analyze CB1 or CB2 phosphorylation, a HisG antibody was used for immunoprecipitation with membrane proteins (500  $\mu$ g). The antibody binding was carried out at 4°C overnight and proteins were visualized by Western blotting with an antibody against phosphorylation of serine residues.

**Nuclear factor- $\kappa$ B activity assay.** The nuclear factor- $\kappa$ B (NF- $\kappa$ B) luciferase reporter plasmid (pGL2-NF- $\kappa$ B-Luc) was transfected into *CB1/2<sup>+/+</sup>* or *CB1/2<sup>-/-</sup>* MEFs together with the phRL-SV40 plasmid. The MEFs were cultured for an additional 36 h, starved for 24 h, and then exposed to UVB (4 kJ/m<sup>2</sup>). The cells were disrupted with lysis buffer 12 or 18 h later, and luciferase activity was measured.

## Results

**CB1 and CB2 are activated by UVA or UVB.** The CB1/2 receptors are seven-transmembrane-helix receptors that belong to the superfamily of GPCRs (5, 6). A GPCR isomerizes between two different states, an inactive (R) and an active (R\*) conformation (23), and the R\* state is appropriate for functional coupling to G proteins. Because an agonist has a high relative affinity for R\* forms compared with the inactive form, receptor binding of the agonist results in a shift of the equilibrium toward the active state and G protein coupling can occur virtually concomitantly (23). Activation of the CB1/2 receptors by UV treatment produced a similar shift in affinity for the agonist, (*R*)-(+)-WIN55212-2, compared with the untreated control, which was detected by an agonist-specific increase in binding affinity. Competition binding experiments using membrane fractions from cells overexpressing CB1 and exposed to UVB (9 kJ/m<sup>2</sup>) or UVA (60 or 120 kJ/m<sup>2</sup>) showed a 1.9-, 2.1-, or 2.9-fold lower K<sub>i</sub> value, respectively, compared with untreated control membrane fractions (Table 1). Similar increases in affinity were observed in membrane fractions from cells overexpressing the CB2 receptor: a 1.4-fold lower K<sub>i</sub> value with UVB (9 kJ/m<sup>2</sup>) and a 2.6-fold lower K<sub>i</sub> value with UVA (60 kJ/m<sup>2</sup>; Table 1). These findings suggested that the CB1 and CB2 receptors are activated by UVA or UVB treatment.


**CB1/2 double-knockout mice show resistance to UV-induced skin inflammation.** Because of the potential role of cannabinoids in cell growth inhibition and inflammation (24), we investigated

whether  $CB1/2^{-/-}$  mouse skin would exhibit an altered susceptibility to UVB-induced inflammation. At the established peak response time of 24 hours after irradiation (25),  $CB1/2^{+/+}$  and  $CB1/2^{-/-}$  mouse skin was harvested and analyzed. Histologic analysis indicated that UVB-treated  $CB1/2^{+/+}$  mouse skin showed a pronounced inflammation with the emergence of subcorneal pustules and epidermal erosion (Fig. 1A and B), whereas the epidermis of  $CB1/2^{-/-}$  mice was far more resistant to UVB-induced inflammation (Fig. 1C and D).


**The CB1/2 receptors are phosphorylated in response to UV treatment.** Current theories for GPCR regulation predict that the active state conformation is the target for phosphorylation, internalization, and desensitization (26). Phosphorylation hinders further association of the receptor with G proteins, thereby terminating the signaling pathways (27). In the continued presence of agonist, receptors are targeted to lysosomes for degradation (28) and agonist treatment has been shown to result in rapid phosphorylation of the CB1 and CB2 receptors in transfected cell systems (29, 30). The quick internalization of the CB1 and CB2 receptors after agonist exposure has been reported (30, 31). Activation of the CB1/2 receptors by UV irradiation might produce a similar rapid phosphorylation and internalization of these receptors. To test this hypothesis, we transfected His-epitope-tagged CB1 or CB2 into HEK293 cells and treated the cells with UVB followed by immunoprecipitation and Western blotting. Results indicated that treatment with UVB induced a rapid

phosphorylation of the CB1 and CB2 receptors, which was observed at 5 minutes and remained at this level until 30 minutes after UVB treatment (Fig. 2A and B). Treatment with UVB also induced rapid internalization of CB1 and CB2 as indicated by the time-dependent decrease in protein level of CB1 and CB2 in membrane fractions after UVB treatment (Fig. 2C and D). These results indicated that the CB1 and CB2 receptors are phosphorylated and internalized in response to UVB treatment.

**CB1/2 modulate JNK and ERK signaling cascades in response to UV treatment.** The GPCRs can stimulate the mitogen-activated protein (MAP) kinase cascades, which have prominent roles in the control of cell growth, differentiation, and proliferation (32). To show a potential role for the cannabinoid receptors in UV-induced signal transduction, we examined the phosphorylation of ERKs, JNKs, and p38 in  $CB1/2^{+/+}$  and  $CB1/2^{-/-}$  MEFs following UVB irradiation (4 kJ/m<sup>2</sup>). Results of Western blot analysis indicated that phosphorylation of ERKs, JNKs, and p38 was induced in a time-dependent fashion after exposure of  $CB1/2^{+/+}$  MEFs to UVB (Fig. 3A). In marked contrast, UVB-induced phosphorylation of ERKs and JNKs in  $CB1/2^{-/-}$  cells was almost abolished (Fig. 3A). Although p38 phosphorylation was somewhat attenuated, the decrease was not as substantial as the decrease in ERK and JNK phosphorylation (Fig. 3A). These data provide direct evidence supporting the involvement of the CB1/2 receptors in mediating the activation of the MAP kinase pathways in the cellular response to UVB.




**Figure 1.** CB1/2 deficiency reduces UV-induced skin inflammation.  $CB1/2^{+/+}$  (A and B) or  $CB1/2^{-/-}$  (C and D) mice were untreated or treated with UVB (4 kJ/m<sup>2</sup>). Dorsal skin from irradiated (B and D) or nonirradiated (A and C) mice was harvested 24 h after UVB exposure. For assessment of inflammation, tissues were fixed in 10% neutral-buffered formalin and stained with H&E. All UV irradiation experiments were done in triplicate for each condition and representative images are shown. Bar, 25  $\mu$ m; magnification,  $\times 200$ .



**Figure 2.** UVB induces CB1 or CB2 phosphorylation and internalization. HEK293 cells transfected with pcDNA4-HisG-CB1 (A) or pcDNA4-HisG-CB2 (B) were irradiated with UVB (4 kJ/m<sup>2</sup>) followed by incubation at 37°C in a 5% CO<sub>2</sub> incubator for 5, 15, or 30 min. The transfected cells not exposed to UVB were used as a negative control. Cells were harvested at the indicated time and the membrane protein was extracted and subjected to immunoprecipitation with a HisG monoclonal antibody. CB1 or CB2 serine phosphorylation was detected by Western blotting with a phospho-serine antibody. The total protein levels of CB1 (C) or CB2 (D) using the same sample as in A and B, respectively, were determined by Western blotting with the HisG monoclonal antibody to show the localization of the CB1 or CB2 receptor in the membrane fraction after UVB treatment.

Both CB1 and CB2 receptors are coupled to G<sub>i/o</sub> proteins, through which they stimulate the activity of the MAP kinases (33). The involvement of G<sub>i/o</sub> proteins in coupling the CB1 and/or CB2 receptor to JNK activation in this system was assessed with pertussis toxin (100 ng/mL), a compound that has widely been used as a reagent to characterize the involvement of heterotrimeric G proteins in signaling (34). Pertussis toxin catalyzes the ADP-ribosylation of specific G-protein α subunits of the G<sub>i</sub> family, which prevents the occurrence of the receptor and G-protein interaction (34). Pertussis toxin pretreatment before UVB treatment partially prevented UVB-induced JNK activation in *CB1/2*<sup>+/+</sup> cells, indicating the involvement of G<sub>i/o</sub> proteins in UVB-induced JNK activation (Fig. 3B). Cannabinoid receptors are also known to stimulate the phosphatidylinositol 3-kinase (PI3K)/Akt survival pathway (35), although the negative coupling of cannabinoid receptors to Akt has also been reported (36). In *CB1/2*<sup>+/+</sup> and *CB1/2*<sup>-/-</sup> cells, LY294002, a PI3K inhibitor, was used to determine whether the PI3K/Akt survival pathway is involved in the UVB-induced JNK activation. Results indicated that UVB-induced JNK phosphorylation is not mediated through PI3K (Fig. 3B). Notably, neither compound affected JNK phosphorylation in *CB1/2*<sup>-/-</sup> cells, which was still markedly decreased compared with wild-type cells.

**CB1/2 are involved in TNF-α-mediated NF-κB activation.** Manipulation of the cannabinoid receptors has been useful in the pain management and treatment of osteoporosis, inflammation, and cancer (4). Human keratinocytes constitutively secrete only a small amount of TNF-α and other cytokines, including IL-1, IL-6, IL-8, colony-stimulating factor, transforming growth factors, and platelet-derived growth factor. However, UV exposure significantly induces the production of cytokines (37, 38). TNF-α is an important member of the cytokine cascade and might be involved in the mediation of the inflammatory response to UV exposure, either directly or through the induction of other cytokines (39). We therefore examined the effect of UVB on the expression of TNF-α in *CB1/2*<sup>+/+</sup> and *CB1/2*<sup>-/-</sup> mouse skin. Interestingly, compared with adult *CB1/2*<sup>-/-</sup> mice, adult *CB1/2*<sup>+/+</sup> mouse skin showed a more elevated inflammatory cytokine TNF-α level (Fig. 4A and B) following UVB treatment, corresponding with the observed elevated inflammatory response (Fig. 1A-D). Thus, the CB1/2 receptors are involved in orchestrating events leading to the inflammation observed in UVB-treated mouse skin. Furthermore, CB1 or CB2 activation has been reported to result in enhanced NF-κB transactivation (40). To test whether CB1/2 deficiency is associated with suppression of NF-κB activity, we transfected a NF-κB-linked luciferase promoter plasmid into *CB1/2*<sup>+/+</sup> or *CB1/2*<sup>-/-</sup> MEFs and stimulated the cells with UVB (4 kJ/m<sup>2</sup>). The results indicated that NF-κB activation was induced in *CB1/2*<sup>+/+</sup> MEFs, but not in *CB1/2*<sup>-/-</sup> MEFs (Fig. 4C), suggesting that the CB1/2 receptors are involved in UVB-induced NF-κB activation.




**Figure 3.** UVB-induced MAP kinase activation is markedly attenuated in *CB1/2*-deficient (*CB1/2*<sup>-/-</sup>) MEFs. A, *CB1/2* wild-type (*CB1/2*<sup>+/+</sup>) or *CB1/2*<sup>-/-</sup> MEFs were irradiated with UVB (4 kJ/m<sup>2</sup>) and then incubated at 37°C for 5, 15, or 30 min. The cells not exposed to UVB were used as a negative control. Cells were then harvested and phosphorylation of ERKs, JNKs, and p38 was detected by immunoblotting with the respective specific phospho-antibodies. Total protein levels of ERKs, JNKs, and p38 were used as loading controls. B, *CB1/2*<sup>+/+</sup> and *CB1/2*<sup>-/-</sup> MEFs were treated with pertussis toxin (PTX; 100 ng/mL) or LY294002 (25 μmol/L) before UVB (4 kJ/m<sup>2</sup>) exposure. Cells were then incubated at 37°C for 15 min and harvested. Phosphorylation of JNKs was detected by immunoblotting with a specific phospho-antibody. Total protein levels of JNKs were used as a loading control.


**CB1/2 deficiency reduces UV-induced skin papilloma formation.** Studies in  $CB1^{-/-}$ ,  $CB2^{-/-}$ , or double-knockout mice have revealed responses to cannabinoids that are not mediated by the CB1/2 receptors (41). Candidates for the observed responses included other membrane-bound receptors (42) and the peroxisome proliferator-activated receptor family of nuclear receptor transcription factors (43). These results emphasize the importance of identifying the function of the CB1/2 receptors in carcinogenesis. To elucidate the function of the CB1/2 receptors in skin cancer development, we exposed mice to UVB in the two-stage skin carcinogenesis mouse model. Mice were initiated with DMBA and then treated with increasing doses of UVB beginning 2 weeks after initiation. Notably,  $CB1/2^{+/+}$  mice showed substantially greater overall papilloma development compared with  $CB1/2^{-/-}$  mice. Forty percent of wild-type mice developed papillomas compared with <10% for  $CB1/2^{-/-}$  mice over the same period of time (Fig. 5A). Furthermore, the papillomas found in  $CB1/2^{+/+}$  mice were more numerous (Fig. 5B) and larger (Fig. 5C) compared with  $CB1/2^{-/-}$  mice (Fig. 5D). DMBA alone had no effect on tumor development in either wild-type or CB1/2-deficient mice (data not shown). These results suggested that the CB1/2 receptors might have an enhancing effect on DMBA/UVB-induced skin tumor development.

## Discussion

Solar radiation induces acute and chronic reactions in human and animal skin. Whereas acute UV irradiation causes apoptosis, long-term exposure to UV irradiation induces activation of oncogenes and inactivation of tumor suppressor genes, resulting in abnormal proliferation of keratinocytes that may contain DNA damage leading to the onset of skin cancer (1, 44). In addition, UV irradiation stimulates cell proliferation through signal transduction initiated on the cell membrane. Activation, clustering, and internalization of cell-surface receptors for EGF, TNF- $\alpha$ , and IL-1 after exposure to UVB have been reported (2, 3). The effects of UV may be produced by a perturbation of membrane structure or a conformational change in membrane proteins resulting from energy absorption (3). Whether the CB1 and CB2 receptors are activated by UV radiation is not clear. Our *in vitro* binding studies indicated that UVB or UVA increased the affinity of the CB1 and CB2 receptors for a selected agonist, and UVB also induced CB1 and CB2 phosphorylation and internalization in the membrane fragments of cells overexpressing these proteins. These data indicated that UV activates the CB1 and CB2 receptors directly. Furthermore, from our *in vivo* studies, we found that tumorigenesis was significantly increased in wild-type ( $CB1/2^{+/+}$ ) mice compared with mice lacking the CB1/2 receptors ( $CB1/2^{-/-}$ ) after chronic or long-term exposure to UVB. These data showed that,



**Figure 4.** CB1/2 deficiency suppresses UV-induced NF- $\kappa$ B activation.  $CB1/2^{+/+}$  (A) or  $CB1/2^{-/-}$  (B) mice were untreated or treated with UVB (4 kJ/m<sup>2</sup>). Dorsal skin from irradiated (A and B, right) or nonirradiated (A and B, left) mice was harvested 24 h after UVB exposure. TNF- $\alpha$  was detected in formalin-fixed and paraffin-embedded sections of skin by the avidin-biotin complex immunohistochemistry staining method. TNF- $\alpha$  is visualized as the brown 3,3'-diaminobenzidine staining. All UV irradiation experiments were done in triplicate for each condition and representative images are shown. Bar, 25  $\mu$ m; magnification,  $\times 200$ . C,  $CB1/2^{+/+}$  or  $CB1/2^{-/-}$  MEFs were transiently transfected with 2  $\mu$ g of NF- $\kappa$ B luciferase reporter plasmid together with 0.1  $\mu$ g of the pRL-SV40 plasmid. At 24 h after transfection, cells were starved in 0.1% FBS/DMEM for another 24 h and then were untreated or treated with UVB (4 kJ/m<sup>2</sup>) and incubated at 37°C. After 12 or 18 h, the firefly luciferase activity was determined in cell lysates and normalized against *Renilla* luciferase activity. The relative activity is shown as compared with the activity of  $CB1/2^{+/+}$  MEFs without UVB exposure. Columns, means of triplicate experiments; bars, SD. \*,  $P < 0.05$ , Student's *t* test.



**Figure 5.** CB1/2 deficiency suppresses skin papilloma formation in the two-stage skin carcinogenesis mouse model. *A*, percentage of mice developing papillomas over a total of 34 wk. *B*, average number of papillomas per mouse. *C*, average papilloma volume per mouse as measured by caliper and calculated according to the following formula: [(short diameter)<sup>2</sup> / (long diameter)] / 2. *B* and *C*, points, mean; bars, SD. \*, *P* < 0.05, significantly greater multiplicity (*B*) or volume (*C*) of papillomas in wild-type mice compared with CB1/2 knockout (KO) mice. *D*, representative photographs of UVB-treated *CB1/2<sup>+/+</sup>* and *CB1/2<sup>-/-</sup>* mice.

besides the cell-surface receptors for EGF, TNF- $\alpha$ , or IL-1, the CB1/2 receptors are also the target of UV irradiation and have a marked effect on UV-induced skin carcinogenesis.

Our results also indicated that after acute exposure to UVB, *CB1/2<sup>+/+</sup>* mice displayed a much higher incidence of inflammation compared with *CB1/2<sup>-/-</sup>* mice, coinciding with an increased inflammatory cytokine TNF- $\alpha$  level. These are known effects of acute UV radiation (1). Cannabinoids have been reported to suppress the production of cytokines in animal models and in human cell culture, but they have also been shown to increase the production of cytokines, including TNF, IL-1, IL-6, and IL-10, when they are administered alone or together with bacteria or other antigens (45). Thus, cannabinoids might either suppress or enhance the production of these proinflammatory agents, depending on either the nature of the proinflammatory stimulus or the type of cannabinoids used. Our results showed that activation of the CB1/2 receptors by UV irradiation enhanced the production of TNF- $\alpha$  and inflammation in CB1/2 wild-type mouse skin. One of the crucial mediators of inflammation-induced tumor growth and progression is activation of NF- $\kappa$ B by a classic pathway, which may promote

tumor development (46, 47). Our data also indicated that NF- $\kappa$ B activation was decreased in *CB1/2<sup>-/-</sup>* mice compared with *CB1/2<sup>+/+</sup>* mice. Overall, these observations suggested that the CB1/2 receptors are required in the induction of the proinflammatory cascade-dependent tumor development in response to UVB.

### Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

### Acknowledgments

Received 12/10/2007; revised 1/30/2008; accepted 3/6/2008.

**Grant support:** The Hormel Foundation and NIH grants CA77646, CA27502, and CA111356.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

We thank Dr. Andreas Zimmer (Laboratory of Molecular Biology Clinic of Psychiatry, University Hospital, Bonn, Germany) for the CB1/2-deficient mice, Dr. Beat Lutz (Molecular Genetics of Behavior, Max-Planck-Institute of Psychiatry, Munich, Germany) for the mouse CB1 plasmid, and Dr. Ruud Delwel (Department of Hematology, Erasmus MC, Rotterdam, the Netherlands) for the mouse CB2 plasmid.

### References

- Melnikova VO, Ananthaswamy HN. Cellular and molecular events leading to the development of skin cancer. *Mutat Res* 2005;571:91-106.
- Sachsenmaier C, Radler-Pohl A, Zinck R, Nordheim A, Herrlich P, Rahmsdorf HJ. Involvement of growth factor receptors in the mammalian UVC response. *Cell* 1994;78:963-72.
- Rosette C, Karin M. Ultraviolet light and osmotic

- stress: activation of the JNK cascade through multiple growth factor and cytokine receptors. *Science* 1996;274:1194-7.
- Marx J. Drug development. Drugs inspired by a drug. *Science* 2006;311:322-5.
- Matsuda LA, Lolait SJ, Brownstein MJ, Young AC, Bonner TI. Structure of a cannabinoid receptor and functional expression of the cloned cDNA. *Nature* 1990;346:561-4.
- Munro S, Thomas KL, Abu-Shaar M. Molecular

- characterization of a peripheral receptor for cannabinoids. *Nature* 1993;365:61-5.
- Shim JY, Welsh WJ, Howlett AC. Homology model of the CB1 cannabinoid receptor: sites critical for non-classical cannabinoid agonist interaction. *Biopolymers* 2003;71:169-89.
- Xie XQ, Chen JZ, Billings EM. 3D structural model of the G-protein-coupled cannabinoid CB2 receptor. *Proteins* 2003;53:307-19.
- Palczewski K, Kumasaka T, Hori T, et al. Crystal

- structure of rhodopsin: a G protein-coupled receptor. *Science* 2000;289:739-45.
10. Hirsch JA, Schubert C, Gurevich VV, Sigler PB. The 2.8 Å crystal structure of visual arrestin: a model for arrestin's regulation. *Cell* 1999;97:257-69.
  11. Craft CM, Whitmore DH, Wiechmann AF. Cone arrestin identified by targeting expression of a functional family. *J Biol Chem* 1994;269:4613-9.
  12. Lohse MJ, Benovic JL, Codina J, Caron MG, Lefkowitz RJ.  $\beta$ -Arrestin: a protein that regulates  $\beta$ -adrenergic receptor function. *Science* 1990;248:1547-50.
  13. Attramadal H, Arriza JL, Aoki C, et al.  $\beta$ -arrestin2, a novel member of the arrestin/ $\beta$ -arrestin gene family. *J Biol Chem* 1992;267:17882-90.
  14. Bakshi K, Mercier RW, Pavlopoulos S. Interaction of a fragment of the cannabinoid CB1 receptor C-terminus with arrestin-2. *FEBS Lett* 2007;581:5009-16.
  15. Sarfaraz S, Afaq F, Adhami VM, Mukhtar H. Cannabinoid receptor as a novel target for the treatment of prostate cancer. *Cancer Res* 2005;65:1635-41.
  16. Sarnataro D, Pisanti S, Santoro A, et al. The cannabinoid CB1 receptor antagonist rimonabant (SR141716) inhibits human breast cancer cell proliferation through a lipid raft-mediated mechanism. *Mol Pharmacol* 2006;70:1298-306.
  17. Jorda MA, Lowenberg B, Delwel R. The peripheral cannabinoid receptor Cb2, a novel oncoprotein, induces a reversible block in neutrophilic differentiation. *Blood* 2003;101:1336-43.
  18. Sarfaraz S, Afaq F, Adhami VM, Malik A, Mukhtar H. Cannabinoid receptor agonist-induced apoptosis of human prostate cancer cells LNCaP proceeds through sustained activation of ERK1/2 leading to G<sub>1</sub> cell cycle arrest. *J Biol Chem* 2006;281:39480-91.
  19. Guzman M. Cannabinoids: potential anticancer agents. *Nat Rev* 2003;3:745-55.
  20. Casanova ML, Blazquez C, Martinez-Palacio J, et al. Inhibition of skin tumor growth and angiogenesis *in vivo* by activation of cannabinoid receptors. *J Clin Invest* 2003;111:43-50.
  21. Blazquez C, Carracedo A, Barrado L, et al. Cannabinoid receptors as novel targets for the treatment of melanoma. *FASEB J* 2006;20:2633-5.
  22. Wilkinson JD, Williamson EM. Cannabinoids inhibit human keratinocyte proliferation through a non-CB1/CB2 mechanism and have a potential therapeutic value in the treatment of psoriasis. *J Dermatol Sci* 2007;45:87-92.
  23. D'Antona AM, Ahn KH, Kendall DA. Mutations of CB1 T210 produce active and inactive receptor forms: correlations with ligand affinity, receptor stability, and cellular localization. *Biochemistry* 2006;45:5606-17.
  24. Guzman M. Effects on cell viability. In: Pertwee R, editor. *Cannabinoids. Handbook of Experimental Pharmacology*. Heidelberg: Springer-Verlag; 2005. p. 627-42.
  25. Schwarz A, Mahnke K, Luger TA, Schwarz T. Pentoxifylline reduces the formation of sunburn cells. *Exp Dermatol* 1997;6:1-5.
  26. Leurs R, Smit MJ, Alewijnse AE, Timmerman H. Agonist-independent regulation of constitutively active G-protein-coupled receptors. *Trends Biochem Sci* 1998;23:418-22.
  27. Abood ME. Molecular biology of cannabinoid receptors. In: Pertwee R, editor. *Cannabinoids. Handbook of Experimental Pharmacology*. Heidelberg: Springer-Verlag; 2005. p. 81-115.
  28. von Zastrow M, Kobilka BK. Ligand-regulated internalization and recycling of human  $\beta$ 2-adrenergic receptors between the plasma membrane and endosomes containing transferrin receptors. *J Biol Chem* 1992;267:3530-8.
  29. Garcia DE, Brown S, Hille B, Mackie K. Protein kinase C disrupts cannabinoid actions by phosphorylation of the CB1 cannabinoid receptor. *J Neurosci* 1998;18:2834-41.
  30. Bouaboula M, Dussosoy D, Casellas P. Regulation of peripheral cannabinoid receptor CB2 phosphorylation by the inverse agonist SR 144528. Implications for receptor biological responses. *J Biol Chem* 1999;274:20397-405.
  31. Rinaldi-Carmona M, Le Duigou A, Oustric D, et al. Modulation of CB1 cannabinoid receptor functions after a long-term exposure to agonist or inverse agonist in the Chinese hamster ovary cell expression system. *J Pharmacol Exp Ther* 1998;287:1038-47.
  32. Howlett AC. Cannabinoid receptor signaling. In: Pertwee R, editor. *Cannabinoids. Handbook of Experimental Pharmacology*. Heidelberg: Springer-Verlag; 2005. p. 53-79.
  33. Di Marzo V, Bifulco M, De Petrocellis L. The endocannabinoid system and its therapeutic exploitation. *Nat Rev Drug Discov* 2004;3:771-84.
  34. Fields TA, Casey PJ. Signalling functions and biochemical properties of pertussis toxin-resistant G-proteins. *Biochem J* 1997;321:561-71.
  35. Sanchez MG, Ruiz-Llorente L, Sanchez AM, Diaz-Laviada I. Activation of phosphoinositide 3-kinase/PKB pathway by CB(1) and CB(2) cannabinoid receptors expressed in prostate PC-3 cells. Involvement in Raf-1 stimulation and NGF induction. *Cell Signal* 2003;15:851-9.
  36. Gomez del Pulgar T, Velasco G, Sanchez C, Haro A, Guzman M. *De novo*-synthesized ceramide is involved in cannabinoid-induced apoptosis. *Biochem J* 2002;363:183-8.
  37. Kock A, Schwarz T, Kirnbauer R, et al. Human keratinocytes are a source for tumor necrosis factor  $\alpha$ : evidence for synthesis and release upon stimulation with endotoxin or ultraviolet light. *J Exp Med* 1990;172:1609-14.
  38. Oxholm A, Oxholm P, Staberg B, Bendtzen K. Immunohistological detection of interleukin I-like molecules and tumour necrosis factor in human epidermis before and after UVB-irradiation *in vivo*. *Br J Dermatol* 1988;118:369-76.
  39. Munker R, Gasson J, Ogawa M, Koeffler HP. Recombinant human TNF induces production of granulocyte-macrophage colony-stimulating factor. *Nature* 1986;323:79-82.
  40. Daaka Y, Zhu W, Friedman H, Klein TW. Induction of interleukin-2 receptor  $\alpha$  gene by  $\Delta^9$ -tetrahydrocannabinol is mediated by nuclear factor  $\kappa$ B and CB1 cannabinoid receptor. *DNA Cell Biol* 1997;16:301-9.
  41. Howlett AC, Barth F, Bonner TI, et al. International Union of Pharmacology. XXVII. Classification of cannabinoid receptors. *Pharmacol Rev* 2002;54:161-202.
  42. Sanchez C, Galve-Roperh I, Canova C, Brachet P, Guzman M.  $\Delta^9$ -Tetrahydrocannabinol induces apoptosis in C6 glioma cells. *FEBS Lett* 1998;436:6-10.
  43. Sun Y, Alexander SP, Kendall DA, Bennett AJ. Cannabinoids and PPAR $\alpha$  signalling. *Biochem Soc Trans* 2006;34:1095-7.
  44. Ichihashi M, Ueda M, Budiyo A, et al. UV-induced skin damage. *Toxicology* 2003;189:21-39.
  45. Klein TW. Cannabinoid-based drugs as anti-inflammatory therapeutics. *Nat Rev Immunol* 2005;5:400-11.
  46. Karin M, Greten FR. NF- $\kappa$ B: linking inflammation and immunity to cancer development and progression. *Nat Rev Immunol* 2005;5:749-59.
  47. Coussens LM, Werb Z. Inflammation and cancer. *Nature* 2002;420:860-7.