

# Perspectives in Diabetes

## Insulin, Prostaglandins, and the Pathogenesis of Hypertension

LLOYD AXELROD

**Hypertension is associated with hyperinsulinemia in the presence or absence of obesity or glucose intolerance. Physiological concentrations of insulin decrease the catecholamine-induced production of prostaglandin I<sub>2</sub> (PGI<sub>2</sub>; prostacyclin) and PGE<sub>2</sub>, two potent vasodilators, in adipose tissue, one of the largest organs in the body. This finding suggests that hyperinsulinemia increases peripheral vascular resistance and blood pressure by inhibiting the stimulatory effect of adrenergic agonists (and perhaps other agonists) on the production of PGI<sub>2</sub> and PGE<sub>2</sub> in adipose tissue (and perhaps other tissues). This concept is supported by evidence that PGI<sub>2</sub> and PGE<sub>2</sub> modulate vascular reactivity in states of health and disease. For example, during insulin deficiency, i.e., in diabetic ketoacidosis, PGI<sub>2</sub> and PGE<sub>2</sub> production by adipose tissue are increased, and peripheral vascular resistance and blood pressure are decreased. This hypothesis is also supported by evidence that blood flow through rat and human adipose tissue is decreased in obesity and that insulin decreases the blood flow through adipose tissue in nonobese rats. Thus, insulin may regulate PGI<sub>2</sub> and PGE<sub>2</sub> production by adipose tissue (and possibly other tissues) through a wide range of concentrations with important physiological and clinical consequences. *Diabetes* 40:1223–27, 1991**

**H**ypertension is associated with hyperinsulinemia in the presence or absence of obesity or glucose intolerance (1–4). Physiological concentrations of insulin decrease the catecholamine-induced production of prostaglandin I<sub>2</sub> (PGI<sub>2</sub>; prostacyclin) and PGE<sub>2</sub>, two potent vasodilators, in adipose tissue (5–8). This sug-

gests that an elevation of the circulating insulin level inhibits the stimulatory effect of adrenergic agonists (and perhaps other agonists) on the production of these vasodilative eicosanoids in adipose tissue (and perhaps other tissues) and thereby increases peripheral vascular resistance and blood pressure (Fig. 1). This perspective reviews information that pertains to this hypothesis and compares it with alternative explanations for the relationship between increased circulating insulin levels and hypertension.

### HYPERINSULINEMIA AND HYPERTENSION

Several studies have demonstrated an association between hyperinsulinemia and hypertension (1–4). In one study of 2475 people, insulin resistance and hyperinsulinemia were present in most hypertensive subjects (1), suggesting that insulin resistance and hyperinsulinemia are common pathophysiological characteristics of obesity, glucose intolerance, and hypertension and may be linked to the increased peripheral vascular resistance of hypertension. Although the prevalence of hypertension is increased in obesity, not all obese individuals are hypertensive. Instead, hypertension correlates with upper-body (central or android) obesity, which in turn is associated with insulin resistance, i.e., diminished sensitivity to the actions of insulin that promote glucose uptake by peripheral tissues such as skeletal muscle (9). The consequent hyperinsulinemia, which is compensatory with regard to glucose homeostasis and perhaps also with regard to regulation of body weight (10), may exert effects on other systems that are not resistant to the action of insulin and lead to deleterious consequences such as hypertension, hypertriglyceridemia, and atherosclerosis (9,11–14). Experimentally, modest hyperinsulinemia causes an increase in blood pressure. The infusion of human insulin into rats at a rate sufficient to raise the plasma insulin level from 180 to 288 pM produces an increase in blood pressure within 3 days (15).


### ROLE OF PGI<sub>2</sub> IN BLOOD PRESSURE REGULATION

A considerable amount of evidence supports the view that PGI<sub>2</sub> modulates vascular reactivity and blood pressure in

From the Diabetes Unit and Medical Services, Massachusetts General Hospital, and the Department of Medicine, Harvard Medical School, Boston, Massachusetts.

Address correspondence and reprint requests to Lloyd Axelrod, MD, Diabetes Unit, Massachusetts General Hospital, Boston, MA 02114.

Received for publication 31 May 1991 and accepted 24 June 1991.



**FIG. 1.** Role of insulin and prostaglandins in regulation of blood pressure. Effect of insulin on epinephrine-stimulated (and norepinephrine-stimulated) prostaglandin I<sub>2</sub> (PGI<sub>2</sub>; prostacyclin) and PGE<sub>2</sub> production by adipose tissue in normal state (A), during insulin deficiency (B), and during insulin excess (C). Also depicted are effects of adrenergic stimulation on prostaglandin production by adipose tissue and on peripheral vascular resistance in each circumstance. Each panel depicts effect of circulating insulin level (1) on PGI<sub>2</sub> and PGE<sub>2</sub> production by adipose tissue (2) and peripheral vascular resistance (3). When insulin levels are normal (A), PGI<sub>2</sub> and PGE<sub>2</sub> production by adipose tissue and peripheral vascular resistance are normal. During insulin deficiency (i.e., in diabetic ketoacidosis; B), PGI<sub>2</sub> and PGE<sub>2</sub> production by adipose tissue increases, resulting in decreased peripheral vascular resistance. When insulin levels are elevated (C), PGI<sub>2</sub> and PGE<sub>2</sub> production by adipose tissue decreases, resulting in increased peripheral vascular resistance and hypertension. Note close anatomic relationship between adipocytes and capillaries of adipose tissue. PGI<sub>2</sub> and PGE<sub>2</sub> production by adipose tissue requires interaction between adipocytes and endothelial cells of adipose tissue, in which adipocytes provide arachidonic acid substrate to endothelial cells, which convert it to PGI<sub>2</sub> and PGE<sub>2</sub>.

states of health and disease (16). PGE<sub>2</sub> is also a vasodilator but is less potent than PGI<sub>2</sub>. PGE<sub>2</sub> probably plays a subsidiary role to PGI<sub>2</sub> in the regulation of vascular tone and blood pressure. Of course, the two agents, which are produced in comparable quantities in adipose tissue in response to epinephrine (17), may produce an additive effect.

PGI<sub>2</sub> is produced locally in blood vessels (16). It is the principal product of arachidonic acid in all arteries and veins tested (18) and is also produced by capillary endothelium (19). PGI<sub>2</sub> is also produced by cultured human, bovine, and porcine endothelial cells, cultured human and bovine vascular smooth muscle cells, and other cells (20–28).

Catecholamines and other pressors stimulate PGI<sub>2</sub> production in humans (29,30). Angiotensin II stimulates PGI<sub>2</sub> production by cultured endothelial cells (23), stimulates the production of PGI<sub>2</sub> and other prostaglandins in various tissues (31), and appears to release PGI<sub>2</sub> into the circulation in animals (32,33) and humans (34).

Inhibition of PGI<sub>2</sub> production enhances vascular responsiveness to vasoconstrictors. Indomethacin, an inhibitor of prostaglandin synthesis, prevents the appearance of a PGI<sub>2</sub>-like substance into the circulation and increases the pressor response to angiotensin II in dogs (33). Indomethacin also enhances the pressor response to the intravenous infusion of angiotensin II in humans (35) and causes an increase in blood pressure in animals (36–38) and humans (39–41) under various circumstances.

Rats that are genetically susceptible to salt-induced hypertension exhibit impaired generation of vascular PGI<sub>2</sub> (and

increased generation of vascular thromboxane A<sub>2</sub>) compared with appropriate control animals (42).

PGI<sub>2</sub> modifies vascular resistance and blood pressure in several disease states. The infusion of PGI<sub>2</sub> at low rates, e.g., 5–10 ng · kg<sup>-1</sup> · min<sup>-1</sup>, in healthy subjects produces tachycardia, hypotension, decreased peripheral vascular resistance, nausea, vomiting, and, occasionally, abdominal pain (43–46). This suggests the possibility that increased endogenous PGI<sub>2</sub> production produces similar effects in certain disorders. This appears to be the case.

Direct hemodynamic evidence indicates that the decreased peripheral vascular resistance observed in diabetic ketoacidosis (DKA), despite profound volume depletion (which ordinarily causes increased peripheral vascular resistance), is a result of the vasodilative effects of increased PGI<sub>2</sub> (and possibly PGE<sub>2</sub>) production by adipose tissue as a consequence of insulin deficiency (47; Fig. 1B). Treatment with insulin decreases the increased PGI<sub>2</sub> and PGE<sub>2</sub> production and increases the peripheral vascular resistance toward a normal level (47). Increased PGI<sub>2</sub> and PGE<sub>2</sub> production by adipose tissue in DKA may also account for the nausea, vomiting, and abdominal pain that occur in this disorder, because the PGI<sub>2</sub> infusion rate that decreases vascular resistance and blood pressure in humans also causes nausea, vomiting, and, occasionally, abdominal pain (43–46).

The permissive effect of glucocorticoids on vascular tone and blood pressure, i.e., the enhancement of vascular responsiveness to pressors such as catecholamines without

necessarily having an effect when given alone, may be mediated by the ability of these agents to inhibit PGI<sub>2</sub> production by vascular endothelium and perhaps other cells (16). The inhibition of PGI<sub>2</sub> production by glucocorticoids may explain, at least in part, the hypertension of Cushing's syndrome (16). Conversely, the production of PGI<sub>2</sub> may be raised in Addison's disease and other states of glucocorticoid deficiency (16). Patients with untreated Addison's disease typically have tachycardia, hypotension, nausea, vomiting, and abdominal pain. Thus, increased PGI<sub>2</sub> production may also explain many of the cardiovascular and gastrointestinal features of Addison's disease (16).

In short, PGI<sub>2</sub> and PGE<sub>2</sub> production are inhibited by glucocorticoids and by insulin, possibly at different anatomic locations. When there is a deficiency of either hormone, the production of these vasodilative prostaglandins increases, causing decreased peripheral vascular resistance and hypotension and possibly nausea, vomiting, and abdominal pain. Thus, certain similarities in the clinical features of adrenal insufficiency and DKA may reflect the role of increased prostaglandin production in the pathogenesis of both disorders. When the level of either hormone is excessive, the production of vasodilative prostaglandins decreases, causing increased peripheral vascular resistance and hypertension.

#### ROLE OF PROSTAGLANDIN PRODUCTION BY ADIPOSE TISSUE IN THE PATHOGENESIS OF HYPERTENSION

Adipose tissue is one of the largest organs in the body. It weighs ~11.8 kg in a nonobese young woman and ~4.8 kg in a nonobese young man (48). The contribution of adipose tissue to total body mass is even greater in obesity and in nonobese individuals as they age (49). As noted above, hypertension correlates with upper-body obesity, i.e., with an increase in the size of the abdominal adipose depot.

Enlarged adipose tissue mass may contribute to increased peripheral vascular resistance in obesity. Blood flow through human adipose tissue measured by the clearance rate of radioactive xenon decreases significantly with increasing thickness of the fatty tissue (50). Blood flow through adipose tissue is also decreased in obese Zucker rats (51). Of course, the increase in peripheral vascular resistance in hypertension may also occur in other tissues such as skeletal muscle, but there is no direct evidence to support this possibility (49).

PGI<sub>2</sub> and PGE<sub>2</sub> play important roles in the normal function of adipose tissue and are produced in comparable quantities by adipose tissue (17). Catecholamine-stimulated PGI<sub>2</sub> and PGE<sub>2</sub> production by adipose tissue involves a cooperative interaction between the adipocyte and the endothelial cell of adipose tissue (17,52). Neither cell alone produces PGI<sub>2</sub> or PGE<sub>2</sub> in response to epinephrine. The presence of both cell types is required for the production of PGI<sub>2</sub> and PGE<sub>2</sub> (17,52). The adipocyte provides the arachidonic acid substrate to the endothelial cell, which utilizes the arachidonic acid to produce PGI<sub>2</sub> and PGE<sub>2</sub> (17,52). Although insulin inhibits PGI<sub>2</sub> and PGE<sub>2</sub> production by adipose tissue, the cellular site of action is not known.

One of the major physiological functions of endogenous prostaglandins in adipose tissue is the regulation of the local blood flow (53,54). Both PGE<sub>2</sub> and PGI<sub>2</sub> are potent vasodi-

lators in most vascular beds (55). The functional vasodilation of adipose tissue that accompanies lipolysis is due at least in part to the vasodilative effect of these prostaglandins (53,56). This concept is supported by evidence that the functional vasodilation is abolished by indomethacin and aspirin (53). Reincorporation of free fatty acids released during triglyceride lipolysis is a mechanism for regulation of the net rate of triglyceride breakdown in adipose tissue (57–59). The functional vasodilation of adipose tissue appears to provide a mechanism for the rapid removal of free fatty acids from adipose tissue and their rapid delivery to the systemic circulation. In this way, the functional vasodilation of adipose tissue may increase the efficiency of the lipolytic process and minimize the activity of a futile cycle of triglyceride lipolysis and reesterification of free fatty acids into neutral lipids.

PGI<sub>2</sub>, a potent lipolytic substance, and PGE<sub>2</sub>, a potent antilipolytic substance, provide a mechanism for the coordinate control of catecholamine-stimulated lipolysis in adipose tissue (17). Of course, epinephrine stimulates lipolysis in the absence of PGE<sub>2</sub> and PGI<sub>2</sub>, e.g., in isolated adipocytes. However, when adipocytes are mixed with endothelial cells or intact adipose tissue, epinephrine also stimulates the production of PGE<sub>2</sub> and PGI<sub>2</sub> during the activation of lipolysis (17,52). When both prostaglandins are produced, there appears to be no net effect on the rate of lipolysis (17). Thus, production of PGI<sub>2</sub> and PGE<sub>2</sub> may increase the efficiency of the lipolytic process by their vasodilative effects, whereas concurrent production of both a lipolytic and an antilipolytic prostaglandin prevents any alteration of the lipolytic cascade per se. It is possible that, in some circumstances, PGI<sub>2</sub> and PGE<sub>2</sub> production by adipose tissue is dissociated and that the ensuing imbalance alters the rate of hormone-stimulated lipolysis.

Insulin decreases prostaglandin production and blood flow in adipose tissue. Insulin in physiological concentrations inhibits catecholamine-stimulated PGI<sub>2</sub> production (5–7) and PGE<sub>2</sub> production (5,8) by adipose tissue *in vitro* (6,7) and *in vivo* (5,8). Infusion of insulin into lean Zucker rats produces a significant reduction in blood flow to the retroperitoneal, inguinal, and mesenteric fat depots but does not reduce blood flow to any nonadipose organ studied (60). After a meal, rat adipose tissue blood flow decreases in inverse relation to the circulating insulin levels (61). A glucose infusion sufficient to increase the circulating insulin level also decreases adipose tissue blood flow (62). Therefore, hyperinsulinemia may decrease the rate of blood flow and the efficiency of lipolysis in adipose tissue and increase systemic blood pressure by decreasing PGI<sub>2</sub> and PGE<sub>2</sub> production in adipose tissue (Fig. 1C).

Insulin may regulate prostaglandin production in adipose tissue through a wide range of concentrations with important physiological and clinical consequences. As indicated above, during insulin deficiency, i.e., in DKA, PGI<sub>2</sub> and PGE<sub>2</sub> production by adipose tissue is increased, and peripheral vascular resistance and blood pressure are decreased (47; Fig. 1B). Conversely, hyperinsulinemia may decrease PGI<sub>2</sub> and PGE<sub>2</sub> production by adipose tissue. This could contribute to the hypertension associated with hyperinsulinemia and could decrease the efficiency of lipolysis (Fig. 1C).

It may be argued that systemic production of PGI<sub>2</sub> is so

low that its inhibition by insulin is not likely to engender hypertension. In fact, the basal total-body production of PGI<sub>2</sub> in nonobese humans is small, <100 pg · kg<sup>-1</sup> · min<sup>-1</sup> (63). However, the production rate increases in response to catecholamines and angiotensin II in animals and humans (see above). Also, the basal or stimulated production rate has not been measured in obesity, especially upper-body obesity, and may be higher in obese than in nonobese subjects. Furthermore, increased or decreased PGI<sub>2</sub> or PGE<sub>2</sub> production in the vascular bed of a specific organ such as adipose tissue may not be detected by methods that assess total-body production rates.

#### ALTERNATIVE MECHANISMS BY WHICH INSULIN MAY CAUSE HYPERTENSION

Several mechanisms have been proposed by which hyperinsulinemia may cause hypertension (9,11–14). First, by virtue of its ability to promote renal tubular reabsorption of sodium, insulin causes an increase in total-body sodium and extracellular fluid volume so that higher renal (and therefore arterial) perfusion pressures are necessary to maintain sodium balance (9,11). Second, insulin increases sympathetic nervous system activity in humans (64), and resistance to the action of insulin on the sympathetic nervous system does not appear to develop in obese subjects (9,14). The increased sympathetic activity may then contribute to hypertension by increasing sodium reabsorption, cardiac output, and peripheral vascular resistance (9,64). Third, insulin stimulates the activity of the Na<sup>+</sup>-H<sup>+</sup> exchanger, which is linked to Ca<sup>2+</sup> exchange (11). Intracellular accumulation of Na<sup>+</sup> and Ca<sup>2+</sup> presumed to enhance the sensitivity of vascular smooth muscle to the pressor effects of norepinephrine, angiotensin II, and volume expansion. Fourth, insulin may cause vascular hypertrophy, thereby increasing peripheral vascular resistance (11).

These mechanisms may or may not explain the ability of hyperinsulinemia to cause hypertension. Although evidence can be marshaled in support of each of these proposed mechanisms, none of them is established beyond doubt, and serious reservations may be raised about each of them. For example, it is not established unequivocally that the ability of insulin to cause renal tubular sodium reabsorption increases blood pressure. In one carefully controlled study, the ability of insulin to cause proximal renal tubular sodium reabsorption did not raise blood pressure in healthy subjects or patients with insulin-dependent diabetes mellitus (65). Furthermore, total-body sodium content is increased to a comparable degree in normotensive and hypertensive patients with diabetes mellitus so that increased total-body sodium may contribute to the development of hypertension in susceptible individuals but is not sufficient to explain the relationship between hyperinsulinemia and hypertension (12,65). Similarly, the physiological relevance of insulin's ability to stimulate sympathetic nervous system activity is open to question, because the circulating insulin level that stimulates sympathetic nervous system activity as measured by the circulating norepinephrine level is not usually attained even in obesity (64). The data used to support a role for the Na<sup>+</sup>-H<sup>+</sup> exchanger as a mediator of the hypertensive effect of insulin were not obtained in vascular tissue of patients

with hypertension. Although the Na<sup>+</sup>-H<sup>+</sup> exchanger is a useful marker for hypertension, its role in the pathogenesis of hypertension has not been proven. Finally, the ability of insulin to stimulate vascular hypertrophy does not necessarily translate into increased peripheral vascular resistance. Nevertheless, our hypothesis and the alternative hypotheses proposed previously are not mutually exclusive. In fact, the several proposed mechanisms may be additive or synergistic.

#### CONCLUSIONS

The inhibitory effect of insulin on catecholamine-induced PGI<sub>2</sub> and PGE<sub>2</sub> production by adipose tissue (and possibly other tissues) can explain the association between hyperinsulinemia and hypertension. This concept is already supported by diverse experimental and clinical findings. Further tests of this hypothesis should include studies of the effects of hyperinsulinemia on blood flow and vascular resistance in adipose tissue (and other vascular beds) and on PGI<sub>2</sub> and PGE<sub>2</sub> production by adipose tissue (and other vascular beds) in normotensive and hypertensive subjects with and without obesity and glucose intolerance.

#### ACKNOWLEDGMENTS

This work was supported by grants from The Juvenile Diabetes Foundation International and the Weight Watchers Foundation, Inc.

I am indebted to Drs. David M. Nathan and Randall M. Zusman for reviewing this manuscript, Dr. Jerzy Kolaczynski for helpful discussions, and Carol Bovest and Christine Guterman for excellent secretarial assistance.

#### REFERENCES

1. Modan M, Halkin H, Almog S, Lusky A, Eshkol A, Shefi M, Shitrit A, Fuchs Z: Hyperinsulinemia: a link between hypertension, obesity and glucose intolerance. *J Clin Invest* 75:809–17, 1985
2. Manicardi V, Camellini L, Bellodi G, Coscelli C, Ferrannini E: Evidence for an association of high blood pressure and hyperinsulinemia in obese man. *J Clin Endocrinol Metab* 62:1302–304, 1986
3. Fournier AM, Gadia MT, Kubrusly DB, Skyler JS, Sosenko JM: Blood pressure, insulin, and glycemia in non-diabetic subjects. *Am J Med* 80:861–64, 1986
4. Bauman W, Maimon M, Langer O: An association between hyperinsulinemia and hypertension during the third trimester of pregnancy. *Am J Obstet Gyn* 159:446–50, 1988
5. Axelrod L, Levine L: Plasma prostaglandin levels in rats with diabetes mellitus and diabetic ketoacidosis. *Diabetes* 31:994–1001, 1982
6. Axelrod L, Levine L: Inhibitory effect of insulin on prostacyclin production by isolated rat adipocytes. *Prostaglandins* 25:571–79, 1983
7. Axelrod L, Ryan CA, Shaw JL, Kieffer JD, Ausiello DA: Prostacyclin production by isolated rat adipocytes: evidence for cyclic adenosine 3',5'-monophosphate-dependent and independent mechanisms and for a selective effect of insulin. *Endocrinology* 119:2233–39, 1986
8. Axelrod L, Shulman GI, Blackshear PJ, Bornstein W, Roussel AM, Aoki TT: Plasma level of 13,14-dihydro-15-keto-PGE<sub>2</sub> in patients with diabetic ketoacidosis and in normal fasting subjects. *Diabetes* 35:1004–10, 1986
9. Landsberg L: Diet, obesity and hypertension: an hypothesis involving insulin, the sympathetic nervous system and adaptive thermogenesis. *J Med* 61:1081–90, 1986
10. Clark MG, Rattigan S, Colquhoun EQ: Hypertension in obesity may reflect a homeostatic thermogenic response. *Life Sci* 48:939–47, 1991
11. DeFronzo RA, Ferrannini E: Insulin resistance: a multifaceted syndrome responsible for NIDDM, obesity, hypertension, dyslipidemia, and atherosclerotic cardiovascular disease. *Diabetes Care* 14:173–94, 1991
12. Nosadini R, Fioretto P, Trevisan R, Crepaldi G: Insulin-dependent diabetes mellitus and hypertension. *Diabetes Care* 14:210–19, 1991
13. Weidmann P, Ferrari P: Central role of sodium in hypertension in diabetic subjects. *Diabetes Care* 14:220–32, 1991
14. Daly PA, Landsberg L: Hypertension in obesity and NIDDM: role of insulin and sympathetic nervous system. *Diabetes Care* 14:240–48, 1991

15. Meehan WP, Buchanan TA, Shargill NS, Hsueh W: Chronic insulin administration elevates blood pressure in rats (Abstract). *Circulation* 82 (Suppl. 3):86, 1990
16. Axelrod L: Inhibition of prostacyclin production mediates permissive effect of glucocorticoids on vascular tone: perturbations of this mechanism contribute to pathogenesis of Cushing's syndrome and Addison's disease. *Lancet* 1:904-906, 1983
17. Chatzipanteli K, Rudolph S, Axelrod L: Coordinate control of lipolysis by prostaglandin E<sub>2</sub> and prostacyclin in rat adipose tissue (Abstract). *Clin Res* 39:277A, 1991
18. Dusting GJ, Moncada S, Vane JR: Prostacyclin: its biosynthesis, actions, and clinical potential. In *Prostaglandins and the Cardiovascular System*. Oates JA, Ed. New York, Raven, 1982, p. 59-106
19. Coughlin SR, Moskowitz MA, Zetter BR, Antoniadou HN, Levine L: Platelet-dependent stimulation of prostacyclin synthesis by platelet-derived growth factor. *Nature (Lond)* 288:600-602, 1980
20. Weksler BB, Marcus AJ, Jaffe EA: Synthesis of prostaglandin I<sub>2</sub> (prostacyclin) by cultured human and bovine endothelial cells. *Proc Natl Acad Sci USA* 74:3922-26, 1977
21. Baenziger NL, Becherer PR, Majerus PW: Characterization of prostacyclin synthesis in cultured human arterial smooth muscle cells, venous endothelial cells and skin fibroblasts. *Cell* 16:967-74, 1979
22. Baenziger NL, Fogarty FJ, Mertz LF, Chernuta LF: Regulation of histamine-mediated prostacyclin synthesis in cultured human vascular endothelial cells. *Cell* 24:915-23, 1981
23. MacIntyre DE, Pearson JD, Gordon JL: Localization and stimulation of prostacyclin production in vascular cells. *Nature (Lond)* 271:549-51, 1978
24. Levine L, Alam I: Arachidonic acid metabolism by cells in culture: analyses of culture fluids for cyclooxygenase products by radioimmunoassay before and after separation by high pressure liquid chromatography. *Prostaglandins Leukotrienes Med* 3:295-304, 1979
25. Raisz LG, Vanderhoek JY, Simmons HA, Kream BE, Nicolaou KC: Prostaglandin synthesis by fetal rat bone in vitro: evidence for a role of prostacyclin. *Prostaglandins* 17:905-14, 1979
26. Hassid A: Transport-active renal tubular epithelial cells (MDCK and LLC-PK<sub>1</sub>) in culture. Prostaglandin biosynthesis and its regulation by peptide hormones and ionophore. *Prostaglandins* 21:985-1001, 1981
27. Kreisberg JJ, Karnovsky MJ, Levine L: Prostaglandin production by homogeneous cultures of rat glomerular epithelial and mesangial cells. *Kidney Int* 22:355-59, 1982
28. Mitchell MD, Carr BR, Mason JJ, Simpson ER: Prostaglandin biosynthesis in the human fetal adrenal gland: regulation by glucocorticosteroids. *Proc Natl Acad Sci USA* 79:7547-51, 1982
29. Nadler JL, Zipser RD, Coleman R, Horton R: Stimulation of renal prostaglandins by pressor hormones in man: comparison of PGE<sub>2</sub> and prostacyclin (6-keto-PGF<sub>1α</sub>). *J Clin Endocrinol Metab* 56:1260-65, 1983
30. Nadler J, Zipser RD, Horton R: The effect of adrenergic stimulation on urinary prostaglandin E<sub>2</sub> and 6-keto-PGF<sub>1α</sub> in man. *Prostaglandins* 26:519-30, 1983
31. Jackson EK, Branch RA, Oates JA: Participation of prostaglandins in the control of renin release. In *Prostaglandins and the Cardiovascular System*. Oates JA, Ed. New York, Raven, 1982, p. 255-76
32. Swies J, Radomski M, Gryglewski RJ: Angiotensin-induced release of prostacyclin (PGI<sub>2</sub>) into circulation of anaesthetized cats. *Pharmacol Res Commun* 11:649-55, 1979
33. Mullane KM, Moncada S: Prostacyclin release and the modulation of some vasoactive hormones. *Prostaglandins* 20:25-49, 1980
34. Wennmalm A: Participation of prostaglandins in the regulation of peripheral vascular resistance. In *Prostaglandins and the Cardiovascular System*. Oates JA, Ed. New York, Raven, 1982, p. 303-31
35. Negus P, Tannen RL, Dunn MJ: Indomethacin potentiates the vasoconstrictor actions of angiotensin II in normal man. *Prostaglandins* 12:175-80, 1976
36. Anggard E, Larsson C: Arachidonic acid lowers and indomethacin increases the blood pressure of the rabbit. *J Pharm Pharmacol* 25:653-55, 1973
37. Lonigro AJ, Itskovitz HD, Crowshaw K, McGiff JC: Dependency of renal blood flow on prostaglandin synthesis in the dog. *Circ Res* 32:712-17, 1973
38. Colina-Chourio J, McGiff JC, Nasjletti A: Effect of indomethacin on blood pressure in the normotensive unanaesthetized rabbit: possible relation to prostaglandin synthesis inhibition. *Clin Sci* 57:359-65, 1979
39. Wennmalm A: Hypertensive effect of the prostaglandin synthesis inhibitor indomethacin. *ICRS J Med Sci* 2:1099, 1974
40. Martin K, Zipser R, Horton R: Effect of prostaglandin inhibition on the hypertensive action of sodium-retaining steroids. *Hypertension* 3:622-28, 1981
41. Oates JA, Fitzgerald GA, Branch RA, Jackson EK, Knapp HR, Roberts LJ II: Clinical implications of prostaglandin and thromboxane A<sub>2</sub> formation (second of two parts). *N Engl J Med* 319:761-67, 1988
42. Uehara Y, Tobian L, Iwai J, Ishii M, Sugimoto T: Alterations of vascular prostacyclin and thromboxane A<sub>2</sub> in Dahl genetic strain susceptible to salt-induced hypertension. *Prostaglandins* 33:727-38, 1987
43. Moncada S, Vane JR: Arachidonic acid metabolites and the interactions between platelets and vessel walls. *N Engl J Med* 300:1142-47, 1979
44. O'Grady J, Warrington S, Moti MJ, Bunting S, Flower R, Fowle ASE, Higgs EA, Moncada S: Effects of intravenous infusion of prostacyclin (PGI<sub>2</sub>) in man. *Prostaglandins* 19:319-32, 1980
45. Data JL, Molony BA, Meinzinger MM, Gorman RR: Intravenous infusion of prostacyclin sodium in man: clinical effects and influence on platelet adenosine diphosphate sensitivity and adenosine 3':5'-cyclic monophosphate levels. *Circulation* 64:4-12, 1981
46. Chaignon M, Ruffie ML, Lucsko M, Cuche JL, Guedon J, O'Grady J: Prostacyclin infusion in hypertension: acute hemodynamic and hormonal effects (Letter). *N Engl J Med* 307:559, 1982
47. Quyyumi A, Iaffaldano R, Guerrero JL, Ryan CA, Powell WJ Jr, Axelrod L: Prostacyclin and pathogenesis of hemodynamic abnormalities of diabetic ketoacidosis in rats. *Diabetes* 38:1585-94, 1989
48. Sjöström L, Smith U, Krotkiewski M, Björntorp P: Cellularity in different regions of adipose tissue in young men and women. *Metabolism* 21:1143-53, 1972
49. Bray GA: *The Obese Patient*. Philadelphia, PA, Saunders, 1976
50. Larsen OA, Lassen NA, Quaade F: Blood flow through human adipose tissue determined with radioactive xenon. *Acta Physiol Scand* 66:337-45, 1966
51. West DB, Prinz WA, Francendese AA, Greenwood MRC: Adipocyte blood flow is decreased in obese Zucker rats. *Am J Physiol* 253:R228-33, 1987
52. Parker J, Lane J, Axelrod L: Cooperation of adipocytes and endothelial cells required for catecholamine stimulation of PGI<sub>2</sub> production by rat adipose tissue. *Diabetes* 38:1123-32, 1989
53. Bowery JN, Lewis GP: Inhibition of functional vasodilation and prostaglandin formation in rabbit adipose tissue by indomethacin and aspirin. *Br J Pharmacol* 47:305-14, 1973
54. Axelrod L, Levine L: Prostacyclin production by isolated adipocytes. *Diabetes* 30:163-67, 1981
55. Moncada S, Vane JR: Pharmacology and endogenous roles of prostaglandins, endoperoxides, thromboxane A<sub>2</sub>, and prostacyclin. *Pharmacol Rev* 30:293-333, 1979
56. Bowery NG, Lewis GP, Matthews J: The relationship between functional vasodilation in adipose tissue and prostaglandins. *Br J Pharmacol* 40:437-45, 1970
57. Bülow J, Madsen J: Influence of blood flow on fatty acid mobilization from lipolytically active adipose tissue. *Pflügers Arch Eur J Physiol* 390:169-74, 1981
58. Leibel RL, Hirsch J: A radioisotopic technique for analysis of free fatty acid reesterification in human adipose tissue. *Am J Physiol* 248:E140-47, 1985
59. Edens NK, Leibel RL, Hirsch J: Mechanism of free fatty acid reesterification in human adipocytes in vitro. *J Lipid Res* 31:1423-31, 1990
60. Prinz WA, West DB, Cohen A, Silverberg S, Greenwood MRC: Intraarterial insulin infusion reduces adipose tissue blood flow (ATBF) (Abstract). *Fed Proc* 45:601, 1986
61. West DB, Prinz WA, Greenwood MRC: Regional changes in adipose tissue blood flow and metabolism in rats after a meal. *Am J Physiol* 257:R711-16, 1989
62. Madsen R, Malchow-Moller A: Effects of glucose, insulin and nicotinic acid on adipose tissue blood flow in rats. *Acta Physiol Scand* 118:175-80, 1983
63. Fitzgerald GA, Brash AR, Falardeau P, Oates JA: Estimated rate of prostacyclin secretion into the circulation of normal man. *J Clin Invest* 68:1272-76, 1981
64. Rowe JW, Young JB, Minaker KL, Stevens AL, Pallotta J, Landsberg L: Effect of insulin and glucose infusions on sympathetic nervous system activity in normal man. *Diabetes* 30:219-25, 1981
65. Trevisan R, Fioretto P, Semplicini A, Opocher G, Mantero F, Rocco S, Remuzzi G, Morocutti A, Zanette G, Donadon V, Perico N, Giorato C, Nosadini R: Role of insulin and atrial natriuretic peptide in sodium retention in insulin-treated IDDM patients during isotonic volume expansion. *Diabetes* 39:289-98, 1990