

# Effects of Naltrexone Sustained-Release/Bupropion Sustained-Release Combination Therapy on Body Weight and Glycemic Parameters in Overweight and Obese Patients With Type 2 Diabetes

PRISCILLA HOLLANDER, MD, PHD<sup>1</sup>  
 ALOK K. GUPTA, MD<sup>2,3</sup>  
 RAYMOND PLODKOWSKI, MD<sup>4</sup>  
 FRANK GREENWAY, MD<sup>2</sup>  
 HAROLD BAYS, MD<sup>5</sup>

COLLEEN BURNS, PHD<sup>6</sup>  
 PRESTON KLASSEN, MD<sup>6</sup>  
 KEN FUJIOKA, MD<sup>4</sup>  
 FOR THE COR-DIABETES STUDY GROUP\*

**OBJECTIVE**—To assess the efficacy and safety of 32 mg naltrexone sustained-release (SR)/360 mg bupropion SR (NB) in overweight/obese individuals with type 2 diabetes with or without background oral antidiabetes drugs.

**RESEARCH DESIGN AND METHODS**—This was a 56-week, double-blind, placebo-controlled study in which 505 patients received standardized lifestyle intervention and were randomized 2:1 to NB or placebo. Coprimary end points were percent weight change and achievement of  $\geq 5\%$  weight loss. Secondary end points included achievement of HbA<sub>1c</sub>  $< 7\%$  (53 mmol/mol), achievement of weight loss  $\geq 10\%$ , and change in HbA<sub>1c</sub>, waist circumference, fasting blood glucose, and lipids.

**RESULTS**—In the modified intent-to-treat population (54% female, 80% Caucasian, and mean age 54 years, weight 106 kg, BMI 37 kg/m<sup>2</sup>, and HbA<sub>1c</sub> 8.0% [64 mmol/mol]), NB resulted in significantly greater weight reduction ( $-5.0$  vs.  $-1.8\%$ ;  $P < 0.001$ ) and proportion of patients achieving  $\geq 5\%$  weight loss (44.5 vs. 18.9%,  $P < 0.001$ ) compared with placebo. NB also resulted in significantly greater HbA<sub>1c</sub> reduction ( $-0.6$  vs.  $-0.1\%$  [6.6 vs. 1.1 mmol/mol];  $P < 0.001$ ), percent of patients achieving HbA<sub>1c</sub>  $< 7\%$  (53 mmol/mol) (44.1 vs. 26.3%;  $P < 0.001$ ), and improvement in triglycerides and HDL cholesterol compared with placebo. NB was associated with higher incidence of nausea (42.3 vs. 7.1%), constipation (17.7 vs. 7.1%), and vomiting (18.3 vs. 3.6%). No difference was observed between groups in the incidence of depression, suicidal ideation, or hypoglycemia.

**CONCLUSIONS**—NB therapy in overweight/obese patients with type 2 diabetes induced weight loss, which was associated with improvements in glycemic control and select cardiovascular risk factors and was generally well tolerated with a safety profile similar to that in patients without diabetes.

*Diabetes Care* 36:4022–4029, 2013

Type 2 diabetes is a chronic disease characterized by elevated blood glucose and is often associated with a collection of metabolic abnormalities including low HDL cholesterol (HDL-C), elevated triglyceride concentrations, and high blood pressure. Excess body weight, particularly visceral adiposity, may contribute

to the development of both diabetes and the associated metabolic abnormalities, perhaps due to pathologic effects of excessive adipose tissue (1). Based on the National Health and Nutrition Examination Survey (NHANES) data from 1999–2010, the combined prevalence of overweight and obesity in the U.S. is 69% (2), while the prevalence of overweight and obesity in people with type 2 diabetes has been estimated to be even higher at  $\sim 85\%$  (3). Survey and clinical trial evidence supports a strong association between obesity (characterized by a BMI  $\geq 30$  kg/m<sup>2</sup> and central adiposity) and the development of insulin resistance, prediabetes, and type 2 diabetes (4).

In patients with type 2 diabetes, the best achievable glycemic control for the individual is recommended to reduce the risk of microvascular and possibly macrovascular disease. While lifestyle modifications can improve glycemic control, most patients with type 2 diabetes will typically require the sequential addition of one or more medications to achieve and sustain glycemic control. Unfortunately, many glucose-lowering medications lead to an increase in body weight (5). This presents challenges in an overweight or obese patient population often already struggling with weight control, particularly given that even modest intentional weight loss can improve cardiovascular risk factors (6,7) and decrease mortality (8).

Bupropion is a dopamine and norepinephrine reuptake inhibitor approved for use in the U.S. as an antidepressant and smoking cessation agent (9,10), with variable and modest effects on body weight as monotherapy (11). Naltrexone is an opioid receptor antagonist approved for use in the management of alcohol and opioid dependence (12). Animal studies suggest that when used together, bupropion and naltrexone increase the firing rate of proopiomelanocortin neurons, in part due to naltrexone-induced reduction of an autoinhibitory effect of  $\beta$ -endorphins

From the <sup>1</sup>Baylor Medical Center, Dallas, Texas; the <sup>2</sup>Pennington Biomedical Research Center, Louisiana State University System, Baton Rouge, Louisiana; <sup>3</sup>Our Lady of the Lake Physician Group, Baton Rouge, Louisiana; the <sup>4</sup>Scripps Clinic, La Jolla, California; the <sup>5</sup>Louisville Metabolic and Atherosclerosis Research Center Inc., Louisville, Kentucky; and <sup>6</sup>Orexigen Therapeutics, Inc., La Jolla, California.

Corresponding author: Priscilla Hollander, priscilh@baylorhealth.edu.

Received 28 January 2013 and accepted 15 July 2013.

DOI: 10.2337/dc13-0234. Clinical trial reg. no. NCD00474630, clinicaltrials.gov.

This article contains Supplementary Data online at <http://care.diabetesjournals.org/lookup/suppl/doi:10.2337/dc13-0234/-DC1>.

\*A full list of members of the COR-Diabetes Study Group can be found in the APPENDIX.

© 2013 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

(13). Increased proopiomelanocortin neuronal firing increases secretion of melanocortins such as  $\alpha$ -melanocyte stimulating hormone, which is thought to mediate anorectic effects and regulate energy balance (14). Furthermore, naltrexone and bupropion therapy are believed to influence the mesolimbic dopaminergic reward system, which can modulate reward behaviors such as food intake (14,15). Thus, this novel combination treatment is believed to act on the central nervous system to elicit sustained appetite reduction and enhance control of eating behavior.

While the development and clinical use of bupropion and naltrexone have spanned more than two decades, only recently have studies investigated the combined use of these agents in sustained-release (SR) form as a potential therapy for obesity. All three phase 3 trials of naltrexone SR plus bupropion SR (NB) in overweight or obese patients without diabetes demonstrated significant weight loss with an acceptable safety profile (16–18). This report details a 56-week, double-blind, placebo-controlled phase 3 clinical trial of NB for the treatment of overweight or obese patients with type 2 diabetes.

## RESEARCH DESIGN AND METHODS

This 56-week, randomized, double-blind, placebo-controlled study was conducted at 53 sites in the U.S. between May 2007 and June 2009. The protocol was approved by each site's Institutional Review Board. All patients provided written informed consent, and the study was conducted according to the guidelines and principles of Good Clinical Practices standards and the Declaration of Helsinki (19).

Participants were smoking or non-smoking men and women with type 2 diabetes, aged 18–70 years, with a BMI  $\geq 27$  and  $\leq 45$  kg/m<sup>2</sup>, HbA<sub>1c</sub> between 7% (53 mmol/mol) and 10% (86 mmol/mol), and fasting blood glucose  $< 270$  mg/dL. Participants were either not taking a diabetes medication or were on stable doses of oral antidiabetes drugs (OADs) for  $\geq 3$  months prior to randomization. Systolic and diastolic blood pressure were to be  $< 145$  and  $< 95$  mmHg, respectively. Most medications for the treatment of dyslipidemia and hypertension were allowed, as long as the doses were stable for at least 4 weeks prior to randomization. A complete list of inclusion and exclusion criteria is provided in the Supplementary Data.

Participants were randomized via a computer-generated randomization schedule in a 2:1 ratio (stratified by baseline HbA<sub>1c</sub> [ $\leq 8$  or  $> 8$ %;  $\leq 64$  or  $> 64$  mmol/mol] and sulfonylurea use) to 32 mg/day naltrexone SR combined with 360 mg/day NB or placebo. NB was provided as a single tablet containing 8 mg naltrexone SR and 90 mg bupropion SR. Study medication was initiated at one-quarter of the daily maintenance dose (one tablet) and increased weekly over the first 4 weeks such that the maintenance dose (2 tablets twice daily) was reached at the beginning of the fourth week. At week 2, participants received a telephone call to assess compliance with study medication and to document adverse events. Participant assessments were undertaken at screening, baseline, and every 4 weeks thereafter.

At baseline and weeks 4, 16, 28, and 40, all participants were instructed by study site personnel to follow a hypocaloric diet (500 kcal deficit/day, based on the World Health Organization algorithm for calculating resting metabolic rate). Participants received dietary counseling and the “Exchange Lists for Weight Management” booklets in accordance with the American Diabetes Association and American Dietetic Association guidelines. Participants also received advice on behavioral modification, including written instructions, to increase physical activity (to walking for at least 30 min most days of the week).

Participants were instructed to monitor their blood glucose twice daily for at least the first 4 weeks after randomization and during suspected hypoglycemia. In the case of loss of glucose control, defined as HbA<sub>1c</sub>  $> 9.5$ % (80 mmol/mol) at week 16 or subsequent visits, any postbaseline HbA<sub>1c</sub>  $\geq 10.0$ % (86 mmol/mol) or two or more successive postbaseline fasting glucose levels  $\geq 270$  mg/dL, the participant's dose of current OADs was to be increased (if not at maximal doses) or an additional OAD was to be added using a protocol-defined algorithm.

## Outcome measures

The coprimary end points were 1) percent change in body weight from baseline to week 56 compared with placebo and 2) percentage of participants achieving  $\geq 5$ % reduction in body weight from baseline to week 56 compared with placebo. Secondary end points included percentage of patients achieving  $\geq 10$ % reduction in body weight and change in

selected cardiovascular risk factors including waist circumference, triglycerides, HDL-C, LDL-C, and high-sensitivity C-reactive protein (hs-CRP). Secondary end points related to glycemic control included change from baseline in HbA<sub>1c</sub>, fasting blood glucose, fasting insulin, and homeostasis model assessment of insulin resistance (HOMA-IR), as well as the percentage of patients achieving HbA<sub>1c</sub>  $< 7$ % (53 mmol/mol) and  $< 6.5$ % (48 mmol/mol), changes in OADs, percentage of patients requiring rescue medications for their diabetes, and percentage of patients discontinuing from the study owing to poor glycemic control.

Safety assessments included evaluation of treatment-emergent adverse events, vital signs, electrocardiograms, and clinical laboratory measures. Depressive symptoms, evaluated using the Inventory of Depressive Symptomatology–Self Report (IDS-SR) ([www.ids-qids.org](http://www.ids-qids.org)) and blood pressure were also evaluated for both efficacy and safety.

## Statistical analyses

To obtain the targeted number of participant exposures at 1 year, we estimated that 350 participants would need to be randomized to NB, with an assumed 33% attrition rate (20). It was estimated that 525 participants randomized 2:1 ( $\sim 350$  to NB and  $\sim 175$  to placebo) would provide  $\sim 99$ % power to detect a difference in mean weight loss of  $\geq 5$ % between NB and placebo (assuming an SD of 5%, comparison between groups using a two-sample *t* test and two-sided significance level of 0.05).

Unless otherwise specified, primary efficacy analyses were performed using the modified intent-to-treat (mITT) population, prospectively defined as randomized participants with a baseline and one or more postbaseline measurements of body weight while on the study drug. Missing data were imputed by carrying forward the last observation on study drug (last observation carried forward [LOCF]). Analyses were also performed on the completer population, which comprised all mITT patients who had a body weight measurement at week 56 while on the study drug. Safety analyses were performed on all randomized participants who took one or more tablets of the study drug and had at least one contact or assessment with the investigative site at any time after beginning the study.

General linear models (ANCOVA) including terms for treatment, HbA<sub>1c</sub>

strata  $\leq 8$  or  $> 8\%$ , pharmacotherapy with or without sulfonylurea, and baseline value as covariate were used to analyze continuous end points. Categorical end points were analyzed with a logistic regression model using the same covariates as the continuous end points. To minimize skewness, values for triglycerides, hs-CRP, insulin, and HOMA-IR were log<sub>10</sub> transformed before running the general linear models. The least squares (LS) percent change from baseline was calculated by back-transforming the LS mean in log<sub>10</sub> scale.

To control for multiple comparisons, secondary end points were analyzed in a predetermined sequence, beginning with the change from baseline to end point in HbA<sub>1c</sub>, fasting triglycerides, HDL-C, blood glucose, and waist circumference; proportion of participants with  $\geq 10\%$  decrease in total body weight; proportion of participants with HbA<sub>1c</sub>  $< 7\%$  (53 mmol/mol); proportion of participants requiring rescue medications for diabetes or changing the doses of their OADs; changes from baseline to end point in HOMA-IR and fasting insulin; proportion of participants with HbA<sub>1c</sub>  $< 6.5\%$  (48 mmol/mol); changes from baseline to end point in hs-CRP; percentage of participants discontinuing owing to poor glycemic control; and changes from baseline to end point in LDL-C, systolic blood pressure, diastolic blood pressure, and IDS-SR total score. Testing proceeded in a sequential step-down manner until any end point failed to reach  $P < 0.05$ , after which nominal  $P$  values are reported and findings deemed exploratory. Continuous data are presented as LS mean  $\pm$  SE unless otherwise indicated. All statistical analyses were performed using Windows SAS, version 9.1 (SAS Institute, Cary, NC).

**RESULTS**—Of 1,625 patients who entered screening, 505 were randomly assigned to receive NB ( $n = 335$ ) or placebo ( $n = 170$ ). The most frequent ( $\geq 5\%$ ) reasons for screen failure were laboratory abnormality (58%, including HbA<sub>1c</sub>  $< 7$  or  $> 10\%$ ), concomitant medication (7%), vital sign abnormality (5%), and withdrawal of consent (5%). For NB, the mITT, completer, and safety populations included 265, 175, and 333 patients, respectively. For placebo, the mITT, completer, and safety populations included 159, 100, and 169 patients, respectively (details in Supplementary Fig. 1). Baseline characteristics of the randomized

and the mITT populations were similar (Table 1). Over the 56-week trial, 47.8% of participants in the NB group discontinued the study drug compared with 41.2% in the placebo group. A greater percentage of participants who received NB compared with placebo discontinued owing to an adverse event (29.3 vs. 15.3%). Conversely, a greater percentage of placebo than NB participants were lost to follow-up (8.8 vs. 6.6%), withdrew consent (8.8 vs. 6.3%), or withdrew because of self-perceived insufficient weight loss (3.5 vs. 1.5%).

Participants treated with NB lost significantly more weight than placebo-treated participants (mITT analysis:  $-5.0 \pm 0.3$  vs.  $-1.8 \pm 0.4\%$ ,  $P < 0.001$ ) (Fig. 1A). The difference between the groups was significant at the first time point assessed (Week 4) and sustained throughout 56 weeks of treatment ( $P < 0.001$  for all visits). A similar pattern was observed in the completer population, with numerically greater weight loss relative to the mITT population seen in both NB and placebo patients (completers:  $-5.9 \pm 0.5$  vs.  $-2.2 \pm 0.6\%$ ,  $P < 0.001$ ) (Fig. 1B). Consistent with the mean


percent changes in body weight, more patients treated with NB than placebo achieved  $\geq 5$  and  $\geq 10\%$  reduction in body weight at week 56 (all  $P < 0.001$ ) (Fig. 1A and B).

Patients treated with NB exhibited a greater improvement in HbA<sub>1c</sub> ( $-0.6\%$  [ $-6.6$  mmol/mol]) than in placebo-treated patients ( $-0.1\%$  [ $-1.1$  mmol/mol]), with a placebo-corrected difference of  $-0.5\%$  ( $-5.5$  mmol/mol) for the mITT population at the end of 56 weeks ( $P < 0.001$ ) (Fig. 1C). Treatment with NB also resulted in a greater percentage of patients achieving an HbA<sub>1c</sub> of  $< 7.0\%$  (53 mmol/mol) compared with placebo (44.1 vs. 26.3%;  $P < 0.001$ ) and  $< 6.5\%$  (48 mmol/mol) (20.7 vs. 10.2%;  $P = 0.004$ ) (Table 2). Exploratory analyses demonstrated that HbA<sub>1c</sub> change was significantly correlated with change in body weight with both treatments (NB mITT:  $r = 0.509$ ,  $P < 0.001$ ; placebo:  $r = 0.168$ ,  $P < 0.05$ ). Patients with higher baseline HbA<sub>1c</sub> values also exhibited significantly greater HbA<sub>1c</sub> reductions with NB vs. placebo: baseline  $> 8\%$  (64 mmol/mol)  $-1.1$  vs.  $-0.5\%$ ; baseline  $> 9\%$  (75 mmol/mol)  $-1.2$  vs.  $-0.3\%$ ; both  $P < 0.05$  (Supplementary

**Table 1—Demographics and baseline characteristics**

	Randomized		mITT	
	Placebo	NB	Placebo	NB
N	170	335	159	265
Age (years)	53.5 $\pm$ 9.8	54.0 $\pm$ 9.1	53.8 $\pm$ 9.7	53.9 $\pm$ 9.2
Sex (% female)	52.9	58.2	52.8	54.3
Race (% white/black/other) <sup>†</sup>	82.4/10.6/7.0	77.9/18.8/3.3	83.0/10.7/6.3	78.1/19.2/2.7
Weight (kg)	105.1 $\pm$ 17.0	104.2 $\pm$ 18.9	105.0 $\pm$ 17.1	106.3 $\pm$ 19.1
BMI (kg/m <sup>2</sup> )	36.4 $\pm$ 4.5	36.4 $\pm$ 4.8	36.3 $\pm$ 4.5	36.7 $\pm$ 4.8
HbA <sub>1c</sub> (%)	8.0 $\pm$ 0.9	8.0 $\pm$ 0.8	8.0 $\pm$ 0.9	8.0 $\pm$ 0.8
Dyslipidemia (% yes) <sup>††</sup>	85.3	83.6	85.5	82.6
Sulfonylurea use (%) <sup>†††</sup>	48.8	46.6	49.1	49.1
Thiazolidinedione use (%) <sup>†††</sup>	30.6	30.7	31.4	31.3
Metformin use (%)	76.5	78.5	76.7	79.6
Systolic blood pressure (mmHg)	124.7 $\pm$ 9.7	125.3 $\pm$ 10.9	124.5 $\pm$ 9.6	125.0 $\pm$ 11.0
Diastolic blood pressure (mmHg)	77.6 $\pm$ 7.0	77.3 $\pm$ 7.5	77.4 $\pm$ 7.1	77.5 $\pm$ 7.5
Pulse rate (bpm)	73.2 $\pm$ 9.1	73.0 $\pm$ 8.7	73.1 $\pm$ 9.1	72.9 $\pm$ 8.3
IDS-SR total score	7.7 $\pm$ 5.7	8.1 $\pm$ 5.7	7.8 $\pm$ 5.7	8.2 $\pm$ 5.9

Data are means  $\pm$  SD unless otherwise indicated. Percentages may not add up to 100 owing to rounding. <sup>†</sup>A significant difference between treatment groups existed for race, although the % of white patients did not significantly differ between treatment groups. There were no significant differences between placebo and NB treatment arms (randomized and mITT) for any other baseline/demographic value. <sup>††</sup>Diagnosed at baseline with dyslipidemia, hypercholesterolemia, hypertriglyceridemia, hyperlipidemia, or low HDL-C or had at least one of the following values prior to first dose of study drug: triglyceride  $\geq 200$  mg/dL, LDL-C  $\geq 160$  mg/dL, total cholesterol  $\geq 240$  mg/dL, and HDL cholesterol  $< 40$  mg/dL. <sup>†††</sup>Sulfonylurea medications included avandaryl, glibenclamide, glimepiride, glipizide, and tolazamide; thiazolidinedione medications included avandaryl, pioglitazone, rosiglitazone, troglitazone.



**Figure 1**—A: Body weight change from baseline in the mITT LOCF population. Data are LS mean ± SE. Placebo, n = 159; NB, n = 265. \*\*\*P < 0.001 vs. placebo; except for change from baseline to week 56 end point for mITT LOCF population, P values are nominal (i.e., not adjusted for multiple comparisons) and associated with exploratory analyses. B: Body weight change from baseline to week 56 end point; placebo, n = 100; NB, n = 175. \*\*\*P < 0.001 vs. placebo; except for ≥5% weight loss category for the mITT-LOCF

Fig. 2). Changes in fasting glucose concentration and HOMA-IR were not significantly different compared with placebo (Table 2).

Over the course of the study, fewer NB-treated patients required an increase in dose or the addition of another OAD owing to deterioration of glycemic control (22.3% NB vs. 35.2% placebo; P < 0.01). Because alterations in medications for diabetes can obscure the impact of treatment on glycemic-related secondary end points, an exploratory analysis was performed to investigate changes in glycemic parameters accounting for rescue medication by carrying forward the last observation prior to introduction of the rescue medication. In this analysis, larger placebo-corrected improvements in HbA<sub>1c</sub> (−0.7% [7.7 mmol/mol]; P < 0.001) and fasting blood glucose (−17.5 mg/dL; P < 0.001) were observed; differences in fasting insulin and HOMA-IR were not significant between NB and placebo groups (Supplementary Table 2).

This study also examined the effect of NB treatment on cardiovascular risk factors. Compared with placebo-treated patients, NB-treated patients had significantly greater reductions in waist circumference and serum triglyceride concentration and significant increases in HDL-C concentration (Table 2). No significant differences were observed between the groups in LDL-C or hs-CRP. Systolic and diastolic blood pressure tended to be reduced from baseline in both NB and placebo-treated patients, with numerically but not statistically greater reductions in placebo-treated patients (Table 2).

**Safety and tolerability**

The most common adverse events that were more prevalent in the NB-treated patients were nausea, constipation, vomiting, and diarrhea (Table 3). Nausea led to study withdrawal in 9.6% of NB-treated patients, with the vast majority (28 of 32) of these withdrawals occurring as the result of nausea with an onset during the first 4 weeks of treatment. Adverse events that led to medication discontinuation during the first 4 weeks of treatment were the primary reason that relatively fewer randomized NB patients were included in the mITT population. Nausea occurred at a greater incidence in NB-treated patients taking metformin at baseline (46.2%) compared with those not on metformin (28.2%). The incidence of subjects with serious adverse events was low (3.9% for NB and 4.7% for placebo)

Downloaded from <http://diabetesjournals.org/care/article-pdf/36/12/4022/166084022.pdf> by guest on 18 January 2025

Table 2—Changes in secondary end points

	Placebo	NB	P (NB vs. placebo)
N	159	265	
HbA <sub>1c</sub> (%)			
Baseline	8.0 ± 0.9	8.0 ± 0.8	
Change	−0.1 ± 0.1	−0.6 ± 0.1	<0.001*
% patients with HbA <sub>1c</sub> <7%	26.3	44.1	<0.001
% patients with HbA <sub>1c</sub> <6.5%	10.2	20.7	0.004
Fasting blood glucose (mg/dL)			
Baseline	163.9 ± 44.5	160.0 ± 41.3	
Change	−4.0 ± 3.4	−11.9 ± 2.7	0.065
Fasting insulin (μIU/mL)†			
Baseline	13.8 ± 1.9	15.1 ± 1.9	
% change (95% CI)	−10.4 (−18.8 to −1.1)	−13.5 (−19.7 to −6.8)	0.563
HOMA-IR†			
Baseline	5.2 ± 2.0	5.7 ± 2.0	
% change (95% CI)	−14.7 (−24.7 to −3.3)	−20.6 (−27.8 to −12.6)	0.361
Waist circumference (cm)			
Baseline	114.3 ± 12.4	115.6 ± 12.6	
Change	−2.9 ± 0.6	−5.0 ± 0.5	0.006
Triglycerides (mg/dL)†			
Baseline	165.6 ± 1.6	143.3 ± 1.7	
% change (95% CI)	−0.8 (−7.0 to 5.8)	−11.2 (−15.6 to −6.6)	0.007*
HDL-C (mg/dL)			
Baseline	46.1 ± 11.5	46.2 ± 10.2	
Change	−0.3 ± 0.6	+3.0 ± 0.5	<0.001*
LDL-C (mg/dL)			
Baseline	101.0 ± 33.9	100.2 ± 34.2	
Change	0.0 ± 2.4	−1.4 ± 1.9	0.641
hs-CRP (mg/L)†			
Baseline	3.3 ± 2.8	3.6 ± 3.0	
% change (95% CI)	−13.3 (−24.9 to 0.2)	−20.9 (−29.3 to −11.5)	0.312
Systolic blood pressure (mmHg)			
Baseline	124.5 ± 9.6	125.0 ± 11.0	
Change	−1.1 ± 0.9	0.0 ± 0.7	0.297
Diastolic blood pressure (mmHg)			
Baseline	77.4 ± 7.1	77.5 ± 7.5	
Change	−1.5 ± 0.6	−1.1 ± 0.5	0.582

Baseline values are mean ± SD; change values are LS mean ± SE unless otherwise indicated. Data are for the mITT population, week 56 end point. †Baseline values are geometric mean ± SD; percent change values are LS mean percent change (95% CI); P values are based on log-transformed values. \*Secondary end points that were significant according to the prespecified sequential closed testing procedure conducted to correct for multiple comparisons.

and similar to that previously reported for patients without type 2 diabetes (16–18).

Analyses were performed to examine the percentage of patients who had blood

pressure and pulse rate increases that met prespecified criteria (Supplementary Table 3). The proportions of patients who had increases meeting these criteria were

similar between groups but tended to be numerically higher in the NB treatment group. Importantly, ≤1% of patients in each treatment group had a systolic or diastolic blood pressure value corresponding to stage 2 hypertension (≥160 mmHg or ≥100 mmHg, respectively [21]) for two consecutive visits at any time during the study (or once if it occurred on the last visit).

Effects of NB on depression and mood was examined using the IDS-SR questionnaire and adverse event reports. At baseline, mean IDS-SR total scores in the safety analysis set were similar in the NB and placebo groups (8.1 and 7.7, respectively; scores can range from 0 to 84, with higher scores indicating more severe depression; <14 = nondepressed). At end point, the LS mean ± SE change from baseline in the total score was 0.53 ± 0.37 for NB and −1.41 ± 0.49 for placebo (P = 0.001). IDS-SR scores of ≥25 (indicating moderate depression) at any time during the study were observed in 6.2% of patients receiving placebo and 7.2% of NB-treated patients (NS). There were no differences between NB- and placebo-treated patients in incidence of adverse events of either depression or depressed mood during the study (Table 3).

**CONCLUSIONS**—Treatment of overweight or obese individuals with type 2 diabetes with NB resulted in a 5.0% reduction in body weight, compared with a reduction of 1.8% with placebo. Patients treated with NB were more than twice as likely to lose ≥5% of their initial body weight as patients treated with lifestyle intervention alone. The weight loss in the NB-treated group was evident as early as the first visit (week 4), and was maintained for the 56-week period, with no evidence of weight regain. This weight loss was accompanied by a significant reduction in HbA<sub>1c</sub> (placebo-corrected difference −0.5%) along with other favorable glycemic effects. Modest improvements in waist circumference, HDL-C, and triglycerides were also observed. The most frequently reported adverse event with NB was nausea.

Adipose tissue is an active endocrine and immune organ whose dysfunction contributes to metabolic diseases (e.g., diabetes) and increases cardiovascular disease (22). Thus, the objective of weight loss in overweight patients with metabolic disease is not solely cosmetic but also to improve the health of patients. In terms of

population, P values are nominal (i.e., not adjusted for multiple comparisons) and associated with exploratory analyses. C: Change in HbA<sub>1c</sub> from baseline to week 56 end point; placebo, n = 137; NB, n = 222. \*\*P < 0.01 vs. placebo; \*\*\*P < 0.001 vs. placebo; the mITT LOCF population consists of patients who had a baseline measurement and at least one postbaseline measurement while on study drug. The last observation on study drug was carried forward. The completer population comprised all randomized patients with a baseline measurement and a 56-week measurement while on the study drug.

**Table 3—Adverse events and adverse events leading to discontinuation**

	Placebo	NB
N	169	333
Patients reporting		
any adverse event	85.2	90.4
Nausea	7.1	42.3
Constipation	7.1	17.7
Vomiting	3.6	18.3
Diarrhea	9.5	15.6
Headache	8.9	13.8
Dizziness	5.3	11.7
Insomnia	5.3	11.1
Nasopharyngitis	13.6	8.4
Hypertension	4.1	9.9
Upper-respiratory tract infection	9.5	7.8
Hypoglycemia	7.1	7.5
Tremor	2.4	6.6
Dry mouth	3.0	6.3
Anxiety	1.2	5.4
Upper abdominal pain	1.8	5.1
Patients discontinued		
due to adverse event	15.4	29.4
Gastrointestinal disorders		
Nausea	0	9.6
Vomiting	0	3.0
Nervous system disorders		
Headache	0	1.8
Psychiatric disorders		
Depression	1.8	0.6
Metabolism and nutrition disorders		
Diabetes	1.2	0.3
Hyperglycemia	1.2	0

Safety analysis set. Any adverse event with >5% incidence in the NB group. Adverse events leading to discontinuation with incidence >1% of patients in any treatment group, listed by system organ class.

clinically meaningful weight loss, human studies suggest that 5% weight loss significantly improves glucose control and other cardiovascular parameters and results in reduced mortality in patients with diabetes (6–8). This magnitude of weight loss has also been associated with a delay in the progression to type 2 diabetes in patients with prediabetes (23–25).

The magnitude of weight loss (both absolute and placebo corrected) in the current study of overweight and obese individuals with type 2 diabetes was lower than that observed in NB trials in patients without type 2 diabetes. In the COR-I and COR-II studies (16,17), which used a lifestyle intervention similar to the current study, mean body weight changes

(mITT) were –6.1 and –6.4% with NB vs. –1.3 and –1.2% with placebo. In the COR-BMOD study (18), which included a more intensive, group-based lifestyle intervention, mean mITT body weight changes were –9.3% (NB) and –5.1% (placebo). The somewhat more modest weight loss in the current study is not surprising, as it is typical that less weight loss is observed in patients with diabetes taking OADs compared with patients without diabetes (26,27). While the reason for the attenuated magnitude of weight loss in patients with diabetes is not well understood, it may relate to differences in insulin resistance, adipose cell metabolism, concomitant medications for the treatment of diabetes, and glucose metabolism. Nevertheless, of study participants who completed 1 year of treatment, more than one-half of NB patients lost  $\geq 5\%$  of body weight and more than one-quarter achieved  $\geq 10\%$  weight loss.

Consistent with the observed improvement in body weight, NB-treated patients exhibited clinically and statistically significant improvement in HbA<sub>1c</sub> and generally beneficial changes in other markers related to glycemic control. The extent of reduction in HbA<sub>1c</sub> (–0.5% [–5.5 mmol/mol] vs. lifestyle intervention alone) compares well with the reported efficacy of several drugs currently used for treating diabetes (28). The magnitude of reduction was even greater in patients with higher baseline HbA<sub>1c</sub> values. This degree of improvement is meaningful in relation to long-term risk for microvascular complications and may contribute to a favorable environment with regard to macrovascular complications (29–32). The clinical impact of the HbA<sub>1c</sub> reduction with NB was demonstrated by the fact that patients on placebo were more likely to require an increased dose or number of OADs to maintain glycemic control (which lessened the apparent treatment difference with NB on both HbA<sub>1c</sub> and fasting glucose) (Supplementary Table 2).

NB treatment was also accompanied by improvements in a number of additional markers of cardiovascular risk, such as waist circumference, triglycerides, and HDL-C. In normoglycemic obese patients, weight loss with NB treatment has been demonstrated to be largely due to reductions in subcutaneous and visceral adipose tissue (33), the latter of which is associated with improvements in glycemia and lipidemia (34–36). Thus, treatment with NB in overweight/obese

patients with (and without) diabetes was associated with an improvement in the cardiovascular risk profile. However, bupropion has a documented pressor effect (37,38) that was associated with a numerically, but not statistically, greater reduction in blood pressure in the placebo group compared with NB. Although the hemodynamic effects of NB were demonstrated to be small and transient, the overall cardiovascular effects of NB could not be adequately assessed in the NB phase 3 clinical program because the patient population had an extremely low CV event rate (~0.2/100 patient-years). In order to assess the clinical effect of the NB-related improvements in weight and other cardiovascular risk factors, along with the documented pressor effect of bupropion (a transient elevation in pulse rate and blood pressure), a cardiovascular outcomes trial of NB is currently ongoing (clinical trial reg. no. NCT01601704, clinicaltrials.gov). This outcomes trial is designed to assess the incidence of major cardiovascular events in patients receiving NB compared with placebo.

The safety profile of NB in this population was generally consistent with the phase 3 trials conducted in patients without type 2 diabetes (16–18). Importantly, NB was not associated with an increase in hypoglycemia. The safety profile of the two agents used in combination is also similar to the postmarketing surveillance data available for the two agents used individually. The most common adverse event reported with NB was nausea, which led to withdrawal in ~10% of NB patients. Much of the patient withdrawal occurred within the first 4 weeks of treatment, during which patients started treatment at a lower initial dose and were required to escalate weekly. Patients unable to tolerate the study medication were discontinued from study drug treatment. The rate of nausea in NB-treated patients in the current study (42%) tended to be higher than previous phase 3 studies (29–34%) (16–18). This may be related to interaction with the background medications used by patients with diabetes in the current study, as NB-treated patients taking metformin (~75% of the population) reported a higher incidence of nausea than those not on metformin. This result was not unexpected, as gastrointestinal side effects are commonly associated with metformin treatment (39).

Limitations of this study include the exclusion of patients with type 2 diabetes who were receiving insulin therapy

(which can promote weight gain) or glucagon-like peptide-1 receptor agonist therapy (which is associated with mild weight loss). Thus, these results may not be applicable to all patients with type 2 diabetes. Furthermore, a relatively high dropout rate was observed in this study. Although the dropout rate was similar to the other NB phase 3 trials (16–18) and does not differ from other reported studies in obesity drug development (40), it should be considered when evaluating the treatment effect. Finally, the 1-year study duration, while common, is not adequate to fully assess the consequences of improved body weight on long-term outcomes.

The current epidemic of obesity and obesity-related diabetes continues to be of increasing concern. Unfortunately, a number of the drugs used to treat diabetes are associated with weight gain, and new approaches to prevention and treatment are needed. In this study of overweight/obese patients with type 2 diabetes, NB treatment, in conjunction with a standardized lifestyle intervention, resulted in a clinically meaningful improvement in body weight, which was associated with improvement in HbA<sub>1c</sub> and other cardiovascular risk factors. Thus, a combination drug such as NB has the potential to become a useful agent for the treatment of overweight/obese patients diagnosed with prediabetes or type 2 diabetes.

**Acknowledgments**—P.H., R.P., and H.B. received research funding from Orexigen Therapeutics, Inc. F.G. is a member of the Orexigen Therapeutics, Inc., Advisory Board and is a consultant for Orexigen Therapeutics, Inc. C.B. and P.K. are employees and shareholders of Orexigen Therapeutics, Inc. K.F. is a member of the Orexigen Therapeutics, Inc. Advisory Board and is a consultant for Orexigen Therapeutics, Inc. Brandon Walsh, PhD, and Amy Halseth, PhD, both employees of Orexigen Therapeutics, Inc., assisted in the writing of the manuscript. No other potential conflicts of interest relevant to this article were reported.

P.H. participated in data interpretation and writing and reviewing of the manuscript, approved the final version of the manuscript, participated in the study design, and was an investigator of the study. A.K.G., R.P., and H.B. participated in data interpretation and writing and reviewing of the manuscript, approved the final version of the manuscript, and were investigators of the study. F.G. participated in data interpretation and writing and reviewing of the manuscript, approved the final version of the manuscript, and

participated in study design. C.B. participated in data interpretation and writing and reviewing of the manuscript, approved the final version of the manuscript, and performed statistical analysis of data. P.K. participated in data interpretation and writing and reviewing of the manuscript and approved the final version of the manuscript. K.F. participated in data interpretation and writing and reviewing of the manuscript, approved the final version of the manuscript, and participated in study design. P.H. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Parts of this study were presented in abstract form at the 70th Scientific Sessions of the American Diabetes Association, Orlando, Florida, 25–29 June 2010; the 71st Scientific Sessions of the American Diabetes Association, San Diego, California, 24–28 June 2011; and the 11th International Congress on Obesity, Stockholm, Sweden, 11–15 July 2010.

The authors acknowledge contributions of Eduardo Dunayevich, MD, Amgen, to the design, implementation, and interpretation of the study.

## APPENDIX

### COR-Diabetes Study Group participating investigators

Laura Akright, Mira Baron, Harold Bays, Jolene Berg, Bruce Berwald, James Borders, Robert Buynak, Marc Capobianco, Harold Cathcart, Joseph Cleaver, Milissa Cooper, Martin Conway, Adnan Dahdul, Matthew Davis, Steven Duckor, John Ervin, David Fitz-Patrick, Larry Gilderman, Gopalakrishna Gollapudi, Alok Gupta, Forrest Hanke, Howard Harris, Lydie Hazan, Priscilla Hollander, Adesh Jain, Boris Kerzner, Lawrence Koehler, Ildiko Kovacs, Richard A. Krause, Diane Krieger, Judy Loper, Norman Lunde, Barbara McGuire, Richard Mills, Sunder Mudaliar, Troy Oxner, Naynesh Patel, Angel Pietri, Sanford N. Plevin, Raymond Plodkowski, George Raad, Eric Ross, Robert Schrieman, Sherwyn Schwartz, Stan Self, Stephan Sharp, Diane Smith, Timothy Smith, Philip Snell, Joseph Soufer, Louise Taber, Andrew Thieneman, Joseph Tibaldi, Paul Tung, James Vogt, Claire Waltman, Mervyn Weerasinghe, Daniel Weiss, Troy Williams, Kevin Wingert, Douglas Young, and Douglas Zmolek.

### References

1. Bays HE, González-Campoy JM, Bray GA, et al. Pathogenic potential of adipose tissue and metabolic consequences of adipocyte hypertrophy and increased visceral adiposity. *Expert Rev Cardiovasc Ther* 2008;6:343–368
2. Flegal KM, Carroll MD, Kit BK, Ogden CL. Prevalence of obesity and trends in the distribution of body mass index among US adults, 1999–2010. *JAMA* 2012;307:491–497
3. Centers for Disease Control and Prevention (CDC). Prevalence of overweight and obesity among adults with diagnosed diabetes—United States, 1988–1994 and 1999–2002. *MMWR Morb Mortal Wkly Rep* 2004;53:1066–1068
4. Niswender K. Diabetes and obesity: therapeutic targeting and risk reduction - a complex interplay. *Diabetes Obes Metab* 2010;12:267–287
5. Hollander PA, Kushner P. Type 2 diabetes comorbidities and treatment challenges: rationale for DPP-4 inhibitors. *Postgrad Med* 2010;122:71–80
6. Goldstein DJ. Beneficial health effects of modest weight loss. *Int J Obes Relat Metab Disord* 1992;16:397–415
7. Wing RR, Lang W, Wadden TA, et al.; Look AHEAD Research Group. Benefits of modest weight loss in improving cardiovascular risk factors in overweight and obese individuals with type 2 diabetes. *Diabetes Care* 2011;34:1481–1486
8. Williamson DF, Thompson TJ, Thun M, Flanders D, Pamuk E, Byers T. Intentional weight loss and mortality among overweight individuals with diabetes. *Diabetes Care* 2000;23:1499–1504
9. Ascher JA, Cole JO, Colin JN, et al. Bupropion: a review of its mechanism of antidepressant activity. *J Clin Psychiatry* 1995;56:395–401
10. Foley KF, DeSanty KP, Kast RE. Bupropion: pharmacology and therapeutic applications. *Expert Rev Neurother* 2006;6:1249–1265
11. Anderson JW, Greenway FL, Fujioka K, Gadde KM, McKenney J, O'Neil PM. Bupropion SR enhances weight loss: a 48-week double-blind, placebo-controlled trial. *Obes Res* 2002;10:633–641
12. Bouza C, Angeles M, Muñoz A, Amate JM. Efficacy and safety of naltrexone and acamprosate in the treatment of alcohol dependence: a systematic review. *Addiction* 2004;99:811–828
13. Greenway FL, Whitehouse MJ, Guttadauria M, et al. Rational design of a combination medication for the treatment of obesity. *Obesity (Silver Spring)* 2009;17:30–39
14. Morton GJ, Cummings DE, Baskin DG, Barsh GS, Schwartz MW. Central nervous system control of food intake and body weight. *Nature* 2006;443:289–295
15. Sinnayah P, Wallingford NM, Evans AE, Cowley MA. Bupropion and naltrexone interact synergistically to decrease food

- intake in mice (Abstract). *Obesity* (Silver Spring) 2007;15:A179
16. Apovian CM, Aronne LJ, Rubino DM, et al.; COR-II Study Group. A randomized, phase 3 trial of naltrexone SR/bupropion SR on weight and obesity-related risk factors (COR-II). *Obesity* (Silver Spring) 2013;21:935–943
  17. Greenway FL, Fujioka K, Plodkowski RA, et al.; COR-I Study Group. Effect of naltrexone plus bupropion on weight loss in overweight and obese adults (COR-I): a multicentre, randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet* 2010;376:595–605
  18. Wadden TA, Foreyt JP, Foster GD, et al. Weight loss with naltrexone SR/bupropion SR combination therapy as an adjunct to behavior modification: the COR-BMOD trial. *Obesity* (Silver Spring) 2011;19:110–120
  19. World Medical Association General Assembly. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *J Int Bioethique* 2004;15:124–129
  20. Scheen AJ, Finer N, Hollander P, Jensen MD, Van Gaal LF; RIO-Diabetes Study Group. Efficacy and tolerability of rimonabant in overweight or obese patients with type 2 diabetes: a randomised controlled study. *Lancet* 2006;368:1660–1672
  21. Chobanian AV, Bakris GL, Black HR, et al.; Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. National Heart, Lung, and Blood Institute; National High Blood Pressure Education Program Coordinating Committee. Seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Hypertension* 2003;42:1206–1252
  22. Bays HE. Adiposopathy is “sick fat” a cardiovascular disease? *J Am Coll Cardiol* 2011;57:2461–2473
  23. Knowler WC, Barrett-Connor E, Fowler SE, et al.; Diabetes Prevention Program Research Group. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002;346:393–403
  24. Knowler WC, Fowler SE, Hamman RF, et al.; Diabetes Prevention Program Research Group. 10-year follow-up of diabetes incidence and weight loss in the Diabetes Prevention Program Outcomes Study. *Lancet* 2009;374:1677–1686
  25. Lindström J, Louheranta A, Mannelin M, et al.; Finnish Diabetes Prevention Study Group. The Finnish Diabetes Prevention Study (DPS): Lifestyle intervention and 3-year results on diet and physical activity. *Diabetes Care* 2003;26:3230–3236
  26. Hollander PA, Elbein SC, Hirsch IB, et al. Role of orlistat in the treatment of obese patients with type 2 diabetes. A 1-year randomized double-blind study. *Diabetes Care* 1998;21:1288–1294
  27. Torgerson JS, Hauptman J, Boldrin MN, Sjöström L. XENical in the prevention of diabetes in obese subjects (XENDOS) study: a randomized study of orlistat as an adjunct to lifestyle changes for the prevention of type 2 diabetes in obese patients. *Diabetes Care* 2004;27:155–161
  28. Bennett WL, Maruthur NM, Singh S, et al. Comparative effectiveness and safety of medications for type 2 diabetes: an update including new drugs and 2-drug combinations. *Ann Intern Med* 2011;154:602–613
  29. UK Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet* 1998;352:837–853
  30. Turnbull FM, Abraira C, Anderson RJ, et al.; Control Group. Intensive glucose control and macrovascular outcomes in type 2 diabetes. *Diabetologia* 2009;52:2288–2298
  31. Ray KK, Seshasai SR, Wijesuriya S, et al. Effect of intensive control of glucose on cardiovascular outcomes and death in patients with diabetes mellitus: a meta-analysis of randomised controlled trials. *Lancet* 2009;373:1765–1772
  32. Mannucci E, Monami M, Lamanna C, Gori F, Marchionni N. Prevention of cardiovascular disease through glycemic control in type 2 diabetes: a meta-analysis of randomized clinical trials. *Nutr Metab Cardiovasc Dis* 2009;19:604–612
  33. Smith SR, Fujioka K, Gupta AK, et al. Combination therapy with naltrexone and bupropion for obesity reduces total and visceral adiposity: effects of the naltrexone/bupropion combination on body composition. *Diabetes Obes Metab*. 12 March 2013 [Epub ahead of print]
  34. Ross R, Dagnone D, Jones PJ, et al. Reduction in obesity and related comorbid conditions after diet-induced weight loss or exercise-induced weight loss in men. A randomized, controlled trial. *Ann Intern Med* 2000;133:92–103
  35. Katsoulis K, Blaudeau TE, Roy JP, Hunter GR. Diet-induced changes in intra-abdominal adipose tissue and CVD risk in American women. *Obesity* (Silver Spring) 2009;17:2169–2175
  36. Després JP, Lemieux I. Abdominal obesity and metabolic syndrome. *Nature* 2006;444:881–887
  37. Thase ME, Haight BR, Johnson MC, et al. A randomized, double-blind, placebo-controlled study of the effect of sustained-release bupropion on blood pressure in individuals with mild untreated hypertension. *J Clin Psychopharmacol* 2008;28:302–307
  38. Settle EC, Stahl SM, Batey SR, Johnston JA, Ascher JA. Safety profile of sustained-release bupropion in depression: results of three clinical trials. *Clin Ther* 1999;21:454–463
  39. Glucophage (metformin hydrochloride) tablets [package insert]. Bristol-Meyers Squibb, 2009
  40. Smith SR, Weissman NJ, Anderson CM, et al.; Behavioral Modification and Lorcaserin for Overweight and Obesity Management (BLOOM) Study Group. Multicenter, placebo-controlled trial of lorcaserin for weight management. *N Engl J Med* 2010;363:245–256