

Vitamin C Further Improves the Protective Effect of Glucagon-Like Peptide-1 on Acute Hypoglycemia-Induced Oxidative Stress, Inflammation, and Endothelial Dysfunction in Type 1 Diabetes

ANTONIO CERIELLO, MD^{1,2}
ANNA NOVIALLS, MD^{1,2}
EMILIO ORTEGA, MD^{1,2}
SILVIA CANIVELL, MD^{1,2}
LUCIA LA SALA, PHD^{1,2}

GEMMA PUJADAS, PHD^{1,2}
LOREDANA BUCCIARELLI, MD³
MAURIZIO RONDINELLI, MD³
STEFANO GENOVESE, MD³

OBJECTIVE—To test the hypothesis that acute hypoglycemia induces endothelial dysfunction and inflammation through the generation of an oxidative stress. Moreover, to test if the antioxidant vitamin C can further improve the protective effects of glucagon-like peptide 1 (GLP-1) on endothelial dysfunction and inflammation during hypoglycemia in type 1 diabetes.

RESEARCH DESIGN AND METHODS—A total of 20 type 1 diabetic patients underwent four experiments: a period of 2 h of acute hypoglycemia with or without infusion of GLP-1 or vitamin C or both. At baseline, after 1 and 2 h, glycemia, plasma nitrotyrosine, plasma 8-iso prostaglandin F_{2a} (PGF_{2a}), soluble intracellular adhesion molecule-1a (sICAM-1a), interleukin-6 (IL-6), and flow-mediated vasodilation were measured. At 2 h of hypoglycemia, flow-mediated vasodilation significantly decreased, while sICAM-1, 8-iso-PGF_{2a}, nitrotyrosine, and IL-6 significantly increased. The simultaneous infusion of GLP-1 or vitamin C significantly attenuated all of these phenomena. Vitamin C was more effective. When GLP-1 and vitamin C were infused simultaneously, the deleterious effect of hypoglycemia was almost completely counterbalanced.

RESULTS—At 2 h of hypoglycemia, flow-mediated vasodilation significantly decreased, while sICAM-1, 8-iso-PGF_{2a}, nitrotyrosine, and IL-6 significantly increased. The simultaneous infusion of GLP-1 or vitamin C significantly attenuated all of these phenomena. Vitamin C was more effective. When GLP-1 and vitamin C were infused simultaneously, the deleterious effect of hypoglycemia was almost completely counterbalanced.

CONCLUSIONS—This study shows that vitamin C infusion, during induced acute hypoglycemia, reduces the generation of oxidative stress and inflammation, improving endothelial dysfunction, in type 1 diabetes. Furthermore, the data support a protective effect of GLP-1 during acute hypoglycemia, but also suggest the presence of an endothelial resistance to the action of GLP-1, reasonably mediated by oxidative stress.

Diabetes Care 36:4104–4108, 2013

Recent evidence suggests that hypoglycemia can be considered a new risk factor in favoring diabetes vascular complications (1). It has been reported that hypoglycemia produces endothelial dysfunction and inflammation (2,3), which are well-recognized pathogenic factors for vascular disease, particularly in diabetes

(4). Oxidative stress is considered the key player in the pathogenesis of diabetes complications (5), and it has been suggested that hypoglycemia produces endothelial dysfunction and inflammation through oxidative stress generation both in vitro and in humans (2,3). However, clear evidence in humans is still lacking (6): particularly, what is still missed is the evidence that by using an antioxidant, the elevation of the markers of oxidative stress induced by hypoglycemia can be reversed.

Recently, a possible beneficial effect of glucagon-like peptide 1 (GLP-1) analogs in the management of type 1 diabetes has been suggested (7,8). GLP-1 and its analogs, in addition to their insulin-tropic action in alleviating hyperglycemia, have beneficial effects in protecting progressive impairment of pancreatic β -cell function, preservation of β -cell mass, and suppression of glucagon secretion, gastric emptying, and appetite, all characteristics that could be beneficial for the management of type 1 diabetes (7,8).

Apart from the well-documented incretin effect of GLP-1, its role in the cardiovascular system also arouses interest. GLP-1 effects on the cardiovascular system may include a direct action on the endothelium, where the presence of specific receptors for GLP-1 has been demonstrated (9). Consistently, GLP-1 has been demonstrated to improve endothelial function in diabetes (10,11), possibly increasing the antioxidant defenses of the endothelium (12) and decreasing oxidative stress generation (11). Recently, we have reported that in type 1 diabetes, during induced-hypoglycemia, GLP-1 can partially protect endothelial function and partially decrease the appearance of inflammation, reducing, concomitantly, the level of several markers of oxidative stress (13). However, it is worthy of interest that in type 2 diabetes, hyperglycemia induces an endothelial

From ¹Diabetes and Endocrinology, Institut d'Investigacions Biomèdiques August Pi i Sunyer, Barcelona, Spain; ²Centro de Investigación Biomédica en Red de Diabetes y Enfermedades Metabólicas Asociadas, Hospital Clínic Barcelona, Barcelona, Spain; and the ³Department of Cardiovascular and Metabolic Diseases, Istituto di Ricovero e Cura a Carattere Scientifico Gruppo Multimedica, Sesto San Giovanni, Italy.

Corresponding author: Antonio Ceriello, aceriell@clinic.ub.es.

Received 29 March 2013 and accepted 24 July 2013.

DOI: 10.2337/dc13-0750

© 2013 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

resistance to the action of GLP-1, oxidative stress being the mediator of such a phenomenon (11).

The aim of this study was to test, in patients with type 1 diabetes, whether 1) the concomitant infusion of an antioxidant, vitamin C, can protect endothelial function and reduce the generation of oxidative stress and inflammation during acute induced-hypoglycemia and 2) the effects of GLP-1 and vitamin C on both endothelial dysfunction and oxidative stress induced by acute hypoglycemia were additive.

RESEARCH DESIGN AND METHODS

Subjects

Twenty persons with type 1 diabetes were studied (Table 1). They had normal bedside tests of autonomic function (14) and did not have hypoglycemia unawareness based on the methods of Gold et al. (15) or major macro- or microcomplications of diabetes. They were treated with multiple daily insulin injections. All subjects were non-smokers and had a normal blood count; had normal plasma lipids, plasma electrolytes, and liver and renal function; were normotensive; and were not taking medications, which can influence neuroendocrine responses to hypoglycemia or anti-inflammatory drugs.

Studies were approved by one of the ethical committees of our institutions, and all participants gave written informed consent.

Table 1—Baseline characteristics of the type 1 diabetic patients

Sex	13 male, 7 female
Age (years)	23.2 ± 2.3
BMI (kg/m ²)	22.8 ± 2.4
HbA _{1c} (%)	8.1 ± 0.5
HbA _{1c} (mmol/mol)	65 ± 3.2
Resting diastolic blood pressure (mmHg)	77.3 ± 1.2
Resting systolic blood pressure (mmHg)	115.2 ± 1.4
Total cholesterol (mmol/L)	4.32 ± 0.3
Triglycerides (mmol/L)	1.1 ± 0.4
HDL-C (mmol/L)	1.4 ± 0.2
LDL-C (mmol/L)	2.0 ± 0.2
FMD (%)	6.5 ± 0.4
8-iso-PGF _{2a} (pg/mL)	65.2 ± 4.2
Nitrotyrosine (μmol/L)	0.74 ± 0.04
sICAM-1a (ng/mL)	174.5 ± 13.1
IL-6 (pg/mL)	225.15 ± 11.1

Data are expressed as mean ± SEM. HDL-C, HDL cholesterol; LDL-C, LDL cholesterol.

Patients were asked to avoid any exercise and consume their usual weight-maintaining diet for 3 days before each experiment. All participants were asked to perform intensive home blood glucose monitoring and to avoid hypoglycemia for at least 5 days before a study. On the day prior to a study, intermediate or long-acting insulin was discontinued and replaced by injections of regular insulin before breakfast and lunch. Each subject was admitted to the Research Center the evening before an experiment. At this time, two intravenous cannulas were inserted under 1% lidocaine local anesthesia. One cannula was placed to be used for blood drawing. The other cannula was placed in the contralateral arm for infusions. All subjects received an evening meal, with a continuous low-dose infusion of insulin to normalize plasma glucose. The insulin infusion was adjusted overnight to maintain blood glucose between 4.4 and 7.2 mmol/L.

Acute hypoglycemia experiments

All of the subjects were studied after an overnight, 10-h fast. Four different experiments were planned for each subject in a randomized order: a period of 2 h of hypoglycemia with or without infusion of GLP-1 [Synthetic GLP-1 (7–36)amide; PolyPeptide Laboratories, Wolfenbuttel, Germany] (0.4 pmol · kg⁻¹ · min⁻¹ [16]), vitamin C (30 mg/min [16]), or both. Each subject underwent each experiment with at least 1 week between each.

At time zero, a primed constant (9.0 pmol · kg⁻¹ · min⁻¹) infusion of insulin (Actrapid; NovoNordisk, Copenhagen, Denmark) was started and continued until 120 min. The rate of fall of glucose was controlled (~0.08 mmol/min), and the glucose nadir (2.9 mmol/L) was achieved using a modification of the glucose clamp technique (16). During the clamp period, plasma glucose was measured every 5 min, and a 20% dextrose infusion was adjusted so that plasma glucose levels were held constant at 2.9 ± 0.1 mmol/L (16). Potassium chloride (20 mmol/L) was infused during the clamp to reduce insulin-induced hypokalemia. The experiment was repeated with the infusion of GLP-1, vitamin C, or both.

At baseline and after 1 and 2 h, blood samples were withdrawn for biochemical assays: glycemia, plasma nitrotyrosine (a stable end product of peroxynitrite oxidation) and plasma 8-iso prostaglandin F_{2a} (PGF_{2a}; a biomarker of lipid peroxidation), both markers of oxidative stress, and intercellular adhesion molecule 1 (ICAM-1) and

interleukin-6 (IL-6), both markers of inflammation. Endothelial function was measured by flow-mediated dilation (FMD).

Biochemical and clinical measurements

Cholesterol, triglycerides, HDL cholesterol, LDL cholesterol, and plasma nitrotyrosine were measured according to Ceriello et al. (17). Plasma glucose was measured by the glucose-oxidase method, HbA_{1c} by high-performance liquid chromatography, and insulin by microparticle enzyme immunoassay (Abbott Laboratories, Wiesbaden, Germany). Plasma 8-iso-PGF_{2a} (Cayman Chemical, Ann Arbor, MI), soluble ICAM-1 (sICAM-1; British Bio-technology, Abingdon, Oxon, U.K.), and IL-6 (R&D Systems, Minneapolis, MN) were determined with commercially available kits.

FMD. Endothelial function at macrovascular level was evaluated by measuring FMD of the brachial artery (12,16). The examination was carried out in a temperature- and light-controlled room on subjects who were lying comfortably flat on a couch. Brachial arteries in this study were imaged with a standard ultrasound system (VIVID 7 ECHO machine; GE Vingmed System V) connected with a 12-MHz linear transducer probe. The ultrasound system was connected to a personal computer equipped with a frame grabber and artificial neural network wall detection software (vessel image analysis). Brachial artery FMD was determined using protocol similar to published studies (12–16).

At the end of the study each day, 250 μg of sublingual glyceryl trinitrate was administered in order to assess endothelium-independent vasodilatation.

The intraobserver variability for repeated measurements of resting arterial diameter was 0.02 ± 0.02 mm.

Statistical analysis

The sample size was selected according to previous studies (12,16,18).

Data are expressed as means ± SE. The Kolmogorov-Smirnov algorithm was used to determine whether each variable had a normal distribution. Comparisons of baseline data among the groups were performed using unpaired Student *t* test or Mann-Whitney *U* test, where indicated. The changes in variables during the tests were assessed by two-way ANOVA with repeated-measures or Kolmogorov-Smirnov test, where indicated. If differences

reached statistical significance, post hoc analyses with two-tailed paired *t* test or Wilcoxon signed-rank test for paired comparisons were used to assess differences at individual time periods in the study. Statistical significance was defined as $P < 0.05$.

RESULTS—After 2 h of acute hypoglycemia, FMD significantly decreased, while sICAM-1, 8-iso-PGF2a, nitrotyrosine,

and IL-6 significantly increased, compared with basal values, (Fig. 1). When hypoglycemia was accompanied by the simultaneous infusion of GLP-1, all of these phenomena were significantly attenuated: FMD decreased less, while sICAM-1, 8-iso-PGF2a, nitrotyrosine, and IL-6 were less increased (Fig. 1). Vitamin C infusion had an even better effect than GLP-1 (Fig. 1). However, when GLP-1 and vitamin C were simultaneously

infused, the deleterious effect of hypoglycemia on FMD, sICAM-1, 8-iso-PGF2a, nitrotyrosine, and IL-6 was almost completely counterbalanced (Fig. 1). Endothelial-independent vasodilatation was not affected in any of the experiments. The amount of insulin infused during the experiments was similar (area under the curve $75,200 \pm 350$ vs. $75,320 \pm 370$ vs. $75,150 \pm 330$ vs. $75,300 \pm 320$ pmol).




Figure 1—Glycemia, FMD, IL-6, sICAM-1, nitrotyrosine, and 8-iso-PGF2a changes during hypoglycemia (white triangle), hypoglycemia plus GLP-1 (black circle), hypoglycemia plus vitamin C (white circle), and hypoglycemia plus GLP-1 and vitamin C (black square). Data are mean \pm SEM. * $P < 0.01$ vs. basal, £ $P < 0.05$ vs. hypoglycemia plus GLP-1 and vitamin C, § $P < 0.05$ vs. hypoglycemia plus GLP-1, # $P < 0.05$ vs. hypoglycemia plus vitamin C.

CONCLUSIONS—This study confirms that acute hypoglycemia induces endothelial dysfunction and inflammation in people with type 1 diabetes (13). However, in our opinion, it adds two important findings to our previous report showing that GLP-1 can reduce these phenomena (13).

It has been suggested that hypoglycemia can produce an oxidative stress (3,13,19), which in turn can favor the appearance of an endothelial dysfunction and inflammation (3–13,19). However, this hypothesis has been, until now, just linked to the evidence that several markers of oxidative stress increase during hypoglycemia (3–13,19), and, to date, it has not been confirmed showing that an antioxidant can counterbalance this phenomenon or, better, that the antioxidant given during hypoglycemia, reducing oxidative stress, can simultaneously reduce endothelial dysfunction and inflammation. This is what, for the first time, our study addresses. The infusion of vitamin C, during acute hypoglycemia, produced an expected improvement in oxidative stress, which was accompanied by the simultaneous improvement of both endothelial function and inflammation. In our opinion, this means a causal role of oxidative stress in favoring the appearance of endothelial dysfunction and inflammation during hypoglycemia.

However, another possibility is that acute hypoglycemia causes activation of the sympathetic system (hence the increase in 8-iso-PGF_{2a}), which causes vasoconstriction (hence the decrease in FMD). The endothelial cells try to respond to this by increasing their nitric oxide production (hence the increase in nitrotyrosine) but cannot overcome the autonomic overactivity. Sympathetic activity is known to influence inflammatory signals (20), hence the increase in IL-6 levels. Therefore, the observed acute hypoglycemia-induced alterations could simply be due to a physiological reflex response, not necessarily a pathological oxidative stress-induced endothelial dysfunction.

The second finding is related to the additive effects of vitamin C and GLP-1.

In this study, a protective effect of GLP-1 on hypoglycemia-induced endothelial dysfunction, inflammation, and oxidative stress is confirmed. Studies are accumulating showing that GLP-1 and its analogs used in clinical practice have an antioxidant activity (12,13). Therefore, it is reasonable that GLP-1 should, by reducing oxidative-stress generation,

improve endothelial dysfunction and inflammation generated by hypoglycemia. However, it is worthy of interest that vitamin C was more efficacious than GLP-1 in counterbalancing the effects of hypoglycemia and that only when both vitamin C and GLP-1 were infused simultaneously were the effects of hypoglycemia abolished completely. These data, altogether, support the hypothesis that the endothelium, as in hyperglycemia (11,21,22), became less sensitive to GLP-1 in hypoglycemia, more than GLP-1 itself loses its activity, and that oxidative stress might be the mediator of such phenomenon.

A possible direct influence of insulin on our results cannot be excluded, particularly because it has recently been reported that GLP-1 enhances the vasodilator effect of insulin (23). However, in our experiments, insulin infusion was accompanied by a decrease of the endothelial function, and the amount of insulin infused was the same during all of the experiments, suggesting that the role of insulin in our results could be not of particular relevance. In the same article by Tesaro et al. (23), GLP-1 did not enhance the effect of vitamin C on endothelial dysfunction. However, our experimental conditions are very different. We studied 20 patients with type 1 diabetes, while the study of Tesaro et al. (23) was focused on only 5 patients with metabolic syndrome (23). Moreover, as underlined by the same authors, no control study with vitamin C alone was performed.

Two mechanisms have been suggested to explain this resistance to the GLP-1 action in diabetes: the activation of protein kinase C β (PKC β), induced by hyperglycemia, able to reduce the expression of the GLP-1 receptors (24), and the generation by hyperglycemia of an oxidative stress (11). Nevertheless, the two proposed mechanisms, PKC β activation with reduction of the expression of GLP-1 receptors and oxidative-stress generation, could be convincingly correlated, because it is well-recognized that PKC β is activated by the free radicals (25). Even at this stage, it is just a hypothesis; as both hyperglycemia and hypoglycemia work through the same pathways and mainly generating an oxidative stress (1,26), it seems reasonable that the same mechanisms involved in the appearance of GLP-1 resistance in hyperglycemia might also be involved in hypoglycemia.

The risk of a cardiovascular disease in type 1 diabetes, even still partly neglected, is very high (27).

The role of hyperglycemia in favoring cardiovascular disease in type 1 diabetes seems to be relevant; even many other classical and less classical risk factors seem to be also involved (26). However, particularly the role of the oxidative stress seems to be very relevant in the pathogenesis of these complications in type 1 diabetes (26), and our finding suggests that hypoglycemia, a frequent event in the life of type 1 diabetes, which is emerging as a cardiovascular risk factor (1), can also produce an oxidative stress.

In conclusion, our study shows that hypoglycemia produces endothelial dysfunction and inflammation through oxidative stress and that an antioxidant, such as vitamin C, can partly prevent this effect. While the chronic use of vitamin C cannot be the solution (28), recent development in the antioxidant treatment of diabetes-related complications might provide a future perspective also in preventing the deleterious effects of hypoglycemia on the cardiovascular system (29).

A possible limitation of this study is that type 1 diabetic subjects enrolled in the current study were free of micro- or macrovascular complications and may not represent the entire spectrum of type 1 diabetes.

Acknowledgments—No potential conflicts of interest relevant to this article were reported.

A.C. researched data, contributed to discussion, wrote the manuscript, and reviewed and edited the manuscript. A.N., E.O., and L.B. researched data, contributed to discussion, and reviewed and edited the manuscript. S.C., M.R., and S.G. contributed to discussion and reviewed and edited the manuscript. L.L.S. and G.P. researched data and contributed to discussion. A.C. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

References

1. Wright RJ, Frier BM. Vascular disease and diabetes: is hypoglycaemia an aggravating factor? *Diabetes Metab Res Rev* 2008;24: 353–363
2. Wang J, Alexanian A, Ying R, et al. Acute exposure to low glucose rapidly induces endothelial dysfunction and mitochondrial oxidative stress: role for AMP kinase. *Arterioscler Thromb Vasc Biol* 2012;32: 712–720
3. Razavi Nematollahi L, Kitabchi AE, Stentz FB, et al. Proinflammatory cytokines in response to insulin-induced hypoglycemic stress in healthy subjects [published

- correction appears in *Metabolism* 2009; 58:1046]. *Metabolism* 2009;58:443–448
4. Nandish S, Wyatt J, Bailon O, Smith M, Oliveros R, Chilton R. Implementing cardiovascular risk reduction in patients with cardiovascular disease and diabetes mellitus. *Am J Cardiol* 2011;108(Suppl.): 42B–51B
 5. Giacco F, Brownlee M. Oxidative stress and diabetic complications. *Circ Res* 2010;107:1058–1070
 6. Snell-Bergeon JK, Wadwa RP. Hypoglycemia, diabetes, and cardiovascular disease. *Diabetes Technol Ther* 2012;14 (Suppl. 1):S51–S58
 7. Issa CM, Azar ST. Possible role of GLP-1 and its agonists in the treatment of type 1 diabetes mellitus. *Curr Diab Rep* 2012;12: 560–567
 8. George P, McCrimmon RJ. Potential role of non-insulin adjunct therapy in Type 1 diabetes. *Diabet Med* 2013;30:179–188
 9. Mudaliar S, Henry RR. Effects of incretin hormones on beta-cell mass and function, body weight, and hepatic and myocardial function. *Am J Med* 2010;123(Suppl.): S19–S27
 10. Nyström T, Gutniak MK, Zhang Q, et al. Effects of glucagon-like peptide-1 on endothelial function in type 2 diabetes patients with stable coronary artery disease. *Am J Physiol Endocrinol Metab* 2004; 287:E1209–E1215
 11. Ceriello A, Esposito K, Testa R, Bonfigli AR, Marra M, Giugliano D. The possible protective role of glucagon-like peptide 1 on endothelium during the meal and evidence for an “endothelial resistance” to glucagon-like peptide 1 in diabetes. *Diabetes Care* 2011;34:697–702
 12. Oeseburg H, de Boer RA, Buikema H, van der Harst P, van Gilst WH, Silljé HH. Glucagon-like peptide 1 prevents reactive oxygen species-induced endothelial cell senescence through the activation of protein kinase A. *Arterioscler Thromb Vasc Biol* 2010;30:1407–1414
 13. Ceriello A, Novials A, Ortega E, et al. Glucagon-like peptide 1 reduces endothelial dysfunction, inflammation and oxidative stress induced by both hyperglycemia and hypoglycemia in type 1 diabetes. *Diabetes Care* 2013;36:2346–2350
 14. Ewing DJ, Martyn CN, Young RJ, Clarke BF. The value of cardiovascular autonomic function tests: 10 years experience in diabetes. *Diabetes Care* 1985;8:491–498
 15. Gold AE, MacLeod KM, Frier BM. Frequency of severe hypoglycemia in patients with type 1 diabetes with impaired awareness of hypoglycemia. *Diabetes Care* 1994;17:697–703
 16. Ceriello A, Novials A, Ortega E, et al. Evidence that hyperglycemia after recovery from hypoglycemia worsens endothelial function and increases oxidative stress and inflammation in healthy control subjects and subjects with type 1 diabetes. *Diabetes* 2012;61:2993–2997
 17. Ceriello A, Mercuri F, Quagliaro L, et al. Detection of nitrotyrosine in the diabetic plasma: evidence of oxidative stress. *Diabetologia* 2001;44:834–838
 18. Gogitidze Joy N, Hedrington MS, Briscoe VJ, Tate DB, Ertl AC, Davis SN. Effects of acute hypoglycemia on inflammatory and pro-atherothrombotic biomarkers in individuals with type 1 diabetes and healthy individuals. *Diabetes Care* 2010;33: 1529–1535
 19. Yousefzade G, Nakhaee A. Insulin-induced hypoglycemia and stress oxidative state in healthy people. *Acta Diabetol* 2012;49 (Suppl. 1):S81–S85
 20. Elenkov IJ, Iezzoni DG, Daly A, Harris AG, Chrousos GP. Cytokine dysregulation, inflammation and well-being. *Neuroimmunomodulation* 2005;12:255–269
 21. Meier JJ, Nauck MA. Is the diminished incretin effect in type 2 diabetes just an epi-phenomenon of impaired beta-cell function? *Diabetes* 2010;59:1117–1125
 22. Green CJ, Henriksen TI, Pedersen BK, Solomon TP. Glucagon like peptide-1-induced glucose metabolism in differentiated human muscle satellite cells is attenuated by hyperglycemia. *PLoS ONE* 2012;7:e44284
 23. Tesouro M, Schinzari F, Adamo A, et al. Effects of GLP-1 on forearm vasodilator function and glucose disposal during hyperinsulinemia in the metabolic syndrome. *Diabetes Care* 2013;36:683–689
 24. Mima A, Hiraoka-Yamamoto J, Li Q, et al. Protective effects of GLP-1 on glomerular endothelium and its inhibition by PKC β activation in diabetes. *Diabetes* 2012;61: 2967–2979
 25. Quagliaro L, Piconi L, Assaloni R, Martinelli L, Motz E, Ceriello A. Intermittent high glucose enhances apoptosis related to oxidative stress in human umbilical vein endothelial cells: the role of protein kinase C and NAD(PH)-oxidase activation. *Diabetes* 2003;52:2795–2804
 26. Ceriello A. Hyperglycaemia and the vessel wall: the pathophysiological aspects on the atherosclerotic burden in patients with diabetes. *Eur J Cardiovasc Prev Rehabil* 2010;17(Suppl. 1):S15–S19
 27. Orchard TJ, Costacou T. When are type 1 diabetic patients at risk for cardiovascular disease? *Curr Diab Rep* 2010;10:48–54
 28. Frei B, Birlouez-Aragon I, Lykkesfeldt J. Authors’ perspective: What is the optimum intake of vitamin C in humans? *Crit Rev Food Sci Nutr* 2012;52:815–829
 29. Miyata T, Suzuki N, van Ypersele de Strihou C. Diabetic nephropathy: are there new and potentially promising therapies targeting oxygen biology? *Kidney Int.* 13 March 2013 [Epub ahead of print]