

## Survey and genetic characterization of wastewater in Tunisia for *Cryptosporidium* spp., *Giardia duodenalis*, *Enterocytozoon bieneusi*, *Cyclospora cayetanensis* and *Eimeria* spp.

Layla Ben Ayed, Wenli Yang, Giovanni Widmer, Vitaliano Cama, Ynes Ortega and Lihua Xiao

### ABSTRACT

The microbial diversity of wastewater used for irrigation and fertilization was assessed using specific polymerase chain reaction (PCR) assays to detect and genotype several pathogenic protists including *Cryptosporidium* spp., *Giardia duodenalis*, *Cyclospora* spp., *Eimeria* spp. and *Enterocytozoon bieneusi*. A total of 220 wastewater samples (110 raw, 110 treated) and 12 sludge samples were collected from 2005 to 2008 from 18 treatment plants located throughout Tunisia. Except for *Cyclospora*, which was detected only once, *E. bieneusi* (61%), *G. duodenalis* (28%), *Cryptosporidium* spp. (27%) and *Eimeria* spp. (45%) were frequently observed in wastewater and sludge. Sequencing of PCR products showed that *C. hominis*, *C. andersoni*, *G. duodenalis* sub-assembly A-II and *E. bieneusi* genotypes D and IV were the most prevalent. An analysis of the distribution of 209 internal transcribed spacer sequences of *E. bieneusi* originating from wastewater at the 18 treatment plants showed a similar genetic diversity, regardless of the geographical location. The identification of these parasite species and genotypes and of host-specific *Eimeria* species indicates that the microbial quality of wastewater was impacted by humans, livestock and rodents. Given the public health risks that some of these parasites represent, guidelines on wastewater usage are needed to minimize human exposure to these pathogens.

**Key words** | *Cryptosporidium*, *Enterocytozoon bieneusi*, genotypes, *Giardia duodenalis*, Tunisia, wastewater

#### Layla Ben Ayed

Laboratoire Sciences et Techniques de l'Eau, Institut National Agronomique de Tunisie, 43, Avenue Charles Nicolle 1082, Tunis, Tunisie

#### Wenli Yang

#### Vitaliano Cama

Lihua Xiao (corresponding author)

Division of Foodborne, Waterborne, and Environmental Diseases, National Center for Emerging and Zoonotic Infectious Diseases, Centers for Disease Control and Prevention, Atlanta, Georgia 30333, USA

E-mail: lxiao@cdc.gov

#### Giovanni Widmer

Division of Infectious Diseases, Tufts Cummings School of Veterinary Medicine, North Grafton, MA 01536, USA

#### Ynes Ortega

Center for Food Safety, University of Georgia, Griffin, GA 30223, USA

### INTRODUCTION

As a result of increased human consumption and irregular or diminishing rainfall, Tunisia is experiencing a growing shortage of water. To address this problem, the government is encouraging the reuse of treated wastewater (TWW). Water reuse for agriculture is increasing, with almost 30% of the total volume of TWW used to irrigate about 9,300 hectares (ONAS 2008). The reuse of sewage or sludge in agriculture may facilitate the dissemination of parasites and the contamination of source water via surface runoff, raising public health concerns (Rimhanen-Finne *et al.*

2004). It is therefore important to assess the safety and microbial quality of wastewater in Tunisia.

Tunisian regulations governing the use of TWW and sludge (NT 106.03 and 106.20, respectively; [www.onas.nat.tn](http://www.onas.nat.tn)) define acceptable levels of specific chemicals, bacteria and helminth ova. However, there are no specifications for protists pathogenic to humans, such as *Cryptosporidium* and *Giardia*, which are responsible for large numbers of gastrointestinal infections (Karanis *et al.* 2007; Yoder & Beach 2010). Results of previous studies of wastewater and sludge

in Tunisia showed a more frequent occurrence of *Giardia* and *Entamoeba* cysts than helminth ova (Ben Ayed et al. 2009, 2010). These observations are in agreement with the high prevalence of these parasites in humans. In the Tunis region, the prevalence of *Giardia duodenalis* and *Cryptosporidium* in diarrheic persons was found to be 46.2 and 8%, respectively (Bouratbine et al. 2000). Approximately 16% of immuno-compromised patients had cryptosporidiosis (Bouratbine et al. 2000). In contrast, the prevalence of helminth infections was estimated to be low; 0.03% for *Ascaris lumbricoides* (Chaker et al. 1995) and 10.3% for *Enterobius vermicularis* (Bouratbine et al. 2000).

*Cryptosporidium* and *Giardia* can cause severe gastrointestinal illnesses. *Cryptosporidium* infections have been associated with stunting and malnutrition in young children (Chekley et al. 1997, 1998). They also have an adverse impact on growth and survival of neonatal calves and on milk production (Estban & Anderson 1995; Olson et al. 2004).

*Cyclospora cayetanensis* is recognized as a food- and waterborne parasite, and is transmitted by environmentally robust oocysts (Ortega & Sanchez 2010). Most outbreaks of *Cyclospora* have been associated with food-borne transmission and have occurred in industrialized countries (Thompson et al. 2003; Yoder & Beach 2007); water has however been identified as a route of transmission in developing countries, particularly in South America and Nepal (Shields & Olson 2003). Although related to *Cyclospora*, *Eimeria* spp. commonly infect domestic and wild animals and are not pathogenic to humans. *Enterocytozoon bieneusi* is the most common microsporidian species infecting humans (Didier 1998), domestic animals, wild mammals and birds (Ghosh & Weiss 2009; Santin & Fayer 2011), and is transmitted through the fecal-oral route. Spores of this fungal pathogen have been detected in surface water (Sperfel et al. 1997; Dowd et al. 1998) and swimming pools (Fournier et al. 2002).

In the present study, polymerase chain reaction (PCR) was used to evaluate the occurrence and geographical distribution of *Cryptosporidium* spp., *Cyclospora* spp., *Eimeria* spp., *G. duodenalis* and *E. bieneusi* in raw wastewater (RWW), TWW and sludge. Results of this survey document a high prevalence and genetic diversity of these parasites in wastewater and sludge in Tunisia.

## MATERIALS AND METHODS

### Study sites


A total of 110 samples of RWW, an equal number of TWW samples and 12 sludge samples (four dry and eight dehydrated) were collected monthly between 2005 and 2008 from 18 wastewater treatment plants located throughout Tunisia. Seven of the plants were located in the district of Tunis (Charguia, Chotrana, Cotière Nord, Grombalia, Kalaat el Andalous, Mornag and Sud Méliane), six in the coastal and tourist region (Kélibia, Mahdia, Nabeul, Slimane, Sousse Nord and Sousse Sud), three in the southern part of the country (Gafsa, Sfax and Sidi Bouzid), one in the center of the country (Kairouan) and one in the Sahel region (Moknine) (Figure 1). The main characteristics and the treatment process for each plant are listed in Table 1. These plants were selected mainly based on: (1) the final destination of the effluents (surface or sea waters, agricultural reuse, green space irrigation, etc.); (2) prior evidence of intestinal parasites in the local population (Bouratbine et al. 2000; Fathallah et al. 2004); (3) treatment process (activated sludge, waste stabilization pond); and (4) diversified geographical representation.

### Sample collection and processing

Samples of 5 L of RWW or TWW were collected over a 24 h period with an automatic sampler, whereas 100 g grab samples of sludge were collected. Upon arrival in the laboratory, wastewater samples were sedimented for 24 h at room temperature. The supernatant was removed and the sediment was centrifuged at 1,000 × g for 15 min. The final volume of sediment was about 10 mL. About 2 mL of sediment was stored in 2.5% potassium dichromate solution at 4 °C for the detection of targeted protists by PCR. Prior to DNA extraction, suspended solids were washed in water by centrifugation to remove the potassium dichromate.

### DNA extraction

DNA was extracted from 0.5 mL of wastewater sediment or 0.5 g of dry or dehydrated sludge using the FastDNA SPIN



**Figure 1** | Geographical location of 18 wastewater treatment plants. 1–6: Tunis metropolitan region; 7–13: coastal/tourist region; 14: Sahel region; 15: center; 16–18: south.

Kit for Soil (Q-Biogene, Irvine, CA) as described by Jiang *et al.* (2005). DNA was eluted into 100  $\mu$ L of reagent-grade water and stored at  $-80^{\circ}\text{C}$  until analyzed by PCR. Each sample was analyzed at least five times using 2  $\mu$ L of extracted DNA for each PCR target. The sample was considered positive if at least one reaction was positive. Following analysis with the nested PCR protocol described below, secondary PCR products were detected by gel electrophoresis on 1.5% agarose. Although a positive PCR outcome is not indicative of infectivity, the fact that DNA

extracted from the precipitate was used supports the view that a positive PCR result indicates the presence of intact spores, oocysts and cysts.

#### ***Cryptosporidium* detection, genotyping and subtyping**

A fragment of approximately 830 bp of the small subunit (SSU) rRNA gene was amplified by nested PCR as described by Jiang *et al.* (2005). To identify the *C. parvum* and *C. hominis* subtypes, an approximately 850-bp product of the 60 kDa

**Table 1** | Characteristics of treatment plants sampled in the study

| Plant              | Loc | Treatment |           |              | Final destination              |
|--------------------|-----|-----------|-----------|--------------|--------------------------------|
|                    |     | Primary   | Secondary | Disinfection |                                |
| Charguia           | DT  | PD        | AS        | –            | Irrigation, sea                |
| Chotrana           | DT  |           |           |              | Irrigation, sea                |
| Cotière Nord       | DT  | –         | WSP       | –            | Irrigation, sea                |
| Grombalia          | DT  | –         | AS        | –            | Surface water                  |
| Kalaat El Andalous | TCA | –         | WSP       | –            | Irrigation, sea                |
| Kélibia            | TCA | –         | AS        | UV           | Surface water                  |
| Mahdia             | TCA | –         | WSP       | UV           | Irrigation, sea                |
| Nabeul SE4         | TCA | PD        | AS        | –            | Surface water, recharge        |
| Sousse Sud         | TCA | PD        | AS + BB   | –            | Surface water                  |
| Gafsa              | SO  | –         | WSP       | –            | Irrigation                     |
| Sfax               | SO  | PD        | AS        | –            | Irrigation, sea                |
| Mornag             | DT  | PD        | AS        | UV           | Irrigation, sea                |
| Sud Méliane        | DT  | –         | AS        | –            | Surface water, sea             |
| Sidi Bouzid        | SO  | –         | WSP       | –            | Surface water                  |
| Slimane I          | TCA | –         | AS        | –            | Irrigation, sea                |
| Kairouan           | C   | –         | AS        | –            | Irrigation, surface water, sea |
| Sousse Nord        | TCA | PD        | AS        | –            | Irrigation, sea                |
| Moknine            | SA  | –         | WSP       | –            | Irrigation, surface water      |

Loc: location; DT: District of Tunis; TCA: tourist and coastal area; C: central; N: north; SO: south; SA: Sahel; PD: primary decantation; AS: activated sludge; BB: bacterial bed; WSP: waste stabilization pond; UV: ultraviolet light.

glycoprotein (gp60) gene was amplified by nested PCR (Alves *et al.* 2003). The secondary PCR products were sequenced using the secondary PCR primers and an intermediary sequencing primer (5'-GAGATATATCTTGTTGCG-3'). The established subtype nomenclature (Sulaiman *et al.* 2005) was used to classify the gp60 subtypes.

#### Detection and genotyping of *G. duodenalis*

To detect and genotype *G. duodenalis*, a 530-bp fragment of the triosephosphate isomerase (tpi) gene was amplified by nested PCR (Sulaiman *et al.* 2003a). Likewise, to detect and genotype *E. bienersi*, a fragment of about 392 bp of the ribosomal internal transcribed spacer (ITS) was amplified by nested PCR (Sulaiman *et al.* 2003b). The secondary PCR products of *G. duodenalis* and *E. bienersi* were sequenced. The ITS genotypes of *E. bienersi* were named according to the established nomenclature (Santin & Fayer 2009).

#### Detection and identification of *C. cayetanensis* and *Eimeria* spp.

To detect *C. cayetanensis*, a 294-bp fragment of the SSU rRNA gene was amplified by nested PCR (Ortega *et al.* 1993). Because these primers also amplify DNA of *Eimeria* spp., the identification of *C. cayetanensis* was confirmed by sequencing the secondary amplicons. *Eimeria* spp. were also targeted because they form monophyletic host-specific groups in sequence analyses, which was good for contamination source tracking.

#### DNA sequencing and sequence analysis

Secondary PCR products were purified using the Multiscreen PCR Plate (Millipore, Bedford, MA) and sequenced in both directions using the BigDye Terminator v. 3.1 cycle sequencing kit (Applied Biosystems, Foster City, CA) and an ABI 3130 automated sequencer (Applied Biosystems, Foster City, CA).

To confirm the accuracy of the sequence and to detect multiple genotypes and subtypes in a sample, all positive PCR products from each sample were sequenced, except for *Cyclospora/Eimeria* for which only two positive products per sample were sequenced. The genetic diversity of *Cryptosporidium* spp., *G. duodenalis*, *E. bieneusi* and *Cyclospora/Eimeria* spp. was examined by aligning the nucleotide sequences with those found in GenBank using ClustalX (<ftp://ftp-igbmc.ustrasbg.fr/pub/ClustalX>). Genotypes were classified by comparing each sequence with reference sequences.

### Meta-analysis of protist sequences

Given that the wastewater survey included treatment plants distributed over most of the country, protist species and genotypes were analyzed to identify geographical patterns in the occurrence and genetic diversity of different species and genotypes. Sequences were grouped into 15 taxa (*C. hominis*, *C. parvum*, *C. andersoni*, other *Cryptosporidium*, *G. duodenalis* A-I, *G. duodenalis* A-II, *G. duodenalis* B, other *Giardia*, *E. bieneusi* Peru8, *E. bieneusi* D, *E. bieneusi* IV, other *E. bieneusi*, ruminant *Eimeria*, poultry *Eimeria* and other *Eimeria*) and the presence/absence of each group in 111 samples tabulated. Pair-wise Jaccard distances (Jaccard 1908) between all pairs of taxa were calculated and taxa hierarchically clustered by similarity. The resulting tree was examined for the possible co-occurrence of protozoan species/genotypes with similar host range.

Parasite presence/absence data were analyzed for the co-occurrence of any *G. duodenalis* genotype and *Cryptosporidium* genotype, as well as the co-occurrence of *Cryptosporidium* and *E. bieneusi*, and *Cryptosporidium* species and *Eimeria* species. A Chi-square test was applied to test for any significant association between these species.

After eliminating ITS sequences with ambiguous positions, 209 *E. bieneusi* sequences were retained and numbered such that each unique sequence was assigned a different number ( $n = 44$ ). Likewise, after removing sequences with ambiguous positions, 191 small-subunit rRNA sequences of *Eimeria/Cyclospora* were aligned and numbered such that each unique sequence ( $n = 43$ ) was assigned an individual number. The diversity of the *E. bieneusi* ITS marker was compared to a previously published global collection of 135 ITS sequences (Widmer & Akiyoshi 2010) with individual-based

rarefaction analysis (Gotelli & Colwell 2001) using the program EstimateS (<http://viceroy.eeb.uconn.edu/estimates>). Heatmaps representing *E. bieneusi* ITS genotype abundance were drawn with a Microsoft Excel macro downloaded from <http://homepage.mac.com/yabyab/my.html>.

The ITS sequence diversity of *E. bieneusi* was displayed as a phylogenetic tree using the Neighbor-Joining clustering method (Saitou & Nei 1987). Sequences from a diversified global collection ( $n = 135$ ) and the present study ( $n = 209$ ) were pooled, aligned using ClustalX and imported into Mega 4.0 software (Tamura *et al.* 2007). A bootstrap consensus tree was drawn based on 500 replicates. The percentage of replicate trees in which a specific cluster was found is indicated in the plot. Rank-abundance plots (Whittaker 1965) were drawn to visualize the diversity of *E. bieneusi* genotypes among nine plants that had provided a minimum of 10 complete ITS sequences, and a same number of plants from which a minimum of nine *Eimeria* SSU rRNA sequences were obtained. The aim of this analysis was to assess whether treatment plants differed with respect to the genetic diversity of parasites. These two parasites were selected because of their extensive genetic diversity. The evenness of *E. bieneusi* ITS diversity was quantified using Pielou's  $J'$  ratio (Pielou 1969). This measure expresses the relative evenness as compared to maximum evenness which is found when all sequences have the same abundance. Treatment plants ( $n = 16$ ) contributing a minimum of four sequences were ranked from north to south to assess whether geography and environment would impact *E. bieneusi* genotype evenness.

### Nucleotide sequence accession numbers

Unique nucleotide sequences were deposited in GenBank under accession numbers EU140501–EU140508 and JF909981–JF910012.

## RESULTS

### Frequency and distribution of *Cryptosporidium* spp. in wastewater and sludge

Based on PCR analysis of the SSU rRNA gene, 42 of 110 (38%) RWW, 14 of 110 (13%) TWW, and five of 12 (33%)



**Table 2** | Parasites detected in raw and treated wastewater by treatment plant

| Plant              | Types of sample | # Sample   | # Cry positive | # <i>Giardia</i> positive | # Eb positive | # Ei positive |
|--------------------|-----------------|------------|----------------|---------------------------|---------------|---------------|
| Charguia           | RWW/TWW         | 8/8        | 3/1            | 4/1                       | 7/2           | 7/0           |
| Chotrana           | RWW/TWW/DeS     | 1/1/3      | 1/0/2          | 1/0/1                     | 0/0/2         | 0/0/1         |
| Cotière Nord       | RWW/TWW         | 5/5        | 3/0            | 5/1                       | 4/3           | 4/0           |
| Grombalia          | RWW/TWW         | 6/6        | 3/0            | 2/1                       | 5/2           | 3/0           |
| Kalaat El Andalous | RWW/TWW         | 28/28      | 7/5            | 8/2                       | 24/12         | 22/2          |
| Kélibia            | RWW/TWW         | 4/4        | 0/0            | 1/1                       | 3/1           | 2/0           |
| Mahdia             | RWW/TWW         | 4/4        | 3/1            | 0/0                       | 3/2           | 2/0           |
| Nabeul SE4         | RWW/TWW/DeS     | 25/25/1    | 10/2/0         | 9/4/0                     | 18/13/1       | 17/7          |
| Sousse Sud         | RWW/TWW/DeS     | 4/4/1      | 2/0/0          | 2/0/0                     | 3/2/1         | 4/2           |
| Gafsa              | RWW/TWW         | 5/5        | 2/2            | 3/0                       | 3/2           | 5/2           |
| Sfax               | RWW/TWW         | 6/6        | 0/0            | 3/1                       | 4/3           | 4/1           |
| Mornag             | RWW/TWW/DrS     | 4/4/4      | 1/1/1          | 2/2/1                     | 3/2/1         | 4/1/4         |
| Sud Méliane        | RWW/TWW/DeS     | 3/3/3      | 2/1/2          | 2/2/1                     | 2/1/3         | 1/3/2         |
| Sidi Bouzid        | RWW/TWW         | 2/2        | 0/1            | 1/0                       | 2/1           | 1/1           |
| Slimane I          | RWW/TWW         | 1/1        | 1/0            | 0/0                       | 1/0           | 1/0           |
| Kairouan           | RWW/TWW         | 2/2        | 2/0            | 2/0                       | 2/1           | 2/0           |
| Sousse Nord        | RWW/TWW         | 1/1        | 1/0            | 1/0                       | 1/1           | 1/0           |
| Moknine            | RWW/TWW         | 1/1        | 1/0            | 1/0                       | 1/1           | 1/0           |
| TOTAL              | RWW/TWW/sludge  | 110/110/12 | 42/14/5        | 47/15/3                   | 86/49/8       | 81/19/7       |

Cry: *Cryptosporidium* spp; DeS: Dehydrated sludge; DrS: Dry sludge; Eb: *Enterocytozoon bieneusi*, Ei: *Eimeria*.

sludge samples were positive for *Cryptosporidium* spp. (Tables 2 and 3). Altogether, 16 of the 18 plants sampled were positive for *Cryptosporidium* spp., including RWW samples from 15 plants and TWW samples from eight plants (Table 2). Sequencing of PCR products revealed the presence of *C. parvum*, *C. muris*, *C. andersoni*, *C. hominis*, *C. ubiquitum*, rat genotype IV, an unknown *Cryptosporidium*, sheep genotype, *C. meleagridis* and avian genotype II (Table 4). With 47 positive PCR products from nine wastewater treatment plants, *C. andersoni* was the most prevalent

species. Both the A and B copy of the gene, which differ by one T indel (Nagano *et al.* 2007), were found in RWW and TWW samples. The second-most common species was *C. hominis* (22 positive products), which was detected in RWW samples from 10 different plants and in one TWW sample collected from the plant of Charguia. In addition, 19 PCR products were positive for *C. parvum*, 11 for *C. muris*, six for *C. ubiquitum*, two for *C. meleagridis* and one each for the rat genotype IV, sheep genotype and a new genotype (Table 4). The sequence of the new *Cryptosporidium* genotype

**Table 3** | Frequency of *Cryptosporidium* spp., *Giardia duodenalis*, *Enterocytozoon bieneusi* and *Eimeria* spp./*Cyclospora cayetanensis* in wastewater and sludge samples in Tunisia

| Sample type        | No. of plants sampled | Sample size | <i>Cryptosporidium</i> spp. |                  | <i>Giardia duodenalis</i> |                  | <i>Enterocytozoon bieneusi</i> |                  | <i>Eimeria/Cyclospora</i> spp. |                  |
|--------------------|-----------------------|-------------|-----------------------------|------------------|---------------------------|------------------|--------------------------------|------------------|--------------------------------|------------------|
|                    |                       |             | Positive plants             | Positive samples | Positive plants           | Positive samples | Positive plants                | Positive samples | Positive plants                | Positive samples |
| Raw wastewater     | 18                    | 110         | 15                          | 42               | 16                        | 47               | 17                             | 86               | 17                             | 81               |
| Treated wastewater | 18                    | 110         | 8                           | 14               | 9                         | 15               | 16                             | 49               | 8                              | 19               |
| Dry sludge         | 1                     | 8           | 1                           | 1                | 1                         | 1                | 1                              | 1                | 1                              | 4                |
| Dehydrated sludge  | 4                     | 4           | 2                           | 4                | 2                         | 2                | 4                              | 7                | 2                              | 3                |

**Table 4** | *Cryptosporidium* species and genotypes in 232 wastewater and sludge samples

| Species, genotype and gp60 subtype | No. of positive plants | No. of positive PCR products |
|------------------------------------|------------------------|------------------------------|
| Humans and ruminants               |                        | 95                           |
| <i>C. hominis</i> (AF093491)       | 10                     | 22                           |
| IaA26R3                            | 1                      | 1                            |
| IaA27R4                            | 1                      | 1                            |
| IdA14                              | 2                      | 3                            |
| <i>C. parvum</i> (AF093490)        | 10                     | 19                           |
| IlaA15G2R1                         | 1                      | 1                            |
| IlaA17G2R1                         | 1                      | 1                            |
| IlaA18G3R1                         | 1                      | 1                            |
| IlaA20G2R1                         | 1                      | 1                            |
| IlaA21R1                           | 1                      | 2                            |
| IlaA21G2R1                         | 2                      | 2                            |
| IlaA5G3b                           | 1                      | 1                            |
| <i>C. andersoni</i> (AF093490)     | 9                      | 47                           |
| Sheep genotype (AY898790)          | 1                      | 1                            |
| <i>C. ubiquitum</i> (AF262328)     | 4                      | 6                            |
| Rodents                            |                        | 12                           |
| <i>C. muris</i> (AF093498)         | 4                      | 11                           |
| Rat genotype IV (AY737581)         | 1                      | 1                            |
| Birds                              |                        | 3                            |
| <i>C. meleagridis</i> (AF112574)   | 1                      | 2                            |
| Avian genotype II (DQ8853402)      | 2                      | 4                            |
| New genotype                       | 1                      | 1                            |

was similar to sequences of *C. baileyi* (DQ650339) and the avian genotype I (DQ885340) and could therefore be of avian origin. In 14 samples, more than one *Cryptosporidium* genotype was found; nine had two genotypes and five had three.

### *C. hominis* and *C. parvum* gp60 subtypes

Fourteen of the *C. hominis*- or *C. parvum*-positive samples were successfully subtyped by sequence analysis of the gp60 gene (Table 4). This analysis showed the presence of the *C. hominis* gp60 subtype families Id (3 sequences of 1 subtype) and Ia (two sequences of two subtypes) and *C. parvum* subtype families Ila (eight sequences of six subtypes) and Ila (one sequence).

### *G. duodenalis* genotypes

Sixty-five (28%) of 232 wastewater and sludge samples were positive for *G. duodenalis* by tpi PCR (Table 3), including RWW samples from 16 plants and TWW samples from nine plants (Table 2). For one collection each from Mornag and Kélibia, the TWW sample was positive despite a negative matching RWW sample as also observed with *Cryptosporidium*.

DNA sequence analysis of the PCR products revealed the presence of three known genotypes of *G. duodenalis* including assemblages A, B and E (Table 5). The most prevalent genotype was assemblage A, which was detected in 86 PCR products, followed by assemblage B in 12 PCR products. Assemblage E was detected in one sample (Table 5). Most of the assemblage A samples ( $n = 84$  PCR products) had the A2 subtype; A1 was detected twice.

### *E. bieneusi* genotypes

*E. bieneusi* was detected in 143 (62%) of 232 samples (Tables 2 and 3). In 2005, 61% of the samples were positive (5/7 RWW, 2/7 TWW, 1/2 dry sludge and 3/3 dehydrated sludge). In 2006, 44% of the samples showed the presence of *E. bieneusi* (38/57 RWW, 11/57 TWW and 4/5 dehydrated sludge). By 2007, *E. bieneusi* occurred in 80% of the samples (27/30 RWW and 21/30 TWW); in 2008 this species was present in all 16 RWW samples and in 94% (15/16) of TWW samples. A total of 86 RWW samples

**Table 5** | *Giardia duodenalis* assemblages and genotypes in 232 wastewater and sludge samples

| Genotype and subtype       | No. of positive plants | No. of positive PCR products |
|----------------------------|------------------------|------------------------------|
| Mammals (including humans) |                        | 14                           |
| A1 (L02120)                | 2                      | 2                            |
| B (U564283)                | 4                      | 10                           |
| B7 (EU140501)              | 1                      | 1                            |
| B8 (EU140504)              | 1                      | 1                            |
| Humans                     |                        | 84                           |
| A2 (U57897)                | 14                     | 84                           |
| Livestock                  |                        | 1                            |
| E (EU189356)               | 1                      | 1                            |

from 17 plants and 49 TWW samples from 16 plants were positive for *E. bieneusi* (Tables 2 and 3).

Overall, 44 unique ITS sequences were identified (Figure 2). Genotypes D and IV were the most prevalent with 85 and 82 products from 15 and 16 plants, respectively. These genotypes were also found in 10 and 11 TWW samples, respectively. In order of decreasing frequency, the next genotypes were Peru8 and WL4/WL5. Previously unknown ITS genotypes were detected in 22 samples. These were found in raw and TWW samples collected in 2007 and 2008.

To analyze whether *E. bieneusi* ITS sequences were geographically clustered, genotype counts were normalized by plant and graphically represented as a heat map (Figure 2). The most prevalent genotypes appear as columns of dark grey and grey. The frequency of detection does not show any association with specific plants. Frequent genotypes were detected in most plants, and rare genotypes were geographically scattered. This analysis indicates no geographic partitioning.


A total of 16 PCR amplicons generated a cluster of sequences not reported previously (Figure 3). These sequences formed a branch within a large group of ITS sequences originating from human and animal hosts comprising the majority of known ITS sequences (Thellier & Breton 2008; Widmer & Akiyoshi 2010) and designated group I by Henriques-Gil et al. (2010). In a bootstrap analysis

of 344 ITS sequences with 500 replicates, the cluster of 16 unique ITS sequences was present in 63% of the trees. The samples in which this sequence was found originated from Nabeul, Kalaat el Andalous, Mahdia, Kairouan and Sfax.

Noteworthy was the presence of three Tunisian ITS genotypes which cluster with a small, highly divergent group of three sequences previously identified in raccoons by Sulaiman et al. (2003b) and designated WL1, WL2, WL3. This cluster, designated as group III by Henriques-Gil et al. (Henriques et al. 2010) and group 4 by Thellier & Breton (2008), can therefore no longer be considered specific to raccoons as these animals do not live in North Africa. Consistent with the high prevalence of *E. bieneusi* positive PCR reactions, samples with more than one genotype were relatively common. A total of 35 samples from 14 plants had more than one genotype: 23 had two genotypes, eight had three and one had four different genotypes.


### *E. bieneusi* genetic diversity

To compare the genetic diversity of *E. bieneusi* ITS sequences with a previously analyzed global collection of ITS genotypes (Widmer & Akiyoshi 2010), a rarefaction analysis was performed. From the 292 ITS sequences obtained, 209 that had no ambiguous nucleotide positions were included in the analysis. The sequence diversity of



**Figure 2** | Distribution of *E. bieneusi* ITS sequences in RWW and TWW collected from 18 treatment plants. A total of 44 unique ITS sequences were detected among 209 PCR amplicons. Raw ITS counts for each plant were normalized by row such that each cell represents the percentage of samples by treatment plant (rows add up to 100%). Light shading  $\geq 12\%$  and black = 0 (not detected); lighter shading indicates higher percentages. Note the homogeneous distribution of *E. bieneusi* genotypes among plants. The arbitrarily assigned genotype numbers are shown uppermost. ITS genotype designation according to Santin & Fayer (2009) is shown below. As multiple ITS sequences are  $\geq 99\%$  similar to a named genotype, the same genotype name may apply to different columns.





**Figure 3** | Phylogenetic analysis of 344 *E. bieneusi* ITS sequences. Black and white bars represent samples from the global collection ( $n = 135$ ) and from Tunisia ( $n = 209$ ), respectively. Host species refer to isolates from the global collection. A large cluster of 263 related sequences from Tunisia and other countries is collapsed. Note the cluster of 16 ITS sequences not encountered previously. For branches observed >50% of the time in a bootstrap analysis with 500 replicates, the percentage of times the branch was observed is indicated. Scale: 0.2 nucleotide substitution per site.

the 209 sequences was 'rarefied' to a sample size of 135 to enable a direct comparison of the genetic diversity of the Tunisian and global ITS sequences. Based on this analysis, for a sample size of 135 the ITS diversity in Tunisia was estimated to be 29.2 genotypes with a 95% confidence interval of 23.0–35.4. This interval is less than half the observed global ITS diversity of 70 unique sequences. This finding is consistent with a more limited genetic diversity of *E. bieneusi* in Tunisian wastewater samples when compared to the global diversity.

#### PCR detection of *C. cayetanensis* and *Eimeria* spp.

Of the 232 samples analyzed for *C. cayetanensis*, 107 samples (45%) were PCR positive. Because the PCR also amplified DNA of *Eimeria* spp., amplicons were sequenced to identify the species. A total of 170 full-length sequences were obtained from PCR products originating from the 107 positive samples, including 81 RWW samples from 17 plants and 19 TWW samples from eight plants (Tables 2 and 3). *Cyclospora* DNA was detected in one RWW

**Table 6** | *Eimeria* species and related parasites in 232 wastewater and sludge samples from Tunisia

| Parasite <sup>a</sup>  | No. of positive plants | No. of positive PCR products |
|--|------------------------|------------------------------|
| <i>Eimeria</i> spp. from cattle  |                        | 59                           |
| <i>Eimeria alabamensis</i> (AF291427)  | 34                     | 52                           |
| <i>Eimeria alabamensis</i> -like (one SNP)   | 4                      | 7                            |
| <i>Eimeria</i> spp. from chicken   |                        | 15                           |
| <i>E. tenella</i> (U40264)   | 4                      | 6                            |
| <i>E. acervulina</i> (U67115)  | 2                      | 3                            |
| <i>E. mitis</i> (U40262)   | 1                      | 1                            |
| <i>E. tenella</i> -like 1 (one SNP)  | 2                      | 2                            |
| <i>E. tenella</i> -like 2 (two SNPs)   | 2                      | 2                            |
| Unknown species from birds (EU044785)  | 1                      | 1                            |
| <i>Eimeria</i> spp. from sheep   |                        | 19                           |
| <i>E. ahsata</i> (AF338350)  | 3                      | 5                            |
| <i>E. crandallis</i> (AF336339)  | 2                      | 4                            |
| <i>E. ahsata</i> -like 1 (one SNP)   | 1                      | 2                            |
| <i>E. ahsata</i> -like 2 (one SNP)   | 1                      | 1                            |
| <i>E. ahsata</i> -like 3   | 1                      | 1                            |
| <i>E. ahsata</i> -like 4   | 1                      | 1                            |
| <i>E. crandallis</i> -like 1 (one SNP)   | 1                      | 1                            |
| <i>E. crandallis</i> -like 2 (one SNP)   | 1                      | 1                            |
| <i>E. faurei</i> -like 1 (one SNP comparing to AF345998)                               | 1                      | 1                            |
| <i>E. faurei</i> -like 2 (one SNP and two deletions)                                   | 2                      | 2                            |
| <i>Eimeria</i> spp. from rodents   |                        | 27                           |
| <i>E. nieschulzi</i> (U40263)  | 8                      | 13                           |
| <i>E. telekii</i> (AF246717)   | 7                      | 11                           |
| <i>E. separate</i> -like 1 (one SNP comparing to AF311643)                             | 1                      | 1                            |
| <i>E. separate</i> -like 2 (two SNPs comparing to AF311643)                            | 2                      | 2                            |
| <i>Eimeria</i> spp. from rabbits   | 3                      | 3                            |
| <i>E. magna</i> or <i>E. media</i> (EF694016 or EF694013)                              | 1                      | 1                            |
| <i>E. magna</i> or <i>E. media</i> -like 1 (one SNP)                                   | 1                      | 1                            |
| <i>E. magna</i> or <i>E. media</i> -like 2 (two SNPs)                                  | 1                      | 1                            |
| <i>Isoospora</i> spp. from birds   | 32                     | 38                           |
| <i>Isoospora</i> sp. (AY331571)  | 29                     | 35                           |
| <i>Isoospora</i> sp.-like 1  | 1                      | 1                            |
| <i>Isoospora</i> sp.-like 2  | 2                      | 2                            |
| <i>Cycloospora cayetanensis</i> -like (one SNP and one deletion comparing to AF111183) | 1                      | 1                            |
| New genotype 1   | 3                      | 5                            |
| New genotype 2   | 1                      | 2                            |

<sup>a</sup>Forty-three unique sequences are assigned to 32 *Eimeria*, *Isoospora* and *Cycloospora* species.

sample collected in Kalaat el Andalous on 26 March 2008. The sequence obtained had two nucleotide differences from *C. cayetanensis*. The other PCR amplicons originated mostly from *Eimeria alabamensis* and relates species infecting cattle (59 sequences), *E. tenella*, *E. acervulina*, *E. mitis* and their relatives from chicken (15 sequences), *E. ahsata*, *E. crandallis* and relatives from sheep (19 sequences), *E. nieschulzi*, *E. telekii* and their relatives from rodents (27 sequences), *E. magna* or *E. media* and their relatives infecting rabbits (three sequences), *Isospora* spp. from birds (38 sequences) and two unknown coccidia called new genotypes 1 (JF910008, JF910009 JF910011 and JF910012) and two (JF910004) (seven sequences) (Table 6).

### Meta-analysis of protist sequences from wastewater

Using cluster analysis, no association between species with similar host range was found. For instance, *C. andersoni*, a parasite of cattle, did not cluster with bovine *Eimeria* species, and the anthroponotic species *C. hominis* did not cluster with *E. bienewsi* ITS genotypes found in humans, such as Peru8. The lack of detectable co-occurrence of parasites with similar host range indicates that parasites in different plants originate from multiple sources, including humans and livestock. The same set of presence/absence data was analyzed for the co-occurrence of *G. duodenalis* genotypes and *Cryptosporidium* genotypes, as well as the co-occurrence of *Cryptosporidium* and *E. bienewsi*, and *Cryptosporidium* species and *Eimeria* species. This analysis revealed a significant association among parasites infecting both humans and animals, i.e. *Cryptosporidium*, *G. duodenalis* and *E. bienewsi* (Bonferroni-adjusted  $p = 0.03$ ), whereas the occurrence of animal parasites of the genus *Eimeria* was not associated with *Cryptosporidium* (Bonferroni-adjusted  $p = 0.6$ ). Thus, although hierarchical clustering did not suggest the co-occurrence according to host range, a significant association between three zoonotic genera was found.

As mentioned above, in 209 *E. bienewsi* PCR products, 44 unique ITS sequences were identified. For *Eimeria*, 191 products were positive by PCR and 43 unique SSU rRNA sequences were found. This analysis revealed clear differences in *E. bienewsi* and *Eimeria* genetic diversity among treatment plants (data not shown). For instance, the

*E. bienewsi* ITS sequences obtained from Charguia (District of Tunis) were the most diverse group, whereas those from Sousse Sud (Coastal area) and Sud Meliane (District of Tunis) were the least diverse. In the latter two plants, 62 and 60% of the ITS sequences, respectively, were identical. A similar wide range of genetic diversity among plants was observed in the SSU rRNA sequences of *Eimeria*. Interestingly, no correlation was observed between the diversity of the two parasites. Sousse Sud had the least diverse *E. bienewsi* population but the most diverse *Eimeria* population. Conversely, Charguia had the most diverse *E. bienewsi* population but the least diverse *Eimeria* population. This analysis could not be performed for other protists due to the relatively low genetic diversity.

The analysis of the evenness of *E. bienewsi* ITS genotypes revealed no north-south cline; plants with even ITS distribution were found in the north of the country, for instance Charguia ( $J' = 0.78$ ), as well as in the center (Mahdia,  $J' = 0.87$ ) and south (Sfax,  $J' = 1.00$ ).

### DISCUSSION

Results of the survey document a high prevalence and genetic diversity of pathogenic protists in RWW in Tunisia, and suggest that the various wastewater treatment processes adopted in Tunisia are insufficient in removing these parasites. As sedimentation was a key process in all treatment processes, the reduction in the occurrence of these protists was affected by the size of the environmental stage. As a result, a higher reduction in occurrence was achieved for the large *Eimeria* oocysts (76%) than for the small *E. bienewsi* spores (43%), with the medium-sized *Cryptosporidium* oocysts (66%) and *Giardia* cysts (68%) in between. Nevertheless, there were no apparent differences in the reduction of the occurrence of protists among different treatment practices. For example, the two plants that were sampled most frequently (Kalaat El Andalous and Nabeul SE4) had a similar reduction in the occurrence of *Cryptosporidium* spp., *G. duodenalis*, *E. bienewsi* and *Eimeria* spp., even though the former had no primary treatment and used waste stabilization pond as the secondary treatment, whereas the latter used decantation as the primary treatment and activated sludge as the secondary treatment

(Tables 1 and 2). These data should be useful in the development of measures to reduce the transmission of enteric pathogens and to improve the safety of discharge and reuse of wastewater and sludge.

The use of PCR for analyzing wastewater samples provided an overview of protists circulating in the population at the time of sampling. This information is relevant to understanding transmission cycles and the potential sources of infections affecting humans, livestock and other animals. This type of study could also provide a useful approach for collecting data on the occurrence of enteric diseases in different regions of the country. However, no information on the infectivity of waterborne parasite spores, cysts and oocysts was gained from the detection of DNA. The fact that solids from wastewater samples were sedimented and PCR was applied to DNA extracted from the solid fraction suggests that the DNA was enclosed in intact spores, cysts and oocysts.

The analysis of co-occurrence of parasites with similar host range gave inconsistent results. Hierarchical clustering showed no association, whereas a positive association among zoonotic parasites was found. This outcome could originate from the fact that for hierarchical clustering 15 species/genotypes were considered, whereas association was tested among *Giardia*, *Cryptosporidium*, *Eimeria* and *E. bieneusi* without differentiating between species and genotypes. The lack of consistent outcome of these analyses is in agreement with the wide distribution and frequent co-occurrence of protist species in different plants.

The large number of positive samples enabled an analysis of geographical distribution of species and genotypes. These analyses showed no obvious difference between plants or regions. Due to the large number of water samples positive for *E. bieneusi*, geographical trends were explored by analyzing the distribution of *E. bieneusi* ITS genotypes and, to some extent, *Eimeria* SSU rRNA genotypes. The lack of any geographical pattern is consistent with the relatively small distances between plants and the fact that wastewater is mostly impacted by human and livestock waste. Any geographical partition would be expected to be rapidly eliminated by human mobility and the presence of numerous slaughterhouses throughout the country.

The results of the rarefaction analysis of *E. bieneusi* ITS genotypes is consistent with a globally restricted distribution of certain ITS genotypes; that is, after adjustment for equal

sampling, fewer *E. bieneusi* genotypes are present in Tunisia as compared to a global reference collection. This observation needs to be interpreted with caution. The diversity of the reference collection (Widmer & Akiyoshi 2010) might be higher than the real diversity because this collection is derived from the GenBank database. Investigators who deposit sequences from epidemiological surveys frequently only deposit one sequence representative of multiple identical sequences, and may not deposit a sequence that is already found in GenBank. Although this bias may be reduced by the fact that some investigators submit sequences regardless of what has already been submitted, the possibility that the global reference collection we used as a benchmark is more diverse than in reality needs to be considered.

Pathogens present in wastewater have the potential to be transmitted to humans, especially when water is reused for irrigation. In this study, wastewater was found to frequently carry pathogens of human and animal origin. The presence of species which do not infect humans such as *Eimeria* demonstrates that wastewater, even in urban areas, is contaminated with fecal matter of animal origin. Given the public health risk of these parasites to humans and farm animals, guidelines on the discharge and reuse of wastewater are needed to reduce the transmission potential and transport of these pathogens in Tunisia and other countries with limited water sources.

## ACKNOWLEDGEMENTS

We thank Abdellatif Boudabous (El Manar University, Tunis) for facilitating this project. This work was partially carried out in the laboratory of Fethi Lebdi at the Institut National Agronomique de Tunisie. GW acknowledges financial support from the NIAID (grant R21AI083719). The findings and conclusions in this report are those of the authors and do not necessarily represent the views of the Centers for Disease Control and Prevention.

## REFERENCES

- Alves, M., Xiao, L., Sulaiman, I., Lal, A. A., Matos, O. & Antunes, F. 2005 *Subgenotype analysis of Cryptosporidium isolates*

- from humans, cattle, and zoo ruminants in Portugal. *J. Clin. Microbiol.* **41**, 2744–2747.
- Ben Ayed, L., Schijven, J., Alouini, Z., Jemli, M. & Sabbahi, S. 2009 Presence of parasitic protozoa and helminth in sewage and efficiency of sewage treatment in Tunisia. *Parasitol. Res.* **105**, 393–406.
- Ben Ayed Khouja, L., Cama, V. L. & Xiao, L. 2010 Parasitic contamination in wastewater and sludge samples in Tunisia using three different detection techniques. *Parasitol. Res.* **107**, 109–116.
- Bouratbine, A., Aoun, K., Siala, E., Chahed, M. K., Ben Hassine, L. & Meherzi, A. 2000 Pour une meilleure estimation de la prévalence du parasitisme intestinal dans la région de Tunis. *Bull. Soc. Pathol. Exot.* **93**, 353–355.
- Chaker, E., Bel Hadj, S., Khaled, S., Ben Moussa, M. & Ben Rachid, M. S. 1995 Les parasitoses digestives, problème toujours d'actualité. *La Tunisie médicale* **73**, 53–56.
- Checkley, W., Gilman, R. H., Epstein, L. D., Suarez, M., Diaz, J. F., Cabrera, L., Black, R. E. & Sterling, C. R. 1997 Asymptomatic and symptomatic cryptosporidiosis: their acute effect on weight gain in Peruvian children. *Am. J. Epidemiol.* **145**, 156–163.
- Checkley, W., Epstein, L. D., Gilman, R. H., Black, R. E., Cabrera, L. & Sterling, C. R. 1998 Effects of *Cryptosporidium parvum* infection in Peruvian children: growth faltering and subsequent catch-up growth. *Am. J. Epidemiol.* **148**, 497–506.
- Didier, E. S. 1998 Microsporidiosis. *Clin. Infect. Dis.* **27**, 1–8.
- Dowd, S. E., Gerba, C. P. & Pepper, I. L. 1998 Confirmation of the human-pathogenic microsporidie *Enterocytozoon bienewisi*, *Encephalitozoon intestinalis* and *Vittiforma corneae* in water. *Appl. Environ. Microbiol.* **64**, 3332–3335.
- Estban, E. & Anderson, B. C. 1995 *Cryptosporidium muris*: prevalence, persistency, and detrimental effect on milk production in a drylot dairy. *J. Dairy Sci.* **78**, 1068–1072.
- Fathallah, A., Saghrouni, F., Madani, B., Ben Rejeb, N. & Ben Said, M. 2004 Le parasitisme digestif dans la région de Sousse. Bilan d'une étude rétrospective de 16 années. *La Tunisie Médicale* **82**, 335–343.
- Fournier, S., Dubrou, S., Liguory, O., Gaussin, F., Santillana-Hayat, M., Sarfati, C., Molina, J. M. & Derouin, F. 2002 Detection of microsporidia, cryptosporidia and *Giardia* in swimming pools: a one-year prospective study. *FEMS Immuno. Med. Microbiol.* **33**, 209–213.
- Ghosh, K. & Weiss, L. M. 2009 Molecular diagnostic tests for microsporidia. *Interdisciplinary Perspectives Infect. Dis.* 2009, Article ID 926521.
- Gotelli, N. J. & Colwell, R. K. 2001 Quantifying biodiversity: procedures and pitfalls in the measurement and comparison of species richness. *Ecol. Lett.* **4**, 379–391.
- Henriques-Gil, N., Haro, M., Izquierdo, F., Fenoy, S. & del Aguila, C. 2010 Phylogenetic approach to the variability of the microsporidian *Enterocytozoon bienewisi* and its implication for inter- and intrahost transmission. *Appl. Environ. Microbiol.* **76**, 3333–3342.
- Jaccard, P. 1908 Nouvelles recherches sur la distribution florale. *Bull. Soc. Vaudoise Sci. Nat.* **44**, 223–270.
- Jiang, J., Alderisio, A. K., Singh, A. & Xiao, L. 2005 Development of procedures for direct extraction of *Cryptosporidium* DNA from water concentrates and for relief of PCR inhibitors. *Appl. Environ. Microbiol.* **71**, 1135–1141.
- Karanis, P., Kourenti, C. & Smith, H. 2007 Waterborne transmission of protozoan parasites: a worldwide review of outbreaks and lessons learnt. *J. Water Health* **5**, 1–38.
- Nagano, S., Matsubayashi, M., Kita, T., Narushima, T., Kimata, I., Iseki, M., Hajiri, T., Tani, H., Sasai, K. & Baba, E. 2007 Detection of a mixed infection of a novel *Cryptosporidium andersoni* and its subgenotype in Japanese cattle. *Vet. Parasitol.* **149**, 213–218.
- Office National d'Assainissement (ONAS) 2008 Rapport annuel d'exploitation des stations d'épurations de l'année 2006. Office National d'Assainissement, Tunis, Tunisia, 80.
- Olson, M. E., O'Handley, R. M., Ralston, B. J., McAllister, T. A. & Thompson, R. C. A. 2004 Update on *Cryptosporidium* and *Giardia* infections in cattle. *Trends Parasitol.* **20**, 185–191.
- Ortega, Y. R. & Sanchez, R. 2010 Update on *Cyclospora cayetanensis*, a food-borne and waterborne parasite. *Clin. Microbiol. Rev.* **23**, 218–234.
- Ortega, Y. R., Sterling, C. R., Gilman, R. H., Cama, V. A. & Diaz, F. 1993 *Cyclospora* species-a new protozoan pathogen of humans. *N. Engl. J. Med.* **328**, 1308–1312.
- Pielou, E. C. 1969 *An Introduction to Mathematical Ecology*. Wiley, New York.
- Rimhanen-Finne, R., Vourinen, A., Marmo, S., Malmberg, S. & Hanninen, M. L. 2004 Comparative analysis of *Cryptosporidium* and *Giardia* and indicator bacteria during sewage sludge hygienization in various composting processes. *Letts. Appl. Microbiol.* **38**, 301–305.
- Saitou, N. & Nei, M. 1987 The neighbour-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**, 406–425.
- Santin, M. & Fayer, R. 2009 *Enterocytozoon bienewisi* genotype nomenclature based on the internal transcribed spacer sequence: a consensus. *J. Eukaryot. Microbiol.* **56**, 34–38.
- Santin, M. & Fayer, R. 2011 Microsporidiosis: *Enterocytozoon bienewisi* in domesticated and wild animals. *Res. Vet. Sci.* **90**, 363–371.
- Shields, J. M. & Olson, B. H. 2003 *Cyclospora cayetanensis*: a review of an emerging pathogenic coccidian. *Int. J. Parasitol.* **33**, 371–391.
- Sperfel, J. M., Sarfati, C., Liguory, O., Caroff, B., Dumoutier, N., Gueglio, B., Billaud, E., Raffi, F., Molina, J. M., Miegville, M. & Derouin, F. 1997 Detection of microsporidia and identification of *Enterocytozoon bienewisi* in surface water by filtration followed by specific PCR. *J. Eukaryot. Microbiol.* **44**, 78s.
- Sulaiman, I. M., Fayer, R., Bern, C., Gilman, R. H., Trout, J. M., Schantz, P. M., Das, P., Lal, A. A. & Xiao, L. 2003a Triosephosphate isomerase gene characterization and



- potential zoonotic transmission of *Giardia duodenalis*. *Emerg. Infect. Dis.* **9**, 1444–1452.
- Sulaiman, I. M., Fayer, R., Lal, A. A., Trout, J. M., Schaefer, F. W. & Xiao, L. 2003b Molecular characterization of microsporidia indicates that wild mammals harbour host-adapted *Enterocytozoon* spp. as well as human-pathogenic *Enterocytozoon bieneusi*. *Appl. Environ. Microbiol.* **69**, 4495–4501.
- Sulaiman, I. M., Hira, P. R., Zhou, L., Al-Ali, F. M., Schweiki, H. M., Iqbal, J., Khaled, N. & Xiao, L. 2005 Unique endemicity of cryptosporidiosis in children in Kuwait. *J. Clin. Microbiol.* **43**, 2805–2809.
- Tamura, K., Dudley, J., Nei, M. & Kumar, S. 2007 MEGA4: Molecular evolutionary genetic analysis (MEGA) software version 4.0. *Mol. Biol. Evol.* **24**, 1596–1599.
- Theulier, M. & Breton, J. 2008 *Enterocytozoon bieneusi* in humans and animals, focus on laboratory identification and molecular epidemiology. *Parasite* **17**, 321–328.
- Thompson, R. C. A., Armson, A. & Ryan, U. M. 2003 *Cryptosporidium: From Molecules to Disease*. Elsevier Science Ltd, Amsterdam.
- Whittaker, R. H. 1965 Dominance and diversity in land plant communities. *Science* **147**, 250–260.
- Widmer, G. & Akiyoshi, D. 2010 Host-specific segregation of ribosomal nucleotide sequence diversity in the microsporidian *Enterocytozoon bieneusi*. *Infect. Genet. Evol.* **10**, 122–128.
- Yoder, J. S. & Beach, M. J. 2007 Giardiasis surveillance – United States 2003–2005. *MMWR* **56** (SS07), 11–18.
- Yoder, J. S. & Beach, M. J. 2010 *Cryptosporidium* surveillance and risk factors in the United States. *Exp. Parasitol.* **124**, 31–39.

First received 30 November 2011; accepted in revised form 22 April 2012. Available online 30 May 2012