

3. small white area=2 points
4. brown scar within extremely white point in it =1 point
5. entire brown scar=0 points

The controls should receive a score of zero. Each patch is scored separately and the three are averaged for each concentration.

#### Problems in the Investigation:

1. Some of the *Tenebrio* larvae would not pupate. Be very careful not to chill the larvae. Chilling lengthens the life cycle considerably.
2. After injection some of the pupae died. Use a fine gauge needle (approximately a 25-27) so less damage is done to the insect.
3. Inject slowly. If you inject rapidly the powerful spray may damage tissues excessively.
4. Use sufficient numbers of experimental insects to make valid conclusions. In some of the early work only six insects were used and four died at various stages making it almost impossible to make valid conclusions. In later experiments it was noted that we got nearly 30% recovery or completion of the experiment. (Nearly 70% died before reaching adulthood.

#### Results of the Investigation:

Of the 30% of insects recovered it was found that adults treated with 1:0, 1:1, 1:2, 1:4 and 1:8 farnesol to corn oil concentrations received scores of 4. Insects treated with concentrations of 1:16, 1:32 and 0:1 received scores of 1 and had brown scars. To interpret these results, the farnesol apparently does act as the juvenile hormone, neotenin. It retards normal metamorphosis of tissue and thus leaves white scars. If normal metamorphosis goes on brown scars show at the points of injection

injury.

The adults that received a score of 4 had large sections of pupal cuticle and the wings were deformed. The *Tenebrio* treated with 1:0 concentration of farnesol had wings that were only about half the length of the abdomen.

The controls (treated with pure oil) had brown scars and had no visible pupal cuticle.

This type of investigation has opened up many questions such as why is the hatching rate so low? Is it humidity, temperature, light or what? Could farnesol be applied on the surface and get results without breaking the epidermis? Was there some product in the vials retarding metamorphosis such as found by Dr. Carroll Williams at Harvard? He found some paper contains a natural juvenile hormone which apparently was an evolutionary feature to protect certain trees from injury by insects.

#### ■ BIBLIOGRAPHY

- CUMMINS, K. W., L. D. MILLER, N. A. SMITH, and R. M. FOX, *Experimental entomology*. New York: Reinhold Publishing Corp., 1965.
- FISHER, FRANK M., and SANBORN, RICHARD C. *Nosema* as a source of juvenile hormone in parasitized insects, *Biological Bulletin*, Vol. 126, No. 2, April, 1964, pp. 235-52.
- FISHER, FRANK M., and SANBORN, RICHARD C. Production of insect juvenile hormone by the microsporidian parasite *Nosema*, *Nature*, Vol. 194, No. 4834, June 23, 1962, p. 1193.
- HANSTROM, BERTIL. *Hormones in invertebrates*. Oxford, Great Britain: Oxford University Press, 1939.
- TURNER, C. DONNELL. *General endocrinology*. Philadelphia: W. B. Sanders Company, 1948.
- WATTERSON, RAY. *Endocrines in development*. Chicago: University of Chicago Press, 1959.
- Personal Communications:  
DR. RICHARD C. SANBORN, Biology Department, Purdue University.

## FLEA BEETLES

Some 2,000 Argentine flea beetles have joined the Army Corps of Engineers' battle against the channel-choking alligator weed. The tiny insects feed entirely on the noxious aquatic weed, which is a serious impediment to navigation and stream flow in 3,000 miles of inland waterway in eight southeastern states from North Carolina around to Texas. The beetles were released on an experimental basis and with the consent of the states concerned and the U. S. Fish and Wildlife Service, in the Savannah National Wildlife Refuge near the South Carolina-Georgia border. The move followed four years of research in cooperation with the Agricultural Experiment Service of the U. S. Department of Agriculture to make sure the introduction of the beetles would have no adverse repercussions on other aquatic growth.

## MOUTH PROTECTORS

The football player who wears a mouthguard inside his mouth may be helping to protect himself from a brain concussion.

Mouth protectors aid in reducing skull deformation and compression between the brain. Such injuries are common in contact sports and often result from blows to the head. If the injuries are severe, brain damage may occur.

Dentists have felt for some time that mouth protectors may do much more than prevent injury to the oral tissues.

## FISH REST REGULARLY

Fish seldom appear to be sleeping because they do not have eyelids, but observations reveal that they do rest regularly.