

# Simulations of Natural Selection

By ROBERT PATTERSON, T. CUSTER, and BAYARD H. BRATTSTROM

Natural-selection simulation games have been used in a variety of ways in elementary schools (Klimas, 1969) and secondary schools (Kuhn, 1969). The present model includes color-matching by prey (crypsis), morphologic adaptation to habitat (beak length vs. prey size), flock vs. individual feeding success, and the concept of carrying capacity of the environment. This has been successfully used in a college and a secondary school and with modification could be used in elementary schools.

## Concealing Coloration

The class is divided into two equal groups, each including a person who acts as recorder. The group is taken to a preselected habitat, which consists of two large lawns or weedy fields. Each group walks onto one of the lawns or fields. The students close their eyes while 100 red and 100 green toothpicks (cocktail or food-color-dyed) are scattered at random in each of the habitats. (This can be done before class.) The toothpicks represent insect prey. The students then open their eyes. Pretending to be birds, they collect as many of the toothpick prey as possible in a 30-second trial. (Size of plots and amount of time spent feeding can be varied, with interesting results.) They close their eyes again, and the recorder tallies and collects the toothpicks from each person. The trials are repeated until each group has collected most of the 200 toothpick prey. The red toothpicks are very obvious and are picked up rapidly, but eight or more trials may be required to find all of the green

toothpicks. This agrees with and supports the findings of Kettlewell (1959) on the selective advantage of habitat-matching in moths.

The class is next taken to a brown dirt habitat, where 200 red and green toothpicks are distributed and the hunt for them is repeated. Both colors of prey (toothpicks) are easily seen here and are selected against by the birds (students); usually fewer than four trials are needed to collect most of the toothpicks.

## Morphologic Adaptation

To show morphologic adaptation within a habitat, beak length in birds as seen in the Galápagos Islands finches (Lack, 1953) can be easily investigated. The two groups are taken to the grassy or weedy plots as before. Half of each group is provided with kitchen tongs or long forceps: these students are the long-beaked birds. The other half of each group is asked to pick up prey with only one hand: they are the short-beaked birds. Wooden matches are scattered over the two plots—long-stemmed matches on one plot and short-stemmed matches on the other. In either case the matches are to be picked up with the hand or tongs and transferred to the other hand, for holding. The data show the disadvantage of using tongs (long beaks) to pick up the short-stemmed matches (small prey) but not the long-stemmed matches (larger prey).

## Feeding Efficiency

Flock vs. individual feeding efficiency can be studied by using 200 green matches on the grassy plots. Half the class “feeds” as individuals; the other half feeds as a flock or herd, each student remaining

Authors' address: c/o B. H. Brattstrom, Biology Dept., California State College, 800 N. State College Blvd., Fullerton 92631.

Fig. 1. Data sheet (abbreviated) used by a recorder.

CONDITION:	BIRD NO.	TRIALS AND PREY TYPE													
		1		2		3		4		5		6		7	
Grass		R	G	R	G	R	G	R	G	R	G	R	G	R	G
	1	12	14	7	8	7	3	1	1	0	2	0	0		
	2	9	4	8	5	5	4	0	2	0	1	0	1		
	3	5	2	6	5	3	1	3	1	0	1	0	1		
	4	4	4	3	4	2	3	0	2	0	0	0	0		
	5	4	5	5	3	1	2	0	0	0	0	0	0		
	6	7	6	3	6	1	4	4	4	0	1	0	0		
	Sum	41	35	32	31	19	17	8	10	0	5	0	2		
	Cum f	41	35	73	66	92	83	100	98	—	98	—	100		

within 30 cm of his neighbor. The flock or herd usually will collect more food items, because of its greater efficiency in finding prey in a restricted locality, its cooperative strategy in hunting, and its social facilitation, as was noted by Etkin (1967) in both mammals and birds.

The effect of injury on feeding efficiency can be studied by using 100 red toothpicks on the green plot. In one group the students cover one eye with a hand, simulating an eye injury; in the other group the students simply place the unused hand on top of the head. The data usually show higher efficiency in the binocular birds than in the injured, monocular birds. The role of injury and illness in decreasing the chances of survival is well known for both individuals and groups of animals; for example, Washburn and DeVore (1961) noted this in baboon troops.

### Carrying Capacity

Carrying capacity can be illustrated on two grassy plots of equal size. Scatter 200 toothpicks on each plot. Start with only six people (birds) to a plot. After each trial, while the number of prey items captured is being tallied, cast 26 new toothpicks onto




Fig. 2. Capture of red and green prey in a green grass habitat.

the plot selected for prey growth. The birds on the other plot, which has no prey growth, are soon observed to “starve”; but the competition for prey becomes rigorous on the plot with prey growth. The model can be amplified and altered by changing the numbers of birds on each plot. Data can be suggested to the class for consideration. For example, a species of longspur, which is a tundra bird, requires three prey items every 15 minutes for eight hours every day, on average (Custer, 1971). According to Welty (1962) the European blackbird (a thrush) requires 7.3% of its body weight in food daily; for most birds the figure is 3% to 30%. Data from the Tasmanian sheep populations (Davidson, 1938; Caughley, 1970) can be used to illustrate growth to a fixed value that represents the carrying capacity of the range. Overpopulation and crashes can be illustrated by using data from the Kiabab Plateau population of mule deer, as presented by Caughley (1970).

### Instructional Use of Data

Fig. 1 is a data sheet (abbreviated) used by one of the recorders. The data can be cumulated and graphed after the recorder returns to the laboratory. A graph could show the cumulative number of prey removed from the habitat under a given condition of predation (fig. 2), or it could show the number of prey remaining in the habitat per trial. The first graph would show the number of prey removed under the two conditions; the latter would show the number of prey still to be removed from the habitat. The students can also tally the data for each individual and then rank the birds as to their efficiency in each habitat: Usually they will discover that a bird is more efficient in one habitat than in another. Data on sex can also be tallied: usually males will have collected more prey items than most of the females.

*Acknowledgment.*—Al Muth made helpful suggestions and tested several of the ideas.

### REFERENCES

CAUGHLEY, G. 1970. Eruption of ungulate populations, with Himalayan thar in New Zealand. *Ecology* 51 (1): 53-72.

- CUSTER, T. 1971. *Breeding biology of the Alaskan longspur*. Master's thesis, California State College, Fullerton.
- DAVIDSON, J. 1938. On the growth of the sheep population in Tasmania. *Transactions of the Royal Society of South Australia* 62: 342-346.
- ETKIN, W. 1967. *Social behavior from fish to man*. University of Chicago Press, Chicago.
- KETTLEWELL, H. B. D. 1959. Darwin's missing evidence. *Scientific American* 200 (4): 48-53.
- KLIMAS, J. 1969. A cooperative program for improving elementary science education. *American Biology Teacher* 31: 493-496.
- KUHN, D. J. 1969. A simulation game on natural selection. *Science Teacher* 36 (1): 68.
- LACK, D. 1953. Darwin's finches. *Scientific American* 188 (4): 66-72.
- WASHBURN, S. L., and I. DEVORE. 1961. The social life of baboons. *Scientific American* 204 (6): 62-71.
- WELTY, J. D. 1962. *The life of birds*. W. B. Saunders Co., Philadelphia.

## ANCIENT ALGAE HOLD SECRETS

Fossil evidence that blue-green algae much like those that exist today existed as much as two billion years ago has been found on a knoll near Eveleth, Minn. Blue-green algae are among the world's most primitive organisms; they are not much more advanced than the most primitive bacteria. Yet the Eveleth samples appear to be very similar to the blue-green algae that help clog today's eutrophying ponds. They are among the world's earliest identifiable life forms, and the Minnesota fossils, says Preston Cloud, a biogeologist at the University of California at Santa Barbara, are the oldest demonstrable examples yet found.

He said the Minnesota fossils appear to be "slightly older—probably not more than a few million or tens of millions of years," than the famous Gunflint microfossils from the north shore of Lake Superior, the oldest that could up to now be confidently identified.

"But the real significance is not the age," said Cloud, whose research is supported by the National Science Foundation and the National Aeronautics and Space Administration. "They are not all that much older than the Gunflint fossils." In contrast with the Gunflint microfossils, the fact that they are readily separable from the rock deposits in which they occur permits a comparison with living organisms with a precision and detail not heretofore possible.

If, thanks to fine detail like that seen in these fossils from the Pokegama strata in Minnesota, evolutionary changes can be detected in blue-green algae, he suggests, then perhaps they can be used to identify the relative ages of strata in which they are found.

Older microfossils have been reported; they go back some 3.2 billion years or more. But Cloud feels that those are still open to scientific question; the Pokegama and Gunflint fossils, he said, are the oldest ones that can be called relics of early life with 100% certainty.

## SUGGESTIONS FOR CONTRIBUTORS

STYLE. *American Biology Teacher* would rather receive an ill-written article containing worthwhile ideas than a stylistic masterpiece that says little: our editors can mend bad writing in a good cause. However, we do hope for clear terse prose, free of jargon. Sensible advice for writers will be found in the Conference of Biological Editors' *Style Manual for Biological Journals*, 2nd ed., published by the American Institute of Biological Sciences; and *How to Write Scientific and Technical Papers*, by Sam F. Trelease.

In matters of punctuation, enumeration, and the like we follow generally the above-mentioned C.B.E. manual and the University of Chicago *Manual of Style*. Our spellings are usually those preferred by *Webster's Third New International Dictionary* (G. & C. Merriam Co., 1965) and its abridgments.

Technical measurements are in metric, not English, units.

Avoid footnotes of any kind. References to the literature are made on-line (not by means of superscripts) within the text. If only one, two, or three works are cited, each is given in full, in the form "A. B. Smith, 1969: *Elements of Biology*, 4th ed., Jones Publishing Co., New York" for a book and "W. X. White and Y. Z. Green, 1965: 'The Inquiry Process,' *Journal of Pedagogy* 7 (2): 53-56" for an article. If four or more works are cited, they are presented at the end of the article as a bibliography arranged alphabetically by (principal) authors' last names, in the following forms for books and journals.

SMITH, A. B. 1969. *Elements of biology*, 4th ed. Jones Publishing Co., New York.

WHITE, W. X., and Y. Z. GREEN. 1965. The inquiry process. *Journal of Pedagogy* 7 (2): 53-56.

(Note that publishers' names and addresses are given and that names of periodicals are not abbreviated.) Reference to the bibliography from the text takes the parenthetical form "(Smith, 1969)"; if the same title is cited a second time this short form is repeated or, better, the reference is recast as, for example, "Smith also says. . . ." The aim is to disburden the text of apparatus—*ibid.* and its relatives. Within text or bibliography a reference may be made precise by adding, for example, "pp. 123-145" or "ch. 8." Responsibility for exact quotation lies with the writer, not the editor.

MANUSCRIPT. Double-space *all material*, on one side only of standard (8½-by-11-inch) bond paper, allowing 1½-inch margins all around. Avoid line-end division of words.

ILLUSTRATIONS, ETC. Photos should be glossy prints not less than 5 inches wide. Other kinds of illustrations should be rendered in black ink on heavy paper, preferably with labeling done expertly on a transparent overlay. Key each illustration, on the back, to its legend ("caption") written on a separate sheet—being sure to mention credits, including "photo by author." Tabular material, too, must be presented on separate sheets—regardless of length. Within the body of the manuscript simply indicate relevance at the proper place, as, "see fig. 1" or "see table."

GENERAL CONSIDERATIONS. The editor welcomes letters of inquiry describing, in some detail, articles he may wish to see. Manuscripts that arrive unannounced may be considered, but will not be returned unless accompanied by a stamped, self-addressed envelope.

We acknowledge receipt of manuscript immediately. During preparation of articles for the press we expect authors to answer queries promptly and to observe deadlines rigorously. Authors will be given two opportunities to make changes: substantially on a copy of the manuscript as edited, minimally on galley proofs.

Offprints may be purchased at rates that will be quoted by the printer.