

# Insulin and Cortisol Promote Leptin Production in Cultured Human Fat Cells

Martin Wabitsch, Per Bo Jensen, Werner F. Blum, Claus T. Christoffersen, Piera Englaro, Eberhard Heinze, Wolfgang Rascher, Walter Teller, Hans Tornqvist, and Hans Hauner

The aim of this study was to investigate the regulation of leptin expression and production in cultured human adipocytes using the model of in vitro differentiated human adipocytes. Freshly isolated human preadipocytes did not exhibit significant leptin mRNA and protein levels as assessed by reverse transcriptase (RT)-polymerase chain reaction (PCR) and radioimmunoassay (RIA). However, during differentiation induced by a defined adipogenic serum-free medium, cellular leptin mRNA and leptin protein released into the medium increased considerably in accordance with the cellular lipid accumulation. In fully differentiated human fat cells, insulin provoked a dose-dependent rise in leptin protein. Cortisol at a near physiological concentration of  $10^{-8}$  mol/l was found to potentiate this insulin effect by almost threefold. Removal of insulin and cortisol, respectively, was followed by a rapid decrease in leptin expression, which was reversible after readdition of the hormones. These results clearly indicate that both insulin and cortisol are potent and possibly physiological regulators of leptin expression in human adipose tissue. *Diabetes* 45:1435-1438, 1996

The recent identification of the *ob* gene, which encodes a 167-amino acid protein expressed exclusively in adipose tissue (1), has drawn growing attention to the investigation of body weight control, as injection of this factor, named leptin, into obese and lean animals was found to induce a reduction of food and subsequent decrease of body fat stores (2-4). Recent studies have shown that this protein is also synthesized in human adipose tissue (1,5-8) and is apparently overexpressed in human obesity (6-8). Serum leptin concentrations are correlated not only with body fat but also with serum insulin concentrations (9). To gain a better insight into the regulation of leptin expression in humans, we investigated the effects of insulin and cortisol, two important regulators of gene expression, on leptin

production in adipose tissue using the model of in vitro differentiated human adipocytes.

## RESEARCH DESIGN AND METHODS

**Subjects.** Human adipose tissue samples (wet weight 200-500 g) were obtained from the mammary adipose tissue depot of 10 healthy women (mean age  $28 \pm 7$  years [means  $\pm$  SD], mean BMI  $28.3 \pm 2.3$  kg/m<sup>2</sup>) undergoing mammary reduction. All patients gave their informed consent, and the study protocol was approved by the ethical committee of the University of Ulm.

**Cell culture.** After removal, the adipose tissue samples were stored in phosphate-buffered saline containing 20 mg/ml bovine serum albumin and immediately transferred to the laboratory. The isolation and culture of the adipose tissue derived stromal cells was performed as described previously with minor modifications (10). Cells were cultured in serum-free DME/Ham's F-12 medium (vol/vol, 1:1) supplemented with 15 mmol/l NaHCO<sub>3</sub>, 15 mmol/l HEPES, 33  $\mu$ mol/l biotin, 17  $\mu$ mol/l pantothenate, 10  $\mu$ g/ml human transferrin, 100 U/ml penicillin, and 0.1 mg/ml streptomycin (basal medium). To induce adipose differentiation, 100 nmol/l cortisol, 66 nmol/l insulin, and 200 pmol/l triiodothyronine (T<sub>3</sub>) were added (adipogenic medium). The percentage of differentiated fat cells after 16 days of culture was assessed by microscopically counting the number of lipid-filled cells in relation to total cell number in 10 randomly selected areas per dish, each area representing 1 mm<sup>2</sup>.

**Total RNA extraction and reverse transcriptase (RT)-polymerase chain reaction (PCR).** Total RNA of cultured cells was prepared according to the method of Chomczynski and Sacchi (11). The detailed description of reverse transcriptase (RT)-polymerase chain reaction (PCR) is given elsewhere (12). Briefly, total RNA was diluted to 0.2  $\mu$ g/ $\mu$ l in Tris/EDTA buffer, and first-strand cDNA was prepared using 5  $\mu$ l RNA, Superscript RT kit, and random hexamers (both Gibco/BRL) according to the instructions of the manufacturer. cDNA was diluted 1:2 with H<sub>2</sub>O and PCR carried out using 3  $\mu$ l diluted cDNA and a PCR primer mix containing 2.5 U Taq polymerase in its buffer (Promega), 40  $\mu$ mol/l of dCTP, dGTP, dTTP, 20  $\mu$ mol/l dATP, 2.5  $\mu$ Ci of 1,000-3,000 Ci/mmol ( $\alpha$ -<sup>32</sup>P)-dATP (Amersham), and 10 pmol of each primer in a 50  $\mu$ l volume overlaid with 50  $\mu$ l mineral oil. PCR conditions were a denaturation step at 95°C for 1 min followed by 23 cycles of 94°C, 30 s; 55°C, 1 min; 72°C, 1 min. PCR products were analyzed on a 6% polyacrylamide/7 M urea gel exposed to a Phosphor-Imager screen suitable for quantification.

In this semiquantitative RT-PCR method, two primer sets were used simultaneously in the same tube whereby variation in amount of cDNA and between tubes was taken into account. One primer set was specific for leptin cDNA and the other specific for the transcription factor Sp1, which is ubiquitously expressed and is necessary for the transcription of many housekeeping genes as well as for other genes and, thus, can be used as internal standard (12). When using only 23 cycles of PCR amplification, we made sure that the amplification of both leptin and Sp1 cDNA was within the exponential range, verified by running different numbers of PCR cycles (20-28 cycles). Human leptin gene specific primers: 5' primer 5'-gacttcattcctggctccacc-3', 3'-primer 5'-cctgaagctccaggacacc-3'. Human Sp1 specific primers: 5' primer 5'-gagagt-gctcacagcctgtc-3', 3' primer 5'-gttcagagcatcagaccctg-3' (13).

Since it is essential to demonstrate that only mRNA is measured by RT-PCR and not DNA contamination, we performed the PCR also in the absence of reverse transcriptase. In this case, no PCR products could be detected.

**Measurement of leptin concentration in the culture media.** Leptin was measured in the culture media at the time intervals indicated using a specific radioimmunoassay (RIA), which has been described in detail elsewhere (W.F. Blum, unpublished observation). In brief, recombinant human leptin (gift from Eli Lilly Research Laboratories, Indianapolis, IN) was used for the production of antiserum in rabbits and for the preparation of tracer by the chloramine T method (14) and of standards. The assay buffer was composed of 0.05 mol/l sodium phosphate, pH 7.4, 0.1 mol/l NaCl, 0.05% (wt:vol) NaN<sub>3</sub>, 0.1%

From the Department of Pediatrics I (M.W., E.H., W.T.), University of Ulm, Germany; the Hagedorn Research Institute (P.B.J., C.T.C., H.T.), Gentofte, Denmark; the Department of Pediatrics (W.F.B., P.E., W.R.), University of Gießen, Germany; Lilly Deutschland (W.F.B.), Bad Homburg, Germany; and the Diabetes Research Institute (H.H.), Düsseldorf, Germany.

Address correspondence and reprint requests to Dr. Martin Wabitsch, Department of Pediatrics I, University of Ulm, Prittwitzstr. 43, D-89075 Ulm, Germany.

Received for publication 20 June 1996 and accepted in revised form 1 August 1996.

PCR, polymerase chain reaction; RIA, radioimmunoassay; RT, reverse transcriptase.




FIG. 1. A: leptin mRNA expression measured by RT-PCR in human adipose tissue samples (lane 1), in isolated adipocytes (lane 2), in stromal cells isolated from adipose tissue (lane 3), in stromal cells cultured in basal medium for 16 days (lane 4), and in newly differentiated human fat cells after 16 days of culture in adipogenic medium (lane 5). B: the same data as in Fig. 1A presented as leptin/Sp1 mRNA ratio.

(vol:vol) gelatine from teleost fish (Sigma, München, Germany), 0.1% (vol:vol) triton X 100 and 0.2% normal horse serum, which showed no cross-reaction in the assay. The assay volume was 0.3 ml. After incubation at room temperature overnight, bound and unbound tracer were separated by a second antibody technique. Maximal tracer binding was 37–45% and half-maximal binding occurred at 0.9  $\mu\text{g/l}$  unlabeled leptin. Sensitivity was 0.03  $\mu\text{g/l}$ . The intra- and interassay coefficients of variation were 0.8% and 8.5%, respectively.

## RESULTS

Between 70 and 80% of the stromal cells isolated from human adipose tissue acquired adipocyte morphology within 16 days of culture upon stimulation by insulin, glucocorticoids, and  $T_3$ . Semiquantitative measurement of leptin mRNA by RT-PCR revealed that compared with freshly isolated adipocyte precursor cells substantially higher amounts of specific mRNA were measured in newly differentiated adipocytes that were comparable to the amount found in freshly isolated human adipocytes (Fig. 1). Even after 8 days of culture, a significant increase in leptin mRNA was evident, whereas cells cultured without adipogenic medium were devoid of any significant leptin mRNA expression. In the culture medium of stimulated cells, leptin production was measurable not before 8 days of culture but then increased markedly in accordance with the morphological maturation of the cells (Fig. 2). This observation suggests that the newly synthesized hormone is rapidly secreted by the fat cells and accumulates in the culture medium. On day 16, when adipocytes were fully developed, leptin production reached maximum levels and remained relatively stable at 2  $\text{ng}/10^5$  adipocytes/48 h. When cells were cultured in basal medium without adipogenic factors, adipose differentiation of the cells did not occur and leptin was undetectable in the medium (Fig. 2).

Replacement of the adipogenic medium by a medium without insulin led to a significant and rapid decrease of leptin mRNA by up to 70% and of leptin protein by up to 40% after 24 h. This decrease was more pronounced when cortisol was also removed (data not shown). Readdition of insulin at rising concentrations for another 24 h increased leptin production in a dose-dependent manner being significant at concentrations not lower than  $10^{-9}$  mol/l. If cortisol was present at a concentration of  $10^{-8}$  mol/l, the insulin effect on leptin expression was clearly potentiated. A more than threefold increase in leptin concentration

from 0.35 to 1.40  $\text{ng}/10^5$  adipocytes/24 h was observed, when a combination of  $10^{-7}$  mol/l insulin and  $10^{-8}$  mol/l cortisol was supplemented for 24 h (Fig. 3).

The effect of cortisol alone on leptin expression was also pronounced. Removal of cortisol at a persistent insulin concentration of  $10^{-8}$  mol/l for 24 h resulted in a 80% decrease in both leptin mRNA and leptin protein content in the medium. When cortisol was supplemented again to the medium at a concentration of  $10^{-8}$  mol/l, leptin mRNA and protein returned to the initial values within 24 h (Fig. 4).

## DISCUSSION

The results of this study clearly show that leptin gene expression and production in cultured human preadipocytes and adipocytes is dependent on the stage of differentiation. In fully developed human adipocytes, similar amounts of leptin mRNA were measured as in freshly isolated human fat cells. Both insulin and cortisol were found to stimulate leptin production, the former in a dose-dependent manner. It was also apparent from these experiments that changes of leptin mRNA levels are paralleled by corresponding changes of leptin in the culture medium.




FIG. 2. Leptin production by cultured human preadipocytes during adipose differentiation. Preadipocytes were cultured in adipogenic medium and leptin concentrations were measured every 48 h from day 2 until day 24 in the aspirated medium, which was renewed thereafter. Control cultures were kept in basal medium without adipogenic factors. Means  $\pm$  SE of three experiments in duplicate.




FIG. 3. Effect of rising concentrations of insulin on leptin content in the culture medium of newly differentiated human adipocytes. After differentiation, insulin was removed for 24 h and then readed for another 24 h in the absence or presence of  $10^{-8}$  mol/l cortisol. Leptin was measured in the medium by RIA. Data are expressed as means  $\pm$  SE of three experiments in duplicate.

Recent studies in animal models of obesity as well as in established adipocyte cell lines have indicated that insulin may play an important role as a regulator of leptin gene expression (15–19). This has been shown mainly at the mRNA levels. In one of these studies, it has been clearly worked out that the insulin effect takes place at the transcriptional level and is rapidly reversible (18). Our findings extend this observation by demonstrating that 1) insulin also induces an increased leptin production in adipocytes and 2) insulin is an important regulator of leptin gene expression in humans.

The assumption that insulin may be involved in the regulation of leptin expression is only partly supported by clinical studies. While Considine et al. (9) have demonstrated a clear correlation between serum leptin and insulin concentrations, recent studies using the euglycemic-hyperinsulinemic clamp technique failed to show an increase of serum leptin concentrations at least under short-term hyperinsulinemia (20,21). In addition, 24-h profiles of circulating leptin were not correlated with insulin levels (22). However, this does not exclude the possibility that the chronic hyperinsulinemia, which characterizes obesity, may cause and maintain an elevated leptin expression. On the other hand, fasting and weight loss not only are associated with low insulin levels, but also were found to lead to a substantial decline in leptin mRNA and protein levels (9).

It is also known from animal experiments and in vitro studies that glucocorticoids may be involved in the control of leptin expression. Treatment of normal rats with different types of glucocorticoids at catabolic doses rapidly induced leptin expression in adipose tissue followed by a concordant decrease in body weight gain and food intake (23). Exposure of isolated rat adipocytes to dexamethasone was reported to increase leptin mRNA within hours by four- to eightfold (24) as well as leptin secretion (25). In contrast, in 3T3 L1 adipocytes dexamethasone was found to exert a slightly inhibitory effect on leptin mRNA levels (19), indicating substantial differences in the regulation of




FIG. 4. Effect of cortisol on leptin mRNA and protein in newly differentiated human fat cells. After having developed the adipocyte phenotype (A), cortisol was removed from the culture medium for 24 h (B) and then readed for another 24 h at a concentration of  $10^{-8}$  mol/l (C). Leptin mRNA was determined by RT-PCR and expressed as leptin/Sp1 ratio (■), leptin concentration in the culture medium was measured by RIA (▨). The results of one representative experiment is shown. Leptin/Sp1 mRNA (■); Leptin (▨)

leptin expression among fat cell models. Our in vitro experiments in human adipose cells clearly suggest that cortisol is a potent stimulator of leptin gene expression and may potentiate the effect of insulin. This action is obviously independent of the differentiation-promoting effect of cortisol, since leptin gene expression was also triggered in fully differentiated fat cells and was rapidly reversible after removal of cortisol. The physiological importance of this finding remains to be demonstrated, as the reported 24-h profile of circulating serum leptin (22) does not correspond with the circadian pattern of serum cortisol levels.

In conclusion, the results of this study suggest that both insulin and cortisol potently induce leptin production in cultured human fat cells. Although these data may indicate that both hormones are contributing to the elevated leptin concentrations found in obesity, it is likely that other hormonal and/or metabolic factors are also involved in the regulation of leptin production in vivo.

#### ACKNOWLEDGMENTS

The authors gratefully acknowledge the excellent technical assistance of Simone Schmid and the help of Bettina Wabitsch.

#### REFERENCES

- Zhang Y, Proenca R, Maffei M, Barone M, Leopold L, Friedman JAM: Positional cloning of the mouse obese gene and its human homologue. *Nature* 372:425–432, 1994
- Pelleymounter MA, Cullen MJ, Baker MB, Hecht R, Winters D, Boone T, Collins F: Effects of the obese gene product on body weight regulation in *ob/ob* mice. *Science* 269:540–543, 1995
- Halaas JL, Gajiwala KS, Maffei M, Cohen SL, Chait BT, Rabinowitz D, Lallone R, Burley SK, Friedman JM: Weight-reducing effects of the plasma protein encoded by the obese gene. *Science* 269:543–546, 1995
- Campfield LA, Smith FJ, Guisez Y, Devos R, Burn P: Recombinant mouse OB protein: evidence for a peripheral signal linking adiposity and central

- neural networks. *Science* 269:546–549, 1995
5. Considine RV, Considine EL, Williams CJ, Nyce MR, Magosin SA, Bauer TL, Rosato EL, Colberg J, Caro JF: Evidence against either a premature stop codon or the absence of obese gene mRNA in human obesity. *J Clin Invest* 95:2986–2988, 1995
  6. Masuzaki H, Ogawa Y, Isse N, Satoh N, Okazaki T, Shigemoto M, Mori K, Tamura N, Hosoda K, Yoshimasa Y, Jingami H, Kawada T, Nakao K: Human obese gene expression: adipocyte-specific expression and regional differences in the adipose tissue. *Diabetes* 44:855–858, 1995
  7. Lönnquist F, Arner P, Nordfors L, Schalling M: Overexpression of the obese (*ob*) gene in adipose tissue of human obese subjects. *Nature Med* 1:950–953, 1995
  8. Hamilton BS, Paglia D, Kwan AY, Deitel M: Increased obese mRNA expression in omental fat cells from massively obese humans. *Nature Med* 9:953–956, 1995
  9. Considine RV, Sinha MK, Heiman ML, Kriauciunas A, Stephens TW, Nyce MR, Ohannesian JP, Marco CC, McKee LJ, Bauer TL, Caro JF: Serum immunoreactive leptin concentrations in normal-weight and obese humans. *N Engl J Med* 334:292–295, 1996
  10. Hauner H, Entenmann G, Wabitsch M, Gaillard D, Ailhaud G, Negrel R, Pfeiffer EF: Promoting effects of glucocorticoids on the differentiation of human adipocyte precursor cells cultured in a chemically defined medium. *J Clin Invest* 84:1663–1670, 1989
  11. Chomczynski N, Sacchi N: Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 162:156–159, 1987
  12. Jensen J, Serup P, Carlsen C, Nielsen TF, Madsen OD: mRNA profiling of rat islet tumors reveals Nkx 6.1 as a  $\beta$ -cell specific homeodomain transcription factor. *J Biol Chem*. In press
  13. Kadonaga JT, Carner KR, Masiarz FR, Tijan R: Isolation of cDNA encoding transcription factor Sp1 and functional analysis of the DNA binding domain. *Cell* 51:1079–1090, 1987
  14. Blum WF, Breier BH: Radioimmunoassays for IGFs and IFGBPs. *Growth Regulation* 4 (Suppl. 1):11–19, 1994
  15. MacDougald AO, Hwang CS, Fan H, Lane MD: Regulated expression of the obese gene product (leptin) in white adipose tissue and 3T3-L1 adipocytes. *Proc Natl Acad Sci USA* 92:9034–9037
  16. Saladin R, De Vos P, Guerre-Millo M, Leturque A, Girard J, Staels B, Auwerx J: Transient increase in obese gene expression after food intake or insulin administration. *Nature* 377:527–529, 1995
  17. Cusin I, Sainsbury A, Doyle P, Rohner-Jeanrenaud F, Jeanrenaud B: The *ob* gene and insulin: a relationship leading to clues to the understanding of obesity. *Diabetes* 44:1467–1479, 1995
  18. Leroy P, Dessolin S, Villageois P, Moon BC, Friedman JM, Ailhaud G, Dani C: Expression of *ob* gene in adipose cells: regulation by insulin. *J Biol Chem* 271:2365–2368, 1996
  19. Rentsch J, Chiesi M: Regulation of *ob* gene mRNA levels in cultured adipocytes. *FEBS Lett* 379:55–59, 1996
  20. Dagogo-Jack S, Fanelli C, Paramore D, Brothers J, Landt M: Plasma leptin and insulin relationships in obese and nonobese humans. *Diabetes* 45:695–698, 1996
  21. Kolaczynski JW, Nyce MR, Considine RV, Boden G, Nolan JJ, Henry R, Mudaliar SR, Olefsky J, Caro JF: Acute and chronic effect of insulin on leptin production in humans: studies in vivo and in vitro. *Diabetes* 45:699–701, 1996
  22. Sinha MK, Ohannesian JP, Heiman ML, Kriauciunas A, Stephens TW, Magosin S, Marco C, Caro JF: Nocturnal rise of leptin in lean, obese, and non-insulin-dependent diabetes mellitus subjects. *J Clin Invest* 97:1344–1347, 1996
  23. De Vos P, Saladin R, Auwerx J, Staels B: Induction of *ob* gene expression by corticosteroids is accompanied by body weight loss and reduced food intake. *J Biol Chem* 270:15958–15961, 1995
  24. Murakami T, Iida M, Shima K: Dexamethasone regulates obese expression in isolated rat adipocytes. *Biochem Biophys Res Comm* 214:1260–1267, 1995
  25. Sliker LJ, Sloop KW, Surface PL, Kriauciunas A, LaQuier F, Manetta J, Bue-Valleskey J, Stephens TW: Regulation of expression of *ob* mRNA and protein by glucocorticoids and cAMP. *J Biol Chem* 271:5301–5304, 1996