

Transmission of *Helicobacter pylori* and the role of water and biofilms

Steven L. Percival and John G. Thomas

ABSTRACT

Documented evidence relating to the survival of *Helicobacter pylori* outside the gastric niche is extremely limited. To date the primary transmission routes of *H. pylori* have yet to be confirmed and when this is achieved preventive infection control measures can be implemented to reduce and ultimately prevent human infection from this pathogen. There is mounting evidence which suggests that the prevalence of *H. pylori* infection has a strong correlation with access to clean water, suggesting a transmission route to the host. However, there are no established culture methods for the detection of viable *H. pylori* in the environment, in particular drinking water supplies, preventing the development of true epidemiological and risk assessments. The aim of this review is to highlight the available data to date that suggests drinking water and possible survival in biofilms as a probable transmission mode for *H. pylori*.

Key words | biofilm, drinking water, *Helicobacter pylori*, viable but non-culturable (VBNC)

Steven L. Percival (corresponding author)
John G. Thomas
Department of Pathology,
West Virginia University School of Medicine,
Robert C. Byrd Health Sciences Center-North,
Morgantown,
WV 26506-9203,
USA
E-mail: biofilms@hotmail.co.uk

INTRODUCTION

Helicobacter pylori are micro-aerophilic spiral-shaped bacteria that efficiently colonize the human gastric mucosa (Moreno *et al.* 2007). They were first identified in autopsied rabbits in 1893 (Rothenbacher & Brenner 2003), described in humans in 1906 (Rothenbacher & Brenner 2003), and successfully cultured in 1983 (Warren & Marshall 1983). Initially, research-based studies classified *H. pylori*, based on its 'Campylobacter like' morphology and biochemistry, as *Campylobacter pyloridis* (Warren & Marshall 1983). Later the name of the bacterium was changed to *Campylobacter pylori* and then finally classified as *Helicobacter pylori* some years later (Owen 1995).

Helicobacter pylori is Latin for 'spiral rod of the lower part of the stomach'. It is known to cause inflammation in the form of chronic active gastritis, specifically in certain humans. In addition, *H. pylori* has been linked to a diverse spectrum of gastrointestinal disorders which include peptic ulcer disease, gastric adenocarcinoma and gastric mucosa-associated lymphoid tissue (MALT) lymphoma (Farinha & Gascoyne 2005). MALT lymphoma is a relatively rare

condition caused by *H. pylori* and only very small percentages (1–2%) of infected individuals develop a malignant disease.

Clinical disease typically occurs decades after initial acquisition of infection. However, gastritis may progress over time from an initially superficial nonatrophic form to a more severe atrophic gastritis, with intestinal metaplasia, leading to duodenal ulceration, gastric adenocarcinoma and gastric MALT lymphoma (Farinha & Gascoyne 2005). Because of the vast array of conditions and public health concerns that surround *H. pylori*, the World Health Organization International Agency for Research on Cancer classified *H. pylori* as a class I carcinogen in humans (Sherman 2004). While only a small percentage of individuals carrying *H. pylori* ever develop clinical sequelae, asymptomatic carriage is common (Peterson & Krogfelt 2003) and if left untreated *H. pylori* infection is lifelong (Czinn 2005).

Although it is estimated that approximately 50% of the world's population is colonized with *H. pylori*, prevalence varies widely by age as well as country, ethnic background and socio-economic conditions (Czinn 2005).

The developing world carries the greatest burden, with over 70% of children infected by age 15 (Czinn 2005). In contrast the developed world has, since the 1950s, experienced a decrease in the prevalence of *H. pylori* with each successive generation; hence at present approximately 20% to 30% of individuals harbour *H. pylori* (Rothenbacher & Brenner 2003). This reduction has been attributed to improved socio-economic status and personal hygiene but may equally have transpired because of improvements in drinking water quality.

The aim of this review is to critically evaluate published data and determine whether water and biofilms may constitute possible transmission and environmental niches for *H. pylori*. Consequently, by establishing water and biofilms as a transmission route for *H. pylori* the development of appropriate control measures to reduce the incidence and prevalence of *H. pylori* even further is of great significance, particularly in developing countries.

TRANSMISSION ROUTES OF *H. PYLORI* AND ACQUISITION

Although the natural niche for *H. pylori* is the human stomach, for widespread infection to occur the organism may need to survive in the external environment (Brown 2000). To date the precise mechanism/s involved in the transmission of *H. pylori* is/are unknown, but clearly any approach that introduces the organism into the stomach of a susceptible person may lead to that individual acquiring an infection. Many transmission routes for *H. pylori* have been proposed and have included gastric-oral (Raymond *et al.* 2008), oral-oral (Mégraud 1995), faecal-oral (Raymond *et al.* 2008), zoonotic (Fox 1995) and water/food-borne (Hulten *et al.* 1996; Herrera 2004). Clearly these proposed transmission routes indicate that *H. pylori* infection occurs through multiple acquisition pathways (Goodman & Correa 1995; Velazquez & Feirtag 1999). Despite these proposals, contamination of food by human faecal material has been found to be one of the major risk factors for the acquisition of *H. pylori* (Hopkins *et al.* 1993).

H. pylori has been cultured from the faeces of infected individuals (Thomas *et al.* 1992; Kelly *et al.* 1994) and specific DNA sequences have been amplified from raw

sewage (Forrest *et al.* 1998) providing possible evidence for the faecal-oral route of transmission. However, the faecal-oral route of transmission and its interpretation has been questioned, as the concentration of *H. pylori* cells present in faecal material is considered to be low, particularly when compared with other faecal pathogens of public health significance (Vincent 1995). Furthermore, the prevalence of *H. pylori* IgG antibodies in sewage workers, compared with a control group, matched for age and socio-economic status, has demonstrated no increased risk of infection when exposure to human faecal material has occurred (Friis *et al.* 1996). Despite controversy in this area there is growing evidence that suggests a faecal-oral transmission route for *H. pylori* (Mladenova *et al.* 2006). Consequently, if *H. pylori* is excreted within faeces they may well go on to colonize surfaces present in water sources. Such surfaces subsequently then become transmittable sources of *H. pylori* (Xia & Talley 1997).

The age at which humans are exposed to *H. pylori* may influence its route of transmission. For example, in an Argentinean study, the key risk factor for acquisition of *H. pylori*, specifically in childhood, was the nature of the water source (Olmos *et al.* 2000). In developing countries, many children are infected with *H. pylori* by the age of ten, and relapses are known to occur (Dooley *et al.* 1989; Gurel *et al.* 1999). In Peruvian children *H. pylori* prevalence was found to strongly correlate with socio-economic status: children were found to be three times more likely to be infected with *H. pylori* when they drank from an external water source than those exposed to water from an internal water source (Klein *et al.* 1991). However, no difference was found when those children from high- and low-income families with an internal water source were compared (Klein *et al.* 1991). Children born into high-income families supplied with municipal water are considered 12 times more likely to become colonized with *H. pylori* than those supplied from community wells (Frenck & Clemens 2003). This suggests that municipal water is a possible risk factor in the transmission and acquisition of *H. pylori*. It is plausible to suggest that breaks in municipal pipes allow for infiltration of contaminated surface water (Frenck & Clemens 2003). A group of 3,289 residents in Italy were screened for the prevalence of *H. pylori* IgG antibodies (Dominici *et al.* 1999). The conclusion drawn from this

study was that there was a common source of exposure to *H. pylori* in the environment (Dominici *et al.* 1999).

Some research findings have shown that hands and fingernails contaminated with *H. pylori* may transfer bacteria into a water source (Dowsett *et al.* 1999; Frenck & Clemens 2003). Evidence of *H. pylori* presence in dental plaque has led to the suggestion by a number of researchers that the oral cavity may be a potential reservoir for this bacteria in the adult population (Luman *et al.* 1996; Cave 1997; Kamat *et al.* 1998; Oshowo *et al.* 1998a,b). Souton & Colombo (2008) found that the prevalence of *H. pylori* can be as high as 33.3% in subgingival biofilms and 20% in saliva. Despite this it still remains unclear whether the oral cavity acts as a permanent reservoir for *H. pylori* specifically as within this oral ecosystem *H. pylori* is considered to be transient.

In addition to humans, domestic cats and Old World macaques have been found to be colonized with *H. pylori* but it is doubtful whether these animals provide an important reservoir for human infection (Osata *et al.* 1997; Baker & Hegarty 2001). Flies have also been considered as having a potential role in the vectorial spread of *H. pylori* from human faeces to food (Grubel *et al.* 1998).

TRANSMISSION OF *H. PYLORI* IN WATER

In developing countries, water rather than person-to-person spread plays a significant role in the transmission of *H. pylori* (Akcem *et al.* 2000). Water from streams, rivers and wells has been considered as a common source (Hulten *et al.* 1991, 1996; Klein *et al.* 1991; Mackay *et al.* 1998; Hegarty *et al.* 1999; Engstrand 2001; Mackay *et al.* 2001; Mazari-Hiriart *et al.* 2001; Lu *et al.* 2002; Imanishi *et al.* 2003; Karita *et al.* 2003; Azevedo *et al.* 2004; Gomes & De Martinis 2004; Rolle-Kampczyk *et al.* 2004). In Brazil, Zaterka & colleagues (2007) confirmed that the source of drinking water in childhood was a risk factor for *H. pylori* infection and that the prevalence of *H. pylori* infection was higher when a local river was the source of drinking water and lower when this water was filtered or boiled. In addition, Goodman & colleagues (1996) found that swimming in streams, using streams as a drinking water source, and frequent consumption of raw vegetables, cleaned in contaminated water, increased the likelihood of infection

with *H. pylori*. Other research also supports an association between *H. pylori* infection and consumption of untreated well or spring water (Carballo *et al.* 1997; Benson *et al.* 2004; Reavis 2005). Further evidence concerning the importance of water as a transmission route of *H. pylori* was stressed by Fujimura & colleagues (2004) who collected and analysed a total of 24 water samples from the upper, middle and downstream reaches of four Japanese rivers for evidence of *H. pylori* by nested polymerase chain reaction (PCR). The conclusion from this study suggested that water, in the natural environment, could be a risk factor for *H. pylori* transmission (Fujimura *et al.* 2004).

PERSISTENCE, DETECTION AND CULTURABILITY OF *H. PYLORI*

Despite the numerous research findings identifying *H. pylori* in water, it is important to consider the fact that the use of PCR and other molecular methods for the detection of pathogens in environmental samples has limitations. This is principally due to the inability of PCR to differentiate between naked DNA from dead and living cells. Furthermore, the natural environment contains many microorganisms which have not yet been identified or cultured, which may interfere with molecular technologies. Consequently, to scientifically interpret data regarding the epidemiology of *H. pylori*, cultured bacteria from appropriate water sources are necessary.

The first successful isolation of *H. pylori*, by culturable methods, occurred in a municipal wastewater canal on the US–Mexico border (Lu *et al.* 2002). This canal was found to be heavily contaminated with untreated raw sewage, in an area known to have a high *H. pylori* prevalence (Lu *et al.* 2002). Aside from this study, positive culture of *H. pylori* from drinking water has not been successful, despite efforts to produce a culture-specific media sensitive and selective enough to isolate and grow this organism. A simple plating medium for the detection of *H. pylori* in the environment was investigated by Degnan *et al.* (2003) and Fernández *et al.* (2007). However, the culturable methods employed were unsuccessful in the culturing of *H. pylori*.

H. pylori rapidly transforms into a viable but non-culturable state (VBNC). This state is induced by low

nutrient and hyperosmotic conditions (Mizoguchi *et al.* 1999; Zheng *et al.* 1999; Moreno *et al.* 2003). Such stressed conditions are commonly found in water and the environment. Fluorescent *in situ* hybridization (FISH) with rRNA oligonucleotide probes has been used for detection and identification of VBNC forms of bacteria (Rowan 2004). The role of the VBNC of *H. pylori*, and its associated forms, in infection and transmission remains unclear (Rowan 2004). However, what is significant is that within the VBNC state *H. pylori* cells are still alive (Moreno *et al.* 2003; Rolle-Kampczyk *et al.* 2004; Rowan 2004). Ultimately this may, when evidence becomes available, have important implications for the survival and therefore the potential infectivity of *H. pylori*. To date, little or no evidence exists regarding the resuscitation of VBNC cells of *H. pylori* or on the ability of the VBNC cells to cause infection.

Initial evidence of *H. pylori* presence in environmental water samples has come from PCR amplification of samples obtained from Colombia, where infection rates are over 90% (Handwerker *et al.* 1995). In addition to this a number of PCR assays have been utilized over the years for the detection of *H. pylori* in water (Engstrand *et al.* 1992; Weiss *et al.* 1994; Hulten *et al.* 1996; Sasaki *et al.* 1999; Benson *et al.* 2004; Gomes & De Martinis 2004; Shahamat *et al.* 2004). Furthermore, in the United States, actively respiring *H. pylori* from surface and well water has been detected using fluorescent antibody-tetrazolium reduction (FACTC) microscopy (Hegarty *et al.* 1999) and confirmed using species-specific PCR (Azevedo *et al.* 2006a). Sen *et al.* (2007) investigated the development of internal controls for PCR assays by spiking drinking water with 100 cells of *H. pylori* and demonstrated similar cycle thresholds to those of recombinant *Escherichia coli* during chlorine disinfection. In addition to PCR, FISH was validated as a quick and sensitive method for detection of *H. pylori* in environmental samples (Moreno *et al.* 2003). These findings suggest the presence of *H. pylori* in the natural environment and a possible waterborne route of transmission. Nayak & Rose (2007) demonstrated that quantitative polymerase chain reaction (qPCR) could determine *H. pylori* concentrations in water. In this study real time qPCR was shown to be a specific, sensitive and rapid method to quantify *H. pylori* in sewage. Prior to these studies a two-stage *in vitro* method for detection of *H. pylori* in spiked water and faecal samples

using immunomagnetic separation followed by PCR detection (IMS/PCR) has been described (Enroth & Engstrand 1995).

Numerous studies have shown that *H. pylori* may survive for prolonged periods in water over a range of physical variables (West *et al.* 1992). In fact *H. pylori* strains have been shown to survive for long periods under physiological saline concentrations, low temperatures and a pH range of 5.8 to 6.9. A study by Shahamat & colleagues (1993) has found that *H. pylori* were able to remain viable for periods ranging from 48 hours to between 20 and 30 days when exposed to different temperatures.

A study in Leipzig, Germany, showed a positive correlation between the drinking of *H. pylori*-contaminated well water and the acquisition of a *H. pylori* infection (Rolle-Kampczyk *et al.* 2004). *H. pylori* DNA has been amplified from drinking water samples in Japan (Sasaki *et al.* 1999), Mexico (Mazari-Hiriart *et al.* 2001) and Peru (Hulten *et al.* 1996), from water samples taken from a delivery truck in the Canadian Arctic (McKeown *et al.* 1999) and from drinking water storage pots in the Gambia (Bunn *et al.* 2002). A study by Hulten *et al.* (1998) used two PCR assays to examine municipal treated and well water samples from all 25 counties of Sweden for the presence of *Helicobacter* DNA: 37.5% of wells, 12% of municipal sources and 12% of wastewater samples were found to be positive for *Helicobacter* DNA.

The argument for a waterborne route of *H. pylori* transmission is supported by the maintenance of viability in spiked natural water (West *et al.* 1990, 1992; Shahamat *et al.* 1993; Hunter 1997; Fan *et al.* 1998; Jiang & Doyle 1998; Sato *et al.* 1999; MMWR 1999). While attempts to culture *H. pylori* from environmental water samples have been largely unsuccessful, closely related microaerophilic organisms, *Campylobacter jejuni* and *Arcobacter butzleri*, have been cultured from ground and surface waters (Arvanitidou *et al.* 1994; Stanley *et al.* 1998; Rice *et al.* 1999) and associated with waterborne outbreaks (MMWR 1999). As mentioned previously, viable *H. pylori* cells could be transmitted through faecal material (Thomas *et al.* 1992), which may well provide a route for contaminating drinking water. *H. pylori* have been shown to survive for short periods in water when present in their coccoid morphology (Mizoguchi *et al.* 1999; She *et al.* 2003). This coccoid form of *H. pylori*, because of its increased tolerance to outside

perturbations, may allow the bacteria to survive the extremes of conditions associated with drinking water and water distribution systems. In addition, it has been speculated that *H. pylori* coccoid cells may be able to tolerate the levels of disinfectant normally used in distribution systems and therefore remain viable (Azevedo *et al.* 2008). To date the survival of *H. pylori* is poorly understood in the aquatic environment; likewise we do not know how this environment affects its viability.

DISINFECTION DATA

Present data concerning the effectiveness of standard drinking water disinfection processes on *H. pylori* are limited by the number of published articles in this area. Baker & colleague (2002) found that *H. pylori* were more resistant to low levels of free chlorine than *E. coli* or *C. jejuni*. Conclusions from this research have highlighted the fact that it was possible, under conditions of inadequate disinfection, for *H. pylori* to persist in water. Based on the studies by Johnson *et al.* (1997) and Baker *et al.* (2002) it is possible that reduced chlorine residuals might not provide adequate inactivation of *H. pylori*. Consequently this would not prevent the entry and persistence of *H. pylori* in drinking water systems. This may be particularly so if the bacterium grows within a biofilm state.

A number of drinking water studies have identified *H. pylori* in water pre- and post-chlorination (Mazari-Hiriart *et al.* 2003). Moreno & colleagues (2007) have shown that *H. pylori* could survive disinfection practices that are normally used in drinking water treatment when *H. pylori* are found in the VBNC state. However, they did find that culture of *H. pylori* was lost after 5 min in water despite free chlorine levels of 0.96 mg l⁻¹.

BIOFILMS

It is possible that there are environmentally adapted forms of *H. pylori* within a biofilm community. *H. pylori* readily form biofilms (Carron *et al.* 2006) and in so doing produce a novel antibacterial peptide, which may confer increased persistence in a heterogeneous biofilm environment (Putsep *et al.* 1999). *H. pylori* is also known to produce a water-insoluble

biofilm when grown under high carbon:nitrogen ratio conditions (Stark *et al.* 1999). However, environmentally adapted *H. pylori* have to be isolated and cultured for an evaluation of their true environmental persistence, virulence and ability to be transmitted in water (Vincent 1995).

It is well known that waterborne bacteria can attach to surfaces by aggregating in a hydrated exopolymer known as a biofilm (Costerton *et al.* 1999; Percival *et al.* 2000). The association of bacteria, particularly pathogens, with biofilm communities within a water distribution system may offer vulnerable and susceptible bacteria protection from disinfection and protozoan predation (Sibille *et al.* 1998). In fact microorganisms in drinking water are predominantly associated with biofilms rather than in the planktonic state (Costerton *et al.* 1999; Percival *et al.* 2000, 2004). Consequently, for *H. pylori* to survive the extremes of water it is probable that it would have to reside within a biofilm. In addition to this, *H. pylori* has recently been shown to remain viable and proliferate inside *Acanthamoeba castellanii* for up to 8 weeks when evident in a co-culture (Winiecka-Krusnell *et al.* 2002). Various species of Acanthamoebae are common components of drinking water biofilms and have been shown to shield bacteria from disinfectants and enhance their proliferation when evident in biofilms (Greub & Raoult 2004).

There is evidence that biofilms in water distribution systems may harbour *H. pylori* (Mackay *et al.* 1999; Mackay *et al.* 2001; Park *et al.* 2001). In addition a study undertaken in Western Africa, utilizing 16S rDNA sequences, has shown evidence that *H. pylori* can be detected in natural biofilms (Bunn *et al.* 2002). A more recent study by Watson & colleagues (2004) showed a close link between *Helicobacter* DNA in showerhead biofilm used in domestic households. Furthermore *H. pylori* has been found to have the ability to incorporate itself into lab-grown biofilms (Mackay *et al.* 1999; Azevedo *et al.* 2003). The study by Mackay & colleagues (1999) concluded that *H. pylori* could incorporate itself, and persist in a laboratory-based, mixed-species heterotrophic biofilm, for over 8 days. However, Watson *et al.* (2004) concluded from their study that monitoring the cold water supply to a particular property did not appear to be a reliable means of predicting the *Helicobacter* status of the water distribution system in that property. Azevedo *et al.* (2006a,b) and Bragança *et al.* (2007) have also shown that *H. pylori* may

be present as biofilms on pipe work in drinking water systems. In the study by Azavedo & colleagues (2006a) *H. pylori* was shown to have the ability to adhere to different plumbing materials, namely copper and stainless steel. Copper surfaces were found to be especially suitable for the maintenance of the bacteria in the spiral form.

CONCLUSION

Although inconclusive, epidemiological studies strongly suggest person-to-person transmission of *H. pylori* (Raymond *et al.* 2008). However, recent experimental findings suggest that *H. pylori* transmission may involve the consumption of contaminated drinking water (Bellack *et al.* 2006). Although *H. pylori* is not classified as a food or waterborne pathogen, it is acknowledged as an important pathogen. The ability of *H. pylori* to survive in a coccoid VBNC form and its survival correlated with water should be of interest and possible concern to epidemiologists and public health microbiologists.

Outside the developing world the evidence for waterborne transmission of *H. pylori* is sparse. However, recent evidence has shown that *H. pylori* can survive in water for over 96 hours at 25°C (Azevedo *et al.* 2008). While this timescale is relatively short, it is evident in the developing world that this would be long enough for *H. pylori* transmission and possible infection of a compromised individual to occur.

H. pylori remains a significant problem in the developing world and will continue to be a concern as a result of the poor level of sanitation and hygiene that exists in these regions of the world. Subsequently the importance of *H. pylori* as a primary pathogen is unlikely to diminish in the foreseeable future in these vulnerable areas. Consequently, it remains important that we acquire a better understanding of the risk factors associated with the acquisition of *H. pylori* infection but at the same time we should not overlook the increasing potential that *H. pylori* can be transmitted via a waterborne pathway. Accordingly the 'environmental' biofilm in which *H. pylori* may reside, resuscitate, proliferate, interact socially with other sessile microorganisms and then disseminate warrants further investigation as a source of infection (Gião *et al.* 2008).

REFERENCES

- Akcam, Y., Ersan, S., Alper, M., Bicik, Z. & Aytug, N. 2000 The transmission of *Helicobacter pylori* via exposure to common sources outweighs the person-to-person contact among spouses in developing countries. *Am. J. Gastroenterol.* **95**, 317–319.
- Arvanitidou, M., Stathopoulos, G. A. & Katsouyannopoulos, V. C. 1994 Isolation of *Campylobacter* and *Yersinia* spp. from drinking waters. *J. Travel Med.* **119**, 156–159.
- Azevedo, N. F., Vieira, M. J. & Keevil, C. W. 2003 Establishment of a continuous model system to study *Helicobacter pylori* survival in potable water biofilms. *Water Sci. Technol.* **47**, 155–160.
- Azevedo, N. F., Pacheco, A. P., Vieira, M. J. & Keevil, C. W. 2004 Nutrient shock and incubation atmosphere influence recovery of culturable *Helicobacter pylori* from water. *Appl. Environ. Microbiol.* **70**, 490–493.
- Azevedo, N. F., Pacheco, A. P., Keevil, C. W. & Vieira, M. J. 2006a Adhesion of water stressed *Helicobacter pylori* to abiotic surfaces. *J. Appl. Microbiol.* **101**, 718–724.
- Azevedo, N. F., Pinto, A. R., Reis, N. M., Vieira, M. J. & Keevil, C. W. 2006b Shear stress, temperature, and inoculation concentration influence the adhesion of water-stressed *Helicobacter pylori* to stainless steel 304 and polypropylene. *Appl. Environ. Microbiol.* **72**, 2936–2941.
- Azevedo, N. F., Almeida, C., Fernandes, I., Cerqueira, L., Dias, S., Keevil, C. W. & Vieira, M. J. 2008 Survival of gastric and enterohepatic *Helicobacter* spp. in water: implications for transmission. *Appl. Environ. Microbiol.* **74**, 1805–1811.
- Baker, K. H. & Hegarty, J. P. 2001 Presence of *Helicobacter pylori* in drinking water is associated with clinical infection. *Scand. J. Infect. Dis.* **33**, 744–746.
- Baker, K. H., Hegarty, J. P., Redmond, B., Reed, N. A. & Herson, D. S. 2002 Effect of oxidizing disinfectants (chlorine monochloramine and ozone) on *Helicobacter pylori*. *Appl. Environ. Microbiol.* **68**, 981–984.
- Bellack, N. R., Koehoorn, M. W., MacNab, Y. C. & Morshed, M. G. 2006 A conceptual model of water's role as a reservoir in *Helicobacter pylori* transmission: a review of the evidence. *Epidemiol. Infect.* **134**, 439–449.
- Benson, J. A., Fode-Vaughan, K. A. & Collins, M. L. 2004 Detection of *Helicobacter pylori* in water by direct PCR. *Lett. Appl. Microbiol.* **39**, 221–225.
- Brown, L. M. 2000 *Helicobacter pylori*: epidemiology and routes of transmission. *Epidemiol. Rev.* **22**, 283–297.
- Bragança, S. M., Azevedo, N. F., Simões, L. C., Keevil, C. W. & Vieira, M. J. 2007 Use of fluorescent in situ hybridisation for the visualisation of *Helicobacter pylori* in real drinking water biofilms. *Water Sci. Technol.* **55**(8–9), 387–393.
- Bunn, J. E., MacKay, W. G., Thomas, J. E., Reid, D. C. & Weaver, L. T. 2002 Detection of *Helicobacter pylori* DNA in drinking water biofilms: implications for transmission in early life. *Lett. Appl. Microbiol.* **34**, 450–454.
- Carballo, F., Caballero, P., Parra, T., Aldeguez, J. M. & Pajares, J. M. 1997 Untreated drinking water is a source of *H. pylori*

- infection. European *Helicobacter pylori* Study Group, *10th International Workshop Abstracts*, 12–14 September, Lisbon, Portugal, p. A39.
- Carron, M. A., Tran, V. R., Sugawa, C. & Coticchia, J. M. 2006 Identification of *Helicobacter pylori* biofilms in human gastric mucosa. *J. Gastrointest. Surg.* **10**, 712–717.
- Cave, D. R. 1997 How *Helicobacter pylori* transmitted? *Gastroenterology* **113**, S14.
- Costerton, J. W., Stewart, P. S. & Greenberg, E. P. 1999 Bacterial biofilms: a common cause of persistent infections. *Science* **284**, 1318–1322.
- Czinn, S. J. 2005 *Helicobacter pylori* infection: detection investigation and management. *J. Pediatr.* **146**, S21–S26.
- Degnan, A. J., Sonzogni, W. C. & Standridge, J. H. 2003 Development of a plating medium for selection of *Helicobacter pylori* from water samples. *Appl. Environ. Microbiol.* **69**, 2914–2918.
- Dominici, P., Bellentani, S., Di Biase, A. R., Saccoccio, G., Le Rose, A., Masutti, F., Viola, L., Balli, F., Tiribelli, C., Grilli, R., Fusillo, M. & Grossi, E. 1999 Familial clustering of *Helicobacter pylori* infection: population based study. *Brit. Med. J.* **319**, 537–541.
- Dooley, C. P., Cohen, H. & Fitzgibbons, P. L. 1989 Prevalence of *Helicobacter pylori* infection and histologic gastritis in asymptomatic persons. *New Engl. J. Med.* **321**, 1562–1566.
- Dowsett, S. A., Archila, L., Segreto, V. A., Gonzalez, C. R., Silva, A., Vastola, K. A., Bartizek, R. D. & Kowolik, M. J. 1999 *Helicobacter pylori* infection in indigenous families of Central America: serostatus and oral and fingernail carriage. *J. Clin. Microbiol.* **37**, 2456–2460.
- Engstrand, L. 2001 *Helicobacter* in water and waterborne routes of transmission. *J. Appl. Microbiol.* **90**, 80S–84S.
- Engstrand, L., Nguyen, A. M., Graham, D. Y. & el-Zaatari, F. A. 1992 Reverse transcription and polymerase chain reaction amplification of rRNA for detection of *Helicobacter* species. *J. Clin. Microbiol.* **30**, 2295–2301.
- Enroth, H. & Engstrand, L. 1995 Immunomagnetic separation and PCR for detection of *Helicobacter pylori* in water and stool specimens. *J. Clin. Microbiol.* **33**, 2162–2165.
- Fan, X. G., Chua, A., Li, T. G. & Zeng, Q. S. 1998 Survival of *Helicobacter pylori* in milk and tap water. *J. Gastroenterol. Hepatol.* **13**, 1096–1098.
- Farinha, P. & Gascoyne, R. D. 2005 *Helicobacter pylori* and MALT Lymphoma. *Gastroenterology* **128**, 1579–1605.
- Fernández, M., Contreras, M., Suárez, P., Gueneau, P. & García-Amado, M. A. 2007 Use of HP selective medium to detect *Helicobacter pylori* associated with other enteric bacteria in seawater and marine molluscs. *Lett. Appl. Microbiol.* **45**, 213–218.
- Forrest, K., Stinson, M. & Wright, S. M. 1998 The presence of *Helicobacter pylori* in sewage. *Abstracts of the 98th General Meeting of the American Society for Microbiology*, Atlanta, Georgia, 17–20 May 1998, p. 445.
- Fox, J. G. 1995 Non-human reservoirs of *Helicobacter pylori*. *Aliment. Pharm. Therap.* **9**(Suppl. 2), 93–103.
- Frencik, R. W. & Clemens, J. 2003 *Helicobacter* in the developing world. *Microbes Infect.* **5**, 705–713.
- Friedman, B. 1998 European *Helicobacter Pylori* study group. XIth international workshop on gastroduodenal pathology and *Helicobacter pylori*. *Gut* **43**(Suppl. 2), 2.
- Friis, L., Engstrand, L. & Edling, C. 1996 Prevalence of *Helicobacter pylori* infection among sewage workers. *Scand. J. Work Environ. Health* **22**, 364–365.
- Fujimura, S., Kato, S. & Kawamura, T. 2004 *Helicobacter pylori* in Japanese river water and its prevalence in Japanese children. *Lett. Appl. Microbiol.* **38**, 517–521.
- Gião, M. S., Azevedo, N. F., Wilks, S. A., Vieira, M. J. & Keevil, C. W. 2008 Persistence of *Helicobacter pylori* in heterotrophic drinking water biofilms. *Appl. Environ. Microbiol.* **74**(19), 5898–5904.
- Gomes, B. G. & De Martinis, E. C. P. 2004 The significance of *Helicobacter pylori* in water food and environmental samples. *Food Control* **15**, 397–403.
- Goodman, K. J. & Correa, P. 1995 The transmission of *Helicobacter pylori*. A critical review of the evidence. *Int. J. Epidemiol.* **24**, 875–887.
- Goodman, K. J., Correa, P., Tengana Aux, H. J., Ramirez, H., DeLany, J. P., Guerrero Pepinosa, O., Lopez Quinones, M. & Collazos Parra, T. 1996 *Helicobacter pylori* infection in the Colombian Andes: a population-based study of transmission pathways. *Am. J. Epidemiol.* **144**, 290–299.
- Greub, G. & Raoult, D. 2004 Microorganisms resistant to free-living amoebae. *Clin. Microbiol. Rev.* **17**, 413–433.
- Grubel, P., Huang, L., Masubuchi, N., Stutzenberger, F. J. & Cave, D. R. 1998 Detection of *Helicobacter pylori* DNA in houseflies (*Musca domestica*) on three continents. *Lancet* **352**, 788–789.
- Gurel, S., Besisk, F., Demir, K., Mungan, Z., Kaymakoglu, S., Boztas, G., Cakaloglu, Y., Yeginsu, O. & Okten, A. 1999 After the eradication of *Helicobacter pylori* infection, relapse is a serious problem in Turkey. *J. Clin. Gastroenterol.* **28**, 241–244.
- Handwerker, J., Fox, J. G. & Schauer, D. B. 1995 Detection of *Helicobacter pylori* in drinking water using polymerase chain reaction amplification. *Abstracts of the 95th General Meeting of the American Society for Microbiology*, Washington, DC, 21–25 May 1995, p. 435.
- Hegarty, J. P., Dowd, M. T. & Baker, K. H. 1999 Occurrence of *Helicobacter pylori* in surface water in the United States. *J. Appl. Microbiol.* **87**, 697–701.
- Herrera, A. G. 2004 *Helicobacter pylori* and food products: a public health problem. *Method. Mol. Biol.* **268**, 297–301.
- Hopkins, R. J., Vial, P. A., Ferreccio, C., Ovalle, J., Prado, P., Sotomayor, V., Russell, R. G., Wasserman, S. S. & Morris, J. G., Jr 1993 Seroprevalence of *Helicobacter pylori* in Chile: vegetables serve as one route of transmission. *J. Infect. Dis.* **168**, 222–226.
- Hulten, K., Han, S. W. & El-Zatari, F. A. K. 1991 Detection of *Helicobacter pylori* in Peruvian water sources by two PCR assays based on independent genes. *Lancet* **227**, 15–23.
- Hulten, K., Han, S. W., Enroth, H., Klein, P. D., Opekun, A. R., Gilman, R. H., Evans, D. G., Engstrand, L., Graham, D. Y. &

- El-Zaatari, F. A. K. 1996 *Helicobacter pylori* in the drinking water in Peru. *Gastroenterology* **110**, 1031–1035.
- Hulten, K., Enroth, H., Nyström, T. & Engstrand, L. 1998 Presence of *Helicobacter* species DNA in Swedish well water. *J. Appl. Microbiol.* **85**, 282–286.
- Hunter, P. R. 1997 *Waterborne Disease*. John Wiley & Sons, New York, pp. 202–205.
- Imanishi, Y., Ogata, T., Matsuzuka, A., Tasaki, T., Fujioka, T., Akashi, M., Makino, Y. & Nishizono, A. 2003 Possibility for the presence of *Helicobacter pylori* in drinking well water Kansenshogaku Zasshi. *J. Jpn Assoc. Infect. Dis.* **77**, 18–23.
- Jiang, X. & Doyle, M. P. 1998 Effect of environmental and substrate factors on survival and growth of *Helicobacter pylori*. *J. Food Prot.* **61**, 929–933.
- Johnson, C. H., Rice, E. W. & Reasoner, D. J. 1997 Inactivation of *Helicobacter pylori* by chlorine. *Appl. Environ. Microbiol.* **63**, 4969–4970.
- Kamat, A. H., Mehta, P. R., Natu, A. A., Phadke, A. Y., Vora, I. M., Desai, P. D. & Koppikar, G. V. 1998 Dental plaque: an unlikely reservoir of *Helicobacter pylori*. *Indian J. Gastroenterol.* **17**, 138–140.
- Karita, M., Teramukai, S. & Matsumoto, S. 2003 Risk of *Helicobacter pylori* transmission from drinking well water is higher than that from infected intrafamilial members in Japan. *Digest. Dis. Sci.* **48**, 1062–1067.
- Kelly, S. M., Pitcher, M. C., Farmery, S. M. & Gibson, G. R. 1994 Isolation of *Helicobacter pylori* from feces of patients with dyspepsia in the United Kingdom. *Gastroenterology* **107**, 1671–1674.
- Klein, P. D., Graham, D. Y., Gaillour, A., Opekun, A. R. & Smith, E. O. 1991 Water source as a risk factor for *Helicobacter pylori* in Peruvian children. *Lancet* **337**, 1503–1506.
- Lu, Y., Redlinger, T. E., Avitia, R., Galindo, A. & Goodman, K. 2002 Isolation and genotyping of *Helicobacter pylori* from untreated municipal wastewater. *Appl. Environ. Microbiol.* **68**, 1436–1439.
- Luman, W., Alkout, A. M., Blackwell, C. C., Weir, D. M. & Plamer, K. R. 1996 *Helicobacter pylori* in the mouth: negative isolation from dental plaque and saliva. *Eur. J. Gastroenterol. Hepatol.* **8**, 11–14.
- Mackay, W. G., Gribbon, L. T., Barer, M. R. & Reid, D. C. 1998 Biofilms in drinking water systems: a possible reservoir for *Helicobacter pylori*. *Water Sci. Technol.* **38**, 181–185.
- Mackay, W. G., Gribbon, L. T., Barer, M. R. & Reid, D. C. 1999 Biofilms in drinking water systems: a possible reservoir for *Helicobacter pylori*. *J. Appl. Microbiol.* **85**, 52s–59s.
- Mackay, W. G., Bunn, J. E., Thomas, J. E., Reid, D. C. & Weaver, L. T. 2001 Molecular evidence of *Helicobacter pylori* in biofilms of containers used for storing water. *Arch. Dis. Child.* **84**, A24–A27.
- Mazari-Hiriart, M., Lopez-Vidal, Y. & Calva, J. J. 2001 *Helicobacter pylori* in water systems for human use in Mexico City. *Water Sci. Technol.* **43**, 93–98.
- Mazari-Hiriart, M., Lopez-Vidal, Y., Ponce de Leon, S., Castillo-Rojas, G., Hernandez-Eugenio, C. & Rojo, F. 2003 Bacteria and disinfection byproducts in water from southern Mexico City. *Arch. Environ. Health* **58**, 233–237.
- McKeown, I., Orr, P., McDonald, S., Kabani, A., Brown, R., Coghlan, G., Dawood, M., Embil, J., Sargent, M., Smart, G. & Bernstein, C. N. 1999 *Helicobacter pylori* prevalence in the Canadian arctic: seroprevalence and detection in community water samples. *Am. J. Gastroenterol.* **94**, 1823–1829.
- Mégraud, F. 1995 Transmission of *Helicobacter pylori*: faecal-oral versus oral-oral route. *Aliment. Pharm. Therap.* **9**(Suppl. 2), 85–91.
- Mizoguchi, H., Fujioka, T. & Nasu, M. 1999 Evidence for viability of coccoid forms of *Helicobacter pylori*. *J. Gastroenterol.* **34**(Suppl. 11), 32–36.
- Mladenova, I., Durazzo, M. & Pellicano, R. 2006 Transmission of *Helicobacter pylori*: are there evidences for a fecal-oral route? *Minerva Med.* **97**, 15–18.
- MMWR 1999 Outbreak of *Escherichia coli* O157:H7 and campylobacter among attendees of the Washington County Fair-New York. *Morb. Mortal. Wkly Rep.* **48**, 803–805.
- Moreno, Y., Ferrus, M. A., Alonso, J. L., Jimenez, A. & Hernandez, J. 2003 Use of fluorescent *in situ* hybridization to evidence the presence of *Helicobacter pylori* in water. *Water Res.* **37**, 2251–2256.
- Moreno, Y., Piqueres, P., Alonso, J. L., Jiménez, A., González, A. & Ferrús, M. A. 2007 Survival and viability of *Helicobacter pylori* after inoculation into chlorinated drinking water. *Water Res.* **41**, 3490–3496.
- Nayak, A. K. & Rose, J. B. 2007 Detection of *Helicobacter pylori* in sewage and water using a new quantitative PCR method with SYBR green. *J. Appl. Microbiol.* **103**, 1931–1941.
- Olmos, J. A., Rios, H. & Higa, R. 2000 Prevalence of *Helicobacter pylori* infection in Argentina: results of a nationwide epidemiological study. *J. Clin. Gastroenterol.* **31**, 33–37.
- Osata, M. S., Le, H. H., Ayoub, K. & Graham, D. Y. 1997 Houseflies are an unlikely reservoir for *Helicobacter pylori*. European *Helicobacter pylori* Study Group, *10th International Workshop Abstracts*, 12–14 September, Lisbon, Portugal.
- Oshowo, A., Gillam, D., Botha, A., Tunio, M., Holton, J., Boulos, P. & Hobsley, M. 1998a *Helicobacter pylori*: the mouth, stomach, and gut axis. *Ann. Periodontol.* **3**, 276–280.
- Oshowo, A., Tunio, M., Gillam, D., Botha, A. J., Holton, J., Boulos, P. & Hobsley, M. 1998b Oral colonization is unlikely to play an important role in *Helicobacter pylori* infection. *Brit. J. Surg.* **85**, 850–852.
- Owen, R. J. 1995 Bacteriology of *Helicobacter pylori*. *Bailliere's Clinical Gastroenterology* **9**, 415–445.
- Park, S. R., Mackay, W. G. & Reid, D. C. 2001 *Helicobacter* sp. recovered from drinking water biofilm sampled from a water distribution system. *Water Res.* **35**, 1624–1626.
- Percival, S. L., Walker, J. & Hunter, P. R. 2000 *Microbiological Aspects of Biofilms and Drinking Water*. CRC Press, New York.
- Percival, S. L., Chalmers, R. M., Embrey, M., Hunter, P. R., Sellwood, J. & Wyn-Jones, P. 2004 *Microbiology of Waterborne Diseases*. Elsevier Academic Press, London.
- Peterson, A. M. & Krogfelt, K. A. 2003 *Helicobacter pylori*: an invading micro-organism? A review. *FEMS Immunol. Med. Microbiol.* **36**, 117–126.

- Putsep, K., Branden, C. I., Boman, H. G. & Normark, S. 1999 Antibacterial peptide from *H. pylori*. *Nature* **398**, 671–672.
- Raymond, J., Bergeret, M. & Kalach, N. 2008 *Helicobacter pylori* infection in children. *Presse Med.* **37**, 513–518.
- Reavis, C. 2005 Rural health alert: *Helicobacter pylori* in well water. *J. Am. Acad. Nurse Pract.* **17**, 283–289.
- Rice, E. W., Rodgers, M. R., Wesley, I. V., Johnson, C. H. & Tanner, S. A. 1999 Isolation of *Arcobacter butzleri* from ground water. *Lett. Appl. Microbiol.* **28**, 31–35.
- Rolle-Kampczyk, U. E., Fritz, G. J., Diez, U., Lehmann, I., Richter, M. & Herbarth, O. 2004 Well water: one source of *Helicobacter pylori* colonisation. *Int. J. Hyg. Environ. Health* **207**, 363–368.
- Rothenbacher, D. & Brenner, H. 2003 Burden of *Helicobacter pylori* and *H. pylori*-related diseases in developed countries: recent development and future implications. *Microbes Infect.* **5**, 693–703.
- Rowan, N. J. 2004 Viable but non-culturable forms of food and waterborne bacteria: quo vadis? *Trends Food Sci. Technol.* **15**, 462–467.
- Sasaki, K., Tajiri, Y., Sata, M., Fujii, Y., Matsubara, F., Zhao, M., Shimizu, S., Toyonaga, A. & Tanikawa, K. 1999 *Helicobacter pylori* in the natural environment. *Scand. J. Infect. Dis.* **31**, 275–279.
- Sato, F., Saito, N., Shouji, E., Rani, A., Takeda, H., Sugiyama, T. & Asaka, M. 1999 The maintenance of viability and spiral morphology of *Helicobacter pylori* in mineral water. *J. Med. Microbiol.* **48**, 971.
- Sen, K., Schable, N. A. & Lye, D. J. 2007 Development of an internal control for evaluation and standardization of a quantitative PCR assay for detection of *Helicobacter pylori* in drinking water. *Appl. Environ. Microbiol.* **73**, 7380–7387.
- Shahamat, M., Mai, U., Paszko-Kovla, C., Kessel, M. & Colwell, R. R. 1993 Use of autoradiography to assess viability of *Helicobacter pylori* in water. *Appl. Environ. Microbiol.* **59**, 1231–1235.
- Shahamat, M., Alavi, M., Watts, J. E., Gonzalez, J. M., Sowers, K. R., Maeder, D. W. & Robb, F. T. 2004 Development of two PCR-based techniques for detecting helical and coccoid forms of *Helicobacter pylori*. *J. Clin. Microbiol.* **42**, 3613–3619.
- She, F. F., Lin, J. Y., Liu, J. Y., Huang, C. & Su, D. H. 2003 Virulence of water-induced coccoid *Helicobacter pylori* and its experimental infection in mice. *World J. Gastroenterol.* **9**, 516–520.
- Sherman, P. M. 2004 Appropriate strategies for testing and treating *Helicobacter pylori* in children: when and how? *Am. J. Med.* **117**, 30S–35S.
- Sibille, I., Sime-Ngando, T., Mathieu, L. & Block, J. C. 1998 Protozoan bacterivory and *Escherichia coli* survival in drinking water distribution systems. *Appl. Environ. Microbiol.* **64**, 197–202.
- Souto, R. & Colombo, A. P. 2008 Detection of *Helicobacter pylori* by polymerase chain reaction in the subgingival biofilm and saliva of non-dyspeptic periodontal patients. *J. Periodontol.* **79**, 97–103.
- Stanley, K., Cunningham, R. & Jones, K. 1998 Isolation of *Campylobacter jejuni* from groundwater. *J. Appl. Microbiol.* **85**, 187–191.
- Stark, R. M., Gerwig, G. J., Pitman, R. S., Potts, L. F., Williams, N. A., Greenman, J., Weinzweig, I. P., Hirst, T. R. & Millar, M. R. 1999 Biofilm formation by *Helicobacter pylori*. *Lett. Appl. Microbiol.* **28**, 121–126.
- Thomas, J. E., Gibson, B. R., Darboe, M. K., Dale, A. & Weaver, L. T. 1992 *Helicobacter pylori* from human faeces. *Lancet* **340**, 1194–1195.
- Velazquez, M. & Feirtag, J. M. 1999 *Helicobacter pylori*: characteristics pathogenicity detection methods and mode of transmission implicating foods and water. *Int. J. Food Microbiol.* **53**, 95–104.
- Vincent, P. 1995 Transmission and acquisition of *Helicobacter pylori* infection: evidences and hypothesis. *Biomed. Pharmacother.* **49**, 11–18.
- Warren, J. R. & Marshall, B. 1983 Unidentified curved bacillus on gastric epithelium in active chronic gastritis. *Lancet* **1**, 1273–1275.
- Watson, C. L., Owen, R. J., Said, B., Lai, S., Lee, J. V., Surman-Lee, S. & Nichols, G. 2004 Detection of *Helicobacter pylori* by PCR but not culture in water and biofilm samples from drinking water distribution systems in England. *J. Appl. Microbiol.* **97**, 690–698.
- Weiss, J., Mecca, J., da Silva, E. & Gassner, D. 1994 Comparison of PCR and other diagnostic techniques for detection of *Helicobacter pylori* infection in dyspeptic patients. *J. Clin. Microbiol.* **32**, 1663–1668.
- West, A. P., Millar, M. R. & Tompkins, D. S. 1990 Survival of *Helicobacter-pylori* in water and saline. *J. Clin. Pathol.* **43**, 609.
- West, A. P., Millar, M. R. & Tompkins, D. S. 1992 Effect of physical environment on survival of *Helicobacter pylori*. *J. Clin. Pathol.* **45**, 228–231.
- Winiacka-Krusnell, J., Wreiber, K., von Euler, A., Engstrand, L. & Linder, E. 2002 Free-living amoebae promote growth and survival of *Helicobacter pylori*. *Scand. J. Infect. Dis.* **34**, 253–256.
- Xia, H. H. & Talley, N. J. 1997 Natural acquisition and spontaneous elimination of *Helicobacter pylori* infection: clinical implications. *Am. J. Gastroenterol.* **92**, 1780–1787.
- Zaterka, S., Eisig, J. N., Chinzon, D. & Rothstein, W. 2007 Factors related to *Helicobacter pylori* prevalence in an adult population in Brazil. *Helicobacter* **12**, 82–88.
- Zheng, P. Y., Hua, J., Ng, H. C. & Ho, B. 1999 Unchanged characteristics of *Helicobacter pylori* during its morphological conversion. *Microbios* **98**, 51–64.

First received 2 July 2008; accepted in revised form 6 December 2008. Available online May 2009