

# Metallothionein Induction by Hypoxia Involves Cooperative Interactions between Metal-Responsive Transcription Factor-1 and Hypoxia-Inducible Transcription Factor-1 $\alpha$

Brian J. Murphy,<sup>1</sup> Tomoki Kimura,<sup>2</sup> Barbara G. Sato,<sup>1</sup> Yihui Shi,<sup>1</sup> and Glen K. Andrews<sup>3</sup>

<sup>1</sup>Biosciences Division, SRI International, Menlo Park, California; <sup>2</sup>Department of Toxicology, Faculty of Pharmaceutical Sciences, Setsunan University, Hirakata, Osaka, Japan; and <sup>3</sup>Department of Biochemistry and Molecular Biology, University of Kansas Medical Center, Kansas City, Kansas

## Abstract

Mammalian *metallothionein* (*MT*) genes are transcriptionally activated by the essential metal zinc as well as by environmental stresses, including toxic metal overload and redox fluctuations. In addition to playing a key role in zinc homeostasis, *MT* proteins can protect against metal- and oxidant-induced cellular damage, and may participate in other fundamental physiologic and pathologic processes such as cell survival, proliferation, and neoplasia. Previously, our group reported a requirement for metal-responsive transcription factor-1 (MTF-1) in hypoxia-induced transcription of mouse *MT-I* and human *MT-IIA* genes. Here, we provide evidence that the protumorigenic hypoxia-inducible transcription factor-1 $\alpha$  (HIF-1 $\alpha$ ) is essential for induction of *MT-1* by hypoxia, but not zinc. Chromatin immunoprecipitation assays revealed that MTF-1 and HIF-1 $\alpha$  are both recruited to the mouse *MT-I* promoter in response to hypoxia, but not zinc. In the absence of HIF-1 $\alpha$ , MTF-1 is recruited to the *MT-I* promoter but fails to activate *MT-I* gene expression in response to hypoxia. Thus, HIF-1 $\alpha$  seems to function as a coactivator of *MT-I* gene transcription by interacting with MTF-1 during hypoxia. Coimmunoprecipitation studies suggest interaction between MTF-1 and HIF-1 $\alpha$ , either directly or as mediated by other factors. It is proposed that association of these important transcription factors in a multiprotein complex represents a common strategy to control unique sets of hypoxia-inducible genes in both normal and diseased tissue. (*Mol Cancer Res* 2008;6(3):483–90)

## Introduction

The cysteine-rich metallothioneins (*MT*) bind metals, including zinc, copper, and cadmium, with high affinity and

capacity (1, 2). In mice, *MT-I* and *MT-II* are the most prevalent of the four known *MT* isoforms, and they are recognized to regulate heavy metal metabolism and detoxification and to participate in oxidant scavenging (1, 2). Other studies suggest that these proteins are pluripotent, contributing to a number of other fundamental processes, including proliferation, survival, metabolism, inflammation, and extracellular remodeling (3-18). It is therefore not surprising that aberrant expression patterns of *MTs* can correlate with a number of pathologic conditions, including malignant progression. For example, high levels of *MT-I* and *MT-IIA* have been detected in many (but not all) human tumors, including those of the breast, prostate, cervix, testes, kidney, bladder, brain, and oral epithelium (19-25). Underlying molecular mechanisms are believed to include the ability of *MT* to activate survival pathways and confer resistance against drug and radiation therapy (5, 26-28). Collectively, these data suggest that an understanding of the control mechanisms regulating *MT* isoform expression will provide unique therapeutic insights.

The metal-responsive transcription factor-1 (MTF-1) is a central regulator of metal-inducible expression of *MT-I* and *MT-II*. Current models of MTF-1 activation suggest that zinc directly and reversibly modulates highly specific, linker-mediated zinc finger interactions in MTF-1, resulting in translocation of this transcription factor to the nucleus and rate-limiting binding to metal response elements [MRE; TGC(G/C)CNC(G)] within proximal *MT* promoters (29-34). In addition to zinc, other heavy metals (e.g., cadmium), hypoxia, oxidative stress, nitric oxide, and high temperature induce the transcriptional activity of MTF-1 (1, 20, 30, 35-43). Although the precise mechanisms of action of these other MTF-1 activators remain to be determined, a common underlying theme seems to involve a displacement of labile zinc, perhaps from the *MT* protein, zinc-sensitive activation of MTF-1, and altered expression of MTF-1 target genes (20, 40, 44, 45). Studies by our group, and collaborators, suggest that MTF-1 can contribute to tumorigenic processes, in part, through its transcriptional action as either a positive or a negative regulator of numerous genes, including *zinc transporter 1* (*ZnT1*), *PIGF*, *transforming growth factor- $\beta$ 1* (*TGF- $\beta$ 1*), *tissue transglutaminase-2* (*TG2*), and *Zip10* (20, 46-49).

A number of studies designed to characterize transcription factor interactions at the *MT-I* promoter have been reported (e.g., refs. 31, 32, 43, 50-54). However, a limited number of these studies have focused on the chromatin-packaged *MT-I* promoter, and those only in the context of activation by metal

Received 7/20/07; revised 10/26/07; accepted 11/14/07.

Grant support: NIH grants CA057692 (B.J. Murphy) and ES05704 (G.K. Andrews).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Note: Supplementary data for this article are available at Molecular Cancer Research Online (<http://mcr.aacrjournals.org/>).

Requests for reprints: Brian J. Murphy, Biosciences, SRI International, 333 Ravenswood Avenue, Menlo Park, CA 94025. Phone: 650-859-4213. E-mail: brian.murphy@sri.com

Copyright © 2008 American Association for Cancer Research.

doi:10.1158/1541-7786.MCR-07-0341

and oxidative stresses (31, 32). The proximal *MT-I* promoter is occupied by several transcription factors on at least 10 sites in response to zinc overload. In addition to MTF-1, upstream stimulatory factor (USF)-1, USF-2, Sp1, c-Fos, and c-Jun are bound to the proximal promoter (32). USF-1 and USF-2 strongly interact with the *MT-I* promoter regardless of the zinc concentration or the presence of MTF-1. The heterodimers of USF-1 and USF-2 most likely facilitate the functions of MTF-1 through the maintenance of an open promoter configuration, allowing direct binding of MTF-1 to specific MREs within the proximal promoter in response to metals (32). MTF-1 seems to be only weakly associated with the promoter under normal zinc conditions, whereas increased zinc levels result in binding of MTF-1 to all five MREs within the proximal -250 bp promoter (relative to the transcription start site). C-Fos is also associated with the *MT-I* promoter during zinc exposure, and this MTF-1-dependent recruitment coincides with robust activation of the gene (32). Our recent studies<sup>4</sup> show that metals cause the formation of a protein complex containing (at least) MTF-1, Sp1, and the histone acetyltransferase p300. In this study, the acidic transactivation domain of MTF-1 was found to be essential for *MT-I* gene activation and the binding of p300. MTF-1-containing transcription factor complexes involved in regulation by stresses other than metals remain to be identified.

The cellular response to hypoxia is regulated, in part, by the hypoxia-inducible factor-1 (HIF-1), a basic helix-loop-helix transcription factor composed of two subunits, HIF-1 $\alpha$  and HIF-1 $\beta$  (55). Whereas the HIF-1 $\beta$  subunit is constitutively expressed, HIF-1 $\alpha$  expression is regulated by oxygen tension through a protein stabilization process involving inhibition of posttranslational ubiquitination-triggered proteolysis (56). Our previous studies have placed MTF-1 in as yet unidentified pathway required for stabilization/accumulation of HIF-1 $\alpha$  protein levels in response to hypoxia (57). Under low oxygen tensions, HIF-1 $\alpha$  forms heterodimers with HIF-1 $\beta$  that mediate nuclear translocation and binding to hypoxic response elements (core consensus sequence RCGTG) within the proximal promoters of target genes. HIF-1 regulates a wide range of genes, including many involved in metabolism, angiogenesis, cell survival, and invasion. Specific gene expression patterns depend on the degree and duration of hypoxic insult, cell type, and tissue background (58, 59). Other transcription factors (e.g., c-Jun, nuclear factor- $\kappa$ B, signal transducer and activator of transcription 3, Sp1, and early growth response-1) have been shown to interact with HIF-1 $\alpha$  and cooperate in regulating target gene expression. Some of these proteins can also act independently of HIF-1 $\alpha$  and control unique sets of hypoxia-responsive target genes (e.g., see refs. 20, 60, 61). In addition, the general transcriptional coactivator p300/cyclic AMP-responsive element binding protein-binding protein directly binds to the COOH terminus transcriptional activation domain of HIF-1 $\alpha$  and synergistically enhances its transcriptional activity in response to hypoxia (60, 62).

We previously found MTF-1 to be transactivated by hypoxia and required for the hypoxia-inducible transcription of the *MT-I*

and *MT-IIa* genes (43). Transfection and gel shift analyses identified hypoxic-responsive MREs within these promoters (43). In addition, our group identified the angiogenic *placenta growth factor (Plgf)* as another MTF-1-dependent hypoxic target gene (47). Subsequent collaborative studies found a corequirement for the redox-sensitive nuclear factor- $\kappa$ B in transcriptional control of the *Plgf* (63). Herein, we show that HIF-1 $\alpha$  is also required for the hypoxia-inducible transcription of *MT-I*. Moderate hypoxia caused the recruitment of MTF-1, HIF-1 $\alpha$ , and p300 to the mouse *MT-I* proximal promoter. Furthermore, MTF-1 and HIF-1 $\alpha$  physically interact under these conditions in a mechanism yet to be elucidated.

## Results

### *HIF-1 $\alpha$ and MTF-1 Regulate Hypoxia Induction of MT-I mRNA*

*MT-I* transcription is increased during hypoxia in a MTF-1 (and MRE)-dependent manner; increases in *MT-I* mRNA levels occur within 4 hours after exposure, and maximum induction occurs within 8 to 12 hours (43, 64). MTF-1 is also required in the hypoxia accumulation of HIF-1 $\alpha$ , a central transcriptional regulator of the cellular response to hypoxia (57). Herein, we assessed a possible role of HIF-1 $\alpha$  in the hypoxia-inducible induction of *MT-I*. Northern blot analyses revealed that loss of *HIF-1 $\alpha$*  in both TAG and TAG/*ras* mouse embryonic fibroblasts [MEF; *HIF-1 $\alpha$*  knockout (HIF-KO)] attenuated the induction of *MT-I* mRNA levels during hypoxia but had no effect on zinc induction (Fig. 1A and B, respectively). To confirm this finding, we used RNA interference to silence HIF-1 $\alpha$  expression in *ras*-NIH 3T3 cells. Figure 1C shows a real-time reverse transcription-PCR (RT-PCR) analysis of *MT-I* mRNA levels in aerobic and hypoxic *ras*-NIH 3T3 cells that were transiently transfected with either mouse *HIF-1 $\alpha$*  or control, noncoding, small interfering RNA (siRNA) oligomers. Although the inductions by both hypoxia and zinc were modest, compared with the MEF model, the induction of *MT-I* mRNA levels by hypoxia was nevertheless completely reversed by *HIF-1 $\alpha$*  RNA interference. The ability of the siRNA sequence to inhibit HIF-1 $\alpha$  expression, and thus accumulation, was confirmed by Western analysis (Fig. 1C). Our data also showed that, in contrast to MTF-1 control over HIF-1 $\alpha$  accumulation, loss of *HIF-1 $\alpha$*  had no effect on either whole-cell or nuclear MTF-1 protein levels from aerobic or hypoxic cells (Fig. 1D). This Western analysis also indicated that hypoxia had no apparent inducing effect on MTF-1 nuclear translocation, compared with significant translocation in response to zinc exposure (e.g., see Fig. 3; ref. 31). These data suggest that MTF-1 and HIF-1 $\alpha$  are both essential for induction of *MT-I* gene expression in response to hypoxia and that HIF-1 $\alpha$  probably directly acts at the *MT-I* promoter. Furthermore, hypoxia may affect that transactivation capacity of MTF-1 that resides in the nucleus.

### *Hypoxia Induces Recruitment of MTF-1 and HIF-1 $\alpha$ to the MT-I Promoter*


Chromatin immunoprecipitation (ChIP) assays were done to more fully address the mechanisms by which MTF-1 and HIF-1 $\alpha$  regulate hypoxia-inducible transcription of the *MT-I* gene *in vivo*. TAG/*ras*-transformed wild-type (WT), MTF-KO,

<sup>4</sup> Y. Li et al., in preparation.


and HIF-KO MEFs were exposed to hypoxia (1% O<sub>2</sub>; 6 hours) or ZnSO<sub>4</sub> (100  $\mu$ mol/L; 1 hour), or left untreated before ChIP assays. Cross-linked chromatin was isolated from each culture, sheared, and immunoprecipitated with antibodies against MTF-1, HIF-1 $\alpha$ , and p300. The precipitated DNA was analyzed by PCR using primers specific for the proximal region (-264 to +43 bp) of the *MT-I* gene (Fig. 2). The results show that MTF-1 is recruited to the *MT-I* promoter in response to hypoxia or zinc. The magnitude of recruitment in response to hypoxia was evident but clearly less than that observed after zinc exposure (Fig. 2A and B, *top*), and was

not dependent on HIF-1 $\alpha$  (Fig. 2B). Hypoxia also caused the recruitment of HIF-1 $\alpha$  and p300 to the *MT-I* promoter in WT MEFs (Fig. 2A and B).

The *in vivo* kinetics of binding of MTF-1 and HIF-1 $\alpha$  to the *MT-I* promoter in response to hypoxia was examined (Fig. 2C). MTF-1 was recruited within the first hour of hypoxia and remained associated with this promoter for at least 8 hours. In contrast, HIF-1 $\alpha$  binding to the *MT-I* promoter was not detected until 4 hours of hypoxia. The intensity of the HIF-1 $\alpha$  signal was always modest relative to that of MTF-1; this pattern may be due to a number of possibilities, including



**FIGURE 1.** HIF-1 $\alpha$  contributes to hypoxia-induced *MT-I* transcription. **A** and **B.** Northern blot analysis of *MT-I* mRNA levels as a function HIF-1 $\alpha$  expression. TAG- and TAG/ras-WT and HIF-KO MEFs were examined for *MT-I* mRNA levels as functions of hypoxia (Hx; 1% O<sub>2</sub>; 2-8 h) or ZnSO<sub>4</sub> (Zn; 100  $\mu$ mol/L; 1 h) exposures. **C.** Real-time RT-PCR analysis of *MT-I* mRNA levels as a function of HIF-1 $\alpha$  silencing in hypoxic *ras*-NIH 3T3 cells. Near-confluent cultures were transfected with HIF-1 $\alpha$  siRNA oligonucleotides (see Materials and Methods) and allowed to recover for 16 h before exposure to hypoxia (8 h). Relative *MT-I* mRNA levels were determined by real-time RT-PCR. Western blotting of nuclear HIF-1 $\alpha$  protein was also done to confirm HIF-1 $\alpha$  silencing. **D.** Western analysis of whole-cell extract (WCE) and nuclear MTF-1 protein levels as a function of HIF-1 $\alpha$  loss. KO in **D** denotes HIF-KO. GAPDH, glyceraldehyde-3-phosphate dehydrogenase. A full-length Western blot is presented in Supplementary Data. The position of the MTF-1 protein band, under these electrophoresis conditions, was previously established through the use of WT and *MTF-1* KO MEFs (57) and through reintroduction of *MTF-1* cDNA into KO cells (ref. 31; also see Fig. 3, *bottom*, and Supplementary Data).



**FIGURE 2.** ChIP analysis of HIF-1 $\alpha$ , p300, and MTF-1 with the *MT-I* promoter. WT and MTF-KO (**A**) or HIF-KO MEFs (**B**) were exposed to 100  $\mu$ M ZnSO<sub>4</sub> (Zn) for 1 h or hypoxia (Hx; 1% O<sub>2</sub>; 8 h). C, control. **C.** A time course analysis of MTF-1 and HIF-1 $\alpha$  recruitment to the *MT-I* promoter using WT MEFs. Chromatin was fixed, sonicated, and then precipitated using a polyclonal against Flag or MTF-1. The relative amounts of *MT-I* promoter DNA precipitated were determined by PCR and normalized to amplification of input DNA (bottom blot in **A** and **B**).

antibody affinity and efficiency of recovery in the immunoprecipitation steps, the inability to cross-link HIF-1 $\alpha$  efficiently, or the lower number of HIF-1 $\alpha$  molecules bound to the *MT-I* promoter.

#### *MTF-1 Associates with HIF-1 $\alpha$ in Hypoxia-Treated Cells*


*Ras*-MTF-KO cells stably transformed with a FLAG-tagged *MTF-1* cDNA (MTF-1<sub>flag</sub> MEFs) and MTF-KO MEFs (as a control) were used to assess a possible association between MTF-1 and HIF-1 $\alpha$ . Earlier studies showed that stable introduction of a FLAG-tagged *MTF-1* cDNA vector into MTF-KO cells rescued metal responsiveness of the *MT-I* gene (31). Immunoprecipitation of FLAG-MTF protein was done on nuclear extracts from aerobic and hypoxic (1% O<sub>2</sub>; 7 hours) MTF-KO and MTF-1<sub>flag</sub> cells. The immunoprecipitates were analyzed by Western blot detection of the HIF-1 $\alpha$  protein (Fig. 3). HIF-1 $\alpha$  protein does coprecipitate with FLAG-MTF protein in hypoxic MTF-1<sub>flag</sub> cells, but not under zinc stress, indicating a specific association of MTF-1 with HIF-1 $\alpha$  under hypoxic conditions (Fig. 3). Similar patterns were obtained with TAG-immortalized MEFs (data not shown). Figure 3 also includes Western analysis of FLAG-MTF from the coimmunoprecipitated extracts. As expected, the nuclear level of MTF-1 protein in the MTF-1<sub>flag</sub> cells was greatly enhanced in response to zinc treatment (31). However, we detected no discernible increase in nuclear MTF-1 protein from hypoxic MTF-1<sub>flag</sub> MEFs. These data are consistent with our analysis of total MTF-1 protein in whole-cell and nuclear extracts (Fig. 1B). However, immunofluorescence analysis of vesicular stomatitis virus-tagged MTF-1 protein indicated that at least a fraction of cytoplasmic MTF-1 stores is transported into the nucleus under hypoxic conditions (63).<sup>5</sup> Regardless, these new findings show for the first time that MTF-1 forms a complex that includes HIF-1 $\alpha$  and that these proteins are both recruited to the promoter *in vivo* under hypoxia exposure.

#### Discussion

The data presented herein confirm and extend our original findings regarding the involvement of MTF-1 in cellular responses to hypoxic stress, and specifically in its interactions with HIF-1 $\alpha$ ? Our group had previously reported that hypoxia induces the expression of MT-I and MT-IIA in a MTF-1/MRE-dependent manner (43). In subsequent studies, we found that MTF-1 participates in a more general hypoxia-inducible mechanism, contributing to the stabilization and nuclear accumulation of the HIF-1 $\alpha$  protein, a central regulator of the cellular response to hypoxia (57). This new study provides evidence for recruitment of MTF-1, HIF-1 $\alpha$ , and p300 to the *MT-I* promoter and indicates that MTF-1 and HIF-1 $\alpha$  are components of a transcriptional complex in response to hypoxia.

Cellular hypoxia is a common stress characterizing a number of physiologic and pathogenic conditions, including development, wound healing, ischemia, inflammation, and tumorigenesis (20, 65-67). For example, the establishment of hypoxia microenvironments within many solid tumor types correlates with the phenotypes of increased angiogenesis, survival, and metastasis (see ref. 20 for a review). HIF-1 $\alpha$ , which is stabilized and highly expressed in many solid tumors, is considered a master transcriptional regulator of many of the genes that contribute to these malignant phenotypes and also to normal tissue responses to hypoxia (65, 68).

<sup>5</sup> B. Sato and B. Murphy, unpublished results.



**FIGURE 3.** Coimmunoprecipitation of mouse MTF-1 and HIF-1 $\alpha$ . MTF-KO and MTF-1<sub>flag</sub> MEFs were incubated under hypoxia (Hx; 1% O<sub>2</sub>; 8 h) or ZnSO<sub>4</sub> (Zn; 100  $\mu$ mol/L; 1 h). Nuclear extracts were immunoprecipitated (IP) with an antibody against FLAG followed by Western blot (WB) analysis on the immunoprecipitates for HIF-1 $\alpha$  (**A**) and MTF-1 (**B**) proteins. Full-length blots are presented in Supplementary Data.

Our preliminary Northern analysis of TAG-immortalized and TAG/*ras*-transformed fibroblasts showed that HIF-1 $\alpha$ , in addition to MTF-1, is essential for hypoxia-associated induction of *MT-I* mRNA levels. Importantly, alternative knockdown of HIF-1 $\alpha$  expression (by siRNA), in *ras*-NIH 3T3 cells, confirmed the validity of the data generated by genetic knockout of the gene. A search of the proximal mouse *MT-I* promoter indicates at least two putative HIF-1 $\alpha$  binding sites, hypoxia response elements, that flank the USF element/antioxidant response element fragment (see ref. 32 for description of the complex USF/antioxidant response element site). ChIP analysis was used to examine the occupancy of the *MT-I* promoter under hypoxia. The recruitment of MTF-1 was suggested in our earlier studies (using Northern blot, promoter/reporter, and gel shift analyses), linking MTF-1 activation to the increased transcription of *MT-I* and *MT-III* by hypoxia (43). Hypoxic stress resulted in markedly increased MTF-1 binding to the *MT-I* promoter, and this intensity was comparable with that seen for zinc induction. Although recruitment of MTF-1 occurred within the first hour of hypoxic exposure, detection of HIF-1 $\alpha$  binding was apparently delayed (up to 4 hours of hypoxia) and the binding was consistently less intense than that seen for MTF-1 binding. We previously reported the appearance of detectable increases in *MT-I* mRNA levels within this time frame (4-6 hours) of HIF-1 $\alpha$  binding (43). The weaker binding of HIF-1 $\alpha$  to the *MT-I* promoter may simply reflect a relatively lower HIF-1 $\alpha$  antibody affinity and efficiency, compared with the FLAG antibody. The weak HIF-1 $\alpha$  signal may also reflect a difficulty in stably cross-linking the MTF-1/HIF-1 complex. However, it is also possible that lower numbers of HIF-1 molecules actually bind the *MT-I* promoter under hypoxia or that HIF-1 $\alpha$  must associate with other proteins for recruitment to the promoter rather than directly to the DNA itself. MTF-1 is considered the master regulator of MT genes and a requirement of MTF-1 for recruitment of other transcription factors to the *MT-I* promoter is most likely a common mechanism. For example, the presence of MTF-1 is necessary for zinc-induced c-Fos recruitment to the *MT-I* promoter (32). In sum, the available data imply that recruitment of both MTF-1 and HIF-1 $\alpha$  to the *MT-I* promoter is required for hypoxia-inducible transcriptional activation of *MT-I*.

On stabilization, HIF-1 $\alpha$  forms a heterodimer with HIF-1 $\beta$ , resulting in functional HIF-1 (69). In addition, the HIF-1 $\alpha$

protein subunit physically interacts with a number of other transcription-related proteins, resulting in tissue-specific expression patterns. HIF-1 complexes are usually composed of a number of other transcription factors and coactivators such as p300. Under hypoxia, p300/cyclic AMP-responsive element binding protein-binding protein acetylates H3 histone, through its intrinsic histone acetyltransferase activity, and synergistically enhances HIF-1 $\alpha$  transactivation, in part through direct physical binding (60). Our ChIP data clearly show that p300 is recruited to the HIF-1 complex at the *MT-I* promoter in response to hypoxia. Although loss of HIF-1 $\alpha$  expression markedly attenuates p300 recruitment, MTF-1 does not affect p300 binding to the *MT-I* promoter. More detailed studies will be required to define the complex interactions between MTF-1, p300, and HIF-1 $\alpha$ .

The coprecipitation experiments show an association between the MTF-1 and HIF-1 $\alpha$  proteins during hypoxic stress. These findings are consistent with the formation of a MTF-1/HIF-1 $\alpha$  transcriptional complex that forms at the *MT-I* promoter in response to hypoxia. The precise interaction between these two hypoxia-inducible transcription factors and their interactions with p300 remain to be elucidated.

Another potentially important finding involves the lack of detectable MTF-1 protein translocation into the nucleus under hypoxia. More sensitive immunofluorescence studies previously detected some nuclear translocation of the MTF-1 protein under hypoxia in at least two human cancer lines (63).<sup>5</sup> This apparent lack of nuclear translocation in the MEFs is in direct opposition to the robust nuclear accumulation of MTF-1 seen under zinc treatment. However, our ChIP data showed relatively similar binding of MTF-1 to the *MT-I* promoter in response to both hypoxia and zinc. It is certainly possible that the level of MTF-1 translocation detected by immunofluorescence is sufficient for this hypoxia-inducible recruitment. On the other hand, hypoxia may induce other signaling pathways that activate preexisting pools of nuclear MTF-1, perhaps through posttranslational modification of the transcription factor as described elsewhere (70, 71).

These new studies have uncovered the existence of a potentially central hypoxia-inducible complex that includes MTF-1 and HIF-1 $\alpha$  (and p300). It is likely that MTF-1 is involved in other hypoxia-inducible transcriptional complexes. For example, MTF-1 cooperates with nuclear factor- $\kappa$ B to control the expression of the proangiogenic and prosurvival

PIGF (63). We therefore propose that MTF-1 and HIF-1 $\alpha$  are components of a transcriptional complex that along with other hypoxia-inducible transcription factors and coactivators form combinatorial transcriptional complexes that are activated in a cell- and gene-specific manner. The cooperative interaction, and potential for gene therapy, between MTF-1 and HIF-1 $\alpha$  has been independently verified by at least one other report showing that a chimeric promoter consisting of multiple copies of MREs and hypoxia response elements (and early growth response-1 binding elements) greatly enhanced expression of a reporter in response to hypoxia compared with that seen for any one element (72). Taken together, these studies offer a completely unique insight into the cooperative roles of MTF-1 and HIF-1 $\alpha$  as transcriptional regulators of a subset of hypoxia-sensitive genes. The potential for the development of combined MTF-1/HIF-1 $\alpha$  therapeutic strategies against diseases such as cancer is promising but remains to be tested.

## Materials and Methods

### Materials

Blasticidin S hydrochloride was obtained from Calbiochem; Polybrene (hexadimethrine bromide) was from Sigma; protein G and A/G Plus-agarose beads were from Santa Cruz Biotechnology; and Phoenix-Eco cells were from Orbigen. The gWZl-Blast H-*ras* expression vector was a generous gift from Dr. David Dankort (University of California, San Francisco, CA). The polyclonal antibody against recombinant mouse MTF-1 was generated (by G.K.A.) as previously described (73). The antibody against Flag peptide was purchased from Sigma, and the polyclonal antibody against p300 was from Santa Cruz Biotechnology. Lipofectamine 2000 was purchased from Invitrogen Life Technologies.

### Cell Culture, Transfection, and Hypoxia Treatments

*Ras*-transformed NIH 3T3 cells were purchased from the American Type Culture Collection. The SV40 large T antigen TAg/*ras*-transformed WT and *HIF-1 $\alpha$*  null MEFs (HIF-KO) were a generous gift of Dr. Keith Laderoute (SRI International, Menlo Park, CA). The HIF-KO cells were derived from the immortalized WT MEFs and there was no selection for individual clones (74). The TAg immortalized/Harvey (H)-*ras*-transformed WT and *MTF-1* null (MTF-KO) MEFs used in the Northern and ChIP analyses were a generous gift from Dr. Walter Schaffner (University of Zürich, Zurich, Switzerland). For detailed descriptions of these cell lines, see refs. (47, 57, 75). All cell lines (and new thaws) were genotyped by PCR analysis to determine the presence of the *MTF-1* WT and null alleles, as well as for genomic integration of *TAg* and oncogenic H-*ras*. The MEFs were cultured in DMEM with 10% fetal bovine serum. For coimmunoprecipitation studies, MTF-KO cells stably transfected with a mouse FLAG-tagged *MTF-1* expression vector were used. These cells are designated as MTF-1<sub>flag</sub>. The generation of these immortalized SV40-TAg stably transfected cell lines is described in ref. (31). The MTF-KO and MTF-1<sub>flag</sub> MEFs were then transformed with H-*ras*. Briefly, retrovirus-containing gWZl-Blast H-*ras* was prepared in Phoenix-Eco cells. Approximately  $5 \times 10^5$  cells were plated in 100-mm dishes and infected the next day with retrovirus containing supernatant (1 mL) that also contained

5 mg/mL Polybrene (3  $\mu$ L) in 2 mL of DMEM + 10% fetal bovine serum. After 5 h incubation at 37°C, an additional 7 mL of DMEM + 10% fetal bovine serum were added. The cells were incubated for 36 h, and then transferred into medium containing 2  $\mu$ g/mL blasticidin. Selection was continued until all the mock-infected cells died. Frozen stocks of the surviving cells were prepared. *Ras* transformation was verified by soft agar colony formation. Large colonies were observed in the plates of transformed cells and not in the plates of the mock-infected cells.

Monolayer cultures were incubated in a hypoxic chamber held at a constant oxygen tension: 1% O<sub>2</sub>; 5% CO<sub>2</sub> (InVivo<sub>2</sub> Hypoxic Workstation, Ruskin, Inc.). All media and solutions, such as lysis buffer, were incubated in the chamber before addition to cells.

### Northern Blotting, RNA Interference, and Quantitative Real-time RT-PCR Analysis

Total RNA was extracted from cells using an RNeasy Mini kit (Qiagen) and Northern blot analysis was done as previously described (64). Briefly, RNA was separated by denaturing agarose-formaldehyde gel electrophoresis and transferred (and UV cross-linked) to nylon membranes. Blots were probed with an *EcoRI/HindIII* fragment of the mouse *MT-I* genomic DNA labeled by the random primer method. Signals were scanned on an Alphamager 3400 (Alpha Innotech).

For the RNA interference studies, cells were seeded into 60-mm<sup>3</sup> plates with fresh medium, without antibiotics, 24 h before transfection. The high-performance liquid chromatography-purified siRNAs (nonsilencing control and mouse *HIF-1 $\alpha$*  siRNAs; >97% pure) were purchased from Qiagen-Xeragon. The control siRNA sequence does not target any known mammalian gene (sequence 5'-AATTCTCCGAACGTGTACAGT-3'). The siRNA specific for mouse *HIF-1 $\alpha$*  was derived from the mouse *HIF-1 $\alpha$*  mRNA sequence (5'-TAGCCACAATTGCACAATATA-3'). Lipofectamine 2000 (Invitrogen) was used for all transfections. Cells were subjected to 8 h hypoxia (or normoxia) following treatment with 200 pmol control or mouse *HIF-1 $\alpha$*  siRNA. Zinc exposure (ZnSO<sub>4</sub>; 100  $\mu$ mol/L for 2 h) was used as a positive control of *MT-I* induction. Total RNA from cells was isolated using the RNeasy method and DNA from the RNA preparations was removed by treatment with DNase (TURBO DNA-free Kit). First-strand cDNA was synthesized from total RNA using iScript cDNA synthesis kit (Bio-Rad) according to the manufacturer's instructions. Transcript analysis was done by real-time RT-PCR using the Taqman assay. Hybridization probes and primers (Inventory, chosen from the online catalog) were purchased from Applied Biosystems. All samples were amplified simultaneously in triplicate and amplifications were run in a 7300 Real-time PCR System (Applied Biosystems). Each value was normalized to the corresponding *GAPDH* mRNA levels.

### Chromatin Immunoprecipitation

ChIP assays of the proximal *MT-I* promoter were carried out as previously described (31, 32, 34, 76). MEFs exposed to hypoxia or aerobic controls were fixed in 1% formaldehyde and sheared, chromatin prepared, precleared with protein

G-agarose, and immunoprecipitated with specific antibody overnight at 4°C. The flag antibody (F3165) was purchased from Sigma and the HIF-1 $\alpha$  antibody (NB100-105) was from Novus. Protein G-agarose was used to capture the immune complexes, and the formaldehyde cross-links were reversed. The DNA was analyzed by PCR using primers that span the -264 to +43 bp segment of the mouse *MTF-1* promoter. The PCR products (after 28 cycles) were separated by agarose electrophoresis and stained with SYBR green I, quantified using the ChemiImager System (Alpha Innotech), and normalized to input products. All PCR products were obtained in the linear range of amplification as determined from analysis at various cycles (27-33 cycles).

#### Preparation of Nuclear Extract, Coimmunoprecipitation, and Western Analysis

Nuclei from either aerobic or hypoxic MEF cultures (MTF-KO and MTF-1<sub>flag</sub>) were isolated using the method of Ryan et al. (77) with modifications described elsewhere (57). Cells were washed with ice-cold PBS before addition of lysis buffer that also contained protease inhibitor cocktail. Nuclei from hypoxic MEFs were harvested in the InVivo<sub>2</sub> Hypoxic Workstation.

Protein concentrations were determined using the BCA reagent (Pierce Biotechnology). Equal nuclear protein extracts (200  $\mu$ g) from hypoxia-treated and aerobic MEFs (MTF-KO and MTF-1<sub>flag</sub>) were incubated with 2  $\mu$ g of anti-Flag antibody and 20  $\mu$ L of protein A/G Plus-agarose beads (constant mixing at 4°C overnight). The beads were centrifuged (2,500 rpm  $\times$  5 min) and washed in NP40 lysis buffer five times before Western blotting for HIF-1 $\alpha$  (Novus) and Flag (Sigma).

#### Acknowledgments

We thank Dr. Keith R. Laderoute (SRI International, Menlo Park, CA) for the generous gifts of HIF-KO and WT MEFs, and Joy Calaoagan for some timely technical assistance.

#### References

- Andrews GK. Regulation of metallothionein gene expression by oxidative stress and metal ions. *Biochem Pharmacol* 2000;59:95-104.
- Palmiter RD. The elusive function of metallothioneins. *Proc Natl Acad Sci U S A* 1998;95:8428-30.
- Penkowa M. Metallothioneins are multipurpose neuroprotectants during brain pathology. *FEBS J* 2006;273:1857-70.
- Jin R, Chow VT, Tan PH, Dheen ST, Duan W, Bay BH. Metallothionein 2A expression is associated with cell proliferation in breast cancer. *Carcinogenesis* 2002;23:81-6.
- Lazo JS, Pitt BR. Metallothioneins and cell death by anticancer drugs. *Annu Rev Pharmacol Toxicol* 1995;35:635-53.
- Shimoda R, Achanzar WE, Qu W, et al. Metallothionein is a potential negative regulator of apoptosis. *Toxicol Sci* 2003;73:294-300.
- Cai L. Suppression of nitrate damage by metallothionein in diabetic heart contributes to the prevention of cardiomyopathy. *Free Radic Biol Med* 2006;41:851-61.
- Cai L, Wang J, Li Y, et al. Inhibition of superoxide generation and associated nitrosative damage is involved in metallothionein prevention of diabetic cardiomyopathy. *Diabetes* 2005;54:1829-37.
- Cai L, Cherian MG. Zinc-metallothionein protects from DNA damage induced by radiation better than glutathione and copper- or cadmium-metallothioneins. *Toxicol Lett* 2003;136:193-8.
- Ye B, Maret W, Vallee BL. Zinc metallothionein imported into liver mitochondria modulates respiration. *Proc Natl Acad Sci U S A* 2001;10.
- Beattie JH, Wood AM, Newman AM, et al. Obesity and hyperleptinemia

- in metallothionein (-I and -II) null mice. *Proc Natl Acad Sci U S A* 1998;95:358-63.
- Beattie JH, Wood AM, Trayhurn P, et al. Metallothionein is expressed in adipocytes of brown fat and is induced by catecholamines and zinc. *Am J Physiol Regul Integr Comp Physiol* 2000;278:R1082-9.
- Inoue K, Takano H, Shimada A, et al. Role of metallothionein in coagulatory disturbance and systemic inflammation induced by lipopolysaccharide in mice. *FASEB J* 2006;20:533-5.
- Inoue K, Takano H, Yanagisawa R, et al. Role of metallothionein in antigen-related airway inflammation. *Exp Biol Med (Maywood)* 2005;230:75-81.
- Jiang Y, Kang YJ. Metallothionein gene therapy for chemical-induced liver fibrosis in mice. *Mol Ther* 2004;10:1130-9.
- Seth D, Leo MA, McGuinness PH, et al. Gene expression profiling of alcoholic liver disease in the baboon (*Papio hamadryas*) and human liver. *Am J Pathol* 2003;163:2303-17.
- Wang L, Zhou Z, Saari JT, Kang YJ. Alcohol-induced myocardial fibrosis in metallothionein-null mice: prevention by zinc supplementation. *Am J Pathol* 2005;167:337-44.
- Cherian MG, Kang YJ. Metallothionein and liver cell regeneration. *Exp Biol Med (Maywood)* 2006;231:138-44.
- Bay BH, Jin R, Huang J, Tan PH. Metallothionein as a prognostic biomarker in breast cancer. *Exp Biol Med (Maywood)* 2006;231:1516-21.
- Murphy BJ. Regulation of malignant progression by the hypoxia-sensitive transcription factors HIF-1 $\alpha$  and MTF-1. *Comp Biochem Physiol B Biochem Mol Biol* 2004;139:495-507.
- Jin R, Bay B, Tan P, Tan BK. Metallothionein expression and zinc levels in invasive ductal breast carcinoma [In Process Citation]. *Oncol Rep* 1999;6:871-5.
- Jasani B, Schmid KW. Significance of metallothionein overexpression in human tumors. *Histopathology* 1997;31:211-4.
- Datta J, Majumder S, Kutay H, et al. Metallothionein expression is suppressed in primary human hepatocellular carcinomas and is mediated through inactivation of CCAAT/enhancer binding protein { $\alpha$ } by phosphatidylinositol 3-kinase signaling cascade. *Cancer Res* 2007;67:2736-46.
- Cherian MG, Jayasurya A, Bay BH. Metallothioneins in human tumors and potential roles in carcinogenesis. *Mutat Res* 2003;533:201-9.
- Lee SS, Yang SF, Ho YC, Tsai CH, Chang YC. The upregulation of metallothionein-1 expression in areca quid chewing-associated oral squamous cell carcinomas. *Oral Oncol* 2008;44:180-6.
- Lazo JS, Kuo SM, Woo ES, Pitt BR. The protein thiol metallothionein as an antioxidant and protectant against antineoplastic drugs. *Chem Biol Interact* 1998;111-2:255-62.
- Yang M, Kroft SH, Chitambar CR. Gene expression analysis of gallium-resistant and gallium-sensitive lymphoma cells reveals a role for metal-responsive transcription factor-1, metallothionein-2A, and zinc transporter-1 in modulating the antineoplastic activity of gallium nitrate. *Mol Cancer Ther* 2007;6:633-43.
- Cai L, Satoh M, Tohyama C, Cherian MG. Metallothionein in radiation exposure: its induction and protective role. *Toxicology* 1999;132:85-98.
- Laitly JH, Andrews GK. Understanding the mechanisms of zinc-sensing by metal-response element binding transcription factor-1 (MTF-1). *Arch Biochem Biophys* 2007;463:201-10.
- Westin G, Schaffner W. A zinc-responsive factor interacts with a metal-regulated enhancer element (MRE) of the mouse metallothionein-I gene. *EMBO J* 1988;7:3763-70.
- Jiang H, Daniels PJ, Andrews GK. Putative zinc-sensing zinc fingers of metal-response element-binding transcription factor-1 stabilize a metal-dependent chromatin complex on the endogenous metallothionein-I promoter. *J Biol Chem* 2003;278:30394-402.
- Daniels PJ, Andrews GK. Dynamics of the metal-dependent transcription factor complex *in vivo* at the mouse metallothionein-I promoter. *Nucleic Acids Res* 2003;31:6710-21.
- Potter BM, Feng LS, Parasuram P, et al. The six zinc fingers of metal-responsive element-binding transcription factor-1 form stable and quasi-ordered structures with relatively small differences in zinc affinities. *J Biol Chem* 2005;280:28529-40.
- Li Y, Kimura T, Laitly JH, Andrews GK. The zinc-sensing mechanism of mouse MTF-1 involves linker peptides between the zinc fingers. *Mol Cell Biol* 2006;26:5580-7.
- Radtke F, Heuchel R, Georgiev O, et al. Cloned transcription factor MTF-1 activates the mouse metallothionein I promoter. *EMBO J* 1993;12:1355-62.
- Wang Y, Lorenzi I, Georgiev O, Schaffner W. Metal-responsive transcription factor-1 (MTF-1) selects different types of metal response elements at low vs. high zinc concentration. *Biol Chem* 2004;385:623-32.

37. Selvaraj A, Balamurugan K, Yepiskoposyan H, et al. Metal-responsive transcription factor (MTF-1) handles both extremes, copper load and copper starvation, by activating different genes. *Genes Dev* 2005;19:891–6.
38. Dalton T, Palmiter RD, Andrews GK. Transcriptional induction of the mouse metallothionein-I gene in hydrogen peroxide-treated Hepa cells involves a composite major late transcription factor/antioxidant response element and metal response promoter elements. *Nucleic Acids Res* 1994;22:5016–23.
39. Dalton T, Fu K, Enders GC, Palmiter RD, Andrews GK. Analysis of the effects of overexpression of metallothionein-I in transgenic mice on the reproductive toxicology of cadmium. *Environ Health Perspect* 1996;104:68–76.
40. Stitt MS, Wasserloos KJ, Tang X, Liu X, Pitt BR, St Croix CM. Nitric oxide-induced nuclear translocation of the metal responsive transcription factor, MTF-1 is mediated by zinc release from metallothionein. *Vasc Pharmacol* 2006;44:149–55.
41. Saydam N, Steiner F, Georgiev O, Schaffner W. Heat and heavy metal stress synergize to mediate transcriptional hyperactivation by metal-responsive transcription factor MTF-1. *J Biol Chem* 2003;278:31879–83.
42. Cousins RJ, Liuzzi JP, Lichten LA. Mammalian zinc transport, trafficking, and signals. *J Biol Chem* 2006;281:24085–9.
43. Murphy BJ, Andrews GK, Bittel D, et al. Activation of metallothionein gene expression by hypoxia involves metal response elements and metal transcription factor-1. *Cancer Res* 1999;59:1315–22.
44. Zhang B, Georgiev O, Hagmann M, et al. Activity of metal-responsive transcription factor 1 by toxic heavy metals and H<sub>2</sub>O<sub>2</sub> *in vitro* is modulated by metallothionein. *Mol Cell Biol* 2003;23:8471–85.
45. Castello PR, David PS, McClure T, Crook Z, Poyton RO. Mitochondrial cytochrome oxidase produces nitric oxide under hypoxic conditions: implications for oxygen sensing and hypoxic signaling in eukaryotes. *Cell Metab* 2006;3:277–87.
46. Lichtlen P, Wang Y, Belsler T, et al. Target gene search for the metal-responsive transcription factor MTF-1. *Nucleic Acids Res* 2001;29:1514–23.
47. Green CJ, Lichtlen P, Huynh NT, et al. Placenta growth factor gene expression is induced by hypoxia in fibroblasts: a central role for metal transcription factor-1. *Cancer Res* 2001;61:2696–703.
48. Haroon ZA, Amin K, Lichtlen P, et al. Loss of metal transcription factor-1 suppresses tumor growth through enhanced matrix deposition. *FASEB J* 2004;18:1176–84.
49. Wimmer U, Wang Y, Georgiev O, Schaffner W. Two major branches of anti-cadmium defense in the mouse: MTF-1/metallothioneins and glutathione. *Nucleic Acids Res* 2005;33:5715–27.
50. Andrews GK, Lee DK, Ravindra R, et al. The transcription factors MTF-1 and USF1 cooperate to regulate mouse metallothionein-I expression in response to the essential metal zinc in visceral endoderm cells during early development. *EMBO J* 2001;20:1114–22.
51. Wan M, Heuchel R, Radtke F, Hunziker PE, Kagi JH. Regulation of metallothionein gene expression in Cd- or Zn-adapted RK-13 cells. *Experientia* 1995;51:606–11.
52. Mueller PR, Salser SJ, Wold B. Constitutive and metal-inducible protein: DNA interactions at the mouse metallothionein I promoter examined by *in vivo* and *in vitro* footprinting. *Genes Dev* 1988;2:412–27.
53. Dalton T, Qingwen L, Bittel D, Liang L, Andrews GK. Oxidative stress activates metal-responsive transcription factor-1 binding activity. *J Biol Chem* 1996;271:26233–41.
54. Ghoshal K, Datta J, Majumder S, et al. Inhibitors of histone deacetylase and DNA methyltransferase synergistically activate the methylated metallothionein I promoter by activating the transcription factor MTF-1 and forming an open chromatin structure. *Mol Cell Biol* 2002;22:8302–19.
55. Wang GL, Jiang BH, Rue EA, Semenza GL. Hypoxia-inducible factor 1 is a basic-helix-loop-helix-PAS heterodimer regulated by cellular O<sub>2</sub> tension. *Proc Natl Acad Sci U S A* 1995;92:5510–4.
56. Semenza GL. Hydroxylation of HIF-1: oxygen sensing at the molecular level. *Physiology (Bethesda)* 2004;19:176–82.
57. Murphy BJ, Sato BG, Dalton TP, Laderoute KR. The metal-responsive transcription factor-1 contributes to HIF-1 activation during hypoxic stress. *Biochem Biophys Res Commun* 2005;337:860–7.
58. Greijer A, van der Groep P, Kemming D, et al. Up-regulation of gene expression by hypoxia is mediated predominantly by hypoxia-inducible factor 1 (HIF-1). *J Pathol* 2005;206:291–304.
59. Semenza GL, Shimoda LA, Prabhakar NR. Regulation of gene expression by HIF-1. *Novartis Found Symp* 2006;272:2–8; discussion -14, 33–6.
60. Jung JE, Lee HG, Cho IH, et al. STAT3 is a potential modulator of HIF-1-mediated VEGF expression in human renal carcinoma cells. *FASEB J* 2005;19:1296–8.
61. Liao H, Hyman MC, Lawrence DA, Pinsky DJ. Molecular regulation of the PAI-1 gene by hypoxia: contributions of Egr-1, HIF-1{α}, and C/EBP{α}. *FASEB J* 2007;21:935–49.
62. Ema M, Hirota K, Mimura J, et al. Molecular mechanisms of transcription activation by HLF and HIF1α in response to hypoxia: their stabilization and redox signal-induced interaction with CBP/p300. *EMBO J* 1999;18:1905–14.
63. Cramer M, Nagy I, Murphy BJ, et al. NF-κB contributes to transcription of placenta growth factor and interacts with metal responsive transcription factor-1 in hypoxic human cells. *Biol Chem* 2005;386:865–72.
64. Murphy BJ, Laderoute KR, Chin RJ, Sutherland RM. Metallothionein IIA is up-regulated by hypoxia in human A431 squamous carcinoma cells. *Cancer Res* 1994;54:5808–10.
65. Semenza GL. Targeting HIF-1 for cancer therapy. *Nat Rev Cancer* 2003;3:721–32.
66. Semenza GL. Regulation of physiological responses to continuous and intermittent hypoxia by hypoxia-inducible factor 1. *Exp Physiol* 2006;91:803–6.
67. Semenza GL. Development of novel therapeutic strategies that target HIF-1. *Expert Opin Ther Targets* 2006;10:267–80.
68. Semenza GL. Hypoxia-inducible factor 1: control of oxygen homeostasis in health and disease. *Pediatr Res* 2001;49:614–7.
69. Semenza GL. Expression of hypoxia-inducible factor 1: mechanisms and consequences. *Biochem Pharmacol* 2000;59:47–53.
70. LaRochelle O, Gagne V, Charron J, Soh JW, Seguin C. Phosphorylation is involved in the activation of metal-regulatory transcription factor 1 in response to metal ions. *J Biol Chem* 2001;276:41879–88.
71. Saydam N, Adams TK, Steiner F, Schaffner W, Freedman JH. Regulation of metallothionein transcription by the metal-responsive transcription factor MTF-1: identification of signal transduction cascades that control metal-inducible transcription. *J Biol Chem* 2002;277:20438–45.
72. Lee JY, Lee YS, Kim JM, et al. A novel chimeric promoter that is highly responsive to hypoxia and metals. *Gene Ther* 2006;13:857–68.
73. Smirnova IV, Bittel DC, Ravindra R, Jiang H, Andrews GK. Zinc and cadmium can promote rapid nuclear translocation of metal response element-binding transcription factor-1. *J Biol Chem* 2000;275:9377–84.
74. Laderoute KR, Calaoagan JM, Knapp M, Johnson RS. Glucose utilization is essential for hypoxia-inducible factor 1 α-dependent phosphorylation of c-Jun. *Mol Cell Biol* 2004;24:4128–37.
75. Gunes C, Heuchel R, Georgiev O, et al. Embryonic lethality and liver degeneration in mice lacking the metal-responsive transcriptional activator MTF-1. *EMBO J* 1998;17:2846–54.
76. Jiang H, Fu K, Andrews GK. Gene- and cell-type-specific effects of signal transduction cascades on metal-regulated gene transcription appear to be independent of changes in the phosphorylation of metal-response-element-binding transcription factor-1. *Biochem J* 2004;382:33–41.
77. Ryan HE, Poloni M, McNulty W, et al. Hypoxia-inducible factor-1α is a positive factor in solid tumor growth. *Cancer Res* 2000;60:4010–5.