

Fumonisin B₁ as a Urinary Biomarker of Exposure in a Maize Intervention Study Among South African Subsistence Farmers

Liana van der Westhuizen¹, Gordon S. Shephard¹, Hester M. Burger^{1,2}, John P. Rheeder¹, Wentzel C.A. Gelderblom^{1,2}, Christopher P. Wild³, and Yun Yun Gong⁴

Abstract

Background: The consumption of maize highly contaminated with carcinogenic fumonisins has been linked to high oesophageal cancer rates. The aim of this study was to validate a urinary fumonisin B₁ (UFB₁) biomarker as a measure of fumonisin exposure and to investigate the reduction in exposure following a simple and culturally acceptable intervention.

Methods: At baseline home-grown maize, maize-based porridge, and first-void urine samples were collected from female participants ($n = 22$), following their traditional food practices in Centane, South Africa. During intervention the participants were trained to recognize and remove visibly infected kernels, and to wash the remaining kernels. Participants consumed the porridge prepared from the sorted and washed maize on each day of the two-day intervention. Porridge, maize, and urine samples were collected for FB₁ analyses.

Results: The geometric mean (95% confidence interval) for FB₁ exposure based on porridge (dry weight) consumption at baseline and following intervention was 4.84 (2.87–8.14) and 1.87 (1.40–2.51) $\mu\text{g FB}_1/\text{kg body weight/day}$, respectively, (62% reduction, $P < 0.05$). UFB_{1C}, UFB₁ normalized for creatinine, was reduced from 470 (295–750) at baseline to 279 (202–386) pg/mg creatinine following intervention (41% reduction, $P = 0.06$). The UFB_{1C} biomarker was positively correlated with FB₁ intake at the individual level ($r = 0.4972$, $P < 0.01$). Urinary excretion of FB₁ was estimated to be 0.075% (0.054%–0.104%) of the FB₁ intake.

Conclusion: UFB₁ reflects individual FB₁ exposure and thus represents a valuable biomarker for future fumonisin risk assessment.

Impact: The simple intervention method, hand sorting and washing, could positively impact on food safety and health in communities exposed to fumonisins. *Cancer Epidemiol Biomarkers Prev*; 20(3); 483–9. ©2011 AACR.

Introduction

Fumonisin is a group of mycotoxins; mainly produced by *Fusarium* spp. as secondary metabolites, of which fumonisin B₁, B₂, and B₃ (FB₁, FB₂, and FB₃) are the most prevalent (1). Fumonisin is carcinogenic to rodents, and have been associated with oesophageal and liver cancer in subsistence maize farming communities around the world, as well as occurrence of neural tube

defects (2, 3, 4, 5). There are numerous fumonisin analogues of which FB₁ is the most abundant naturally occurring fumonisin on maize and maize-based food worldwide and was classified as "possibly carcinogenic to humans" by the International Agency for Research on Cancer (6). The Joint FAO/WHO Expert Committee on Food Additives (JECFA) has determined a group provisional maximum tolerable daily intake (PMTDI) for FB₁, FB₂, and FB₃, alone or in combination, of 2 $\mu\text{g/kg body weight/day}$ (7).

In many Sub-Saharan countries, where both maize contamination and maize consumption are high, regulatory mechanisms to control mycotoxin levels, including fumonisin, are either lacking or are not enforced (8). Therefore, reducing exposure levels by intervention, specifically those based on simple low-cost measures acceptable to these communities, becomes critical to protect the population at greatest risk (9). Such an approach gave promising results with respect to aflatoxins in groundnuts (10).

Conventional assessment of mycotoxin exposure, probable daily intake (PDI), is based on the amount of

Authors' Affiliations: ¹PROMEC Unit, Medical Research Council, Tygerberg, South Africa; ²Department of Biochemistry, University of Stellenbosch, Matieland, South Africa; ³International Agency for Research on Cancer, Lyon, France; and ⁴Molecular Epidemiology Unit, Centre for Epidemiology and Biostatistics, Leeds Institute of Genetics, Health and Therapeutics, University of Leeds, United Kingdom

Corresponding Author: Yun Yun Gong, Molecular Epidemiology Unit, Centre for Epidemiology and Biostatistics, Leeds Institute of Genetics, Health and Therapeutics, University of Leeds, LS2 9JT. Phone: 44-113-3437757; Fax: 44-113-3436603. Email: medyg@leeds.ac.uk

doi: 10.1158/1055-9965.EPI-10-1002

©2011 American Association for Cancer Research.

food consumed and the contamination level of the food (11). Accuracy of the PDI can be improved by determining food consumption and the level of the mycotoxin in the food on an individual basis (12). The quality of food consumption information in the subsistence community setting depends on the validity of the assessment methods and measuring of food on the plate prior to and after food consumption, a labour-intensive activity (11). Utilization of validated biomarkers of exposure, taking advantage of measuring individual internal dose, would not require measurement of consumption or contamination level of the food and thus promises to be more accurate than estimations by other assessment methods (13). Aflatoxin, deoxynivalenol, and ochratoxin A, urinary biomarkers, as well as the aflatoxin-albumin adduct in blood, have been implemented in human studies (14–16). The aflatoxin biomarker applications in human health and intervention studies have demonstrated the value of mycotoxin-related biomarkers (13).

Various animal studies have successfully investigated the sphinganine/sphingosine ratio as a biomarker of fumonisin exposure (11, 17). The sphingolipid bases, sphinganine and sphingosine, as well as their ratio, have also been investigated in several human studies in blood and urine, but could not be correlated with fumonisin exposure (11, 12, 18). A recent study has reported a high-performance liquid chromatography–mass spectrometry (HPLC-MS) method for urinary fumonisin B₁ (UFB₁), which was sufficiently sensitive to be positively correlated with fumonisin exposure in a Mexican population consuming different amounts of maize-based tortillas (19).

In a previous study, customary sorting and washing of maize kernels as food preparation procedures were

optimized under laboratory-controlled conditions to achieve optimal fumonisin reduction in contaminated home-grown maize (20). These simple and culturally acceptable food preparation practices reduced fumonisin exposure as assessed by food intake profiles and fumonisin food analysis in a subsistence farming community residing in a high oesophageal cancer incidence area (21). The aim of this study was to validate the UFB₁ biomarker and confirm the reduction in fumonisin exposure at an individual level.

Material and Methods

Participants

The study was approved by the Ethics Committee of the Medical Research Council of South Africa. Following informed consent, apparently healthy females (*n* = 22) aged between 20 and 70 who prepared maize-based meals from home-grown maize, were recruited from the Centane magisterial district, Eastern Cape Province, South Africa.

Baseline phase of the field study

At baseline, participants consumed their customarily prepared maize-based food (~0.5 kg porridge) twice daily for 2 consecutive days (Table 1). Porridge samples (~0.5 kg) were collected on both days for FB₁ analyses. Morning first-void urine samples were collected from each participant on the subsequent days for FB₁ biomarker analysis. The first-void urine collections were done approximately 12 hours after the participants consumed the last porridge meal. Dietary recall questionnaires were recorded to determine maize intake (dry weight) in the 24 hours prior to urine collection (21). Home-grown maize

Table 1. Both the baseline and intervention phases of the study were conducted over 3 consecutive days for each participant. Morning first-void urine samples were collected from the participant individually on each day following the twice daily consumption of the porridge. The training was conducted following the completion of the baseline phase preceding the intervention phase of the study.

Baseline			Training	Intervention		
Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
Consume porridge	Consume porridge	Collect first-void urine	Sorting/washing of maize	Consume porridge	Consume porridge	Collect first-void urine
	Collect first-void urine	Collect first-void urine	Collect sorted/washed maize	Collect first-void urine	Collect first-void urine	Collect first-void urine

samples (~5 kg) were collected from each participant of which subsamples (~0.5 kg) were retained for FB₁ analyses. The remaining kernels were pooled, thoroughly mixed, divided into 4-kg batches ($n = 22$), and set aside for the intervention study.

Intervention phase of the field study

Participants were trained to recognize infected kernels (most likely to be contaminated) and on the washing procedure (10 minutes) for the remaining maize kernels. Following training, each participant first removed visibly infected kernels and subjected the remaining maize kernels to a washing step. Subsamples of the sorted and washed maize of each participant were collected for FB₁ analysis and the remainder of the maize kernels pooled for porridge preparation. On the first day of the intervention phase porridge was prepared and weighed as 0.5 kg portions, which were consumed as 2 separate meals (midday and evening) by the participants. First-void urine was collected on the following morning and 24-hour dietary recall questionnaires completed as described previously. The procedure was repeated on the second day of the intervention.

All the maize and porridge samples were stored at 4°C and -20°C, respectively prior to FB₁ analysis at the PROMEC Unit. Urine samples were stored at -20°C and sent on dry ice for FB₁ analysis at the University of Leeds.

Analyses

The maize and porridge samples were analyzed for FB₁ following solid phase extraction clean-up, derivatization with *o*-phthalaldehyde, and detection by fluorescence HPLC (21, 22). The urine samples were cleaned-up by solid-phase extraction and analyzed for FB₁ by HPLC-MS with a limit of detection (LOD) of 20 pg/mL (19). Urinary creatinine was determined according to the alkaline-picric acid method (23) with minor modifications to adapt to a 96-well plate format. Inter-individual variation in urine concentration was normalized by utilization of urinary creatinine. The biomarker data are presented as both pg FB₁/mL urine (UFB₁) and pg FB₁/mg creatinine (UFB₁C).

Probable daily intake assessment

Porridge (dry weight) consumption was assessed by 24-hour dietary recall questionnaires utilizing full-scale photographs of small, medium, and large porridge portions. Individual maize intakes (dry weight, g/day) were estimated using the local maize porridge recipes and by assigning specific weights to the portions sizes (21). Individual fumonisin PDI was assessed as FB₁ level in the porridge (dry weight) consumed by each participant during the baseline and intervention phases of the study.

FB₁ excretion in urine

The 24-hour urine output was estimated by assuming that the first-void urine volume represented an 8-hour

collection. The urinary FB₁ excretion was calculated as follows:

$$\text{Excretion (\%)} = \frac{\text{UFB}_1 \text{ (pg/mL)} \times (8\text{-hour urine output} \times 3) \text{ (mL/day)}}{\text{PDI} \text{ (}\mu\text{g/kg body weight/day)} \times \text{body weight (kg)} \times 10^6} \times 100$$

Statistical analysis

Individual FB₁ PDI and UFB₁ were assessed as the mean of 2 days. UFB₁C data was not normally distributed and was natural log-transformed to facilitate data analysis. Mean FB₁ levels in urine and maize were expressed geometrically (95% confidence interval) unless otherwise stated. Student's *t* test was used to compare the FB₁ exposure at baseline and following intervention. Correlation and regression analyses were performed to examine the association between the PDI and UFB₁ or UFB₁C level. Urine samples below the LOD were assigned a value of half the LOD. Multiple linear regression modeling was used to investigate the independent contribution of FB₁ PDI and age to the biomarker levels. STATA9.0 (STATA corp) and Number Cruncher Statistical Systems (NCSS) 2004 Dawson Edition (NCSS) were used for the analysis.

Results

Participants

The mean age of the participants was 42 years (range 20–70), their mean body weight was 65 (47–127) kg and their mean maize consumption (dry weight) at baseline was 0.34 (0.28–0.40) kg/day and following intervention it was 0.38 (0.31–0.47) kg/day as reported in the previous study (20). The mean FB₁ level in home-grown maize at baseline [1.16 (0.84–1.56) mg/kg] was significantly reduced (84%, $P < 0.05$) following the sorting and washing of the kernels during intervention [0.19 (0.14–0.25) mg/kg]. The FB₁ represented 72% of the total fumonisins (FB₁ + FB₂ + FB₃) in the maize (FB₂ and FB₃ levels were reported in a previous study; ref. 20). The porridge at baseline had a mean FB₁ of 0.89 (0.55–1.44) mg/kg compared with 0.32 (0.27–0.37) mg/kg following intervention (64% reduction, $P < 0.05$; Table 2).

PDI assessment

The mean PDI of FB₁ at baseline was 4.84 (2.87–8.14) $\mu\text{g/kg}$ body weight/day and following intervention the PDI was significantly reduced (62%, $P < 0.05$) to 1.87 (1.40–2.51) $\mu\text{g/kg}$ body weight/day (Table 2). The mean FB₁ exposure at baseline exceeded the JECFA recommended PMTDI (2 $\mu\text{g/kg}$ body weight/day), and was below this PMTDI following intervention. Before intervention 15 of 21 (71%) participants had PDIs exceeding the recommended PMTDI, whereas this frequency was reduced to 10 of 19 (53%) following intervention ($p > 0.05$). However, FB₁ represented only 72% of the total fumonisins in the maize, and the PMTDI determined

Table 2. The geometric means (95% confidence limits) of FB₁ levels in porridge and urine, as well as PDI in Centane, a rural area from the Eastern Cape Province of South Africa.

	FB ₁ porridge, mg/kg	FB ₁ PDI ^a , µg/kg b.w./day	UFB ₁ , pg/mL	UFB ₁ C ^b , pg/mg creatinine
Baseline	0.89a (0.55–1.44)	4.84a (2.87–8.14)	225a (144–350)	470a (295–750)
Intervention	0.32b (0.27–0.37)	1.87b (1.40–2.51)	109b (85–138)	279a (202–386)
Reduction	64%	62%	52%	41%

^aPDI based on FB₁ levels in porridge (dry weight) consumed as assessed with 24-hour dietary recall questionnaires

^bUFB₁ was normalized with urinary creatinine

Means are significantly ($P < 0.05$) different within columns when followed by different letters

by JECFA was based on FB₁, FB₂, and FB₃, alone or in combination.

Urine

At baseline 43 of 44 (98%) and following intervention 40 of 42 (96%) of the urine samples had FB₁ above the LOD. Following intervention, the mean UFB₁ was reduced from 225 (144–350) to 109 (85–138) pg/mL, a 52% reduction ($P = 0.02$; Table 2). Following normalization with urinary creatinine, UFB₁C at baseline and following intervention was 470 (295–750) and 279 (202–386) pg/mg creatinine, respectively, (41% reduction, $P = 0.06$). Individual FB₁ intake to UFB₁C at baseline and intervention combined is significantly correlated ($r = 0.4972$, $P < 0.01$). In addition, the linear regression relationship between FB₁ intake and UFB₁C, at baseline only and at baseline and intervention combined, is shown in Figure 1. There is a significant positive correlation between FB₁ intake and UFB₁C level (regression coefficient 0.4154, $P < 0.01$). However, the multiple regression model is better fitted when age, a positive contributor (regression coefficient 0.0227, $P < 0.01$), is introduced as

the older women were apparently exposed to higher FB₁ levels.

FB₁ excretion in urine

The urine output as well as the urinary excretion was similar at baseline and following intervention. The estimated mean 24-hour urine output at baseline and during intervention combined was 933 (793–1,099) mL/day. The mean percentage FB₁ excreted in urine, based on FB₁ intake and the estimated mean daily urine output was 0.075%/day (0.054%/day to 0.104%/day).

Discussion

This is the first published study describing a quantitative correlation between a fumonisin biomarker and intake of this mycotoxin measured at the individual level. The investigation provides further validation following a previous report showing a positive correlation between the FB₁ urinary biomarker and different levels of maize-based tortilla consumption in a Mexican population (19). In addition, the current investigation applied an




Figure 1. The relationship between UFB₁C and FB₁ intake (natural log-transformed data) at baseline (left) and at baseline and intervention combined (right) is shown. The lines show the predicted linear regression for UFB₁C and the grey curves indicate the 95% confidence limits.

optimized sorting and washing method to maize, which reduced fumonisin exposure in a subsistence farming community as previously assessed by food analysis and food intake data (21). The biomarker was well correlated with FB₁ exposure at the individual level and confirmed the efficacy of this simple and culturally acceptable intervention method.

Surveys previously conducted in subsistence farming communities in Africa have shown that mycotoxin contamination, such as fumonisin and aflatoxin, can be reduced by hand-sorting and/or washing of maize kernels (24, 25). Featured in this paper and our previous publication (21), the first intervention study to reduce fumonisin exposure by implementing these practices has been conducted recently in a rural subsistence-farming community in South Africa. The customary sorting and washing of maize kernels as food preparation procedures were optimized under laboratory-controlled conditions. The participants were trained to apply the simple 2-step method by identifying the infected maize kernels to ensure proper selection for removal and to follow the correct washing procedures. The intervention method is practical and culturally acceptable as it represents the optimization of existing local practices. The FB₁ level in the maize-based porridge, prepared from the maize sorted and washed by the participants following training, was significantly ($P < 0.05$) reduced by 64% compared with the porridge prepared by the participants prior to training. The intervention study method effectively reduced the PDIs of FB₁ by 62% (21).

The tortillas the participants consumed in the Mexican study were prepared from maize kernels which were boiled in alkaline water followed by prolonged steeping in fresh water (19, 26). Subsequently, the cooked kernels were washed and ground to yield masa from which the tortillas were prepared. The cooking of kernels in alkaline water partially converts the fumonisins to their hydrolyzed analogues, and therefore the tortillas contain both fumonisins and hydrolyzed fumonisins. The geometric mean UFB₁C (134 pg/mg creatinine) in the high tortilla consumption group in the Mexican study (19) was more than 3-fold lower than at baseline in Centane (470 pg/mg creatinine). Based on the UFB₁ level in the high tortilla consumption group, FB₁ exposure was estimated to be 0.37 µg/kg body weight/day assuming a 1,500 mL/day urine output, a 60 kg body weight, and an assumption of a 1% excretion of ingested FB₁ in urine, based on data in swine and nonhuman primates (27, 28). It is notable that in the current study the measured urinary excretion of FB₁ was significantly lower than 1% (see the following text) and hence the intakes in Mexico may have been underestimated if urinary FB₁ excretion in the 2 populations is similar. Re-estimating the PDI in the Mexican study, based on the estimated mean 24-hour urine output and the urine excretion rate estimated for this study, the FB₁ PDI in Mexico (3.04 µg/kg body weight/day) was slightly lower than observed in Centane (4.83 µg/kg body weight/day). A recent study conducted in China with

male and female participants reported median UFB₁C levels of 390 and 3,910 pg/mg creatinine, respectively, in a high-risk hepatocellular carcinoma area (Fusui) and in a high-risk oesophageal cancer area (Huaian; ref. 18). The median level reported for Fusui was similar to Centane (451 pg/mg creatinine), whereas the median level in Huaian was more than 8-fold higher.

The mean UFB₁ excretion in the Centane population, based on individual body weight, an estimated urine output, the actual FB₁ content of the food consumed, and the food consumption (dry weight) was 0.075%. The correlation between FB₁ intake and the urinary biomarker is influenced by the low proportion of FB₁ excreted in urine and the interindividual variation of absorption and excretion of FB₁ (19, 28, 29). This variation was observed in vervet monkeys where toxicokinetic studies revealed individual variations in urinary excretion with values of 0.25%, 0.66%, 1.0%, and 1.5% (28, 29). Experimental studies in rodents and swine reported urinary excretion of 0.4%–2% (27, 30). A more recent study in swine reported a 0.9% urinary excretion with peak urinary excretion between 8 and 24 hours following administration of a single oral dose of 5 mg FB₁/kg (31). In a preliminary human study in the United States, participants consumed maize-based food that approximated maize consumption in urban Guatemala (32). The food was prepared from commercial masa and maize flour, containing 0.8 to 2.5 mg FB₁/kg. The urinary excretion of less than 1% FB₁ measured in the U.S. study was 10-fold higher than the less than 0.075% measured in this current study. It remains possible that there are ethnic differences or food matrix effects, which influence urinary FB₁ excretion in human populations. Future studies to measure FB₁ PDI in population would certainly be useful to improve our understanding on individual and ethnic variation in FB₁ excretion through urine.

Due to the rapid urinary excretion rate of FB₁, first-void urine was collected 12 hours following the last porridge meal consumed by the participants on 2 consecutive days at both baseline and intervention phases of the study. In a rat study following a single FB₁ gavage dose (25 mg/kg body weight) UFB₁ levels peaked at 12 hours, whereafter levels declined rapidly, but FB₁ was still detectable 10 days later (30). This study also suggested that UFB₁ might reflect long-term exposure with chronic FB₁ exposure at low levels in rats (1.0 mg/kg body weight/day). Therefore, residual FB₁ from the previous days' maize consumption in the current study could have contributed to the measured UFB₁. Sources of FB₁ other than maize could also have contributed to the individual exposure. However, as this subsistence community is reliant on maize almost to the exclusion of all other food commodities, the contribution of other sources is expected to be insignificant. This was verified by utilizing a validated questionnaire specifically designed for the local customary diet practices to obtain individual maize-based food consumption (33). The 52% reduction ($P < 0.05$) observed

in UFB₁ (41% in UFB₁C) levels following intervention were comparable to the 62% reduction in FB₁ exposure as previously assessed by food analysis and food intake data (21). Indeed one of the advantages of assessing the intervention using a biomarker is that it provides a measure of the effect of the intervention on FB₁ exposure from all dietary sources, whereas, by definition, the maize and food analysis indicates only the effect on the target commodity itself.

Conclusion

The FB₁ urinary biomarker is quantitatively correlated with FB₁ exposure at the individual level. The biomarker permitted confirmation of the reduction in fumonisin exposure achieved with the culturally acceptable intervention method of hand sorting and washing maize. Utilization of this biomarker will improve assessment of fumonisin exposure and thus contribute to the assessment of the possible health impacts of fumonisin exposure as well as permitting evaluation of intervention strategies to reduce fumonisin exposure. Future intervention studies could be expanded to include larger numbers of both male and female participants including children. Significant advances in food safety and health in subsistence maize farming communities exposed to high

levels of fumonisin could be possible by further development of these approaches.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

The authors thank Mr. John Mokotary for driving expertise and technical assistance, Ms. N Kulati, for leadership and IsiXhosa translations, Mss. N Mpetsheni, N Mbana, and NS Kulati for recruiting and interviewing of the participants, Mr. R Hulley and his family for their invaluable assistance as well as the participants without whom the study would not be possible.

We thank Professor Hans-Ulrich Humpf of University of Münster for his kind support in providing the internal standard for the biomarker analysis.

Grant Support

The authors acknowledge support from the Sir Halley Stewart Trust. Prof. CP Wild and Dr Yun Yun Gong were supported by NIEHS, USA grant no. ES06052, and Ms L van der Westhuizen was in receipt of an ICRET fellowship from the International Union Against Cancer (IUGC).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received September 21, 2010; revised December 21, 2010; accepted January 9, 2011; published OnlineFirst January 25, 2011.

References

- Marasas WF. Discovery, occurrence of the fumonisins: a historical perspective. *Environ Health Perspect* 2001;109:239–43.
- Gelderblom WCA, Abel S, Smuts CM, Marnewick JL, Marasas WFO, Lemmer ER, et al. Fumonisin-induced hepatocarcinogenesis: Mechanisms related to cancer initiation and promotion. *Environ Health Perspect* 2001;109:291–300.
- Marasas WFO, Riley RT, Hendricks KA, Stevens VL, Sadler TW, Gelineau-van Waes J, et al. Fumonisins disrupt sphingolipid metabolism, folate transport, and neural tube development in embryo culture and *in vivo*: a potential risk factor for human neural tube defects among populations consuming fumonisin contaminated maize. *J Nutr* 2004;134:711–6.
- Sun G, Wang S, Hu X, Su J, Huang T, Yu J, et al. Fumonisin B₁ contamination of home-grown corn in high-risk areas for esophageal and liver cancer in China. *Food Addit Contam* 2007;24:181–5.
- Van Der Westhuizen L, Shephard GS, Scussel VM, Costa LL, Vismar HF, Rheeder JP, et al. Fumonisin contamination and *Fusarium* incidence in corn from Santa Catarina, Brazil. *J Agric Food Chem* 2003;51:5574–8.
- IARC. Some traditional herbal medicines, some mycotoxins, naphthalene and styrene. In: Monographs on the evaluation of carcinogenic risks to humans. Lyon (France): International Agency for Research on Cancer Press;2002. p.301–66.
- Bolger M, Coker RD, DiNovi M, Gaylor D, Gelderblom WC, Olsen M, et al. Fumonisins. In: Safety evaluation of certain mycotoxins in food. Food Additives Series No. 47, FAO Food and Nutrition Paper No. 47, Prepared for the 56th Meeting of the Joint FAO/WHO Expert Committee on Food Additives (JECFA). Geneva (Switzerland): World Health Organization (WHO);2001. p.103–279.
- Gelderblom WCA, Riedel S, Burger HM, Abel S, Marasas WFO. Carcinogenesis by the fumonisins: mechanisms, risk analyses, and implications. In: Siantar DP, Trucksess MW, Scott PM, Herman EM, editors. Food Contaminants, mycotoxins and food allergens. ACS Symposium Series 1001;2008. p.80–95.
- Wild CP, Gong YY. Mycotoxins and human disease: a largely ignored global health issue. *Carcinogenesis* 2010;31:71–82.
- Turner PC, Sylla A, Gong YY, Diallo MS, Sutcliffe AE, Hall AJ, et al. Reduction in exposure to carcinogenic aflatoxins by postharvest intervention measures in west Africa: a community-based intervention study. *Lancet* 2005;365:1950–6.
- Shephard GS, Van Der Westhuizen L, Sewram V. Biomarkers of exposure to fumonisin mycotoxins: A review *Food Addit Contam* 2007;24:1196–201.
- Van Der Westhuizen L, Shephard GS, Rheeder JP, Burger HM. Individual fumonisin exposure and sphingoid base levels in rural populations consuming maize in South Africa. *Food Chem Toxicol* 2010a;48:1698–703.
- Wild CP, Turner PC. The toxicology of aflatoxins as a basis for public health decisions. *Mutagenesis* 2002;17:471–81.
- Akdemir C, Ulker OC, Basaran A, Ozkaya S, Karakaya A. Estimation of ochratoxin A in some Turkish populations: an analysis in urine as a simple, sensitive and reliable biomarker. *Food Chem Toxicol* 2010;48:877–82.
- Meky FA, Turner PC, Ashcroft AE, Miller JD, Qiao YL, Roth MJ, et al. Development of a urinary biomarker of human exposure to deoxynivalenol. *Food Chem Toxicol* 2003;41:265–73.
- Wild CP, Hasegawa R, Barraud L, Chutimatawin S, Chapot B, Ito N, et al. Aflatoxin-albumin adducts: a basis for comparative carcinogenesis between animals and man. *Cancer Epidemiol Biomarkers Prev* 1996;5:179–89.
- Riley RT, Enongene E, Voss KA, Norred WP, Meredith FI, Sharma RP, et al. Sphingolipid perturbations as mechanisms for fumonisin carcinogenesis. *Environ Health Perspect* 2001;109:301–8.
- Xu L, Cai Q, Tang L, Wang S, Hu X, Su J, Sun G, Wang J-S. Evaluation of fumonisin biomarkers in a cross-sectional study with two high-risk populations in China. *Food Addit Contam* 2010;27:1161–9.
- Gong YY, Torres-Sanchez L, Lopez-Carrillo L, Peng JH, Sutcliffe AE, White KL, et al. Association between tortilla consumption and human

- urinary fumonisin B₁ levels in a Mexican population. *Cancer Epidemiol Biomarkers Prev* 2008;17:688–94.
20. Van Der Westhuizen L, Shephard GS, Burger HM, Rheeder JP, Gelderblom WCA, Wild CP, et al. Optimising sorting and washing of home-grown maize to reduce fumonisin contamination under laboratory-controlled conditions. *Food Control* 2011;22:396–400.
 21. Van Der Westhuizen L, Shephard GS, Burger HM, Rheeder JP, Gelderblom WCA, Wild CP, et al. Implementation of a simple intervention method to reduce fumonisin exposure in a subsistence maize farming community of South Africa. *Food Addit Contam* 2010b;27:1582–8.
 22. Shephard GS, Leggott NL, Stockenström S, Somdyala NIM, Marasas WFO. Preparation of South African maize porridge: effect on fumonisin mycotoxin levels. *S Afr J Sci* 2002;98:393–6.
 23. Varley H. *Practical clinical biochemistry*. London (NY): William Weimann Medical Books and Interscience Books, Inc. 1967, p.35–6.
 24. Fandohan P, Zoumenou D, Hounhouigan DJ, Marasas WFO, Wingfield MJ, Hell K. Fate of aflatoxins and fumonisins during the processing of maize into food products in Benin. *Int J Food Microbiol* 2005;98:249–59.
 25. Kimanya ME, De Meulenaer B, Tiisekwa B, Ugullum C, Devlieghere F, Van Camp J, et al. Fumonisin exposure from freshly harvested and stored maize and its relationship with traditional agronomic practices in Rombo district, Tanzania. *Food Addit Contam* 2009;26:1199–208.
 26. Dombink-Kurtzman MA, Dvorak TJ. Fumonisin Content in Masa and Tortillas from Mexico. *J Agric Food Chem* 1999;47:622–7.
 27. Fodor J, Meyer K, Riedlberger M, Bauer J, Horn P, Kovacs F, et al. Distribution and elimination of fumonisin analogues in weaned piglets after oral administration of *Fusarium verticillioides* fungal culture. *Food Addit Contam* 2006;23:492–501.
 28. Shephard GS, Thiel PG, Sydenham EW, Alberts JF, Cawood ME. Distribution and excretion of a single-dose of the mycotoxin fumonisin B₁ in a non-human primate. *Toxicon* 1994;32:735–41.
 29. Shephard GS, Thiel PG, Sydenham EW, Savard ME. Fate of a single dose of ¹⁴C-labelled fumonisin B₁ in vervet monkeys. *Nat Toxins* 1995;3:145–50.
 30. Cai Q, Tang L, Wang JS. Validation of fumonisin biomarkers in F344 rats. *Toxicol Appl Pharmacol* 2007;225:28–39.
 31. Dilkin P, Direito G, Simas MM, Mallmann CA, Corrêa B. Toxicokinetics and toxicological effects of single oral dose of fumonisin B₁ containing *Fusarium verticillioides* culture material in weaned piglets. *Chem Biol Interact* 2010;185:157–62.
 32. Riley RT. The kinetics of urinary fumonisin excretion in humans consuming maize-based foods. *PL 1456. The Toxicologist CD – J Soc Toxicol* 2010;114:308–9.
 33. Burger HM, Lombard MJ, Shephard GS, Rheeder JR, Van Der Westhuizen L, Gelderblom WCA. Dietary fumonisin exposure in a rural population of South Africa. *Food Chem Toxicol* 2010;48:2103–8.