

Targeting Colon Luminal Lipid Peroxidation Limits Colon Carcinogenesis Associated with Red Meat Consumption



Océane C.B. Martin^{1,2}, Nathalie Naud¹, Sylviane Taché¹, Laurent Debrauwer¹, Sylvie Chevolleau¹, Jacques Dupuy¹, Céline Chantelauze³, Denis Durand³, Estelle Pujos-Guillot⁴, Florence Blas-Y-Estrada¹, Christine Urbano⁵, Gunter G.C. Kuhnle⁶, Véronique Santé-Lhoutellier⁷, Thierry Sayd⁷, Didier Viala⁷, Adeline Blot⁸, Nathalie Meunier⁸, Pascal Schlich⁹, Didier Attaix^{4,8}, Françoise Guéraud¹, Valérie Scislowski², Denis E. Corpet¹, and Fabrice H.F. Pierre¹

Abstract

Red meat is probably carcinogenic to humans (WHO/IARC class 2A), in part through heme iron-induced lipoperoxidation. Here, we investigated whether red meat promotes carcinogenesis in rodents and modulates associated biomarkers in volunteers, speculating that an antioxidant marinade could suppress these effects via limitation of the heme induced lipid peroxidation. We gave marinated or non-marinated beef with various degrees of cooking to azoxymethane-initiated rats, *Min* mice, and human volunteers (crossover study). Mucin-depleted foci were scored in rats, adenoma in *Min* mice. Biomarkers of lipoperoxidation were mea-

sured in the feces and urine of rats, mice, and volunteers. The organoleptic properties of marinated meat were tested. Fresh beef increased colon carcinogenesis and lipoperoxidation in rats and mice and lipoperoxidation in humans. Without an adverse organoleptic effect on meat, marinade normalized peroxidation biomarkers in rat and mouse feces, reduced peroxidation in human feces and reduced the number of Mucin-depleted foci in rats and adenoma in female *Min* mice. This could lead to protective strategies to decrease the colorectal cancer burden associated with red meat consumption. *Cancer Prev Res*; 11(9); 569–80. ©2018 AACR.

Introduction

Colorectal cancer is the third most common type of cancer and fourth most common cause of death from cancer worldwide (1). In 2007 and confirmed in 2011 and 2017, the World Cancer Research Fund (WCRF) and American Institute for Cancer Research (AICR) panel stated

that there is strong evidence that consuming red meat increases the risk of colorectal cancer (2, 3). In 2015, the International Agency for Research on Cancer (IARC), an agency of the World Health Organization (WHO), classified red meat consumption as "probably carcinogenic to humans" (4).

Epidemiological studies incited the WCRF to recommend limiting red meat consumption to less than 500 g/w (1, 2). These recommendations may reduce the colorectal cancer burden, but sociological studies have demonstrated that people with a lower socioeconomic status are less receptive to nutritional messages and are less likely to change risky behaviors than affluent people (5–9). Simply conveying information on risks and benefits has almost no effect on food choices among less-educated people and tends to enlarge the health inequalities to which colorectal cancer contributes (10, 11). Furthermore, adherence to the WCRF recommendation of limiting red meat intake is low in cancer survivors (8%; ref. 12). On the basis of these observations, we wanted to propose an alternative to the current recommendation of limiting consumption by adding protective additives directly in the commercial product.

¹INRA UMR1331, TOXALIM (Research Centre in Food Toxicology), Université de Toulouse, ENVT, INP-Purpan, UPS, Toulouse, France. ²ADIV, 10 rue Jacqueline Auriol, Clermont-Ferrand, France. ³INRA, UMR1213 Herbivores, Saint-Genès-Champanelle, France. ⁴Clermont Université, Université d'Auvergne, Unité de Nutrition Humaine, Clermont Ferrand, France. ⁵Sensostat, Dijon, France. ⁶Department of Food & Nutritional Sciences, University of Reading, Reading, United Kingdom. ⁷INRA URO370, QuaPA, QuaPA, Saint-Genès-Champanelle, France. ⁸CHU Clermont Ferrand, CRNH Auvergne, France. ⁹Centre des Sciences du Goût et de l'Alimentation, CNRS, INRA, University of Bourgogne Franche-Comté, Dijon, France.

Note: Supplementary data for this article are available at Cancer Prevention Research Online (<http://cancerprevres.aacrjournals.org/>).

Corresponding Author: Fabrice H.F. Pierre, INRA, Toulouse 31070, France. Phone: 335-8206-6370; Fax: 33-56-128-5244; E-mail: fabrice.pierre@inra.fr

doi: 10.1158/1940-6207.CAPR-17-0361

©2018 American Association for Cancer Research.

Different hypotheses have been explored to explain the link between red meat intake and colorectal cancer, including heterocyclic amine formation during high-temperature cooking (13), *N*-nitroso compound formation by endogenous nitrosation, and high heme iron content, principally through the catalysis of dietary lipid oxidation and formation of *N*-nitroso compounds involving gut microbiota (14, 15). We recently showed that, at nutritional doses, heme iron is the major actor in meat-induced promotion of colon cancer without an additive or synergistic effect of heterocyclic amines and endogenous *N*-nitroso compounds (16). The promoting effect of heme iron has been validated in rodent models using purified molecules (hemoglobin or hemin) and lyophilized red meat (16–20). Furthermore, this promoting effect of heme was confirmed at the epidemiological level in the French E3N prospective cohort of women (21). In humans and in rodent models, meat and heme intake have been associated with an increase in two lipoperoxidation biomarkers: thiobarbituric acid reactive substances (TBARS) and 1,4-dihydroxynonane mercapturic acid (DHN-MA; refs. 18–20, 22–24). Heme iron can catalyze dietary lipoperoxidation, a key downstream feature of ferroptosis, an iron-dependent form of non-apoptotic-regulated cell death (25–27) and a reaction leading to the production of genotoxic and/or cytotoxic aldehydes that could be involved in the promotion of carcinogenesis (16, 28–31). Moreover, we have shown that addition of antioxidants to diet is efficient to decrease meat-induced lipoperoxidation and carcinogenesis in animal models (19, 20, 32). The protective effect of the dietary antioxidant capacity was confirmed in humans in the French E3N prospective cohort of women (21).

We designed a project which aimed to limit the heme-induced colonic luminal lipoperoxidation to prevent the risk of colon cancer associated with red meat consumption. The project was divided in six sequential studies, including an industrial task of meat production and analysis, short-term and carcinogenesis animal studies, and clinical and organoleptic acceptability studies in human volunteers (Supplementary Fig. S1).

Materials and Methods

Ethics

All animal studies were performed in an accredited animal facility by approved staff and animal care conducted in accordance with the ARRIVE and European Council on Animals used in Experimental Studies guidelines. The human volunteer crossover study was allowed by a written decision of the ethical committee (French CPP Sud Est VI) and authorized by the French Ministry of Health (No. IDRCB 2013-A01692-43). The study was registered at Clinicaltrials.gov. under the number NCT02473302.

Study design

First, we selected the most effective marinade among the 12 tested (see Supplementary Materials for details) on the basis of a DPPH assay. The selected grape and olive marinade was then incorporated into cooked or uncooked meats to test the modulation of lipoperoxidation and iron content in meats. A 14-days nutritional experiment (see Supplementary Materials for details) was then performed in Fischer 344 male rats to test the effect of different marinated or non-marinated and cooked or uncooked meats on the modulation of lipoperoxidation and iron content in fecal water, two biomarkers associated with the promotion of colon carcinogenesis by red meat. These six meats were then tested in colorectal carcinogenesis studies in chemically induced rats, and four were selected for study in *Apc Min* mice. From these results, the biological effect of the most effective production process was tested in healthy human volunteers, as well as its organoleptic acceptability. A flowchart of the experiments is presented in Supplementary Fig. S1.

Carcinogenesis study in azoxymethane-induced rats

Male Fischer 344 rats ($N = 92$) purchased at 5 weeks of age from Charles River (France) were housed individually in stainless steel wire-bottom cages. The rats were kept at 22°C with a 12–12 hours light–dark cycle and allowed free access to standard AIN76 semipurified diet and tap water. After acclimatization, the rats received a single intraperitoneal injection of azoxymethane (Sigma, 20 mg/kg) in NaCl (9 g/L of water). Seven days later, the rats were randomly allocated to seven groups. Groups of twelve rats were given meats following the same 3×2 factorial protocol as for study 2: meats were raw, rare, or well-done and marinated or not-marinated for each level of cooking. As in the short-term experiment, pieces of meat were completed with low calcium modified AIN76. After evaluating the consumption of the pieces of meat and powder, we calculated that diets contained 47.3 g of meat (dry weight, per 100g total diet) and 52.7 g of low calcium modified AIN76 (See Supplementary Materials for details). One group ($N = 20$) received control diet without meat, composed only of low calcium–modified AIN76. Each diet was stored at -20°C under vacuum and dispensed daily at 5:00 p.m. Rats were fed the experimental diets daily for 98 days before CO_2 euthanasia. Body weight was monitored every week during the first 4 weeks, then every 2 weeks. Food and water intake was measured on days 25 and 60. Feces were collected on days 88 to 91 and kept at -20°C . Each rat was placed in a metabolic cage and urine collected on days 67 to 70 and kept at -20°C .

Carcinogenesis study in *Apc Min* mice

Five- to 8-week-old male and female *Apc Min* mice ($N = 97$; The Jackson Laboratory) were housed 2 to 5 mice per cage under standard laboratory conditions with free access to food and water. After 3 days of acclimatization, the mice

were randomly allocated to five experimental groups balanced for age and sex. Groups of 18 mice were given meats following a 2×2 factorial protocol: meats were raw or rare and marinated or not marinated for each level of cooking. As we could not have the same number of experimental groups as in rats, we decided to eliminate the well-done meat group because it is not as representative of human consumption (33). Unlike for rats, for the mice we had to grind and mix the meat with the modified AIN76 to ensure it was eaten entirely. The meat diets contained 60% (dry weight) meat and 40% low calcium–modified AIN76–base powder (see Supplementary Materials for more details on diets). One group ($N = 25$) received a control diet without meat, composed only of modified AIN76. Each diet was stored at -20°C under vacuum and dispensed daily at 5:00 p.m. Mice were fed the experimental diets for 45 days before euthanasia by cervical dislocation. Body weight was monitored every week during the first 3 weeks, then every 2 weeks. Feces were collected on days 35 to 39 and urine on day 39 and kept at -20°C .

The consumption of raw beef is too unusual to be representative of human consumption; therefore, for studies in humans, the meat was cooked to be consistent with the most common consumption methods (rare and well done).

Cross over study with human volunteers

The human study was performed in the Nutritional Investigation Unit of the Human Nutrition Center of Auvergne (Clermont-Ferrand, France). A single-blind randomized, controlled cross-over trial was performed in 24 human volunteers after obtaining written informed consent (inclusion criteria in Supplementary Materials). During the 1-week run-in period for adaptation, volunteers were asked to eat a diet without beef or pork and low in antioxidant products. Volunteers were then randomly submitted to three alternating 4-day intervention periods with 110 g/d of rare beef, marinated rare beef or marinated well-done beef in a random order. During the intervention periods, the volunteers must not eat meat, fish or eggs outside the provided beef. To facilitate the comparison between human and rodents studies of this project, meats for the human study were obtained from the same production batch as those given to rats and mice. Intervention periods were separated by a wash-out period of at least 3 days with the same diet as the run-in period. Urine and stool were collected during the last 3 days of each intervention period and at the end of each washout period. Each subject came to the Nutrition Investigation Unit four times in addition to the screening visit. Before the first intervention period (visit 1), volunteers were given cooking instructions. At each visit, meats were distributed for the following intervention period. During visits 2, 3, and 4, each volunteer brought back the meat packaging and frozen urine and stool samples. Compliance to diet was assessed after the collection of

feces and urine samples at the end of the intervention periods.

Organoleptic acceptability of marinated meats by a consumer panel

Two consumer studies were organized in INRA-Dijon, France (see Supplementary Materials for details). The first study aimed at assessing a reference beef meat (beef) and the two marinated products (one marinade without grape–olive extracts and one marinade with these extracts) for overall liking by consumers during blind tasting. The second study aimed to replicate the first study (reference vs. grape–olive marinade) but also aimed to assess whether providing the consumers with information about the protective effect of the marinated product could improve their overall liking of the product. Consumers from both groups were regular eaters of beef. The consumers from the treatment group were first (at recruitment and at the beginning of the session) instructed about the risk of colon cancer promotion by excessive red meat consumption. Both consumer studies were designed according to the same rules. Data were collected by the Fizz system in a sensory room equipped with 16 booths and maintained according to ISO Standard (NF EN ISO 8589). Sessions occurred at lunch time. The order of serving the samples was balanced using Williams Latin squares, ensuring that each product was evaluated equally often in the different position orders and preceded each other product equally often. Each consumer had the products served either rare or well done according to his/her usual way of consuming beef. The liking scale was the 9-point hedonic scale with the first tick-box on the left labeled "I don't like it at all" and the ninth on the right labeled "I like it very much."

In the second study, the information about colon cancer risk and the protective effect of new products was provided to the consumer of the informed group (see Supplementary Table S9). This text was sent to them by mail before the experiment and included in a preliminary screen of the sensory session. For each product and subject, the difference of liking scores given in the informed condition (protective effect) versus no information was computed. The 3 differences of each subject were averaged providing individual scores of the effect of information on liking. A number of individual variables (demographics, usages, attitudes, . . .) were screened for differences in information scores among their levels.

Carcinogenesis endpoints

MDF scoring. At euthanasia, the rats' colons were removed, washed with cold Ringer, opened, coded, and fixed flat between two sheets of filter paper in 10% buffered formalin (Sigma). The number of MDF per colon and the number of crypts per MDF, as a measure of their size, were scored by two blinded investigators after high-iron diamine alcian blue staining according to Caderni and colleagues and Femia and colleagues (34, 35).

Tumor scoring. At euthanasia, the small intestine (from duodenum to ileum) and the colon of mice were removed. Sections of the duodenum, jejunum, ileum, and colon were opened along the longitudinal axis and washed in PBS. After fixation, the different sections were stained for 48 hours in a 300 ppm solution of methylene blue in formalin. Two blinded investigators scored tumors and determined their diameters using a binocular microscope at $\times 25$ magnification. All tumors in each section of the intestines were counted; the smallest tumors that could be counted were approximately 0.5 mm in diameter.

Fecal biomarkers

Fecal water preparation. To prepare fecal water of induced rats and mice, 1 mL of distilled water and 50 μ L of butylated hydroxytoluene (BHT, Sigma) 0.45 mol/L were added to 0.4 g of fresh feces. The feces were ground using Fast-Prep (MP Biomedicals) for 30 seconds three times and then centrifuged at 5,500 g for 20 minutes. The fecal water was collected and kept at -20°C until use. For humans, 1 mL of sterilized water and 25 μ L of BHT (0.90 mol/L) were added to 0.25 g of fresh feces and the same protocol followed.

Heme and TBARS assays. Heme was measured by fluorescence in fecal water according to Sesink and colleagues (36) as described previously (20). TBARS were measured in fecal water according to Ohkawa and colleagues as previously described (37).

HNE assay. Samples were prepared for free HNE determination as described previously by Lesgards and colleagues (38) and adapted for the analysis of rat fecal water samples (see Supplementary Materials for details).

Urinary biomarker

Urine preparation. Urine samples were diluted at 1:200 for animal studies and at 1:20 for human study in an EIA/BSA solution containing phosphate buffer (0.1 mol/L at pH 7.4) with NaCl (0.15 mol/L), 0.1% BSA and 0.01% sodium azide.

DHN-MA assay. DHN-MA assay was performed by competitive enzyme immunoassay as described previously using DHN-MA-linked acetylcholinesterase enzyme as tracer (24). Urine samples from non-induced rats study were pooled and analyzed in duplicate, whereas other urine samples were assayed individually.

Statistical analysis

Data were analyzed using Systat 12 software for Windows and reported as mean \pm SEM or mean \pm SD as noted in legends.

In study with induced rats, the effect of meat intake was analyzed by one-way ANOVA comparing the control animals to the meat-fed animals. Next, the control animals

were removed and the effect of marinade and cooking analyzed in two-way ANOVA. The interaction between the two factors (marinade and cooking) was always checked. In study with *Min* mice, the effect of meat intake was analyzed by two-way ANOVA, including the factor sex. Next, the effect of the marinade, cooking, and gender was analyzed in three-way ANOVA. The interaction between the two or three factors was always checked. MDF and tumor scoring was performed in duplicate by two independent readers; therefore, these variables were tested first by 2-factor ANOVA (groups and readers). The group \times reader interaction was never significant, and when total ANOVA was significant ($P < 0.05$), pairwise differences between groups were analyzed.

Human volunteer data were analyzed using the Wilcoxon's signed rank test with each volunteer being his own control. No Bonferroni correction was made for the multiple comparison analysis, as only three pairwise comparisons were made that had been decided beforehand (i.e., rare-cooked meat *vs.* no-meat control period, rare-cooked meat *vs.* rare-cooked meat marinated, and rare-cooked meat marinated *vs.* well-done meat marinated).

Liking data from the first panel and control group of the second panel were analyzed separately using two-way ANOVA model, including the product and the consumer effects. The information (classical/protective) and its interaction with the product effect were added to the model for the informed group of the second panel. Finally, the difference between liking scores with and without the protective information averaged over the two products was used to screen the items on the consumer questionnaires by one-way ANOVAs for the effect of this information.

Results

Associated to fecal and urinary lipoperoxidation biomarkers, consumption of red meat increases the size of colonic preneoplastic lesions in azoxymethane-initiated rats, but marinating meat protects against these modulations

Rats fed meats had significantly larger MDF ($P = 0.028$) and more fecal heme ($P < 0.0001$), TBARS ($P < 0.0001$), free and bound 4-hydroxynonenal (HNE; $P = 0.002$ and $P < 0.0001$, respectively), and urinary DHN-MA ($P < 0.0001$) than rats given the control diet with no meat (Table 1). Proteomic analysis showed that raw no-marinated meat fed rats had significantly more ferritin and annexin in the colon mucosa than control no-meat diet fed rats (Supplementary Table S6). All proteomics results as well as the metabolic pathway affected are presented in Supplementary Data (Supplementary Table S6 and Supplementary Fig. S3). Rats given marinated meats had significantly fewer MDF ($P = 0.018$) and less fecal heme ($P = 0.013$), TBARS ($P = 0.015$), free and bound HNE ($P = 0.003$ and $P = 0.031$, respectively), and urinary DHN-MA ($P = 0.001$)

Table 1. Effect of experimental meats on size and number of preneoplastic Mucin depleted foci (MDF) lesions and on fecal and urinary biomarkers associated with meat-induced promotion in rats previously injected with azoxymethane and fed for 98 days

	Control diet (no meat)	Experimental meats						Statistics ^a		
		No marinade			Marinade			Meat	Marinade	Cooking
		Raw	Rare	WD	Raw	Rare	WD			
MDF size	2.3 ± 0.3	2.5 ± 0.6	2.7 ± 0.9	2.4 ± 0.6	2.6 ± 0.5	2.5 ± 0.5	2.7 ± 0.7	<i>P</i> = 0.028	<i>P</i> = 0.855	<i>P</i> = 0.944
MDF number	19 ± 9	21 ± 11	17 ± 10	15 ± 6	16 ± 9	14 ± 7	13 ± 7	<i>P</i> = 0.067	<i>P</i> = 0.018	<i>P</i> = 0.042
Heme (μmol/L in FW)	2 ± 2	193 ± 72	122 ± 43	90 ± 30	120 ± 58	101 ± 44	99 ± 26	<i>P</i> < 0.0001	<i>P</i> = 0.013	<i>P</i> < 0.0001
TBARS (μmol/L MDA eq. in FW)	13 ± 12	84 ± 21	75 ± 15	91 ± 17	67 ± 19	68 ± 14	81 ± 27	<i>P</i> < 0.0001	<i>P</i> = 0.015	<i>P</i> = 0.034
HNE free (nmol/g FW)	1.1 ± 0.4	1.7 ± 0.4	1.8 ± 0.5	1.7 ± 0.6	1.5 ± 0.3	1.2 ± 0.4	1.3 ± 0.6	<i>P</i> = 0.002	<i>P</i> = 0.003	<i>P</i> = 0.824
HNE bound (nmol/g proteins)	44 ± 17	72 ± 23	97 ± 28	66 ± 32	63 ± 21	62 ± 22	69 ± 28	<i>P</i> < 0.0001	<i>P</i> = 0.031	<i>P</i> = 0.202
DHN-MA (μg/24 h in urine)	178 ± 62	406 ± 173	564 ± 237	485 ± 229	335 ± 86	372 ± 170	310 ± 149	<i>P</i> < 0.0001	<i>P</i> = 0.001	<i>P</i> = 0.172

NOTE: Data are presented as mean ± SD.

Abbreviations: FW, fecal water; WD, well-done.

^aThe effect of meat intake was evaluated by a one-way ANOVA and the effect of marinated and cooked meats intake by a two-way ANOVA.

than rats given non-marinated meats (Table 1). Finally, rats given cooked meats had significantly fewer MDF (*P* = 0.042) and less fecal heme (*P* < 0.0001) and TBARS (*P* = 0.034) than rats given uncooked meats, without a significant difference in free and bound HNE (*P* = 0.8 and *P* = 0.2, respectively) and urinary DHN-MA (*P* = 0.2; Table 1). We identified significant interactions between marinade and cooking factors for fecal heme (*P* = 0.014) and fecal bound HNE (*P* = 0.042).

Red meat intake induces a simultaneous increase in the number of intestinal tumors, the tumor load and the fecal biomarker of peroxidation in *Apc Min* mice. Marinating red meat protects against these increases in female mice

The tumoral character of the enumerated lesions in the intestine after methylene blue staining (Fig. 1A) was evaluated in histological sections (Fig. 1B). Mice given meats had significantly more intestinal tumors and higher tumor load (*P* = 0.048 and *P* = 0.001, respectively; Fig. 1C and D) and more fecal heme (*P* = 0.02, Fig. 2A) and TBARS (*P* < 0.0001, Fig. 2B) than mice given the control diet with no meat. Taking into account the size of the tumors, the significant effect of meat in increasing the number of tumors was confirmed for medium and large tumors (Supplementary Table S7). Furthermore, female mice given meats had significantly more and larger intestinal tumors (*P* = 0.047, Fig. 1C and *P* = 0.009, respectively) than male mice given meats.

We found a significant interaction between marinade and sex factors for tumor number (*P* = 0.015). Female mice given marinated meats had significantly fewer intestinal tumors than female mice given non-marinated meats (*P* = 0.019, Fig. 1E), but no protecting effect of marinade was observed for male mice (*P* = 0.47, Fig. 1F).

Mice given marinated meats had significantly less fecal TBARS (*P* = 0.043) than rats given non-marinated meats without a significant effect on fecal heme (*P* = 0.4), and urinary DHN-MA (*P* = 0.2, Fig. 2). Proteomic analysis showed that rare marinated meat fed mice had significantly

less vimentin in the colon mucosa than control no-meat diet fed mice (Supplementary Table S8). All proteomics results as well as the metabolic pathway affected are presented in Supplementary Data (Supplementary Table S8 and Supplementary Fig. S5). Mice given cooked meats had significantly more intestinal tumors and higher tumor load (*P* = 0.001 and *P* = 0.001, respectively; Fig. 1C and D) than mice given uncooked meats, without a significant difference for fecal heme (*P* = 0.2), TBARS (*P* = 0.1), and urinary DHN-MA (*P* = 0.1, Fig. 2). We found no significant interaction between marinade and cooking factors.

Rare cooked meat consumption increases luminal peroxidation in healthy volunteers and a grape-olive marinade reduces this effect

The nutritional intervention was scheduled with 24 healthy volunteers (see Supplementary Materials for details). Biomarker measurements revealed a significant increase in the heme content of fecal water for the two rare meat periods (Fig. 3A) and a tendency to increase with the well-done meat. We also observed a significant increase in the concentrations of TBARS in fecal water from human volunteers fed 110 g of non-marinated meat for 4 days compared with control periods (*P* = 0.038, Fig. 3B). Marinade significantly decreased fecal TBARS for rare cooking (*P* = 0.046) but not well-done cooking (*P* = 1). We found no significant difference between volunteers in regards to HNE content in fecal water, urinary DHN-MA, or fecal ATNC, and fecal and genotoxicity (Supplementary Fig. S6).

The olive-grape marinade does not alter the organoleptic acceptability of red meat but conversely increases the preference of the meat

To quantify the contribution of antioxidants in the effect of grape-olive marinade on organoleptic acceptability, an antioxidant-free marinade was compared with a marinade with antioxidants. A two-way (product and consumer) ANOVA model of liking scores from the first study demonstrated that the marinated meats were both significantly




Figure 1. Effect of experimental meats on small intestine tumors in *Apc Min* mice fed for 45 days. **A**, Adenoma after methylene blue staining. **B**, Histological section of adenoma (H&E staining). **C**, Number of tumors per mouse. **D**, Tumor load per mouse. **E**, Number of tumors per female mouse; scale bars, 500 μ m. Data are presented in the graphs as mean \pm SEM and the circle are the individuals. The effect of meat intake was evaluated by two-way ANOVA and the effect of marinated and cooked meat intake by three-way ANOVA. #, $P < 0.05$; ##, $P < 0.01$ versus the no-meat-fed mice; *, $P < 0.05$ versus the raw condition with the same marinade status; °, $P < 0.01$ versus the no-marinated condition.

preferred to the reference meat ($P = 0.004$, Fig. 4A). A very slight similar trend was observed in the second study (Fig. 4B), although it did not reach statistical significance ($P = 0.283$). The marinated meat with antioxidants was slightly less appreciated than the marinated meat without antioxidants, though this difference was not significant (Fig. 4A).

The effect of information about colon cancer risk and the protective effect of new products was not significant and had almost ($P = 0.182$) no interaction with the product effect (Fig. 4C). However, looking precisely at the mean liking scores of the four products tested in the test group, a potential positive effect of the information on the liking of the marinated meat was observed ($P = 0.086$).

None of the variables on the usage, attitude, and knowledge questionnaire influenced the effect of the information

on product-liking. However, the level of education and age of the consumers seemed to have impacted that effect. Indeed, the difference between liking scores with and without information is significantly ($P = 0.022$) higher in primary and secondary versus higher school levels (Fig. 4D). Also, Fig. 4E shows that the effect of information would be negative in consumers ages less than 35, about null in consumers between 36 et 55 and positive in consumer older than 56; these differences among the three age categories were almost significant ($P = 0.053$).

Discussion

The present study is the first to show that consumption of fresh red meat for few days increased fecal lipid oxidation




Figure 2. Effect of experimental meats on fecal and urinary biomarkers associated with meat-induced promotion in *Apc Min* mice fed for 45 days. **A**, Heme in fecal water. **B**, TBARS in fecal water. **C**, DHN-MA in urine. Data are presented as mean ± SEM. The effect of meat intake was evaluated by two-way ANOVA and the effect of marinated and cooked meat intake evaluated by three-way ANOVA. *, $P < 0.05$; **, $P < 0.0001$ versus no meat-fed mice, °, $P < 0.05$ versus no-marinated condition.

biomarkers associated with heme-induced promotion of colon carcinogenesis in human volunteers and in two complementary animal models.

These findings agree with the epidemiological data and the conclusions of the WCRF and WHO (3, 4). This study also shows that this increase and the promotion of carcinogenesis in rats and female mice were decreased when meat was treated with an antioxidant marinade.

As people of a lower socioeconomic status are not receptive to nutritional messages and less likely to change risky behaviors than affluent people, we wanted to propose an alternative to the current recommendation of limiting consumption by working directly on the commercial prod-

uct. To achieve this objective, we designed a project (Flow chart in Supplementary Fig. S1) allowing to highlight a beef antioxidant marinade limiting the heme-iron induced lipoperoxidation, a key step in the promotion of colon cancer associated with red meat consumption (11).

For this project, we associated two animal models of carcinogenesis: The AOM-induced rat and the *Apc Min* mice. Indeed, for 30 years, investigators have searched for dietary agents that could impact colon carcinogenesis tumors. For that, the azoxymethane (AOM) model was widely used and more recently associated to *Apc Mice* models. If carcinogenesis in the rat model results of a large bolus of AOM, it has many morphological as well as





Figure 3. Effect of experimental meats on fecal biomarkers associated with meat-induced promotion in human volunteers after 4 days of 110 g/d of meat. **A**, Heme. **B**, TBARS in fecal water. Data are presented as mean ± Min/Max, $N = 21$. *, $P < 0.05$ versus the no-meat control period (Wilcoxon's test). §, $P < 0.05$, versus the no-marinated meat period (Wilcoxon's test).

**Figure 4.**

Product liking means in the first sensory study (A), second sensory study (B), test groups (C). Effect of the information according to education level (D) and age (E). Data are presented as mean \pm SEM. Statistical analysis was three-way ANOVA. *, $P < 0.05$, vs. the reference. °, $P < 0.05$, versus the reference Higher school level; ax.: antioxidants; inform.: after information.

molecular similarities to human sporadic colorectal cancer (39). Similarly, if the main location of tumors in the *Apc Min* mice is the small intestine, this mice is considered to be a model for human familial adenomatous polyposis with applications to better understand the molecular mechanisms of multistage carcinogenesis in the large bowel of humans (40). Furthermore, review of experimental data between diet and colon cancer demonstrated that there is a close agreement between many results obtained in the colons of AOM-initiated rats and in the small intestine of *Min* mice (41) and that there is a reasonable agreement between the results of these animal studies and the more limited clinical studies (42).

First, we defined an antioxidant marinade able to decrease lipoperoxidation in meats and rat fecal water after meat consumption. Several studies have shown that natural antioxidants can be used to reduce lipoper-

oxidation in meat, and that the antioxidant activity of a marinade can be determined by a DPPH assay (43–46). Twelve marinades were prepared to test the antioxidant activity of eight molecules, alone or in mixture, and grape–olive marinade had the highest (Supplementary Table S2). The addition of this marinade to beef sirloin led to a significant decrease in the concentrations of TBARS in meat (study 1, Supplementary Table S3). Although the grape–olive mixture has never been tested, these results are consistent with the literature regarding the properties of grapes and olives (47–49). In a short-term study in rats, we confirmed that the marinade can also decrease meat-induced colonic luminal lipoperoxidation (study 2, Supplementary Table S4), which is consistent with our previous studies (19, 20).

Thus, we wanted to check the effect of marinated and non-marinated meat on two stages of colorectal

carcinogenesis using two complementary animal models: AOM-initiated rats for the preneoplastic stage and *Apc Min* mice for the tumoral stage. Finally, we explored the impact of these meats on fecal and urinary biomarkers in human volunteers given 110 g of meat per day for 4 days.

Our results showed that fresh red meat intake significantly increased the size of pre-cancerous lesion MDF in initiated rats (Table 1) and the number and size of intestinal tumors and tumor load in *Apc Min* mice (Fig. 1). More precisely, fresh red meat intake significantly increased the number of medium and large tumors (Supplementary Table S7). In the two carcinogenesis animal models, meat-induced promotion was associated with a significant increase in fecal heme and lipoperoxidation biomarkers (TBARS, HNE in fecal water, and urinary DHN-MA in rats [Table 1]; TBARS in *Apc Min* mice [Fig. 2]). In mucosa, promotion in initiated rats was positively associated with the level of ferritin and annexin (Supplementary Table S6), two proteins for which expression was positively associated with the degree of dysplasia (50, 51). In our study, the level of ferritin protein in the colon mucosa correlated with the size of MDF ($r = 0.497$, $P = 0.002$), with the TBARS level in fecal water ($r = 0.511$, $P = 0.001$) and with the DHN-NA content in urine ($r = 0.409$, $P = 0.013$). The level of annexin protein in mucosa positively correlated with the size of MDF ($r = 0.452$, $P = 0.006$), with the heme content in fecal water ($r = 0.394$, $P = 0.017$) and with the TBARS level in fecal water ($r = 0.402$, $P = 0.015$). These results on fecal and urinary biomarker modulations are consistent with previous studies showing a promoting effect of hemoglobin or lyophilized red meat on pre-cancerous lesions or on tumors in association with increased fecal lipoperoxidation (16, 18–20). In human volunteers, the present crossover study showed that eating fresh red meat for 4 days was sufficient to increase lipoperoxidation biomarker concentrations in fecal water, compared with the control period without meat (Fig. 4). We did not observe a significant increase in urinary DHN-MA, which is consistent with our previous results with red meats (22). In humans, unlike in rodents, meat intake correlates with endogenous formation of fecal ATNC (52, 53). In this study, we did not observe a significant increase of fecal ATNC (Supplementary Fig. S6) and the level of ATNC was low overall, but this result is consistent with other human studies in which an increase in fecal ATNC was found only in subjects consuming at least 240 g of red meat per day and for a longer consumption period (52, 54, 55). Thus, the intake of fresh red meat can modulate lipoperoxidation biomarkers associated with the promotion of colon carcinogenesis in initiated rats and *Apc Min* mice, which gives experimental support to the epidemiology-based conclusion that red meat may be a cause of colorectal cancer (1, 2, 4).

Grape–olive marinade was efficient in initiated rats to decrease the number of pre-cancerous lesion MDF and was associated with a decrease in heme and lipoperoxidation biomarkers (Table 1). In *Apc Min* mice study, we observed a significant protective effect in female mice (Fig. 1E). To the best of our knowledge, this study is the first in which an antioxidant marinade was used to reduce the promotion of colorectal cancer by fresh red meat. Our results support epidemiological studies showing that adherence to a Mediterranean diet, rich in olive oil and antioxidants, may reduce colorectal cancer risk, mainly in women (56, 57). The fact that we only observed a protective effect of the marinade in female may be consistent with epidemiologic evidences showing a gender-specific associations between antioxidants and rectal cancer risk, pointing a protective effect in women while no effect in men, probably due to the estrogen status (58). In our study, the protection due to the marinade in female *Apc Min* mice was associated with decreased expression of vimentin in the colon mucosa (Supplementary Table S8). Vimentin activates the Wnt signaling pathway, which is detectable as increased β -catenin accumulation in the nucleus with concomitant activation of β -catenin–dependent transcription of Wnt signaling downstream targets (59); thus, a decrease in the activation of this pathway largely involved in the colon carcinogenesis could participate in the observed protection. In human volunteers, as in the two carcinogenesis animal models, the marinade reduced significantly the increase of fecal TBARS when meat was cooked rare (Fig. 3). However, the protective effect was lost when the meat was well-done, likely reflecting an alteration in the antioxidant capacity during cooking at high temperature. In accordance with our initial aim to limit the risk in big meat eaters, in whom the risk of colon cancer increases the most, this absence of effect in well-done meat will not greatly limit the efficiency of the marinade because a majority of big eaters in France (60), in Europe (33), and in the United States (61) consume rare or medium red meat. Moreover, in parallel to the addition of antioxidant *via* a marinade, it may be interesting to test other solutions to provide the antioxidants with the meat: sell antioxidant extracts with the meat for example. These extracts could be used to dress the meat after cooking. However, if it is not certain that a sufficient amount of antioxidant is attained, this solution could be useful to limit the loss of antioxidant activity during "well-done" cooking.

The biological effectiveness of modifying a product is not useful if consumers do not like the product. For the marinated meat, it is important to note that the product had higher liking scores than the unmodified product (Fig. 4). Thus, modifying the product to limit the risk of cancer does not affect the acceptability of the product. Furthermore, we observed a potential positive effect of the information about colon cancer risk and the protective effect of new products on the liking score for the marinated

meat. This suggests that, if the consequences of the modification of the meat product are appreciated, then the consumers can positively value the information. Though the current communication of only limiting consumption does not seem to be well accepted (12), the strategy of associating the modification of the product with the communication on the potential benefit could be very effective in limiting the risk.

We have also evaluated the effect of cooking on meat-induced colorectal cancer. We studied rare and well-done meat in initiated rats and meat cooked rare in *Apc Min* mice because it is the cooking method most used by consumers (62). In a previous study, in which we demonstrated the promotive effect of heme iron (16), we did not observe a positive association between the number of MDF and PhiP and Me-IQx in an AIN-76-based diet. In the present study, cooking had a protective effect on the number of pre-cancerous lesions in initiated-rats. In contrast, in *Apc Min* mice, cooking had a promoting effect on the number and size of intestinal tumors. In our study, this promoting effect did not come from potentially carcinogenic heterocyclic amine formation because concentrations found in meats were very low. Among the 13 tested, 8 heterocyclic amines were not detected or under the quantification limit, only PhiP, IQx, MeIQx, DiMeIQx and DMIP were quantifiable in rare and well-cooked meats, but at low doses and well below the concentrations usually used to promote colorectal carcinogenesis in *Apc Min* mice (Supplementary Table S5; refs. 63, 64). In initiated rats, the protective effect of cooking was associated with a significant decrease in fecal heme, and in *Apc Min* mice the promoting effect of cooking was associated with a tendency of increase in fecal heme. Thus, as observed in our previous studies (16, 19), the promotion of carcinogenesis appears to be associated with the luminal heme content. Differences observed between rats and *Apc Min* mice suggest a difference in heme bioavailability in these two different rodent models. These opposing results make it difficult to come to a conclusion on the effect of cooking on heme bioavailability and the link with lipoperoxidation and the promotion of carcinogenesis. However, in human volunteers, if meat consumption increased the level of heme in fecal water, the cooking method did not significantly affect this level. Thus, the impact of marinated meat on lipoperoxidation in human volunteers could be assigned to the antioxidant capacity of the marinade and not the decrease of heme bioavailability, the catalyzer of lipid peroxidation.

In conclusion, consumption of fresh red meat for few days increased biomarkers associated with heme-induced promotion of colon carcinogenesis. Grape-olive marinade counteracted this promoting effect in carcinogen-injected rats and female *Apc Min* mice. This protection was associated with normalization of fecal biomarkers in the two

animal models and a significant limitation in human volunteers. As in one of our previous studies with processed meat (32), these results suggest that it is possible to modify fresh red meat products by adding antioxidants to reduce the cancer-promoting properties of red meat, which could also improve, or at least not deteriorate, its sensory acceptability. The protective effect of meat antioxidant marinades need now to be validated by epidemiological studies and mechanisms need to be explored more precisely with for example the monitoring of the levels of DNA damage and DNA adducts to lipoperoxidation endproducts. Their effectiveness could lead to protective strategies to decrease the colorectal cancer burden in all populations, especially those who need it most.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: O. Martin, D. Durand, P. Schlich, V. Scislowski, D.E. Corpet, F.H.F. Pierre

Development of methodology: J. Dupuy, C. Chantelauze, C. Urbano, T. Sayd

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): O. Martin, N. Naud, S. Taché, L. Debrauwer, S. Chevolleau, J. Dupuy, C. Chantelauze, E. Pujos-Guillot, F. Blas-Y-Estrada, C. Urbano, V. Santé-Lhoutellier, D. Viala, A. Blot, N. Meunier, D. Attaix

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): O. Martin, S. Chevolleau, J. Dupuy, G.G.C. Kuhnle, V. Santé-Lhoutellier, T. Sayd, D. Viala, P. Schlich, F. Guéraud, D.E. Corpet, F.H.F. Pierre

Writing, review, and/or revision of the manuscript: O. Martin, L. Debrauwer, S. Chevolleau, C. Urbano, V. Santé-Lhoutellier, T. Sayd, D. Viala, A. Blot, N. Meunier, D. Attaix, F. Guéraud, D.E. Corpet, F.H.F. Pierre

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): C. Urbano, T. Sayd, N. Meunier

Study supervision: N. Meunier, D.E. Corpet, F.H.F. Pierre

Other (written the chapter: "Protein identification by mass spectrometry"): D. Viala

Acknowledgments

We thank all of the volunteers who generously gave their time for this research. We thank the ADIV workshop staff for the meat preparation. This work was conducted in the framework of the French network for Nutrition And Cancer Research (NACRe network), www.inra.fr/nacre. This work was supported by the French National Agency of Research (ANR-10-ALIA-14, SecuriViande project), the French National Institute for Agricultural Research (INRA), and by the French Technical Center of Meat (ADIV).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received November 9, 2017; revised March 12, 2018; accepted June 15, 2018; published first June 28, 2018.

References

1. WCRF/AICR. World Cancer Research Fund/American Institute for Cancer Research. Continuous Update Project Report. Food, Nutrition, Physical Activity, and the Prevention of Colorectal Cancer. 2011.
2. WCRF/AICR. Food, nutrition, physical activity, and the prevention of cancer: a global perspective. 2007.
3. WCRF/AICR. Continuous Update Project Report. Diet, Nutrition, Physical Activity, and Colorectal Cancer. 2017.
4. IARC. Carcinogenicity of consumption of red and processed meat. *Lancet Oncol* 2015.
5. Alkerwi A, Sauvageot N, Nau A, Lair ML, Donneau AF, Albert A, et al. Population compliance with national dietary recommendations and its determinants: findings from the ORISCAV-LUX study. *Br J Nutr* 2012;108:2083–92.
6. Estaquio C, Kesse-Guyot E, Deschamps V, Bertrais S, Dauchet L, Galan P, et al. Adherence to the French Programme National Nutrition Sante Guideline Score is associated with better nutrient intake and nutritional status. *J Am Diet Assoc* 2009;109:1031–41.
7. Darmon N, Drewnowski A. Does social class predict diet quality? *Am J Clin Nutr* 2008;87:1107–17.
8. Aston LM, Smith JN, Powles JW. Meat intake in Britain in relation to other dietary components and to demographic and risk factor variables: analyses based on the National Diet and Nutrition Survey of 2000/2001. *J Hum Nutr Diet* 2013;26:96–106.
9. James WP, Nelson M, Ralph A, Leather S. Socioeconomic determinants of health. The contribution of nutrition to inequalities in health. *BMJ* 1997;314:1545–9.
10. Smith MP, Olatunde O, White C. Inequalities in disability-free life expectancy by area deprivation: England, 2001–04 and 2005–08. *Health Stat Q* 2010;36–57.
11. Weiderpass E, Pukkala E. Time trends in socioeconomic differences in incidence rates of cancers of gastro-intestinal tract in Finland. *BMC Gastroenterol* 2006;6:41.
12. Winkels RM, van Lee L, Beijer S, Bours MJ, van Duijnhoven FJ, Geelen A, et al. Adherence to the World Cancer Research Fund/American Institute for Cancer Research lifestyle recommendations in colorectal cancer survivors: results of the PROFILES registry. *Cancer Med* 2016;5:2587–95.
13. Le NT, Michels FA, Song M, Zhang X, Bernstein AM, Giovannucci EL, et al. A prospective analysis of meat mutagens and colorectal cancer in the nurses' health study and health professionals follow-up study. *Environ Health Perspect* 2016;124:1529–36.
14. Martin OC, Lin C, Naud N, Tache S, Raymond-Letron I, Corpet DE, et al. Antibiotic suppression of intestinal microbiota reduces heme-induced lipoperoxidation associated with colon carcinogenesis in rats. *Nutr Cancer* 2015;67:119–25.
15. Ijssennagger N, Belzer C, Hooiveld GJ, Dekker J, van Mil SW, Muller M, et al. Gut microbiota facilitates dietary heme-induced epithelial hyperproliferation by opening the mucus barrier in colon. *Proc Natl Acad Sci U S A* 2015;112:10038–43.
16. Bastide N, Chenni F, Audebert M, Santarelli R, Tache S, Naud N, et al. A central role for heme iron in colon carcinogenesis associated with red meat intake. *Cancer Res* 2015;75:870–9.
17. Sawa T, Akaike T, Kida K, Fukushima Y, Takagi K, Maeda H. Lipid peroxyl radicals from oxidized oils and heme-iron: implication of a high-fat diet in colon carcinogenesis. *Cancer Epidemiol Biomarkers Prev* 1998;7:1007–12.
18. Pierre F, Freeman A, Tache S, Van der Meer R, Corpet DE. Beef meat and blood sausage promote the formation of azoxymethane-induced mucin-depleted foci and aberrant crypt foci in rat colons. *J Nutr* 2004;134:2711–6.
19. Pierre F, Santarelli R, Tache S, Gueraud F, Corpet DE. Beef meat promotion of dimethylhydrazine-induced colorectal carcinogenesis biomarkers is suppressed by dietary calcium. *Br J Nutr* 2008;99:1000–6.
20. Pierre F, Tache S, Petit CR, Van der Meer R, Corpet DE. Meat and cancer: haemoglobin and haemin in a low-calcium diet promote colorectal carcinogenesis at the aberrant crypt stage in rats. *Carcinogenesis* 2003;24:1683–90.
21. Bastide N, Morois S, Cadeau C, Kangas S, Serafini M, Gusto G, et al. Heme iron intake, dietary antioxidant capacity, and risk of colorectal adenomas in a large cohort study of french women. *Cancer Epidemiol Biomarkers Prev* 2016;25:640–7.
22. Pierre F, Peiro G, Tache S, Cross AJ, Bingham SA, Gasc N, et al. New marker of colon cancer risk associated with heme intake: 1,4-dihydroxynonane mercapturic acid. *Cancer Epidemiol Biomarkers Prev* 2006;15:2274–9.
23. Romeu M, Aranda N, Giral M, Ribot B, Nogues MR, Arija V. Diet, iron biomarkers and oxidative stress in a representative sample of Mediterranean population. *Nutr J* 2013;12:102.
24. Gueraud F, Peiro G, Bernard H, Alary J, Creminon C, Debrauwer L, et al. Enzyme immunoassay for a urinary metabolite of 4-hydroxynonanal as a marker of lipid peroxidation. *Free Radic Biol Med* 2006;40:54–62.
25. Gaschler MM, Stockwell BR. Lipid peroxidation in cell death. *Biochem Biophys Res Commun* 2017;482:419–25.
26. Lu B, Chen XB, Ying MD, He QJ, Cao J, Yang B. The role of ferroptosis in cancer development and treatment response. *Front Pharmacol* 2017;8:992.
27. Abete I, Romaguera D, Vieira AR, Lopez de Munain A, Norat T. Association between total, processed, red and white meat consumption and all-cause, CVD and IHD mortality: a meta-analysis of cohort studies. *Br J Nutr* 2014;112:762–75.
28. Tappel A. Heme of consumed red meat can act as a catalyst of oxidative damage and could initiate colon, breast and prostate cancers, heart disease and other diseases. *Med Hypotheses* 2007;68:562–4.
29. Niedernhofer LJ, Daniels JS, Rouzer CA, Greene RE, Marnett LJ. Malondialdehyde, a product of lipid peroxidation, is mutagenic in human cells. *J Biol Chem* 2003;278:31426–33.
30. Awasthi YC, Sharma R, Cheng JZ, Yang Y, Sharma A, Singhal SS, et al. Role of 4-hydroxynonanal in stress-mediated apoptosis signaling. *Mol Aspects Med* 2003;24:219–30.
31. Pierre F, Tache S, Gueraud F, Rerole AL, Jourdan ML, Petit C. Apc mutation induces resistance of colonic cells to lipoperoxide-triggered apoptosis induced by faecal water from haem-fed rats. *Carcinogenesis* 2007;28:321–7.
32. Pierre FH, Martin OC, Santarelli RL, Tache S, Naud N, Gueraud F, et al. Calcium and alpha-tocopherol suppress cured-meat promotion of chemically induced colon carcinogenesis in rats and reduce associated biomarkers in human volunteers. *Am J Clin Nutr* 2013;98:1255–62.
33. de Batlle J, Gracia-Lavedan E, Romaguera D, Mendez M, Castano-Vinyals G, Martin V, et al. Meat intake, cooking methods and doneness and risk of colorectal tumours in the Spanish multicase-control study (MCC-Spain). *Eur J Nutr* 2016.
34. Caderni G, Femia AP, Giannini A, Favuzza A, Luceri C, Salvadori M, et al. Identification of mucin-depleted foci in the unsectioned colon of azoxymethane-treated rats: correlation with carcinogenesis. *Cancer Res* 2003;63:2388–92.
35. Femia AP, Dolara P, Caderni G. Mucin-depleted foci (MDF) in the colon of rats treated with azoxymethane (AOM) are useful

- biomarkers for colon carcinogenesis. *Carcinogenesis* 2004;25:277–81.
36. Sesink AL, Termont DS, Kleibeuker JH, Van der Meer R. Red meat and colon cancer: the cytotoxic and hyperproliferative effects of dietary heme. *Cancer Res* 1999;59:5704–9.
 37. Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal Biochem* 1979;95:351–8.
 38. Lesgards JF, Frayne IR, Comte B, Busseuil D, Rheume E, Tardif JC, et al. Differential distribution of 4-hydroxynonenal adducts to sulfur and nitrogen residues in blood proteins as revealed using Raney nickel and gas chromatography-mass spectrometry. *Free Radic Biol Med* 2009;47:1375–85.
 39. Perse M, Cerar A. Morphological and molecular alterations in 1,2-dimethylhydrazine and azoxymethane induced colon carcinogenesis in rats. *J Biomed Biotechnol* 2011;2011:473964.
 40. Yamada Y, Mori H. Multistep carcinogenesis of the colon in Apc (Min/+) mouse. *Cancer Sci* 2007;98:6–10.
 41. Corpet DE, Pierre F. Point: From animal models to prevention of colon cancer. Systematic review of chemoprevention in min mice and choice of the model system. *Cancer Epidemiol Biomarkers Prev* 2003;12:391–400.
 42. Corpet DE, Pierre F. How good are rodent models of carcinogenesis in predicting efficacy in humans? A systematic review and meta-analysis of colon chemoprevention in rats, mice and men. *Eur J Cancer* 2005;41:1911–22.
 43. Schevey CT, Toshkov S, Brewer MS. Effect of natural antioxidants, irradiation, and cooking on lipid oxidation in refrigerated, salted ground beef patties. *J Food Sci* 2013;78:S1793–9.
 44. Kim SJ, Min SC, Shin HJ, Lee YJ, Cho AR, Kim SY, et al. Evaluation of the antioxidant activities and nutritional properties of ten edible plant extracts and their application to fresh ground beef. *Meat Sci* 2013;93:715–22.
 45. Colindres P, Brewer MS. Oxidative stability of cooked, frozen, reheated beef patties: effect of antioxidants. *J Sci Food Agric* 2011;91:963–8.
 46. Viegas O, Amaro LF, Ferreira IM, Pinho O. Inhibitory effect of antioxidant-rich marinades on the formation of heterocyclic aromatic amines in pan-fried beef. *J Agric Food Chem* 2012;60:6235–40.
 47. Gomez I, Beriain MJ, Sarries MV, Insausti K, Mendizabal JA. Low-fat beef patties with augmented omega-3 fatty acid and CLA levels and influence of grape seed extract. *J Food Sci* 2014;79:S2368–76.
 48. Jongberg S, Skov SH, Torngren MA, Skibsted LH, Lund MN. Effect of white grape extract and modified atmosphere packaging on lipid and protein oxidation in chill stored beef patties. *Food Chem* 2011;128:276–83.
 49. Hayes JE, Stepanyan V, Allen P, O'Grady MN, Kerry JP. Effect of lutein, sesamol, ellagic acid and olive leaf extract on the quality and shelf-life stability of packaged raw minced beef patties. *Meat Sci* 2010;84:613–20.
 50. Wei D, Chen T, Li J, Gao Y, Ren Y, Zhang X, et al. Association of serum gamma-glutamyl transferase and ferritin with the metabolic syndrome. *J Diabetes Res* 2015;2015:741731.
 51. Yang HB, Hsu PI, Lee JC, Chan SH, Lin XZ, Chow NH. Adenoma-carcinoma sequence: a reappraisal with immunohistochemical expression of ferritin. *J Surg Oncol* 1995;60:35–40.
 52. Bingham SA, Hughes R, Cross AJ. Effect of white versus red meat on endogenous N-nitrosation in the human colon and further evidence of a dose response. *J Nutr* 2002;132:3522S–5S.
 53. Silvester KR, Bingham SA, Pollock JR, Cummings JH, O'Neill IK. Effect of meat and resistant starch on fecal excretion of apparent N-nitroso compounds and ammonia from the human large bowel. *Nutr Cancer* 1997;29:13–23.
 54. Cross AJ, Pollock JR, Bingham SA. Haem, not protein or inorganic iron, is responsible for endogenous intestinal N-nitrosation arising from red meat. *Cancer Res* 2003;63:2358–60.
 55. Hughes R, Cross AJ, Pollock JR, Bingham S. Dose-dependent effect of dietary meat on endogenous colonic N-nitrosation. *Carcinogenesis* 2001;22:199–202.
 56. Jones P, Cade JE, Evans CEL, Hancock N, Greenwood DC. The Mediterranean diet and risk of colorectal cancer in the UK Women's Cohort Study. *Int J Epidemiol* 2017;46:1786–96.
 57. Bamia C, Lagiou P, Buckland G, Grioni S, Agnoli C, Taylor AJ, et al. Mediterranean diet and colorectal cancer risk: results from a European cohort. *Eur J Epidemiol* 2013;28:317–28.
 58. Murtaugh MA, Ma KN, Benson J, Curtin K, Caan B, Slattery ML. Antioxidants, carotenoids, and risk of rectal cancer. *Am J Epidemiol* 2004;159:32–41.
 59. Satelli A, Hu J, Xia X, Li S. Potential function of exogenous vimentin on the activation of wnt signaling pathway in cancer cells. *J Cancer* 2016;7:1824–32.
 60. AFSSA. Etude Individuelle Nationale des Consommations Alimentaires 2 (INCA 2) 2006–07. 2009. 225p.
 61. Reicks AL, Brooks JC, Garmyn AJ, Thompson LD, Lyford CL, Miller MF. Demographics and beef preferences affect consumer motivation for purchasing fresh beef steaks and roasts. *Meat Sci* 2011;87:403–11.
 62. INCA-2. National Individual Study of Dietary Consumption-2 2006–2007. 2009.
 63. Andreassen A, Vikse R, Mikalsen A, Adamovic T, Steffensen IL, Hjertholm H, et al. 2-Amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) induces genetic changes in murine intestinal tumours and cells with ApcMin mutation. *Mutat Res* 2006;604:60–70.
 64. Svendsen C, Meinel W, Glatt H, Alexander J, Knutsen HK, Hjertholm H, et al. Intestinal carcinogenesis of two food processing contaminants, 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine and 5-hydroxymethylfurfural, in transgenic FVB min mice expressing human sulfotransferases. *Mol Carcinog* 2012;51:984–92.