

# Recruitment in flatfish, with special emphasis on North Atlantic species: Progress made by the Flatfish Symposia

Henk W. van der Veer, Rüdiger Berghahn,  
John M. Miller, and Adriaan D. Rijnsdorp



Van der Veer, H. W., Berghahn, R., Miller, J. M., and Rijnsdorp, A. D. 2000. Recruitment in flatfish, with special emphasis on North Atlantic species: Progress made by the Flatfish Symposia. – ICES Journal of Marine Science, 57: 202–215.

In summarizing the main results on recruitment that emerged from the series of Flatfish Symposia, two aspects were distinguished: mean level and interannual variability. Recruitment to a stock appears to be related to the quantity of juvenile nursery habitats, suggesting that either larval supply or the carrying capacity of the nurseries is the limiting factor. However, available information on growth of 0-group flatfish suggests that the carrying capacity of nursery areas is never reached. Variability in year-class strength is generated during the pelagic egg and larval stage, probably by variations in the hydrodynamic circulation and in the mortality rates of eggs and larvae. Density-dependent processes seem to occur only during the juvenile stages, particularly in respect of growth. However, no impact on recruitment variability has been found. Density-dependent mortality during the phase shortly after settlement dampens the interannual recruitment variability. There is no evidence of density-dependent effects in the adult phase at present, but these may have been important at lower levels of exploitation. The importance of the factors determining recruitment vary not only among species, but also over the species' range. It is suggested that damping processes can only occur in the demersal stage, implying that variability in year-class strength can only decrease in fish species with a demersal stage. If true, ultimate variability in recruitment in fish species will be related to the relative duration of the pelagic and demersal stages.

© 2000 International Council for the Exploration of the Sea

Key words: control, flatfish, recruitment, regulation, year-class strength.

*H. W. van der Veer: Netherlands Institute for Sea Research, PO Box 59, 1790 AB Den Burg Texel, The Netherlands. R. Berghahn: Federal Environmental Agency, Institute for Water, Soil and Air Hygiene, Field Station Marienfelde, Schichauweg 58, 12307 Berlin, Germany. J. M. Miller: North Carolina State University, Zoology Department, Campus Box 7617, Raleigh NC 27695, USA. A. D. Rijnsdorp: Netherlands Institute for Fisheries Research (RIVO-DLO), PO Box 68, 1970 AB IJmuiden, The Netherlands. Correspondence to H. W. van der Veer: e-mail: [veer@nioz.nl](mailto:veer@nioz.nl)*

## Introduction

Flatfish are distributed over virtually all latitudes (Pauly, 1994) and therefore present an attractive group for the study of geographic variation in recruitment processes. Most species are characterized by pelagic egg and larval stages and demersal juvenile and adult stages. Juveniles are often concentrated in restricted shallow “nursery” areas, where they offer excellent opportunities to carry out quantitative studies.

Studies on the early life history of flatfish started with the work of Pearcy (1962) on winter flounder (*Pseudopleuronectes americanus* (Walbaum)) on the east coast of the United States. This pioneering work was followed up by classical studies in Great Britain, particularly on

plaice (*Pleuronectes platessa* L.). These concentrated initially on the demersal juvenile stage (e.g., Riley *et al.*, 1981; Edwards and Steele, 1968; Lockwood, 1974), but later focused also on the pelagic egg and larval phase (e.g., Talbot, 1978; Harding *et al.*, 1978). Zijlstra (1972) emphasized the importance of the Wadden Sea estuaries along the Dutch, German, and Danish coasts as nursery areas for plaice and sole (*Solea solea* (L.)). His publication formed the first in a series on these species in continental Europe (e.g., Kuipers, 1977; De Vlas, 1979; Zijlstra and Witte, 1985; Berghahn, 1987; Rijnsdorp *et al.*, 1985; Van der Veer, 1986; Pihl, 1990; Hovenkamp, 1992).

In the mid-1980s, extensive investigations indicated that year-class strength in plaice was determined during

the early pelagic stages (Zijlstra and Witte, 1985; Van der Veer, 1986). However, it remained unclear whether similar conclusions were valid for other species and areas. At the 1988 ICES Symposium on “The Early Life History of Fish” in Bergen, the idea originated to organize a flatfish symposium aimed at making an inventory of the present state of knowledge. The late Ray Beverton strongly supported the plan and he took a lead as a member of the organizing committee. The First Flatfish Symposium in 1990 with the theme “Life Cycle”, was followed up by a Second in 1993 on “Recruitment” and a Third in 1996 on “System Dynamics of Flatfish”. Simultaneously, other forums were organized, such as the International Symposium on North Pacific Flatfish in 1994 (Anon., 1995). But even after three decades of investigation, the central question in flatfish biology remains: what causes variability in year-class strength?

We review the progress made over the last decade in solving the recruitment problem, with a main focus on North Atlantic flatfish species. Firstly, we discuss the factors determining the mean level of recruitment. Secondly, the processes affecting variability in recruitment are analysed. And finally, a synthesis of the available information is presented.

## Definitions

Recruitment is defined as the number of individuals that reach a particular age to join a specific part of the population (e.g., the mature population). It is thus a number that varies from year to year. We use recruitment process to cover everything that affects survival between spawning (or even before that: maturation) and the stage of life where year-class strength is more or less fixed. We distinguish two important aspects: the actual number of recruits and the (interannual) variability therein. We use recruitment level to identify the ultimate number of a specific year class that survives to attain sexual maturity and joins the reproductive population. In addition, indices of the number of juveniles of a specific year class surviving between spawning and recruitment are called year-class strength estimates. The interannual variability in recruitment can be standardized by expressing it as coefficient of variation.

Two sets of factors may be distinguished: variability generating or controlling factors and variability damping or regulating factors. Controlling factors enhance the variability in year-class strength, and are reflected by an increase in the coefficient of variation at a particular life stage compared with a previous stage. Regulating factors reduce variability in year-class strength and ultimately in recruitment. These show up as a decrease in the coefficient of variation at a certain life stage compared with a previous stage (Fig. 1). Correlations




Figure 1. Sequence of variability generating or controlling factors (C) and variability damping or regulating factors (R) illustrated by changes in the coefficient of variation in abundance during early life history of plaice (*Pleuronectes platessa*) (data after Van der Veer, 1986).

between abundance estimates at various life stages and recruitment reveal when year-class strength is determined and fixed.

## Recruitment level

Rijnsdorp *et al.* (1992) found a positive relationship between relative recruitment, both mean and maximum, and the approximate surface area of the nursery grounds for a number of sole stocks (Fig. 2). This intraspecific “nursery size hypothesis” appears to be valid also for plaice: the abundance of 0-group for different populations around the North Sea shows a positive relationship to the size of the nursery area (Fig. 3). Recently, Steinarsson and co-workers (unpubl.) carried out a similar analysis for subpopulations of plaice around Iceland and found a positive relationship between size of the nursery area and adult stock size (Fig. 4). Rijnsdorp *et al.* (1992) further suggested that the “nursery size hypothesis” might also be valid for interspecific differences between flatfish species. In examining the abundance of flatfish species in the North Sea, Gibson (1994) indeed found a significant positive correlation between the habitat requirements of juveniles in terms of depth range and their abundance (Fig. 5). The hypothesis might also explain why turbot (*Scophthalmus maximus* (L.)) and brill (*Scophthalmus rhombus* (L.)) are relatively rare species in the North Sea (Daan *et al.*, 1990), since their nursery areas are restricted to a narrow band along sandy beaches with a water depth of less than 1 m (Riley *et al.*, 1981). Obviously, the information available strongly supports the “nursery size hypothesis” not only as explanation of intraspecific but also of interspecific differences in recruitment.




Figure 2. Sole *Solea solea* stocks in ICES Subareas IV and VII. (a) Location of stocks and nursery areas. (b) Relationship between average recruitment (geometric mean  $\pm$  s.e.) and surface area of the nursery grounds (data after Rijnsdorp *et al.*, 1992).

The positive relationship between recruitment and nursery size raises the question whether nursery areas may ever become saturated with settling larvae and reach their “carrying capacity”. In this context, carrying capacity is defined according to MacCall’s theoretical basin model (1990) as the population density of a habitat at which the *per capita* population growth rate is zero. Studies on individual growth of juveniles have yielded contradictory results. Data on growth rate of 0-group plaice and sole generally failed to show an effect of population density. Spatial growth differences could be attributed to different origin of the settlers and its effects on timing of settlement in relation to distance to the spawning grounds (Karakiri *et al.*, 1991), while interannual growth variations could be explained by differences in temperature regimes (Edwards and Steele, 1968; Zijlstra *et al.*, 1982; Van der Veer, 1986; Bergman




Figure 3. Relationship between average year-class strength in 0-group plaice (*Pleuronectes platessa*) in August (mean  $\pm$  s.e.) and surface area of the nursery grounds for different areas around the North Sea (data after [1] Edwards and Steele, 1968; [2] Lockwood, 1974; [3] Zijlstra *et al.*, 1982; [4] Pihl, 1990; and [5] Mengedoh, 1995).





Figure 4. Plaice (*Pleuronectes platessa*) populations around Iceland. (a) Subdivisions used. (b) Relationship between stock biomass (mean  $\pm$  s.e.) estimated from CPUE multiplied by the surface area of the <100 m zone and the depth zone <20 m (as a measure of nursery size), 1985–1997 (data from Steinarrsson, unpubl.).




Figure 5. Relationship between stock numbers and maximum depth distribution as a measure of habitat size in North Sea flatfish species (data after Gibson, 1994).

*et al.*, 1988; Van der Veer *et al.*, 1990; Rogers, 1994; Nash *et al.*, 1994; Jager *et al.*, 1995). Only Modin and Pihl (1994) were able to show that density-dependent growth in 0-group plaice did occur at extremely high densities. Also, I- and II-group plaice have shown depressed growth rates during periods of high abundance (Rijnsdorp and Van Leeuwen, 1992, 1996).

Studies on population growth of juveniles are scarce. Zijlstra *et al.* (1982) and Van der Veer (1986) provide information for 0-group plaice in the Dutch Wadden Sea for a number of years (Fig. 6). Population growth is expressed as the change in total individual metabolic biomasses ( $W^{0.8}$ ), because the relationship between energy requirements and metabolic biomass is linear in contrast to that with biomass (Fonds *et al.*, 1992). In all years, population biomass increases after settlement in spring until emigration starts in autumn and does not level off during the summer period. The increase during the growing season implies that the *per capita* population growth rate remains positive, suggesting that the carrying capacity is never reached.

It should be noted that interpretation of these growth studies may be biased. Firstly, ambient temperatures cannot be estimated with high accuracy and the true effect may therefore be underestimated. Secondly, field studies have not addressed the potential effects of variations in abundance on the distribution of a year class over primary and marginal habitat. If a larger part of the population of a strong year class moves into a marginal habitat where it experiences lower growth rates and higher mortality rates, a study that is confined to the primary habitat may give a biased signal of the population-regulating processes. Thirdly, the lack of evidence for density-dependent growth in juveniles may be due to other disturbing factors. For example, eutrophication may have resulted in an increased production of suitable benthic food. There are indeed indications that growth rates were lower during the




Figure 6. Seasonal pattern in metabolic biomass of 0-group plaice (*Pleuronectes platessa*) at the Balgzand, 1975–1982. For each survey, individual biomass per 0.5 mm size class interval [ $W$  (g)= $0.0019 L^3$  (cm); Fonds *et al.*, 1992] was converted into metabolic biomass ( $MB=W^{0.8}$ ), multiplied by the number of individuals of that size, and summed over all size classes.

pre-eutrophication period (Rijnsdorp and Van Leeuwen, 1992; Gibson, 1994). There is ample evidence that food availability and quality is an important factor in determining growth (Rijnsdorp and Van Beek, 1991; Van der Veer and Witte, 1993; Berghahn *et al.*, 1995). Although the information is restricted to a few species only, the data do support the suggestion by Gibson (1994) that saturation of nursery grounds is rare or non-existent, implying that fluctuations in larval supply might be the key factor. Local physical and morphological conditions will determine the passive transport and the ultimate supply of larvae to the nurseries (Rijnsdorp *et al.*, 1985; Bergman *et al.*, 1989). A larger nursery area may be expected to trap on average a larger number of settling larvae.

On a global scale, Pauly (1994) observes a decrease in biomass – and hence recruitment – of flatfish from temperate to tropical areas. He suggests that flatfish are overadapted to feeding on zoobenthic infauna and epifauna, and that, in combination with a decreasing trend in importance of the benthic system towards the tropics, food-limitation might increase with decreasing latitude. This would explain lower biomass values. In contrast to Gibson (1994), Pauly (1994) implicitly assumes that the carrying capacity of the nursery acts as a limiting factor and that it decreases towards the tropics.

## Recruitment variability

Variability in recruitment is generally much larger than variability in spawning stock biomass or potential egg production, because the adult population incorporates members of several year classes. The implication is that variability-generating processes operate between spawning and recruitment. The evidence for variability-generating and variability-damping factors during different life stages within a population is discussed separately. Subsequently, the single stock approach will be expanded to differences between populations of the same species over its geographical range of distribution, and between species.

## Variability-generating factors

### The adult stage

For individual stocks, first maturation occurs within a specific size range. After spawning, the energy lost by reproductive investment has to be compensated for first before surplus energy may be used for growth. Since fecundity is size-specific, the number of eggs produced during consecutive spawning events will increase. Egg size may also increase with increasing body size and age. Furthermore, female plaice appears to start spawning earlier in the season as they grow older (Rijnsdorp, 1989). There is no indication whether sex ratio or size composition of the males affects spawning success. At the population level, total reproductive investment depends on sex ratio, size composition, and absolute numbers involved, but information on fertilization rate and interannual variability therein is lacking (Rijnsdorp, 1994).

Rijnsdorp (1994) provided evidence that variability-generating processes may occur in the adult phase. The interannual variability in size-specific fecundity varies in different species between 5% and 25%. Furthermore, exploitation may strongly modify the adult population in terms of absolute numbers, sex ratio, size composition, spawning season, and egg quality (size). Although the importance of maternal effects on reproductive potential has often been stressed (Solemdal, 1997), convincing evidence that these contribute strongly to generating interannual variability in recruitment is still lacking. Interannual variability in the number of fertilized eggs might be generated compared to a simple measure of spawning stock biomass by variations in any of the aspects mentioned above, but field evidence for major shifts is lacking. Rijnsdorp (1994) concludes that time series of recruitment so far do not provide evidence that these factors are important.

### Pelagic egg and larval stage

Leggett and DeBlois (1994) provide a review of the timing of year-class formation in marine fish. Their




Figure 7. Relationship between mean daily mortality and water temperature of plaice (*Pleuronectes platessa*) eggs during development in the Southern Bight of the North Sea (data after Harding *et al.*, 1978).

findings were consistent with Gulland's (1965) hypothesis of a coarse control during the period of egg and larval drift, suggesting that variability-generating processes operate during this period. Most studies focused on the possible role of food and predation as controlling factors. Notwithstanding the conflicting and often indirect nature of the information (Leggett and DeBlois, 1994), the hypothesis that year-class strength in marine fish larvae is determined by starvation and predation in the egg and larval stages could not be rejected.

In a zoogeographic study on recruitment variation in dab (*Limanda limanda* (L.)), the available evidence also points to critical events during the pelagic stage, suggesting that year-class strength might even be determined during the egg stage (Henderson, 1998). As in plaice (Van der Veer, 1986), a negative relationship between year-class strength and water temperature at the time of spawning was observed. Henderson (1998) assumed that pathogens may be important in determining egg mortality. Support for his hypothesis is provided by the significant positive correlation between egg mortality in plaice and sole and water temperature (Fig. 7), while prevalence of malformations in flatfish eggs was negatively correlated with water temperature (Dethlefsen *et al.*, 1996). Based on laboratory experiments, Fonds (1995) also suggested that the observed negative relationship between survival of plaice eggs and water temperature might be caused by a reduced activity of pathogenic bacteria at low temperatures.

Recently, the role of hydrodynamics in explaining interannual variability in flatfish recruitment has been given attention (Pepin *et al.*, 1995; Werner *et al.*, 1997). In their review, Werner *et al.* (1997) were not able to provide explicit links between physical variability and recruitment, despite a large number of studies for a variety of marine fish populations. On interannual time




Figure 8. Correlation between year-class strength of turbot (*Scophthalmus maximus*) around the coast of England and Wales and offshore winds. Dashed lines indicate 95% confidence limits (data after Riley *et al.*, 1981).

scales, they listed wind as the most intensively studied abiotic factor contributing to recruitment variability. Using wind as a measure of transport, Riley *et al.* (1981) found a significant positive correlation between the year-class strength of turbot around the coast of England and Wales and offshore winds (Fig. 8). Also in the petrale sole (*Eopsetta jordani* (Lockington)) along the northeast Pacific Ocean, variation in year-class strength was associated with oceanographic conditions during the pelagic larval stages (Castillo *et al.*, 1994). Coombs *et al.* (1990) suggest that weather conditions during embryonic development might significantly influence recruitment strength of North Sea plaice, rough weather supposedly leading to increased egg mortality. Recently, Nielsen *et al.* (1998) showed that year-class strength of plaice along the Danish Kattegat coast was related to prevailing wind conditions, suggesting that recruitment to the area depends on egg and larval transport from the Skagerrak. Further support for the importance of transport in determining year-class strength was provided by Van der Veer *et al.* (1998) in an analysis of the impact of interannual variability in hydrodynamic circulation on egg and larval transport of plaice in the southern North Sea. Simulations with a realistic 2D circulation model were compared with observed larval concentrations in the coastal zone. Results showed that interannual variability in transport was in the same order as observed in year-class strength. The authors concluded that the variability in circulation patterns during the early pelagic stages in the open sea might be a key factor. A similar suggestion was already made a few decades ago by Talbot (1976, 1978).

#### Demersal juvenile stage

Metamorphosis and settlement of flatfish larvae has been considered to be a key process in determining recruitment. The change from a symmetrical pelagic larva to a demersal flatfish requires extensive functional

changes, involving flattening of the body, strong development of the unpaired fins, often a marked reduction of the paired fins, eye migration, and changes in pigmentation (Fuiman, 1997; Osse, 1997). Especially the rewiring and recalibration of vision during eye migration will strongly affect food detection (Osse *et al.*, 1997) and maybe also predator avoidance. Although Creutzberg *et al.* (1978) indeed found that almost 100% of the settling plaice larvae had empty stomachs, starvation was not causing additional mortality but was merely a key factor in the settlement process. Starving larvae remained swimming until favourable benthic food conditions were encountered. Then the larvae settled and remained demersal. Also the positive linear relationship between abundance of pelagic plaice larvae and subsequently settled juveniles (Van der Veer, 1986) suggests the absence of either a starvation- or predation-induced variability-generating process.

Variability-generating processes during this phase can be introduced by variability in either mortality or the length of the period, or by a combination of these two factors. Van der Veer *et al.* (1994) reviewed the impact of juvenile growth on recruitment in flatfish, which might be caused either by size-selective mortality and/or by size-dependent onset of maturation. Although interannual variability existed in both factors, there was no indication that these processes were important. In a recent paper on size-selective predation on 0-group flatfish, this conclusion was confirmed (Van der Veer *et al.*, 1997).

Bailey (1994) discussed the impact of predation on recruitment variability in flatfish. Fluctuating numbers of predators are considered to be the most likely condition leading to increasing variability in recruitment, since predator population dynamics might be influenced by factors other than prey abundance. Iles and Beverton (1991) and Beverton and Iles (1992a, b) could not find any indication of variability-generating factors in a systematic analysis of mortality rates in juvenile flatfish in European waters. Bailey (1994) confirmed this general conclusion, but provided evidence that the situation might be different under exceptional conditions. In plaice, strong year classes are associated with cold winters (Brander and Houghton, 1982; Van der Veer, 1986; Van der Veer and Witte, 1999) and it has been argued that the extremely strong 1963 year class may have been partly caused by a release from predation by crustaceans in the nursery (Van der Veer and Bergman, 1987). Beukema (1991, 1992) observed a delay of about 1 month in the arrival of crustaceans after a cold winter. Other indirect support arises from observations by Pihl (1990) in plaice nurseries along the Swedish west coast. In these areas, with large year-to-year variability in winter temperature and hence in predator abundance, variability in year-class strength increased during the first year of life (Van der Veer *et al.*, 1990). Adverse

environmental conditions may also occur in the demersal stage. In North Sea sole, low winter temperatures caused high mortality in all demersal stages (Woodhead, 1964). Mortality between the 0- and I-group was negatively correlated with winter temperature, generating a larger variability in I-group abundance in the more northerly situated nursery areas only (Van Beek *et al.*, 1989; Rijnsdorp *et al.*, 1992).

Overall, there is no strong evidence that variability-generating processes in the demersal juvenile stage are common and of importance. Only in areas with cold and strongly varying winter temperatures might fluctuating numbers of predators induce interannual variability in year-class strength.

### Variability-damping factors

A first approach to check the presence of regulating factors is to investigate stock-recruitment data. Iles (1994) showed in a review of 20 flatfish stocks that in six cases there was a significant non-linear stock-recruitment relationship. This evidence for density dependence was strengthened by the fact that for eight additional stocks a model of constant recruitment was favoured over a model in which recruitment was directly proportional to stock size. The identification of the form of the relationship has considerable potential for insight in population-regulating mechanisms (Iles and Beverton, 1999). A further analysis by Beverton (1995) in relation to his "concentration hypothesis" yielded strong evidence that there are critical periods in both the adult and juvenile stage that are of primary importance in regulating recruitment.

#### The adult stage

Regulating processes may operate through density-dependent adult mortality, growth, maturation, and/or fecundity. Data on density-dependent adult mortality are virtually absent. Predation would seem to become less important with increasing fish size. The general size rules in predator-prey relationships indicate that mortality through cannibalism in the adult phase is unlikely. Whereas optimum prey size appears to be in general about one-third of the predator size, the length at first maturity is at half the asymptotic maximum length in most flatfish species (Beverton, 1992).

Growth, maturation, and fecundity are intricately linked within the energy budget of a fish, where the amount of surplus energy plays a crucial role (Rijnsdorp, 1989, 1993). Factors affecting surplus energy, such as intra- and interspecific competition and environmental conditions (especially food availability and temperature) will also affect the allocation of energy over somatic growth and reproduction. On an annual

basis, surplus energy is a continuously increasing function of body size, and reproductive investment is a function of body weight. Reproductive effort is not only determined by egg numbers, but also by egg size. Egg size shows in general an inverse trend during the spawning season of an individual fish and a positive trend with increasing fish size (Solemdal, 1997). Annual variations in surplus energy in the adult phase will be reflected in both somatic growth and corresponding reproductive effort. The most extreme trade-off between somatic growth and reproduction exist when adults skip a spawning season (intermittent spawning).

Regulating processes should become particularly evident at high levels of abundance or biomass. However, most information available is derived from commercially important populations which appear to have been reduced by fishing below the threshold at which food supply per fish becomes limiting (Rijnsdorp, 1994). At such reduced levels of abundance, regulatory processes can no longer be detected. The analysis of available information from the early years of exploitation (Rijnsdorp, 1994) suggests that density-dependent growth has occurred in the past. The feature of intermittent spawning has also been suggested, at least in fish inhabiting colder waters. On the other hand, size-specific fecundity appeared to be stable over a broad range of population abundance and may only decrease at very high values. The mechanisms involved are further complicated by the trade-off between egg numbers and egg size. Overall, Rijnsdorp (1994) concluded that density-dependent ripening and absolute fecundity, mediated through density-dependent growth, may regulate recruitment at high levels of population abundance.

#### Pelagic egg and larval stage

Information on regulating processes operating during egg and larval development are scarce. Predation-induced mortality by planktivorous fish has been identified as a main source of mortality (Pommeranz, 1981; Daan *et al.*, 1985; Ellis and Nash, 1997).

No evidence for regulatory processes has been found in the egg stage. Observations on plaice and sole eggs (Harding *et al.*, 1978; Van der Land, 1991) indicate mortality to be independent of density (Fig. 9a). In both species, mortality is related to water temperature (Van der Land, 1991; Rijnsdorp and Vingerhoed, 1994). However, stage-specific cumulative mortality rates will be less dependent on temperature because of the levelling out effect of temperature on daily growth and daily mortality rates (Pepin, 1991).

Also during larval development, data suggesting the existence of major regulatory processes are lacking. The available quantitative information for plaice (Harding *et al.*, 1978) suggests a negative correlation (not significant;  $r_s = -0.40$ ;  $n=4$ ) between larval abundance and




Figure 9. Relationship between daily mortality ( $Z$ ) and abundance of pelagic first egg and larval stages of plaice (*Pleuronectes platessa*) (data after Harding *et al.*, 1978) and sole (*Solea solea*) (data after Van der Land, 1991) in the Southern Bight of the North Sea. (a) Eggs of stage I. (b) Larvae of stage I.

mortality (Fig. 9b), indicating a variability-generating rather than damping process.

### Demersal juvenile stage

Regulating processes operating during the juvenile stage could act either on stage duration through density-dependent growth (for review, see Van der Veer *et al.*, 1994) or on survival through density-dependent mortality. Extensive research on plaice has revealed the general picture that mean length of juveniles is reduced at high densities as observed during the pre-war period and during the time of the two exceptionally strong year classes of 1963 and 1985 (Rijnsdorp and Van Leeuwen, 1992, 1996). In contrast, there is no evidence for density-dependent growth of 0-group plaice in the nursery areas along the European coast (Zijlstra *et al.*, 1982; Van der Veer, 1986; Bergman *et al.*, 1988; Van der Veer *et al.*, 1990; Mengedoht, 1995). However, this lack of evidence may be due to our inability to estimate precisely the

effect of differences in ambient temperatures on growth rate and to possible bias in the field studies (Van der Veer *et al.*, 1997).

Population regulation through density-dependent mortality has also been observed. In Japanese flounder [*Paralichthys olivaceus* (Temminck and Schlegel)], cannibalism of I- and II-group on 0-group is an important source of mortality (Furuta, 1996). A comparison of mortality rates of demersal 0-group plaice, dab, and turbot in European waters yielded significant density-dependent effects (Iles and Beverton, 1991; Beverton and Iles, 1992a, b). Predation is probably the most important source of mortality, since a variety of crustaceans, fish, and birds have been found to prey upon juvenile flatfish (for review, see Van der Veer *et al.*, 1997). At least for predation by crustaceans, density-dependent effects were generated by both a functional and a numerical response of the predator (Van der Veer and Bergman, 1987).

### Intra- and interspecific variation

The analysis of variability-generating and variability-damping processes so far does not take into account that species vary in life history traits over their range of distribution (e.g. Leggett and Carscadden, 1978; Houde, 1989) and, therefore, their likely susceptibility to different control mechanisms. Miller *et al.* (1991), in applying this reasoning, developed a hypothesis, which has been later referred to as the “species range hypothesis” (Leggett and Frank, 1997). The underlying assumptions are: (1) species differ in life history traits, and therefore in susceptibility to different controls on recruitment, and (2) species life history traits vary over their range of distribution, and therefore controlling factors differ over both latitudinal and inshore–offshore gradients. Using latitudinal and inshore–offshore gradients in food, predation, and abiotic factors, these assumptions lead to the hypothesis that factors controlling recruitment of a species vary over its range with the following implications: (1) abiotic factors are most important at the edges of the range; (2) predation plus abiotic factors control recruitment at the polar edge of the range; (3) food plus abiotic factors control recruitment at the equatorial edge.

Miller *et al.* (1991) predicted that recruitment would be most variable at the polar edge of the species range, least near the centre of the range, and be intermediate near the equatorial edge (Fig. 10). However, they emphasized that the latitudinal pattern might be modified or obscured in the case of migratory species and that inshore–offshore environmental gradients may swamp latitudinal effects.

A first attempt to falsify the species range hypothesis was made by Walsh (1994). His analysis of recruitment variability in populations of long rough dab






Figure 10. Schematic representation (redrawn after Leggett and Frank, 1997) of the species range hypothesis of Miller *et al.* (1991) with respect to: (a) coefficient of variation in recruitment, (b) importance of density-dependent processes.




Figure 11. Latitudinal recruitment variability of age-4 long rough dab (American plaice) *Hippoglossoides platessoides* in the western and eastern North Atlantic expressed as the standard deviation of the residuals of detrended log (base 10) recruitment (data after Walsh, 1994).

(=American plaice) (*Hippoglossoides platessoides* (Fabricius)) showed an increasing south to north latitudinal gradient without a valley in the centre (Fig. 11), and on this basis at least part of the hypothesis would have to be rejected. Leggett and Frank (1997) expanded the validation by including a variety of North Atlantic flatfish species, based on stock and recruitment data compiled by Myers *et al.* (1995). The results were generally inconsistent with the species range hypothesis and they concluded that recruitment variability is dominated by density-independent factors operating at a local scale on the egg and larval stages. Neither Walsh (1994) nor Leggett and Frank (1997) considered

inshore–offshore gradients. Philippart *et al.* (1998), in an analysis of latitudinal variation in recruitment in north-west Europe based on other data sources, observed patterns of variation which also did not correspond with the expectation. However, they suggested that other sources of variation might have masked the expected patterns, such as different sampling methods, average age at recruitment, inshore–offshore gradients, and shifts in species ranges over time.

The basic assumption that species differ in susceptibility to different controls on recruitment still seems valid. In a recent paper, Tanaka *et al.* (1997) present an overview of latitudinal differences in life history traits in Japanese flounder, in combination with seasonal differences in food availability and potential predators. The abundance of mysids, one of the main food sources, decreased markedly in the southern range at increasing temperatures. Abundance of predatory shrimps showed a reverse trend with highest densities at the northern edge. Miller *et al.* (1991) based their hypothesis on likely trends in food availability, predation pressure, and abiotic factors. These factors are most likely to act only in the juvenile stage, while year-class strength appears to be established already in the pelagic phase (Leggett and Frank, 1997). This suggests that the species range hypothesis does not apply to the main factors causing the coarse control, but to the subsequent fine control. The dominance of density-independent factors operating at a local scale on the egg and larval stages stresses the importance of hydrodynamic circulation as a key factor in determining year-class strength (Leggett and Frank, 1997). It is noteworthy that the Japanese flounder spawns closer to the inshore nursery area than most species, and thus may escape many of the vagaries caused by long migration ways to the nurseries (Minami and Tanaka, 1992).

If hydrodynamic circulation represents a key factor, it can be hypothesized that recruitment variability depends on pelagic stage duration as a forcing factor. Since egg and larval development is strongly affected by temperature, the implication is that recruitment variability over the range of a species should be inversely correlated with temperature, as observed for long rough dab (Fig. 11). Furthermore, a large portion of the patterns observed by Leggett and Frank (1997) for plaice and sole might be explained this way. Exceptions may be found in areas where temperature is inversely correlated with hydroclimate for migration, e.g., stronger shoreward currents in colder years (Miller *et al.*, 1984).

## Synthesis

In addressing variability, Heath's (1992) statement should not be forgotten that all the attention on recruitment variability masks the underlying stability of marine

populations. Slight variations in mortality over the larval stage would lead to much larger variations in recruitment than actually observed. In fact, recruitment to most populations is remarkably stable.

Variability-generating processes occur in all life stages of flatfish, but only those operating in the 3-D pelagic egg and larval stages appear to be of major importance. After settlement and transfer to the 2-D environment, regulating mechanisms dominate. This general scheme may be modified by at least four factors: fisheries, location within the range of the species, species type, and feeding type.

The main effect of fisheries appears to be that adult biomass may be reduced below the threshold at which food supply per fish is limiting (Rijnsdorp, 1994), in which case intraspecific regulating processes in the adult phase (food limitation, intermittent spawning) are no longer expected to occur. It is unclear whether this has also affected recruitment variability. The impact of the variability-generating processes in the egg and larval stages might well dominate the regulating processes in the adult stage.

Location is obviously important, because food and predators show clear trends over the distribution range of a species. Although detailed information is lacking, especially predation on early juvenile stages varies. With decreasing water temperature, the abundance of mobile epibenthic predators (crustaceans) becomes more variable (Van der Veer *et al.*, 1990), mainly as a result of lethal winter temperatures or avoidance behaviour. The consequence is a shift from regulation to variability generation during the juvenile stage. No information is available concerning the extent to which this is also true for other sources of juvenile mortality, such as predation by larger fish, which appear to be even more capable of avoiding unfavourable environments.

Within the Pleuronectiformes, two major groups can be distinguished: those from cold waters, principally members of the Pleuronectidae, and those from warm-water habitats, including members of the Citharidae, Paralichthyidae, Bothidae, Solidae, and Cynoglossidae. In temperate locations, the groups overlap. Depending on their origin in the cold or warm region, species may show a different physiological response to their environment and as a consequence differences in recruitment patterns between species and differences over the range of species are expected.

With respect to feeding, a specialization into three groups has been recognized: fish feeders, crustacean feeders, and polychaete-mollusc feeders (De Groot, 1971). Most species change their diet from zooplankton to (epi)benthic prey after settlement. The abundance in epibenthic prey (e.g. mysids; Tanaka *et al.*, 1997) is thought to be more variable than that of benthic food items (e.g. regenerating body parts; Kuipers, 1977; De Vlas, 1979). If variable food conditions are important in

generating recruitment variability, this is probably more so in species depending on epibenthic prey than in species depending on benthic prey.

Hjort (1914) put forward the hypothesis that year-class strength is controlled during the early life history, when numbers of individuals are at a maximum. Gulland (1965) suggested that the coarse control during the period of egg and larval drift was followed by a second period of fine regulation later on. The findings of Leggett and DeBlois (1994) supported this view: year-class strength in marine fish appears to be established during the pelagic egg or larval period.

According to Leggett and Frank (1997), regulation principally operates at the local (stock) scale rather than at large geographic scales. Positive correlations in recruitment time series occurred only among stocks in close geographic proximity to each other (Rijnsdorp *et al.*, 1992; Leggett and Frank, 1997), suggesting that oceanographic conditions such as wind or sea surface temperatures are acting as forcing functions over some 100–500 km (Myers *et al.*, 1997). This suggestion is consistent with other independent observations. Recruitment patterns in Northwest Atlantic fish stocks indicate that large-scale physical forcing, rather than local biological interactions, largely determines year-class strength (Koslow, 1984). Other analyses have also suggested that recruitment success of species in the North-east Pacific Ocean and in the Northwest Atlantic is strongly influenced by environmental conditions (Hollowed *et al.*, 1987; Frank, 1991; Planque and Fox, 1998). For instance, year-to-year variability in temperature has been shown to be linked with year-class strength (Frank, 1991; Planque and Fox, 1998). However, the correlations varied over the range of distribution. Positive relationships were found near the northern limits, negative relationships near the southern limits and no correlation was observed in the centre of the distribution (Frank, 1991).

A striking difference between flatfish and most other marine fish species is the relatively low recruitment variability among the former, suggesting that the degree of variation might be inversely related to the degree of concentration during early life phases (Beverton, 1984). Later, Beverton (1995) elaborated on his “concentration hypothesis” and put forward the statement that species which tend to concentrate spatially on nursery grounds during their early life may be “saturating” the carrying capacity of those habitats even when the adult population is still relatively small and not limited by their habitat. According to his view, this saturation results in density-dependent processes and explains the relatively low variability in these species. However, one might wonder whether it is the degree of concentration during early life that is responsible for the observed patterns. Regulating processes in flatfish were only observed to operate during the demersal stage and the concentration

into nursery areas coincides with the transfer from the plankton to the demersal stage. Therefore, instead of the degree of concentration, the underlying mechanism might be that regulating mechanisms only occur during demersal life stages. This might explain the apparent paradox that the lemon sole (*Microstomus kitt* L.) has a low recruitment variability even though this flatfish species does not concentrate spatially (Rae, 1970). In this respect, the results of Sundby *et al.* (1989) for cod are intriguing. Mortality in 2–3 months old juveniles in the transition of becoming demersal (but still at least partly planktonic) was density-dependent. If the implication that year-class strength variability can only decrease in fish species with a demersal stage is true, variability in recruitment in fish species will be related to the relative duration of the pelagic and demersal stages.

## Acknowledgements

This personal account of the progress made during three Flatfish Symposia at NIOZ would not have been possible without the inspiring support of Ray Beverton, as well as his scientific ideas, in setting up these symposia, nor without the stimulating discussions and support of all participants. We also thank the reviewers for improving the manuscript. This is NIOZ publication No 3232. JMM was supported by NOAA, US Dept. of Commerce, through NC Sea Grant NA90AA-D-SG-062.

## References

- Anon. 1995. Proceedings of the International Symposium on North Pacific Flatfish. University of Alaska, Fairbanks. 643 pp.
- Bailey, K. M. 1994. Predation on juvenile flatfish and recruitment variability. *Netherlands Journal of Sea Research*, 32: 175–189.
- Berghahn, R. 1987. Effects of tidal migration on growth of 0-group plaice (*Pleuronectes platessa* L.) in the North Frisian Wadden Sea. *Berichte der Deutschen Wissenschaftlichen Kommission für Meeresforschung*, 31: 209–226.
- Berghahn, R., Lüdemann, K., and Ruth, M. 1995. Differences in individual growth of newly settled 0-group plaice (*Pleuronectes platessa* L.) in the intertidal of neighbouring Wadden Sea areas. *Netherlands Journal of Sea Research*, 34: 131–138.
- Bergman, M. J. N., Van der Veer, H. W., Stam, A., and Zuidema, D. 1989. Transport mechanisms of larval plaice (*Pleuronectes platessa* L.) from the coastal zone into the Wadden Sea nursery areas. *Rapports et Procès-Verbaux des Réunions du Conseil International pour l'Exploration de la Mer*, 191: 43–49.
- Bergman, M. J. N., Van der Veer, H. W., and Zijlstra, J. J. 1988. Plaice nurseries: effects on recruitment. *Journal of Fish Biology*, 33 Supplement A: 201–218.
- Beukema, J. J. 1991. The abundance of shore crabs *Carcinus maenas* (L.) on a tidal flat in the Wadden Sea after cold and mild winters. *Journal of Experimental Marine Biology and Ecology*, 153: 97–113.
- Beukema, J. J. 1992. Dynamics of juvenile shrimp *Crangon crangon* in a tidal-flat nursery of the Wadden Sea after mild and cold winters. *Marine Ecology Progress Series*, 83: 157–165.
- Beverton, R. J. H. 1984. Dynamics of single species. *In* Exploitation of marine communities, vol 32, pp. 13–58. Ed. by R. M. May. Springer Verlag, Berlin.
- Beverton, R. J. H. 1992. Fish resources; threats and protection. *Netherlands Journal of Zoology*, 42: 139–175.
- Beverton, R. J. H. 1995. Spatial limitation of population size; the concentration hypothesis. *Netherlands Journal of Sea Research*, 34: 1–6.
- Beverton, R. J. H., and Iles, T. C. 1992a. Mortality rates of 0-group plaice (*Pleuronectes platessa* L.), dab (*Limanda limanda* L.) and turbot (*Scophthalmus maximus* L.) in European waters II. Comparison of mortality rates and construction of life table for 0-group plaice. *Netherlands Journal of Sea Research*, 29: 49–59.
- Beverton, R. J. H., and Iles, T. C. 1992b. Mortality rates of 0-group plaice (*Pleuronectes platessa* L.), dab (*Limanda limanda* L.) and turbot (*Scophthalmus maximus* L.) in European waters III. Density-dependence of mortality rates of 0-group plaice and some demographic implications. *Netherlands Journal of Sea Research*, 29: 61–79.
- Brander, K., and Houghton, R. G. 1982. Predicting the recruitment of North Sea plaice from egg surveys. *ICES CM* 1982/G:5.
- Castillo, G. C., Li, H. W., and Golden, J. T. 1994. Environmentally induced recruitment in petrale sole, *Eopsetta jordani*. *Fishery Bulletin U.S.*, 92: 481–493.
- Coombs, S. H., Nichols, J. H., and Fosh, C. A. 1990. Plaice eggs (*Pleuronectes platessa* L.) in the southern North Sea: abundance, spawning area, vertical distribution, and buoyancy. *Journal du Conseil International pour l'Exploration de la Mer*, 47: 133–139.
- Creutzberg, F., Eltink, A. Th. G. W., and Van Noort, G. J. 1978. The migration of plaice larvae *Pleuronectes platessa* into the western Wadden Sea. *In* Physiology and behaviour of marine organisms. Proceedings of the 12th European Symposium on Marine Biology, pp. 243–251. Ed. by D. S. McLusky, and A. J. Berry. Pergamon Press, New York.
- Daan, N., Rijnsdorp, A. D., and Van Overbeeke, G. R. 1985. Predation by North Sea herring *Clupea harengus* on eggs of the plaice *Pleuronectes platessa* and the cod *Gadus morhua*. *Transactions of the American Fisheries Society*, 114: 499–506.
- Daan, N., Bromley, P. J., Hislop, J. R. G., and Nielsen, N. A. 1990. Ecology of North Sea fish. *Netherlands Journal of Sea Research*, 26: 343–386.
- De Groot, S. J. 1971. On the interrelationships between morphology of the alimentary tract, food and feeding behaviour in flatfishes (Pisces: Pleuronectiformes). *Netherlands Journal of Sea Research*, 5: 121–196.
- De Vlas, J. 1979. Annual food intake by plaice and flounder in a tidal flat area in the Dutch Wadden Sea, with special reference to consumption of regenerating parts of macrobenthic prey. *Netherlands Journal of Sea Research*, 13: 117–153.
- Dethlefsen, V., Von Westernhagen, H., and Cameron, P. 1996. Malformations in North Sea pelagic fish embryos during the period 1984–1995. *ICES Journal of Marine Science*, 53: 1024–1035.
- Edwards, R., and Steele, J. H. 1968. The ecology of 0-group plaice and common dab at Loch Ewe. I. Population and food. *Journal of Experimental Marine Biology and Ecology*, 2: 215–238.
- Ellis, T., and Nash, R. D. M. 1997. Predation by sprat and herring on pelagic fish eggs in a plaice spawning area in the Irish Sea. *Journal of Fish Biology*, 50: 1195–1202.

- Fonds, M. 1995. Effect of winter temperature on the brood success of flatfish. In Annual Report 1995, p. 59. Netherlands Institute for Sea Research, Texel.
- Fonds, M., Cronie, R., Vethaak, A. D., and Van der Puyl, P. 1992. Metabolism, food consumption and growth of plaice (*Pleuronectes platessa*) and flounder (*Platichthys flesus*) in relation to fish size and temperature. Netherlands Journal of Sea Research, 29: 127–143.
- Frank, K. T. 1991. Predicting recruitment variation from year class specific vertebral counts: an analysis of the potential and a plan for verification. Canadian Journal of Fisheries and Aquatic Science, 48: 1350–1357.
- Fuiman, L. A. 1997. What can flatfish ontogenies tell us about pelagic and benthic lifestyles? Journal of Sea Research, 37: 257–267.
- Furuta, S. 1996. Predation on juvenile Japanese flounder (*Paralichthys olivaceus*) by diurnal piscivorous fish: field observations and laboratory experiments. In Survival strategies in early life stages of marine resources, pp. 285–294. Ed. by Y. Watanabe, Y. Yamashita, and Y. Oozeki. A. A. Balkema, Rotterdam.
- Gibson, R. N. 1994. Impact of habitat quality and quantity on the recruitment of flatfishes. Netherlands Journal of Sea Research, 32: 191–206.
- Gulland, J. A. 1965. Survival of the youngest stages of fish and its relation to year-class strength. International Commission for the Northwest Atlantic Fisheries, 6: 365–371.
- Harding, D., Nicols, J. H., and Tungate, D. S. 1978. The spawning of the plaice (*Pleuronectes platessa* L.) in the Southern North Sea and the English Channel. Rapports et Procès-Verbaux des Réunions du Conseil International pour l'Exploration de la Mer, 172: 102–113.
- Heath, M. R. 1992. Field investigations of the early life stages of marine fish. Advances in Marine Biology, 28: 2–133.
- Henderson, P. 1998. On the variation in dab *Limanda limanda* recruitment: a zoogeographic study. Journal of Sea Research, 40: 131–142.
- Hjort, J. 1914. Fluctuations in the great fisheries of Northern Europe viewed in the light of biological research. Rapports et Procès-Verbaux des Réunions du Conseil International pour l'Exploration de la Mer, 20: 1–228.
- Hollowed, A. B., Bailey, K. M., and Wooster, W. S. 1987. Patterns in recruitment of marine fishes in the Northeast Pacific Ocean. Biological Oceanography, 5: 99–131.
- Houde, E. D. 1989. Comparative growth, mortality, and energetics of marine fish larvae: temperature and implied latitudinal effects. Fishery Bulletin U.S., 87: 471–495.
- Hovenkamp, F. 1992. Growth-dependent mortality of larval plaice *Pleuronectes platessa* in the North Sea. Marine Ecology Progress Series, 82: 95–101.
- Iles, T. C. 1994. A review of stock-recruitment relationships with reference to flatfish populations. Netherlands Journal of Sea Research, 32: 399–420.
- Iles, T. C., and Beverton, R. J. H. 1991. Mortality rates of 0-group plaice (*Pleuronectes platessa* L.), dab (*Limanda limanda* L.) and turbot (*Scophthalmus maximus* L.) in European waters. I. Statistical analysis of the data and estimation of parameters. Netherlands Journal of Sea Research, 27: 217–235.
- Iles, T. C., and Beverton, R. J. H. 1999. The concentration hypothesis: the statistical evidence. ICES Journal of Marine Science (this volume).
- Jager, Z., Kleef, H. L., and Tydeman, P. 1995. Mortality and growth of 0-group flatfish in the brackish Dollard (Ems estuary, Wadden Sea). Netherlands Journal of Sea Research, 34: 119–129.
- Karakiri, M., Berghahn, R., and Van der Veer, H. W. 1991. Variations in settlement and growth of 0-group plaice (*Pleuronectes platessa* L.) in the Dutch Wadden Sea as determined by otolith microstructure analysis. Netherlands Journal of Sea Research, 27: 345–351.
- Koslow, J. A. 1984. Recruitment patterns in Northwest Atlantic fish stocks. Canadian Journal of Fisheries and Aquatic Science, 41: 1722–1729.
- Kuipers, B. R. 1977. On the ecology of juvenile plaice on a tidal flat in the Wadden Sea. Netherlands Journal of Sea Research, 11: 56–91.
- Leggett, W. C., and Carscadden, J. E. 1978. Latitudinal variation in reproductive characteristics of American shad (*Alosa sapidissima*): evidence for population specific life history strategies in fish. Journal of the Fisheries Research Board of Canada, 35: 1469–1478.
- Leggett, W. C., and DeBlois, E. M. 1994. Recruitment in marine fishes: is it regulated by starvation and regulation in the egg and larval stages? Netherlands Journal of Sea Research, 32: 119–134.
- Leggett, W. C., and Frank, K. T. 1997. A comparative analysis of recruitment variability in North Atlantic flatfishes – testing the species range hypothesis. Journal of Sea Research, 37: 281–299.
- Lockwood, S. J. 1974. The settlement, distribution and movements of 0-group plaice *Pleuronectes platessa* (L.) in Filey Bay, Yorkshire. Journal of Fish Biology, 6: 465–477.
- MacCall, A. D. 1990. Dynamic geography of marine fish populations. University of Washington Press, Seattle. 153 pp.
- Mengedoht, D. M. 1995. Growth and mortality of 0-group plaice *Pleuronectes platessa* L. in coastal waters. Internal Report Netherlands Institute for Sea Research, 1995–8. 22 pp.
- Miller, J. M., Burke, J. S., and Fitzhugh, G. R. 1991. Early life history patterns of Atlantic North American flatfish: likely and (unlikely) factors controlling recruitment. Netherlands Journal of Sea Research, 27: 261–275.
- Miller, J. M., Reed, J. P., and Pietrafesa, L. J. 1984. Patterns, mechanisms and approaches to the study of migrations of estuarine-dependent fish larvae and juveniles. In Mechanisms of migration in fishes, pp. 209–225. Ed. by J. D. McCleave, G. P. Arnold, J. J. Dodson, and W. H. Neill. Plenum Press, New York.
- Minami, T., and Tanaka, M. 1992. Life history cycles in flatfish from the northwestern Pacific, with particular reference to their early life histories. Netherlands Journal of Sea Research, 29: 35–48.
- Modin, J., and Pihl, L. 1994. Differences in growth and mortality of juvenile plaice, *Pleuronectes platessa* L., following normal and extremely high settlement. Netherlands Journal of Sea Research, 32: 331–341.
- Myers, R. A., Bridson, J., and Barrowman, N. J. 1995. Summary of worldwide spawner and recruitment data. Canadian Technical Report of Fisheries and Aquatic Science, 2024: 1–274.
- Myers, R. A., Mertz, G., and Bridson, J. 1997. Spatial scales of interannual recruitment variations of marine, anadromous, and freshwater fish. Canadian Journal of Fisheries and Aquatic Science, 54: 1400–1407.
- Nash, R. D. M., Geffen, A. J., and Hughes, G. 1994. Individual growth of juvenile plaice (*Pleuronectes platessa* L.) on a small Irish Sea nursery ground (Port Erin Bay, Isle of Man, UK). Netherlands Journal of Sea Research, 32: 369–378.
- Nielsen, E., Bagge, O., and MacKenzie, B. R. 1998. Wind-induced transport of plaice (*Pleuronectes platessa*): early life-history stages in the Skagerrak-Kattegat. Journal of Sea Research, 39: 11–28.
- Osse, J. W. M., and Van den Boogaart, J. G. M. 1997. Size of flatfish larvae at transformation, functional demands and historical constraints. Journal of Sea Research, 37: 229–239.

- Pauly, D. 1994. A framework for latitudinal comparisons of flatfish recruitment. *Netherlands Journal of Sea Research*, 32: 107–118.
- Pearcy, W. C. 1962. Ecology of an estuarine population of winter flounder, *Pseudopleuronectes americanus* (Walbaum). Part I–IV. *Bulletin of the Bingham Oceanographic Collection*, 18: 1–78.
- Pepin, P. 1991. Effect of temperature and size on development, mortality, and survival rates of the pelagic early life history stages of marine fish. *Canadian Journal of Fisheries and Aquatic Science*, 48: 503–518.
- Pepin, P., Helbig, J. A., Laprise, R., Colbourne, E., and Shears, T. H. 1995. Variations in the contribution of transport to changes in planktonic animal abundance: a study of the flux of fish larvae in Conception Bay, Newfoundland. *Canadian Journal of Fisheries and Aquatic Science*, 52: 1475–1486.
- Philippart, C. J. M., Henderson, P. A., Johannessen, T., Rijnsdorp, A. D., and Rogers, S. I. 1998. Latitudinal variation in fish recruits in Northwest Europe. *Journal of Sea Research*, 39: 69–77.
- Pihl, L. 1990. Year-class strength regulation in plaice (*Pleuronectes platessa* L.) on the Swedish west coast. *Hydrobiologia*, 195: 79–88.
- Planque, B., and Fox C. J. 1998. Interannual variability in temperature and the recruitment of Irish Sea cod. *Marine Ecology Progress Series*, 172: 101–105.
- Pommeranz, T. 1981. Observations on the predation of herring (*Harengus harengus* L.) and sprat (*Sprattus sprattus* L.) on fish eggs and larvae in the southern North Sea. *Rapports et Procès-Verbaux des Réunions du Conseil International pour l'Exploration de la Mer*, 178: 402–404.
- Rae, B. B. 1970. The distribution of flatfishes in Scottish and adjacent waters. *Marine Research*, 2: 1–39.
- Rijnsdorp, A. D. 1989. Maturation of male and female North Sea plaice (*Pleuronectes platessa* L.). *Journal du Conseil International pour l'Exploration de la Mer*, 46: 35–51.
- Rijnsdorp, A. D. 1993. Relationship between juvenile growth and the onset of sexual maturity of female North Sea plaice, *Pleuronectes platessa*. *Canadian Journal of Fisheries and Aquatic Science*, 50: 1617–1631.
- Rijnsdorp, A. D. 1994. Population-regulating processes during the adult phase in flatfish. *Netherlands Journal of Sea Research*, 32: 207–223.
- Rijnsdorp, A. D., and Van Beek, F. A. 1991. The effects of the plaice box on the reduction in discarding and on the level of recruitment of North Sea sole. *ICES CM 1991/G:47*.
- Rijnsdorp, A. D., Van Beek, F. A., Flatman, S., Millner, R. M., Riley, J. D., Giret, M., and De Clerck, R. 1992. Recruitment in sole stocks, *Solea solea* (L.) in the northeast Atlantic. *Netherlands Journal of Sea Research*, 29: 173–192.
- Rijnsdorp, A. D., and Van Leeuwen, P. I. 1992. Density-dependent and density-independent changes in somatic growth of female plaice *Pleuronectes platessa* between 1930 and 1985 as revealed by back-calculation of otoliths. *Marine Ecology Progress Series*, 88: 19–32.
- Rijnsdorp, A. D., and Van Leeuwen, P. I. 1996. Changes in growth of North Sea plaice since 1950 in relation to density, eutrophication, beam-trawl effort, and temperature. *ICES Journal of Marine Science*, 53: 1199–1213.
- Rijnsdorp, A. D., Van Stralen, M., and Van der Veer, H. W. 1985. Selective tidal transport of North Sea plaice *Pleuronectes platessa* in coastal nursery areas. *Transactions of the American Fisheries Society*, 114: 461–470.
- Rijnsdorp, A. D., and Vingerhoed, B. 1994. The ecological significance of geographical and seasonal differences in egg size in sole *Solea solea* (L.). *Netherlands Journal of Sea Research*, 32: 255–270.
- Riley, J. D., Symonds, D. J., and Woolner, L. 1981. On the factors influencing the distribution of 0-group demersal fish in coastal waters. *Rapports et Procès-Verbaux des Réunions du Conseil International pour l'Exploration de la Mer*, 178: 223–228.
- Rogers, S. I. 1994. Population density and growth rate of juvenile sole *Solea solea* (L.). *Netherlands Journal of Sea Research*, 32: 353–360.
- Solemdal, P. 1997. Maternal effects – a link between the past and the future. *Journal of Sea Research*, 37: 213–227.
- Sundby, S., Bjørke, H., Soldal, A. V., and Olsen, S. 1989. Mortality rates during the early life stages and year-class strength of northeast Arctic cod (*Gadus morhua* L.). *Rapports et Procès-Verbaux des Réunions du Conseil International pour l'Exploration de la Mer*, 191: 351–358.
- Talbot, J. W. 1976. The dispersal of plaice eggs and larvae in the Southern Bight of the North Sea. *Journal du Conseil International pour l'Exploration de la Mer*, 37: 221–248.
- Talbot, J. W. 1978. Changes in plaice larval dispersal in the last fifteen years. *Rapports et Procès-Verbaux des Réunions du Conseil International pour l'Exploration de la Mer*, 172: 114–123.
- Tanaka, M., Ohkawa, T., Maeda, T., Kinoshita, I., Seikai, T., and Nishida, M. 1997. Ecological diversities and stock structure of the flounder in the Sea of Japan in relation to stock enhancement. *Bulletin of the National Research Institute of Aquaculture*, Suppl. 3.
- Van Beek, F. A., Rijnsdorp, A. D., and De Clerck, R. 1989. Monitoring juvenile stocks of flatfish in the Wadden Sea and the coastal areas of the southeastern North Sea. *Helgoländer Meeresuntersuchungen*, 43: 275–293.
- Van der Land, M. A. 1991. Distribution of flatfish eggs in the 1989 egg surveys in the southeastern North Sea, and mortality of plaice and sole eggs. *Netherlands Journal of Sea Research*, 27: 277–286.
- Van der Veer, H. W. 1986. Immigration, settlement and density-dependent mortality of a larval and early post-larval 0-group plaice (*Pleuronectes platessa*) population in the western Dutch Wadden Sea. *Marine Ecology Progress Series*, 29: 223–236.
- Van der Veer, H. W., Berghahn, R., and Rijnsdorp, A. D. 1994. Impact of juvenile growth on recruitment in flatfish. *Netherlands Journal of Sea Research*, 32: 153–173.
- Van der Veer, H. W., and Bergman, M. J. N. 1987. Predation by crustaceans on a newly settled 0-group plaice *Pleuronectes platessa* population in the western Wadden Sea. *Marine Ecology Progress Series*, 35: 203–215.
- Van der Veer, H. W., Ellis, T., Miller, J. M., Pihl, L., and Rijnsdorp, A. D. 1997. Size-selective predation on juvenile North Sea flatfish and possible implications for recruitment. *In* Early life history and recruitment in fish populations, pp. 279–303. Ed. by R. C. Chambers, and E. A. Trippel. Chapman and Hall, London.
- Van der Veer, H. W., Pihl, L., and Bergman, M. J. N. 1990. Recruitment mechanisms in North Sea plaice *Pleuronectes platessa*. *Marine Ecology Progress Series*, 64: 1–12.
- Van der Veer, H. W., Ruardy, P., Van den Berg, A. J., and Ridderinkhof, H. 1998. Impact of interannual variability in hydrodynamic circulation on egg and larval transport of plaice *Pleuronectes platessa* L. in the southern North Sea. *Journal of Sea Research*, 39: 29–40.
- Van der Veer, H. W., and Witte, J. IJ. 1993. The “maximum growth/optimal food condition” hypothesis: a test for 0-group plaice *Pleuronectes platessa* L. in the Dutch Wadden Sea. *Marine Ecology Progress Series*, 101: 81–90.
- Van der Veer, H. W., and Witte, J. IJ. 1999. Year-class strength of plaice *Pleuronectes platessa* L. in the Southern Bight of

- the North Sea: a validation and analysis of the inverse relationship with winter seawater temperature. *Marine Ecology Progress Series*, 184: 245–257.
- Walsh, S. J. 1994. Recruitment variability in populations of lough rough dab (American plaice) *Hippoglossoides platessoides* (Fabricius) in the North Atlantic. *Netherlands Journal of Sea Research*, 32: 421–431.
- Werner, F. E., Ouinlan, J. A., Blanton, B. O., and Luettich, R. A., Jr 1997. The role of hydrodynamics in explaining variability in fish populations. *Journal of Sea Research*, 37: 195–212.
- Woodhead, P. M. J. 1964. Changes in the behaviour of the sole, *Solea vulgaris*, during cold winters and the relation between winter catch and sea temperature. *Helgoländer wissenschaftliche Meeresuntersuchungen*, 10: 328–340.
- Zijlstra, J. J. 1972. On the importance of the Waddensea as a nursery area in relation to the conservation of the southern North Sea fishery resources. *Symposia of the Zoological Society of London*, 29: 233–258.
- Zijlstra, J. J., Dapper, R., and Witte, J. IJ. 1982. Settlement, growth and mortality of post-larval plaice (*Pleuronectes platessa* L.) in the western Wadden Sea. *Netherlands Journal of Sea Research*, 15: 250–272.
- Zijlstra, J. J., and Witte, J. IJ. 1985. On the recruitment of 0-group plaice in the North Sea. *Netherlands Journal of Zoology*, 35: 360–376.