

Molecular Pathways: Estrogen Pathway in Colorectal Cancer

Afsaneh Barzi¹, Annika Medea Lenz¹, Melissa J. Labonte², and Heinz-Josef Lenz¹

Abstract

Worldwide, colorectal cancer has a higher incidence rate in men than in women, suggesting a protective role for sex hormones in the development of the disease. Preclinical data support a role for estrogen and its receptors in the initiation and progression of colorectal cancer and establishes that protective effects of estrogen are exerted through ER β . Hormone replacement therapy (HRT) in postmenopausal women as well as consumption of soy reduces the incidence of colorectal cancer. In the Women's Health Initiative trial, use of HRT in postmenopausal women reduced the risk of colon cancer by 56% [95% confidence interval (CI), 0.38–0.81; $P = 0.003$]. A recent meta-analysis showed that in women, consumption of soy reduced the risk of colon cancer by 21% (95% CI, 0.03–0.35; $P = 0.026$). In this review, using the preclinical data, we translate the findings in the clinical trials and observational studies to define the role of estrogen in the prevention of colorectal cancer. We hypothesize that sometime during the tumorigenesis process ER β expression in colonocytes is lost and the estrogen ligand, HRT, or soy products, exerts its effects through preventing this loss. Thus, in the adenoma-to-carcinoma continuum, timing of HRT is a significant determinant of the observed benefit from this intervention. We further argue that the protective effects of estrogen are limited to certain molecular subtypes. Successful development of estrogen modulators for prevention of colorectal cancer depends on identification of susceptible colorectal cancer population(s). Thus, research to better understand the estrogen pathway is fundamental for clinical delivery of these agents. *Clin Cancer Res*; 19(21); 5842–8. ©2013 AACR.

Background

Age is the strongest risk factor for colorectal cancer, with 90% of cases and 94% of deaths reported in individuals older than 50 years. Yet the incidence is lower in women across all age groups (1). This epidemiologic observation can be explained by differences in the colorectal cancer-related risk factors between men and women such as obesity, smoking, dietary exposures, and physical activity. However, persistence of the gender differences across several decades, in different racial groups and across the world, suggest that sex hormones (estrogen and progesterone) play a role in the pathogenic pathways of colorectal cancer and this role is most likely protective.

In addition, studies from relationship between gender and mortality of colorectal cancer consistently show lower mortality for women, especially premenopausal women. Epidemiologic studies over several decades suggest a decline in mortality of women attributed to use of hormone therapy (2, 3).

The protective role of the estrogen pathway in development of colon cancer has been studied and established in animal models. In several studies, exposure of ovariectomized rats to estrogen reduced the rate of colon tumors by 71% even in the presence of an ER α knockout (4, 5). In addition, in the Apc^{Min/+} mice ovariectomy resulted in an increased number of polyps, whereas replacement of estrogen in these mice reduced the number of polyps to the baseline values (6).

Over the past two decades use of hormone replacement therapy (HRT) provided the opportunity to discover the role of this intervention in prevention and outcome of colorectal cancer. Data from multiple prospective and retrospective cohorts support a protective role for HRT in the development and outcome of colon cancer.

Experience with soy products complements the hypothesis that estrogens have a protective effect against colorectal cancer. Lower incidence of colorectal cancer in Asians is attributed to high soy content. In animal models genistein, estrogen in soy, inhibits colon cancer cell proliferation and enhances apoptosis through interaction with several pathways (7, 8).

The estrogen pathway has the potential to be targeted for preventive and therapeutic strategies in colorectal cancer; therefore, its role and relationship with other pathways warrant further investigation. In this review, we provide comprehensive evidence on the role of estrogen and progesterone and how their signaling pathways play a critical role in colorectal cancer development and outcome.

Authors' Affiliations: ¹USC Norris Comprehensive Cancer Center, Los Angeles; and ²Azusa Pacific University, Azusa, California

Corresponding Author: Heinz-Josef Lenz, University of Southern California, Norris Comprehensive Cancer Center, 1441 Eastlake Avenue, Los Angeles, CA 90033. Phone: 1-323-865-3955/3967; Fax: 1-323-865-0061; E-mail: Lenz@med.usc.edu

doi: 10.1158/1078-0432.CCR-13-0325

©2013 American Association for Cancer Research.

Molecular effects of estrogen and progesterone in colorectal cancer

Estrogens are members of the steroid hormone family and are traditionally associated with female reproductive development. The most abundant and most potent estrogen in humans is 17 β -estradiol (E2). Other natural estrogens include estrone (E1) and estriol (E3), metabolites of E2. The estrogen receptor (ER) recognizes a molecule as estrogen based on its three-dimensional configuration and charge, but preferentially binds E2 over E3 (2 \times) and E1 [3 \times (ref. 9)]. Estrogen production and metabolism are tissue specific and are different in colon compared with breast tissue. In the colon, 17 β HSD2 and 4 convert E2 to E1, which is anti-proliferative in human colon cancer cell lines (10, 11).

The discovery of different ERs has revealed their significant role in tissue types other than the female reproductive tract, including the gastrointestinal system (12). Estrogens regulate these cellular effects through their intracellular receptors, ER α and ER β . The two receptors are coded for by separate genes, *ESR1* (ER α) and *ESR2* (ER β), with each gene producing different receptor isoforms from alternative splicing, resulting in three ER α and five ER β variants. The two receptors share DNA homology, coreceptors, and downstream signaling, however; their dissimilarities dictate the differences observed in their effects (13). Estrogen interacts differently with ER α and ER β downstream pathways (14). The differences in the downstream signaling of these two receptors may explain their distinctive intracellular effects. Furthermore, it has been shown that the biologic response of activated ER α and ER β is dependent on the ratio of ER α to ER β in the cell and that ER β inhibits transcriptional activity of ER α (15).

The intracellular effects of estrogen are exerted through two main pathways: genomic and nongenomic.

Genomic effects of estrogen in colorectal cancer. An activated ER activates gene transcription through either direct interaction with specific DNA sequences known as estrogen response elements (ERE) or other transcription factors such as c-Jun and/or c-Fos, resulting in transcription (Fig. 1). While interaction of both ERs with ERE is similar, ER α interacts with other transcription factors such as c-Jun and c-Fos of the activating protein-1 complex (AP1) and SP1, whereas ER β 's interaction with these transcription factors is less significant (13). In the case of AP1, it has been shown that binding of E2 to ER α results in transcription activation and that binding to ER β inhibits transcription through diversion of estrogen away from the ER α pathway (16, 17).

Genomic effects can also be seen in a ligand-independent manner, where ERs are phosphorylated through an activated kinase pathway in the absence of the estrogen ligand (Fig. 1). As an example, the EGF receptor (EGFR) can activate the Ras/Raf/MAPK pathway, which in turn will phosphorylate the ER, resulting in dimerization and ligand-independent activation of gene expression (18, 19).

The genomic-mediated effects of estrogen result in activated transcription of a wide array of genes including genes involved in angiogenesis, such as VEGF, cellular adhesion,

such as *cadherins* and *laminins*, and proliferation and apoptosis, including the TGF β family.

Nongenomic effects of estrogen in colorectal cancer. In addition to genomic effects, estrogen and its receptors can activate several signaling pathways without direct interaction with DNA, resulting in modulation of additional cellular processes. Transmembrane ERs have been shown to activate diverse intracellular pathways, including protein kinase C (PKC; ref. 20), intracellular Ca²⁺ (21, 22), cytosolic cAMP (23), nitric oxide (24), and MAPK (ref. 25; Fig. 1). For example, transmembrane ER α signaling through PI3K can result in cell proliferation and survival, whereas transmembrane ER β signaling causes the influx of Ca²⁺ in the cell, resulting in suppression of PKC signaling (Fig. 1). An additional role of ER β is through modulation of cell-cycle control through interactions with c-Myc, cyclin D1 (26), and cyclin A (27) resulting in inhibition of cell-cycle progression.

In both normal and cancerous colonocytes, ER α expression levels remain low. In contrast, ER β is the predominant ER in the normal colon (28, 29) with higher expression level in the ascending colon (30). The expression level of ER β in tumor tissue compared with normal colon mucosa is decreased and correlates with stage of the disease (31, 32).

Hartman and colleagues showed that transfection of SW480 cell lines with ER β resulted in inhibition of proliferation and cell-cycle arrest. SW480 xenografts with ER β expression had 70% reduction in the tumor weight (33). Furthermore, knockout of ER β in Apc^{Min/+} mice resulted in a dramatic increase in the polyps, and addition of E2 treatment did not prevent development of polyps in these mice (34).

Edvardsson and colleagues had shown that transfection of colon cancer cell lines with ER β affects the MAPK signaling pathway (35). Furthermore, ER β results in down-regulation of interleukin-6 and thus in reducing inflammation (35). Giroux and colleagues postulate that the effects of ER β in Apc^{Min/+} mice are through modulation of the TGF β signaling pathway (34).

Several studies support the cross-talk between Wnt/ β -catenin signaling and ER α . In one experiment, ER α transfection of SW480 and HCT116 resulted in activation of Wnt signaling and addition of an ER antagonist resulted in deactivation of the pathway. Moreover, in this experiment adding an antibody against β -catenin resulted in activation of ERE in an estrogen-dependent manner (36).

Despite the established cellular role for the ERs in colorectal cancer, the exact role of estrogen in colorectal cancer remains unclear. Effects of estrogen on colonocytes may be related to the direct effects of the estrogen ligand and its interaction with the receptors (directly related to ligand) or changes in the ER ratio and function independent of the ligand (indirectly related to ligand). For example, the rise in the incidence of colon cancer after menopause in women, when the circulating amount of estrogen ligand is decreased, could be related to the loss of cell-cycle regulatory effects of ER β (directly related to ligand), or that the decrease in the endogenous estrogen level at the time of




Figure 1. Intracellular effects of estrogen are exerted through two main pathways, genomic and nongenomic. Interaction of the ER and ligand with DNA, through ERE or other transcription factors, such as C-Fos/C-Jun/AP1, results in transcriptional alteration of genes and is the main genomic mechanism of estrogen in cells. The genomic effects can also be achieved from interactions of the receptor, without the ligand, through ligand-independent pathways, where ERs are phosphorylated through the activated kinase pathway. The nongenomic effects are achieved through several intracellular pathways, including PKC, intracellular Ca^{++} , cytosolic CAMP, nitric oxide, and MAPK.

menopause results in a change in the ratio of ER (indirectly related to ligand). In fact, in animal models the transcriptional activity of ERs changes over time and is influenced by estrogen level (37). In APC mouse models, E2 treatment results in an increase in the ratio of ER β and protection against colon cancer (6). Therefore, it is likely that HRT in women protects against colon cancer through an increased ratio of ER β . As ER β expression decreases throughout the process of tumorigenesis the timing of HRT to prevent loss of ER β is critical (38).

As for estrogen metabolism, polymorphism in the metabolic pathway of estrogen may result in a higher level of E2 and a lower level of E1 in colonocytes, thereby stimulating growth. The expression level of 17 β -HSD2 and 17 β -HSD4 is

lower in colon cancer than in normal colon tissue, resulting in lower levels of E1 in the tumor. The imbalance in the level of the estrogen metabolites can result in an increase in proliferation or a decrease in apoptosis (10, 11, 39). In a recent study, the level of E2 was higher in a cohort of Chinese men with colon cancer compared with controls. In this population combination of genetic variations in the ESR2 and E2 level was predictive of the risk of colon cancer, whereas in the group with high E2 level and CT/TT genotypes of ESR2 the risk of colon cancer was 2.3 (95% CI, 1.4–3.9) compared with the risk in those with a low level of E2 and CC genotype (40).

The literature on soy supports that use of isoflavones, a strong ER β agonist, in the DLD-1 human colon cancer cell

line, resulted in G₂ cell-cycle arrest. Use of soy isoflavones after silencing of ER β in this cell line failed to suppress G₂ arrest (41). Pintova and colleagues had shown that genistein reduces proliferation in HT29, DLD1, and RKO cell lines through inhibition of the Wnt/ β -catenin pathway (42). While these findings are exciting, results of a meta-analysis of the relationship between use of soy and risk of colorectal cancer in humans suggests that the protective effects of soy are limited to women (43), thus suggesting that effects of soy depend on the endogenous estrogen level through altering estrogen metabolism (44). Furthermore, use of soy isoflavones in rats resulted in increased expression of ER β and decreased expression of ER α in osteocytes; exposure of colonocytes to soy may also result in increased expression of ER β .

In addition to estrogen, the steroid hormone progesterone has been shown to play a distinct role in the regulation of the normal female reproductive function and regulating the bone matrix, but further has antiestrogen effects in other tissue types (45). Cellular effects of progesterone are mediated through a similar pathway to that of the ER, in which progestins bind either progesterone receptor (PR) A or B and mediate transcription through interaction with progesterone response elements. The only prominent non-genomic effect of progesterone that is reported is a rapid and transient activation of the MAPK pathway (46). The observed cellular effects of progesterone are largely attributed to its ability to oppose the action of estrogen, causing an inhibition of ER expression and through abrogating the induction of estrogen-related genes (47). Cellular effects of progesterone in colonocytes have been minimally studied. In mammary cells, interception of the PR by progesterone results in upregulation of EGFR and Wnt-1 (48–50). In animal breast cancer models, combination of estrogen and progesterone results in increased amphirgulin and augmented proliferation compared with estrogen alone. Assuming that EGFR and Wnt-1 can be upregulated in

colonocytes, endogenous progesterone may have a proliferative effect on the colonocytes. These findings are, presumably, fitting the interactions of progesterone with ER α , and interactions between ER β and progesterone are unexplored. Medroxyprogesterone, the progesterone formulation in HRT, has a direct inhibitory effect on colon cancer cell lines, supporting the observation that HRT is protective against colon cancer (50).

Epidemiologic data

The results of the Women's Health Initiative (WHI) study raised the awareness of the scientific community about the interactions between sex steroids and incidence and outcome of colon cancer. One limitation of the WHI study is that colon cancer was not the primary endpoint of the study and therefore the number of colorectal cancers in the study was small (122 during the trial and 263 during and after follow-up), making the data hypothesis generating. Hartz and colleagues published an analysis inclusive of all WHI enrollees and its observational cohort and found that use of any form of HRT reduces the risk of colon cancer by 30% [95% confidence interval (CI), 0.62–0.80; $P < 0.01$] and the risk of rectal cancer by 43% ($P < 0.0001$; ref. 51).

As the publication of WHI results, several observational studies have examined the role of HRT in the incidence and outcome of colon cancer. Table 1 summarizes the data from a few selected observational studies. The majority of these studies suggest that use of HRT reduces the incidence of colon cancer (52–55), but, this finding is not confirmed in other studies (52, 53, 56, 57). It is important to point out that these studies are planned and analyzed over several decades, with different formulations of hormones being used by participants. There is a large variation in the timing of start of these hormones after menopause and duration of hormonal use. It is unknown whether they are used to alleviate symptoms of menopause or for other reasons.

Table 1. Selected observational studies assessing the interactions between HRT and incidence and mortality of colorectal cancers

Study (pub year; ref.)	Risk	Product	Mean age	Colorectal cancer	HR (95% CI; P)
French E3N prospective cohort (2012; ref. 52)	I	E	51.1	525	0.72 (0.56–0.94) [S]
French E3N prospective cohort (2012; ref. 52)	I	E + P	56		1 (0.83–1.21)
Cancer Prevention Study II Nutrition Cohort (2009; ref. 53)	I	E	61&66	776	0.76 (0.59–0.97; 0.01)
Cancer Prevention Study II Nutrition Cohort (2009; ref. 53)	I	E + P	57&59		0.84 (0.54–1.30; 0.72)
UK General Practice Research Database (GPRD; 2007; ref. 56)	I	E	60.5		1.18 (0.72–1.92; 0.77)
UK General Practice Research Database (GPRD; 2007; ref. 54)	I	E + P	60.5		0.56 (0.35–0.87)
European Prospective Investigation into Cancer and Nutrition (57)	I	HRT	49.7	1,186	1 (0.86–1.16; 0.06)
Nested case control (2012; ref. 55)	I	HRT	73.8	4,708	0.81 (0.73–0.91; 0.001)
Cancer Prevention Study II (1995; ref. 58)	M	HRT	65.8	897	0.71 (0.63–0.81; 0.001)
Utah, California, and Minnesota Cancer Registry Data (59)	M	HRT			0.4
Seattle HMO database (60)	M	HRT		699	0.41
Nurses Health Study (61)	M	HRT	63	834	0.64

Abbreviations: E, estrogen; I, incidence; M, mortality; P, progesterone; [S], significant.

More importantly, due to the observational nature of these studies, compliance with the treatment and indication for treatment is unknown. Furthermore, in all these studies, prior use of hormonal therapy (postmenopausal or contraceptives) may have resulted in a difference in the observed outcome. Although, statistical methods are used to adjust for these variations, one should not ignore the potential role of bias in the results given the multitude of underlying factors.

In addition, four studies evaluated the role of hormones in mortality from colon cancer in patients who were diagnosed with colon cancer. All of these studies reported that HRT reduces the risk of death from colon cancer (Table 1; refs. 58–61). However, Newcomb and colleagues evaluated the risk of death in patients with colon cancer who were users of HRT and found that HRT does not reduce the risk of death from colon cancer (HR, 1.09; 95% CI, 0.81–1.47; ref. 62). These results further support a role for estrogen in reducing the mortality of colon cancer.

Clinical-Translational Advances

Although the estrogen pathway is not the central pathway in colorectal cancer, it plays an important role in the initiation and progression of the disease. Despite expression of ER β in normal colonocytes, none of the existing colon cancer cell lines express a sufficient amount of ER β , thus suggesting that loss of expression of ER β is part of the tumorigenesis process. This review establishes that the ERs play a major role in the fate of the colonocytes and ligands maintain the receptors' balance.

Data from the WHI and observational cohorts collectively support that HRT, estrogen surrogate, reduces the risk of developing colon cancer. Nevertheless, the degree of risk reduction differs among individuals, likely based on the activated pathways leading to colorectal cancer. In the WHI study colon cancer in the hormone replacement group seemed to be more advanced and had a higher number of positive lymph nodes, suggesting that tumors developing while on HRT seem to be biologically more aggressive (63). Interestingly, in the observational studies, obesity and smoking were associated with lesser degree of benefit from HRT. These phenotypic features are linked to aberrant activation of the pathways that are independent from the estrogen pathway and thus HRT may be less effective in preventing the tumors driven by these ER-independent pathways. For example, obesity results in hyperinsulinemia, which in turn results in PI3K activation and increased risk of colorectal cancer (64). As shown in Fig. 1, there is no relationship between PI3K and ER β (Fig. 1), and therefore estrogen may not protect against PI3K-driven tumors. In addition, smoking is linked to increased risk of MSI-high tumors and HRT does not decrease the risk of MSI-high tumors, which emphasizes that HRT may be effective in molecularly defined populations (65, 66).

In the WHI study, women older than 60 years on HRT had a lesser degree of colorectal cancer risk reduction, suggesting that timing of start of HRT (shorter interval from menopause) may play a crucial role in its protective effects on

colon cancer. On the basis of the existing preclinical evidence, early start of HRT prevents changes in the ratio of the receptors and results in a lower rate of colorectal cancer through maintaining ER β receptors. Furthermore, it is likely that start of estrogen before activation of the aberrant pathways will prevent colorectal cancer, while if the precancerous process is already initiated and ER β is lost, estrogens will not be able to reverse the process.

Besides HRT, high estrogen content of soy is implicated to be protective against colorectal cancer. However, recent evidence suggests that higher soy consumption is protective against colon cancer only in women (43).

To maximize the benefit of estrogen for prevention of colorectal cancer, exploring the phenotypic and molecular subgroups of the individuals who achieve maximum benefit from this intervention is essential. Until then, use of estrogens for prevention of colon cancer should remain investigational.

Translational Relevance

Use of estrogens for prevention of colon cancer is an attractive concept in women; however, the increased rates of cardiovascular events with HRT limit use of these agents in clinical practice.

While there are no clinical data to prove that ER β modulation reduces the incidence of colon cancer, preclinical models of ER β modulation are promising. In the study by Schleipen and colleagues, treatment of ovariectomized rats with ER β agonists or genistein increased the apoptosis and reduced proliferation in the ileum and colon, further supporting the role of these agents in the prevention of colon cancer (67). Also, Giroux and colleagues showed that ER β agonist treatment reduces the number of small-intestinal polyps in Apc^{Min/+} male and female mice (68). Although development of selective ER β modulators has proven to be very challenging, two agents are currently in phase III and IV trials. MF101 and raloxifen are being studied for their effects on menopausal symptoms and osteoporosis, respectively (69, 70). Long-term follow-up will show if any of the agents in clinical development have an effect on the incidence of colon cancer in women. The challenge is that benefit from these agents happens primarily in the presence of ER β ; thus, secondary prevention in patients who express ER β is the most logical first step to test the clinical efficacy of these agents.

We conclude by stating that although estrogen modulation has a role in colorectal cancer prevention, the burden is on investigators to identify the appropriate population and product to move the field forward.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: A. Barzi, A.M. Lenz, M.J. Labonte, H.-J. Lenz

Development of methodology: A.M. Lenz, H.-J. Lenz

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): A. Barzi, A.M. Lenz, H.-J. Lenz

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): A.M. Lenz, H.-J. Lenz

Writing, review, and/or revision of the manuscript: A. Barzi, A.M. Lenz, M.J. Labonte, H.-J. Lenz

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): A.M. Lenz, M.J. Labonte, H.-J. Lenz
Study supervision: H.-J. Lenz

Grant Support

The work was supported by NIH support, P30, and Wunderglo Foundation.

Received February 4, 2013; revised August 2, 2013; accepted August 8, 2013; published OnlineFirst August 21, 2013.

References

- Siegel R, Jemal A. Colorectal cancer facts & figures. American Cancer Society; 2011.
- Hendifar A, Yang D, Lenz F, Lurje G, Pohl A, Lenz C, et al. Gender disparities in metastatic colorectal cancer survival. *Clin Cancer Res* 2009;15:6391-7.
- Fernandez E, Bosetti C, La Vecchia C, Levi F, Fioretti F, Negri E. Sex differences in colorectal cancer mortality in Europe, 1955-1996. *Eur J Cancer Prev* 2000;9:99-104.
- Guo JY, Li X, Browning JD Jr, Rottinghaus GE, Lubahn DB, Constantinou A, et al. Dietary soy isoflavones and estrone protect ovariectomized ERalphaKO and wild-type mice from carcinogen-induced colon cancer. *J Nutr* 2004;134:179-82.
- Smirnov P, Liel Y, Gnainky J, Shany S, Schwartz B. The protective effect of estrogen against chemically induced murine colon carcinogenesis is associated with decreased CpG island methylation and increased mRNA and protein expression of the colonic vitamin D receptor. *Oncol Res* 1999;11:255-64.
- Weyant MJ, Carothers AM, Mahmoud NN, Bradlow HL, Remotti H, Bilinski RT, et al. Reciprocal expression of ERalpha and ERbeta is associated with estrogen-mediated modulation of intestinal tumorigenesis. *Cancer Res* 2001;61:2547-51.
- Kim EJ, Shin HK, Park JH. Genistein inhibits insulin-like growth factor-I receptor signaling in HT-29 human colon cancer cells: a possible mechanism of the growth inhibitory effect of Genistein. *J Med Food* 2005;8:431-8.
- Qi W, Weber CR, Wasland K, Savkovic SD. Genistein inhibits proliferation of colon cancer cells by attenuating a negative effect of epidermal growth factor on tumor suppressor FOXO3 activity. *BMC Cancer* 2011;11:219.
- Ruh TS, Katzenellenbogen BS, Katzenellenbogen JA, Gorski J. Estrone interaction with the rat uterus: *in vitro* response and nuclear uptake. *Endocrinology* 1973;92:125-34.
- English MA, Kane KF, Cruickshank N, Langman MJ, Stewart PM, Hewison M. Loss of estrogen inactivation in colonic cancer. *J Clin Endocrinol Metab* 1999;84:2080-5.
- English MA, Stewart PM, Hewison M. Estrogen metabolism and malignancy: analysis of the expression and function of 17beta-hydroxysteroid dehydrogenases in colonic cancer. *Mol Cell Endocrinol* 2001;171:53-60.
- Gustafsson JA. Estrogen receptor beta—a new dimension in estrogen mechanism of action. *J Endocrinol* 1999;163:379-83.
- Deroo BJ, Buensuceso AV. Minireview: estrogen receptor-beta: mechanistic insights from recent studies. *Mol Endocrinol* 2010;24:1703-14.
- Nilsson S, Makela S, Treuter E, Tujague M, Thomsen J, Andersson G, et al. Mechanisms of estrogen action. *Physiol Rev* 2001;81:1535-65.
- Hall JM, McDonnell DP. The estrogen receptor beta-isoform (ERbeta) of the human estrogen receptor modulates ERalpha transcriptional activity and is a key regulator of the cellular response to estrogens and antiestrogens. *Endocrinology* 1999;140:5566-78.
- Foley EF, Jazaeri AA, Shupnik MA, Jazaeri O, Rice LW. Selective loss of estrogen receptor beta in malignant human colon. *Cancer Res* 2000;60:245-8.
- Paech K, Webb P, Kuiper GG, Nilsson S, Gustafsson J, Kushner PJ, et al. Differential ligand activation of estrogen receptors ERalpha and ERbeta at AP1 sites. *Science* 1997;277:1508-10.
- Driggers PH, Segars JH. Estrogen action and cytoplasmic signaling pathways. Part II: the role of growth factors and phosphorylation in estrogen signaling. *Trends Endocrinol Metab* 2002;13:422-7.
- Martin LA, Farmer I, Johnston SR, Ali S, Dowsett M. Elevated ERK1/ERK2/estrogen receptor cross-talk enhances estrogen-mediated signaling during long-term estrogen deprivation. *Endocr Relat Cancer* 2005;12 Suppl 1:S75-84.
- Alzamora R, Harvey BJ. Direct binding and activation of protein kinase C isoforms by steroid hormones. *Steroids* 2008;73:885-8.
- Zhang L, Blackman BE, Schonemann MD, Zogovic-Kapsalis T, Pan X, Tagliaferri M, et al. Estrogen receptor beta-selective agonists stimulate calcium oscillations in human and mouse embryonic stem cell-derived neurons. *PLoS ONE* 2010;5:e11791.
- Lamprecht SA, Lipkin M. Chemoprevention of colon cancer by calcium, vitamin D, and folate: molecular mechanisms. *Nat Rev Cancer* 2003;3:601-14.
- Aronica SM, Kraus WL, Katzenellenbogen BS. Estrogen action via the cAMP signaling pathway: stimulation of adenylate cyclase and cAMP-regulated gene transcription. *Proc Natl Acad Sci U S A* 1994;91:8517-21.
- Kim HP, Lee JY, Jeong JK, Bae SW, Lee HK, Jo I. Nongenomic stimulation of nitric oxide release by estrogen is mediated by estrogen receptor alpha localized in caveolae. *Biochem Biophys Res Commun* 1999;263:257-62.
- Endoh H, Sasaki H, Maruyama K, Takeyama K, Waga I, Shimizu T, et al. Rapid activation of MAP kinase by estrogen in the bone cell line. *Biochem Biophys Res Commun* 1997;235:99-102.
- Liu MM, Albanese C, Anderson CM, Hilty K, Webb P, Uht RM, et al. Opposing action of estrogen receptors alpha and beta on cyclin D1 gene expression. *J Biol Chem* 2002;277:24353-60.
- Treeck O, Latrich C, Springwald A, Ortmann O. Estrogen receptor beta exerts growth-inhibitory effects on human mammary epithelial cells. *Breast Cancer Res Treat* 2010;120:557-65.
- Fiorelli G, Picariello L, Martinetti V, Tonelli F, Brandi ML. Functional estrogen receptor beta in colon cancer cells. *Biochem Biophys Res Commun* 1999;261:521-7.
- Arai N, Strom A, Rafter JJ, Gustafsson JA. Estrogen receptor beta mRNA in colon cancer cells: growth effects of estrogen and genistein. *Biochem Biophys Res Commun* 2000;270:425-31.
- Papaxoinis K, Triantafyllou K, Sasco AJ, Nicolopoulou-Stamati P, Ladas SD. Subsite-specific differences of estrogen receptor beta expression in the normal colonic epithelium: implications for carcinogenesis and colorectal cancer epidemiology. *Eur J Gastroenterol Hepatol* 2010;22:614-9.
- Castiglione F, Taddei A, Rossi Degl'Innocenti D, Buccoliero AM, Bechi P, Garbini F, et al. Expression of estrogen receptor beta in colon cancer progression. *Diagn Mol Pathol* 2008;17:231-6.
- Jassam N, Bell SM, Speirs V, Quirke P. Loss of expression of oestrogen receptor beta in colon cancer and its association with Dukes' staging. *Oncol Rep* 2005;14:17-21.
- Hartman J, Edvardsson K, Lindberg K, Zhao C, Williams C, Strom A, et al. Tumor repressive functions of estrogen receptor beta in SW480 colon cancer cells. *Cancer Res* 2009;69:6100-6.
- Giroux V, Lemay F, Bernatchez G, Robitaille Y, Carrier JC. Estrogen receptor beta deficiency enhances small intestinal tumorigenesis in ApcMin/+ mice. *Int J Cancer* 2008;123:303-11.
- Edvardsson K, Strom A, Jonsson P, Gustafsson JA, Williams C. Estrogen receptor beta induces antiinflammatory and antitumorigenic networks in colon cancer cells. *Mol Endocrinol* 2011;25:969-79.
- Kouzmenko AP, Takeyama K, Ito S, Furutani T, Sawatsubashi S, Maki A, et al. Wnt/beta-catenin and estrogen signaling converge *in vivo*. *J Biol Chem* 2004;279:40255-8.

37. Della Torre S, Biserni A, Rando G, Monteleone G, Ciana P, Komm B, et al. The conundrum of estrogen receptor oscillatory activity in the search for an appropriate hormone replacement therapy. *Endocrinology* 2011;152:2256–65.
38. Konstantinopoulos PA, Kominea A, Vadoros G, Sykietis GP, Andriopoulos P, Varakis I, et al. Oestrogen receptor beta (ERbeta) is abundantly expressed in normal colonic mucosa, but declines in colon adenocarcinoma paralleling the tumour's dedifferentiation. *Eur J Cancer* 2003;39:1251–8.
39. Di Domenico M, Castoria G, Bilancio A, Migliaccio A, Auricchio F. Estradiol activation of human colon carcinoma-derived Caco-2 cell growth. *Cancer Res* 1996;56:4516–21.
40. Wu H, Xu L, Chen J, Hu J, Yu S, Hu G, et al. Association of estrogen receptor beta variants and serum levels of estradiol with risk of colorectal cancer: a case control study. *BMC Cancer* 2012;12:276.
41. Bielecki A, Roberts J, Mehta R, Raju J. Estrogen receptor-beta mediates the inhibition of DLD-1 human colon adenocarcinoma cells by soy isoflavones. *Nutr Cancer* 2011;63:139–50.
42. Pintova S, Planoutis K, Planutiene M, Holcombe RF. Genistein-induced reduction of Wnt/beta-catenin pathway activity in colorectal cancer cell lines correlates with inhibition of proliferation and colony growth in soft agar [abstract]. In: *Proceedings of the Annual Meeting of the American Association for Cancer Research*; 2013; Washington, DC. Philadelphia (PA): AACR; 2013.
43. Yan L, Spitznagel EL, Bosland MC. Soy consumption and colorectal cancer risk in humans: a meta-analysis. *Cancer Epidemiol Biomarkers Prev* 2010;19:148–58.
44. Xu X, Duncan AM, Wangen KE, Kurzer MS. Soy consumption alters endogenous estrogen metabolism in postmenopausal women. *Cancer Epidemiol Biomarkers Prev* 2000;9:781–6.
45. Graham JD, Clarke CL. Physiological action of progesterone in target tissues. *Endocr Rev* 1997;18:502–19.
46. Faivre E, Skildum A, Pierson-Mullany L, Lange CA. Integration of progesterone receptor mediated rapid signaling and nuclear actions in breast cancer cell models: role of mitogen-activated protein kinases and cell cycle regulators. *Steroids* 2005;70:418–26.
47. Takeda A, Leavitt WW. Progesterone-induced downregulation of nuclear estrogen receptor in uterine decidual cells: analysis of receptor synthesis and turnover by the density-shift method. *Biochem Biophys Res Commun* 1986;135:98–104.
48. Faivre EJ, Lange CA. Progesterone receptors upregulate Wnt-1 to induce epidermal growth factor receptor transactivation and c-Src-dependent sustained activation of Erk1/2 mitogen-activated protein kinase in breast cancer cells. *Mol Cell Biol* 2007;27:466–80.
49. Peterson CM. Estrogen and progesterone receptors: an overview from the year 2000. *J Soc Gynecol Invest* 2000;7:S3–7.
50. Tanaka Y, Kato K, Mibu R, Uchida S, Asanoma K, Hashimoto K, et al. Medroxyprogesterone acetate inhibits proliferation of colon cancer cell lines by modulating cell cycle-related protein expression. *Meno-pause* 2008;15:442–53.
51. Hartz A, He T, Ross JJ. Risk factors for colon cancer in 150,912 postmenopausal women. *Cancer Causes Control* 2012;23:1599–605.
52. Morois S, Fournier A, Clavel-Chapelon F, Mesrine S, Boutron-Ruault MC. Menopausal hormone therapy and risks of colorectal adenomas and cancers in the French E3N prospective cohort: true associations or bias? *Eur J Epidemiol* 2012;27:439–52.
53. Hildebrand JS, Jacobs EJ, Campbell PT, McCullough ML, Teras LR, Thun MJ, et al. Colorectal cancer incidence and postmenopausal hormone use by type, recency, and duration in cancer prevention study II. *Cancer Epidemiol Biomarkers Prev* 2009;18:2835–41.
54. Tannen RL, Weiner MG, Xie D, Barnhart K. A simulation using data from a primary care practice database closely replicated the Women's Health Initiative trial. *J Clin Epidemiol* 2007;60:686–95.
55. Green J, Czanner G, Reeves G, Watson J, Wise L, Roddam A, et al. Menopausal hormone therapy and risk of gastrointestinal cancer: nested case-control study within a prospective cohort, and meta-analysis. *Int J Cancer* 2012;130:2387–96.
56. Tannen RL, Weiner MG, Xie D, Barnhart K. Estrogen affects postmenopausal women differently than estrogen plus progestin replacement therapy. *Hum Reprod* 2007;22:1769–77.
57. Tsilidis KK, Allen NE, Key TJ, Sanjoaquin MA, Bakken K, Berrino F, et al. Menopausal hormone therapy and risk of colorectal cancer in the European Prospective Investigation into Cancer and Nutrition. *Int J Cancer* 2011;128:1881–9.
58. Calle EE, Miracle-McMahill HL, Thun MJ, Heath CW Jr. Estrogen replacement therapy and risk of fatal colon cancer in a prospective cohort of postmenopausal women. *J Natl Cancer Inst* 1995;87:517–23.
59. Slattery ML, Anderson K, Samowitz W, Edwards SL, Curtin K, Caan B, et al. Hormone replacement therapy and improved survival among postmenopausal women diagnosed with colon cancer (USA). *Cancer Causes Control* 1999;10:467–73.
60. Mandelson MT, Miglioretti D, Newcomb PA, Harrison R, Potter JD. Hormone replacement therapy in relation to survival in women diagnosed with colon cancer. *Cancer Causes Control* 2003;14:979–84.
61. Chan JA, Meyerhardt JA, Chan AT, Giovannucci EL, Colditz GA, Fuchs CS. Hormone replacement therapy and survival after colorectal cancer diagnosis. *J Clin Oncol* 2006;24:5680–6.
62. Newcomb PA, Chia VM, Hampton JM, Doria-Rose VP, Trentham Dietz A. Hormone therapy in relation to survival from large bowel cancer. *Cancer Causes Control* 2009;20:409–16.
63. Chlebowski RT, Wactawski-Wende J, Ritenbaugh C, Hubbell FA, Ascensao J, Rodabough RJ, et al. Estrogen plus progestin and colorectal cancer in postmenopausal women. *N Engl J Med* 2004;350:991–1004.
64. Carling D. The AMP-activated protein kinase cascade—a unifying system for energy control. *Trends Biochem Sci* 2004;29:18–24.
65. Newcomb PA, Zheng Y, Chia VM, Morimoto LM, Doria-Rose VP, Templeton A, et al. Estrogen plus progestin use, microsatellite instability, and the risk of colorectal cancer in women. *Cancer Res* 2007;67:7534–9.
66. Limsui D, Vierkant RA, Tillmans LS, Wang AH, Weisenberger DJ, Laird PW, et al. Cigarette smoking and colorectal cancer risk by molecularly defined subtypes. *J Natl Cancer Inst* 2010;102:1012–22.
67. Schleipen B, Hertrampf T, Fritzsche KH, Kluxen FM, Lorenz A, Molzberger A, et al. ERbeta-specific agonists and genistein inhibit proliferation and induce apoptosis in the large and small intestine. *Carcinogenesis* 2011;32:1675–83.
68. Giroux V, Bernatchez G, Carrier JC. Chemopreventive effect of ERbeta-Selective agonist on intestinal tumorigenesis in Apc(Min/+) mice. *Mol Carcinog* 2011;50:359–69.
69. Grady D, Sawaya GF, Johnson KC, Koltun W, Hess R, Vittinghoff E, et al. MF101, a selective estrogen receptor beta modulator for the treatment of menopausal hot flashes: a phase II clinical trial. *Meno-pause* 2009;16:458–65.
70. Walsh JM, Cheung AM, Yang D, Litwack S, Grady D. Raloxifene and colorectal cancer. *J Womens Health (Larchmt)* 2005;14:299–305.