

Association of Inflammation-Related and microRNA Gene Expression with Cancer-Specific Mortality of Colon Adenocarcinoma

Aaron J. Schetter,¹ Giang Huong Nguyen,^{1,2} Elise D. Bowman,¹ Ewy A. Mathé,¹ Siu Tsan Yuen,³ Jason E. Hawkes,^{1,2} Carlo M. Croce,⁴ Suet Yi Leung,³ and Curtis C. Harris¹

Abstract **Purpose:** Inflammatory genes and microRNAs have roles in colon carcinogenesis; therefore, they may provide useful biomarkers for colon cancer. This study examines the potential clinical utility of an inflammatory gene expression signature as a prognostic biomarker for colon cancer in addition to previously examined *miR-21* expression. **Experimental Design:** Quantitative reverse transcriptase-PCR was used to measure the expression of 23 inflammatory genes in colon adenocarcinomas and adjacent noncancerous tissues from 196 patients. These data were used to develop models for cancer-specific mortality on a training cohort ($n = 57$), and this model was tested in both a test ($n = 56$) and a validation ($n = 83$) cohort. Expression data for *miR-21* were available for these patients and were compared and combined with inflammatory gene expression. **Results:** *PRG1*, *IL-10*, *CD68*, *IL-23a*, and *IL-12a* expression in noncancerous tissue, and *PRG1*, *ANXA1*, *IL-23a*, *IL-17a*, *FOXP3*, and *HLA-DRA* expression in tumor tissues were associated with poor prognosis based on Cox regression ($|Z\text{-score}| > 1.5$) and were used to generate the inflammatory risk score (IRS). IRS was associated with cancer-specific mortality in the training, test ($P = 0.01$), and validation ($P = 0.02$) cohorts. This association was strong for stage II cases ($P = 0.002$). Expression of *miR-21* was associated with *IL-6*, *IL-8*, *IL-10*, *IL-12a*, and *NOS2a*, providing evidence that the function of this microRNA and these inflammatory genes are linked. Both IRS and *miR-21* expression were independently associated with cancer-specific mortality, including stage II patients alone. **Conclusion:** IRS and *miR-21* expression are independent predictors of colon cancer prognosis and may provide a clinically useful tool to identify high-risk patients. (Clin Cancer Res 2009;15(18):5878–87)

Authors' Affiliations: ¹Laboratory of Human Carcinogenesis, Center for Cancer Research, National Cancer Institute, NIH, Bethesda, Maryland; ²Howard Hughes Medical Institute-NIH Research Scholar, Howard Hughes Medical Institute, Chevy Chase, Maryland; ³Department of Pathology, University of Hong Kong, Queen Mary Hospital, Pokfulam, Hong Kong; ⁴Department of Molecular Virology, Immunology, and Medical Genetics, Ohio State University Comprehensive Cancer Center, Columbus, Ohio

Received 3/12/09; revised 6/9/09; accepted 6/9/09; published OnlineFirst 9/8/09.

Grant support: Intramural Research Program of the National Cancer Institute, Center for Cancer Research, NIH. A.J. Schetter was supported by the Cancer Prevention Fellowship Program, Office of the Director, National Cancer Institute, NIH. G.H. Nguyen and J.E. Hawkes were supported by the Howard Hughes Medical Institute Grant for Graduate Medical Education.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Note: Supplementary data for this article are available at Clinical Cancer Research Online (<http://clincancerres.aacrjournals.org/>).

Requests for reprints: Curtis C. Harris, Laboratory of Human Carcinogenesis, Center for Cancer Research, NCI, NIH, 37 Convent Drive, Building 37, Room 2068A, Bethesda, MD 20892. Phone: 301-496-2048; Fax: 301-496-0497; E-mail: Curtis_Harris@nih.gov.

© 2009 American Association for Cancer Research.
doi:10.1158/1078-0432.CCR-09-0627

Colon adenocarcinoma is a leading cause of cancer mortality worldwide (1) and accounts for ~50,000 deaths annually in the United States (2). Although current adjuvant treatment modalities improve survival for tumor-node-metastasis (TNM) stage III colon cancer patients, whether stage II patients should be given these therapies remains controversial (3, 4). Some stage II patients will benefit from therapy, but therapy for others will harm quality of life with little therapeutic benefit. Therefore, it is important to develop biomarkers to identify high-risk, early-stage patients who may be suitable for therapeutic intervention.

Inflammation plays a key role in tumor initiation, progression, and metastasis (5, 6). Chronic inflammation is associated with increased rates of colon cancer for both ulcerative colitis and Crohn's disease (7–9). Nonsteroidal anti-inflammatory drugs can reduce colon cancer risk (10). Inflammation-modulating cytokines affect tumor development through roles in cell proliferation, angiogenesis, and apoptosis (11). Cytokines can signal changes directly within the tumor or the tumor microenvironment to influence cancer progression (12). Because inflammation contributes to colon carcinogenesis, expression of inflammatory genes may serve as biomarkers for colon cancer.

Translational Relevance

We report that the expression pattern of inflammatory-related genes in tumors and paired noncancerous tissues was an independent prognostic marker for colon adenocarcinoma patients. This gene signature was associated with prognosis in early-stage patients. Therefore, this gene signature may be useful for identifying high-risk, early-stage patients to assist in decisions regarding appropriate therapeutic intervention. We also show that combining independent biomarkers can improve predictions over single biomarkers. The combination of the inflammatory gene signature with available *miR-21* expression data improved predictions with prognosis over either alone. These findings show the potential of inflammatory risk score and/or *miR-21* as prognostic biomarkers for early-stage colon cancer.

Infiltration of inflammatory cells in colorectal cancer has been associated with prognosis (13–15). Polymorphisms in inflammatory genes have been associated with colon cancer incidence and prognosis (16–18). Expression of inflammatory genes has also been associated with TNM staging and prognosis in colon cancer (19, 20). Previous studies identified unique expression signatures composed of a panel of inflammatory/immune-response genes that predict metastatic progression and survival of hepatocellular carcinoma (21) and lung adenocarcinoma (22) patients. Building on these findings in hepatocellular carcinoma and lung adenocarcinoma, we determined if expression of these inflammatory genes in tumors and the surrounding noncancerous tissue can be used as a prognostic biomarker for colon adenocarcinoma.

Combining multiple, independent prognostic biomarkers may improve the ability to identify cancer patients at high risk of disease progression and mortality. Therefore, adding an additional factor to an inflammatory gene biomarker may provide a more clinically useful biomarker than either alone. MicroRNA expression may serve this purpose. MicroRNAs are small, non-coding RNA molecules that have shown potential as biomarkers in cancer (23–26). Expression levels of microRNAs are altered in all cancers that have been studied. Alteration of specific microRNAs can alter tumor progression in mouse models (27), showing their potential to be causal factors in carcinogenesis. We recently reported that patients with tumors expressing high levels of an oncogenic microRNA, *miR-21*, have worse survival prognosis for stage II or stage III colon adenocarcinoma, showing its potential as a prognostic biomarker for colon cancer (28). The expression of *miR-21* has previously been linked to inflammatory responses. Expression of *miR-21* is increased following lipopolysaccharide-induced inflammation (29), and increased *miR-21* expression occurs during T-cell differentiation (30). Interleukin 6 (*IL-6*), a proinflammatory cytokine, can drive *miR-21* expression through a signal transducers and activators of transcription 3 (*STAT3*)-dependent mechanism (31). Because *miR-21* expression is linked to inflammation and both *miR-21* and inflammatory gene expression are linked to colon cancer, combining inflammatory gene biomarkers with *miR-21* may improve their clinical utility.

In this study, we set out to measure the expression of inflammatory genes in tumors and paired noncancerous tissue from 196 colon adenocarcinoma patients, and use these data to develop an inflammatory risk model that could be used as a prognostic biomarker for colon cancer. In addition, we used previously acquired data on *miR-21* to address two specific questions. First, does *miR-21* expression correlate with specific inflammatory genes as is predicted from mechanistic studies in cell culture? Second, does the combination of the inflammatory risk model with *miR-21* expression have improved associations with cancer-specific mortality over either alone?

Materials and Methods

Tissue collection and RNA isolation. Pairs of primary colon tumor and adjacent noncancerous tissues came from 83 patients recruited from the University of Maryland Medical Center or Baltimore Veterans Administration Medical Center from 1993 to 2002, and from 113 patients recruited from Queen Mary Hospital in Hong Kong from 1991 to 2000. These patients have been described in a previous study (28). Cases with familial adenomatous polyposis were excluded. Tissues were grossly dissected and flash-frozen after surgery, prior to any adjuvant therapy. Detailed backgrounds for each tissue donor, including age, gender, clinical staging, tumor location, and survival time from diagnosis were collected. Final dates of follow-up were December 31, 2005 and August 16, 2004 for the NCI-Maryland and the Hong Kong cohorts, respectively. Tumor histopathology was classified according to the WHO Classification of Tumor system (1). Informed consent was given by all participants. This study was approved by the Institutional Review Board of the NIH, the Institutional Review Board of the University of Hong Kong/Hospital Authority Hong Kong West Cluster, and the Institutional Review Board for Human Subject Research at the University of Maryland.

RNA isolation and quantitative reverse transcriptase-PCR of inflammatory genes. RNA from frozen tissue samples was extracted using standard TRIZOL (Invitrogen) methods. RNA was reverse-transcribed using the cDNA Archive Kit (Applied Biosystems) with a 50 ng/ μ L final concentration. Expression levels of inflammatory genes were measured with custom-designed, Taqman low-density-array real-time PCR plates (Applied Biosystems) containing probes to 23 inflammatory genes: Annexin A1 (*ANXA1*; assay ID Hs00167549_m1), colon stimulating factor 1 (*CSF1*; assay ID Hs00174164_m1), MHC class II antigen DR α (*HLA-DRA*; ID Hs00219575_m1), MHC class II antigen DP α 1 (*HLA-DPA1*; ID Hs00410276_m1), *IFN- γ* (ID Hs00174143_m1), interleukin 1 α (*IL-1A*; ID Hs00174092_m1), *IL-1B* (ID Hs00174097_m1), *IL-2* (ID Hs00174114_m1), *IL-4* (ID Hs00174122_m1), *IL-5* (ID Hs00174200_m1), *IL-6* (ID Hs00174131_m1), *IL-8* (ID Hs00174103_m1), *IL-10* (ID Hs00174086_m1), *IL-12A* (ID Hs00168405_m1), *IL-12B* (ID Hs00233688_m1), *IL-15* (ID Hs00542571_m1), *IL-17A* (ID Hs00174383_m1), *IL-23A* (ID Hs00372324_m1), proteoglycan 1 (*PRG1*; ID Hs00160444_m1), nitric oxide synthase 2A (*NOS2A*; ID Hs00167257_m1), forkhead box p3 (*FOXP3*; ID Hs00203958_m1), cluster of differentiation 68 (*CD68*; ID Hs00154355_m1), and tumor necrosis factor α (*TNF- α* ; ID Hs00174128_m1), with 18 s rRNA (ID Hs99999901_s1) as a normalization control. Expression of inflammatory genes was measured while blinded to all clinical outcomes. For quality control, any tissue that had 18 s threshold cycle values >15 were considered poor quality and were removed. A patient was removed from this study if either noncancerous or paired tumor tissues failed quality control.

Measurement of miR-21. In a previous study, microRNA expression levels were measured in all of these patient samples and is described in detail there (28). Briefly, microRNA expression levels in the NCI-Maryland cohort were measured using microRNA microarrays (Ohio State microRNA microarray version 2.0). For the Hong Kong cohort,

expression of *miR-21* was measured using quantitative reverse transcriptase-PCR (qRT-PCR) using Taqman microRNA assays (Applied Biosystems) according to manufacturer's instructions. High-expression cases for *miR-21* were defined based on highest tertile separately for the microarrays and qRT-PCR results.

Statistical analyses. Expression data were imported into Biometric Research Branch Array Tools v3.6.0, and the median was normalized for the Hong Kong cohort and Maryland cohort separately. Paired *t* tests identified differentially expressed genes between tumor and noncancerous tissue for the Hong Kong and the Maryland cohorts separately. To account for multiple comparisons, only differences that were found and validated in each cohort separately ($P < 0.05$) were considered significant. Graphpad Prism v5.0 (Graphpad Software Inc.) was used for correlation analysis.

The Hong Kong cohort was divided randomly into a training cohort and a test cohort to identify a gene expression model associated with cancer-specific mortality. The Maryland cohort was used as the validation cohort for this model. Prior to beginning the analysis, the Hong Kong cohort was selected to divide into a training cohort and a test cohort because it was the larger of the two cohorts and would likely result in a model with improved accuracy compared with the smaller Maryland cohort. Univariate Cox regression analysis on the training cohort was used to select genes associated with cancer-specific mortality ($|Z\text{-score}| > 1.5$; $P < 0.13$) to include in multivariate risk models using previously reported methods (32). All genes were included for these purposes, and expression values for all analyses are continuous variables. For multivariate Cox regression models, missing values for genes were replaced with the average values. In the training cohort, selected genes were used to build multivariate models for tumor and noncan-

cerous tissue separately. Coefficients from these models were multiplied with gene expression values and summed to build risk scores. Individuals were defined as high risk if they had higher than median risk scores for both tumor and noncancerous models. Kaplan-Meier analysis was done with WINSTAT 2007 (R Fitch Software, Bad Krozingen). Cox regression was carried out in Stata 9.2 (StataCorp LP).

Linear regression models identified associations between *miR-21* expression and inflammatory gene expression in noncancerous tissue, tumor tissue, and then a combined analysis of both tumor and noncancerous tissue adjusting for tumor status. Only the qRT-PCR data from the Hong Kong cohort were analyzed for these purposes because the microarray data from the Maryland cohort was considered less reliable. *IL-4*, *IL-5*, and *IL-12b* were excluded because they were missing data for >25% of the samples. The Bonferroni-Holm method (33) was used to adjust for multiple comparisons in the combined tumor and noncancerous regression models.

Results

Expression of inflammatory genes are systematically altered in colon adenocarcinoma. This study used two independent cohorts, one consisting of 113 cases recruited from Hong Kong and a second cohort of 83 cases recruited from Maryland (Table 1). The median follow-up times were 84.6 and 80 months for patients in the Hong Kong and the NCI-Maryland cohorts, respectively. The cohorts were similar in TNM staging ($P = 0.65$, Fisher's exact) and cancer-specific mortality

Table 1. Characteristics of study populations and tumors

	Hong Kong cohort		Maryland cohort
	Training (n = 57)	Test (n = 56)	Validation (n = 83)
Age at enrollment (y)			
Mean (SD)	57.8 (15.5)	53.8 (14.1)	64.4 (10.6)
Range	32-84	30-82	32-87
Gender, no. (%)			
Male	30 (53)	26 (46)	65 (78)
Female	27 (47)	30 (54)	18 (22)
Tumor location,* no. (%)			
Distal	41 (72)	49 (87.5)	48 (59)
Proximal	16 (28)	7 (12.5)	33 (41)
Adenocarcinoma histology, no. (%)			
Adenocarcinoma	53 (93)	52 (93)	74 (89)
Mucinous adenocarcinoma	4 (7)	3 (5)	8 (10)
Adenosquamous	0	0	1 (1)
Signet ring cell and mucinous	0	1 (2)	0
Adjuvant chemotherapy, [†] no. (%)			
Received	22 (39)	18 (32)	22 (38)
Did not receive	35 (61)	38 (57)	36 (62)
TNM staging, [‡] No. (%)			
I	2 (4)	7 (13)	8 (10)
II	19 (33)	18 (32)	29 (35)
III	27 (47)	21 (38)	36 (43)
IV	9 (16)	10 (18)	9 (11)
Removed during quality control [§]	0	0	10 (12)

*Distal includes tumors located in or distal to the descending colon. Proximal tumors include tumors in or proximal to the splenic flexure. Tumor location was available for all cases in the Hong Kong cohort and 81 cases in the Maryland cohort.

[†]Detailed information pertaining to receipt of chemotherapy was available for all patients in the Hong Kong cohort and 58 in the Maryland cohort. Chemotherapy was primarily fluorouracil-based (in forms of either i.v. fluorouracil or oral drugs including tegafur with uracil) with or without levamisole or leucovorin.

[‡]For one patient in the Maryland cohort, it was unclear if that patient had stage III or stage IV colon cancer, therefore this patient was removed from analyses stratifying by TNM stage.

[§]Cases with poor quality data from quantitative RT-PCR for either the tumor or nontumorous tissue were removed.




Fig. 1. Inflammatory genes are consistently altered in colon tumors on both cohorts. *A*, unsupervised hierarchical clustering (correlation, average linkage) using 21 inflammatory genes on 113 pairs of cancerous and noncancerous tissues in the Hong Kong cohort. Clustering separates tissues into two distinct groups: one composed of 97% tumor tissues and one composed of 99% noncancerous tissue. *B*, correlation of the tumor/noncancerous tissue expression ratio comparing the Hong Kong cohort with the NCI-Maryland cohort indicates consistent changes in inflammatory gene expression in both cohorts.

($P = 0.46$, Kaplan-Meier log rank) with 5-year survival rates of 49.5% (Hong Kong cohort) and 59.7% (NCI-Maryland cohort). Along with the racial, cultural, and geographic differences of the two cohorts, the Maryland cohort was considerably older, with a higher percentage of men.

We measured the expression of 23 inflammatory genes in primary colon tumor and paired noncancerous tissues using low-density-array real-time PCR. Eighteen of these genes were selected because they were included in previous studies of hepatocellular carcinoma (21) and lung adenocarcinoma (22). The additional five genes (*IL-17A*, *IL-23A*, *CD68*, *NOS2A*, and *FOXP3*) were selected based on literature supporting their roles in colonic inflammation or cancer (34–36). *IL-4* and *IL-5* were not detectable in the majority of tissues and were removed from all further analyses.

Inflammatory gene expression was systematically altered in tumors. Expression of the 21 inflammatory genes could distinguish tumor from noncancerous tissue pairs with 99% or 100% accuracy based on nearest centroid or 3-nearest neighbors class prediction algorithms, respectively (10-fold cross-validation repeated 100 times) using the Hong Kong cohort. Unsupervised hierarchical clustering of the 21 genes separated tissues into two distinct groups: one composed of 97% tumor tissue and

the other composed of 99% noncancerous tissue (Fig. 1A). Of the 21 inflammatory genes examined, the expression of 18 was altered in tumors in the Hong Kong cohort ($P < 0.05$; paired *t*-test; Supplementary Table S1). Of these, *IL-8* showed the largest fold-increase in tumors at ~13-fold higher levels in tumors whereas *IL-2* showed the largest reduction in tumor with ~80% less in tumors. These results indicate a systematic change in the expression of inflammatory genes during tumorigenesis.

We next analyzed the NCI-Maryland cohort. Fold changes in tumors for these inflammatory genes were consistent with the Hong Kong cohort (Pearson $R = 0.96$; Fig. 1B), indicating that these changes in gene expression are likely representative of the majority of colon adenocarcinomas. Expression of *IL-8*, *IL-23a*, *IL-1a*, *IL-1b*, *FOXP3*, *IL-17a*, *IFN-γ*, and *IL-6* was significantly increased in tumors from both cohorts, whereas expression of *IL-2*, *IL-15*, *IL-10*, *IL-12a*, *CSF1*, *HLA-DPA1*, *HLA-DRA*, *TNF-a*, and *CD68* was significantly decreased in tumors from both cohorts.

Colon adenomas represent an early, precancerous lesion of the colon. Changes in inflammatory gene expression in adenoma tissues may indicate early changes in the inflammatory state that can lead to cancer. We evaluated the expression levels of the 23 inflammatory-related genes in 18 pairs of colon adenomas and nonadenoma tissues. Although there was limited




Fig. 2. Strategy for building inflammatory risk scores. Genes were selected for inclusion in the risk score based on univariate Cox regression on the Hong Kong training cohort. Multivariate Cox regression on the training cohort was used to build the risk models. This model was then tested on the Hong Kong test and Maryland validation cohorts.

power to detect differences in expression of these genes due to using a limited number of tissues, we found similar changes in gene expression in adenoma as compared with the colon cancer tissues (Supplementary Table S1). When using the Hong Kong cohort as a reference, expression changes in colon adenomas were consistent with colon cancer tissues for these inflammatory genes (Pearson $R = 0.91$; $P < 0.0001$). Of the 10 genes significantly decreased in tumors, all showed decreased expression in adenomas and seven of these (*IL-2*, *IL-10*, *IL-12a*, *CSF1*, *HLA-*

DPA1, *HLA-DRA*, and *PRG1*) were significantly reduced. Of the eight genes significantly increased in tumors, all eight were increased in adenomas and six of these (*IL-8*, *IL-23a*, *IL-1a*, *IL-1b*, *FOXP3*, and *IL-17a*) were significantly increased. Similar to colon cancer tissues, *IL-8* showed the greatest increase and *IL-2* showed the greatest decrease in adenoma tissues.

Inflammatory risk score is associated with cancer-specific mortality. We evaluated the expression of these inflammatory genes for associations with cancer-specific mortality. Constructing a

multi-gene signature using several genes with moderate associations can provide more accurate predictions than a model using a single gene. Therefore, we used univariate Cox regression to identify genes with moderate associations with prognosis following previously established methodologies (32). We randomly split the Hong Kong cohort into a training cohort ($n = 57$) and a test cohort ($n = 56$; Fig. 2). These cohorts were similar in clinical characteristics, including age at enrollment, gender, and TNM staging. Based on univariate Cox regression on the training cohort, expression of *PRG1*, *IL-10*, *CD68*, *IL-23a*, and *IL-12a* in noncancerous tissue, and *PRG1*, *ANXA1*, *IL-23a*, *IL-17a*, *FOXP3*, and *HLA-DRA* in tumors was moderately associated with cancer-specific mortality ($|Z\text{-score}| > 1.5$; using criteria from ref. 32; Supplementary

Fig. S1). These genes were selected to construct a multigene risk signature. Using the training cohort, multivariate Cox regression was done on selected genes to develop risk models. The non-cancerous risk model was $[(0.855 \times PRG1) + (0.720 \times IL-10) + (0.458 \times CD68) + (-0.494 \times IL-23a) + (-0.635 \times IL-12a)] = \text{risk score}$. The tumor risk model was $[(1.321 \times PRG1) + (0.840 \times ANXA1) + (0.123 \times IL-23a) + (0.484 \times IL-17a) + (0.367 \times FOXP3) + (-0.373 \times HLA-DRA)] = \text{risk score}$. Individuals having higher than median values for both models were classified as having high inflammatory risk score (IRS). All others were considered low. When evaluated separately, patients classified as high IRS had significantly worse cancer-specific mortality for the Hong Kong training cohort, the Hong Kong test cohort ($P = 0.01$,




Fig. 3. IRS is associated with cancer-specific mortality in TNM stage II patients. A, IRS in Hong Kong training, Hong Kong test, and NCI-Maryland validation cohorts separately. B, combined analysis of Hong Kong test and NCI-Maryland validation cohorts, stratified by TNM stage. For one patient in the Maryland Cohort, it was unclear if that patient had stage III or stage IV colon cancer and therefore was removed from analyses stratifying by TNM stage.




Fig. 4. Combined IRS and *miR-21* expression predicts colon cancer-specific mortality better than either alone. All patients from the Hong Kong test and NCI-Maryland validation cohorts are analyzed. Left, stratified by *miR-21* expression; middle, stratified by IRS and *miR-21* expression; right, combined IRS and *miR-21* expression. *miR-21* expression data were available for 115 of 117 patients and only those patients were used for this analysis. The Hong Kong training cohort was also excluded from this analysis.

Kaplan-Meier log rank), and NCI-Maryland validation cohort ($P = 0.02$, Kaplan-Meier log rank; Fig. 3A).

To evaluate the potential use of IRS as a biomarker, we did a stratified analysis by TNM staging. For these analyses, the Hong Kong training cohort was excluded to prevent overfitting. The Hong Kong test and NCI-Maryland validation cohorts were combined. IRS was associated with TNM stage ($P = 0.03$, Fisher's exact test). Patients with more advanced TNM stage were more likely to be classified as high IRS. Four of 14 (29%) stage I, 9 of 42 (21%) stage II, 16 of 46 (35%) stage III, and 9 of 14 (64%) stage IV cases were classified as high IRS. High IRS was associated with poor cancer-specific mortality for all patients ($P = 0.0003$, Kaplan-Meier log rank; Fig. 3B). When stratified by TNM stage, IRS was associated with cancer-specific mortality in stage II cases ($P = 0.002$, Kaplan-Meier log rank; Fig. 3B). IRS was not associated with prognosis in stage I, stage III, or stage IV patients.

We were unable to analyze associations with therapeutic outcome. Of the 34 stage II patients for whom we had information about receipt of adjuvant therapy, only 7 received it. Only one of these seven patients was classified as high IRS. Therefore, we did not have sufficient power to analyze associations between IRS and therapeutic outcome.

miR-21 expression is associated with the expression of IL-6, IL-8, IL-10, IL-12a, and NOS2a. Expression of *miR-21* was available for these samples from a previous study (28). *miR-21* has previously been shown to be associated with inflammation. Therefore, we used linear regression to examine the associations of *miR-21* expression with inflammatory genes in these tissues. This was evaluated in noncancerous tissues, tumor tissues, and then a combination of all tissues adjusting for tumor status (Supplementary Table S2). In the combined model, expression of *IL-6* ($P < 0.0005$), *IL-8* ($P < 0.0005$), and *IL-10* ($P = 0.002$) was positively associated, and *IL-12a* ($P < 0.0005$) and *NOS2a* ($P = 0.0025$) expression was negatively associated with *miR-21* expression. Only *IL-6* and *IL-12a* expression was

statistically significant ($P < 0.05$) in both the tumor and non-cancerous tissues, separately.

IRS and miR-21 expression are independently associated with cancer-specific mortality, including stage II patients. We previously reported that high *miR-21* expression in tumors was associated with poor prognosis in colon adenocarcinoma (28). That study utilized the same patients as the current study and provides an opportunity to combine *miR-21* and IRS to determine if together they have improved prognostic utility. High *miR-21* expression was defined in our previous publication where the highest tertile (>3.3 -fold higher than average noncancerous tissue) is defined as high (28). Survival information for the NCI-Maryland cohort was updated from that study to include an additional 12 months of available survival information. Consistent with our previous report, high *miR-21* expression is associated with cancer-specific mortality using all cases ($P < 0.0001$, Kaplan-Meier log rank) or stage II cases ($P = 0.006$, Kaplan-Meier log rank; Fig. 4). Due to the association between *IL-6* and *miR-21* expression, we investigated if combining *IL-6* and *miR-21* expression data into survival models would alter the association between *miR-21* and cancer-specific mortality, and found that it did not (data not shown).

Although IRS and *miR-21* expression were each associated with prognosis, they were not associated with one another ($P = 0.83$, Fisher's exact). Therefore, combination of these biomarkers may identify high-risk patients that would be misclassified by a single end point. We did a stratified analysis of *miR-21* and IRS (Fig. 3). Patients with low *miR-21* expression and low IRS had the best prognosis. Patients with high *miR-21*/low IRS or low *miR-21*/high IRS had an intermediate prognosis. Patients with high *miR-21*/high IRS had the worst prognosis. This was true when observing all cases or stage II cases alone. Upon combining intermediate groups, patients classified as high for either *miR-21* or IRS score had significantly worse cancer-specific mortality than those classified as low *miR-21*/low IRS for all cases ($P = 0.0002$, Kaplan-Meier

log rank) or stage II cases ($P = 0.002$, Kaplan-Meier log rank). Patients classified as high for both *miR-21* and IRS had worse survival than patients classified as high for either using all cases ($P = 0.002$, Kaplan-Meier log rank) or stage II alone ($P = 0.02$, Kaplan-Meier log rank). Only two stage II patients were classified as high IRS and high *miR-21*. Therefore, one should be cautious interpreting the poor outcome of these stage II patients and future studies will explore this association.

Univariate Cox regression analysis for all cases found that high IRS [hazard ratio (HR), 2.4; 95% confidence interval (95% CI), 1.4-4.2], high *miR-21* (HR, 3.0; 95% CI, 1.7-5.1), and TNM staging (HR, 4.7; 95% CI, 2.5-8.8) were each associated with poor prognosis (Table 2). Multivariate analyses showed that both high IRS (HR, 2.2; 95% CI, 1.3-3.8) and high *miR-21* (HR, 3; 95% CI, 1.7-5.2) were independent of one another and TNM staging. Additionally, the multivariate model including IRS, TNM staging, and *miR-21* did significantly better than the model without *miR-21* ($P < 0.001$, likelihood ratio test). When restricting the analysis to stage II cases, univariate analyses showed that high IRS (HR, 5.4; 95% CI, 1.7-17.2) and high *miR-21* (HR, 4.8; 95% CI, 1.4-16.1) were each associated with poor prognosis. Multivariate analysis showed that high IRS (HR, 7.5; 95% CI, 2.2-25.6) and high *miR-21* (HR, 6.5; 95% CI, 1.9-21.9) were each associated with prognosis independent of one another. A multivariate model including both IRS and *miR-21* in stage II patients did significantly better than a model including only IRS ($P = 0.004$, likelihood ratio test). Therefore, IRS and *miR-21* expression may be used together as a prognostic biomarker for stage II colon adenocarcinoma.

Discussion

We found systematic changes in inflammatory gene expression in colon tumors. Of the eight inflammatory genes consistently increased in tumors, seven (*IL-8*, *IL-23a*, *IL-1a*, *IL-1b*, *IL-17a*, *INF γ* , and *IL-6*) are proinflammatory cytokines and the other is *FOXP3*, a marker for regulatory T cells. These results are consistent with other reports evaluating their expression in colon cancer (37). Therefore, there are predictable changes in inflammatory gene expression in colon tumors, consistent for a role for these genes in carcinogenesis. We found similar changes in gene expression in colon adenomas. This indicates that changes in the inflammatory state may be an early event in colon carcinogenesis.

Expression of inflammatory genes was associated with *miR-21* expression. The association of *IL-6* and *IL-12a* expression was statistically significant in both the tumor and noncancerous tissues, separately. *IL-6* is thought to drive the expression of *miR-21* in a *STAT3*-dependent mechanism (31). Our results are consistent with that model and provide evidence that this mechanism may be relevant to colon cancer. There is also a predicted binding site for *miR-21* in the 3' untranslated region of *IL-12a* as indicated by Targetscan 5.0 (38) and miRanda (39). *IL-12a* has a negative correlation with *miR-21*, which is consistent with a pattern for a *miR-21* target. Based on this finding, mechanistic studies should be done to determine if *IL-12a* is a target of *miR-21*. The interaction between *miR-21* and inflammatory genes may play an important role in colon carcinogenesis. Although the associations between *miR-21*, *IL-6*, and *IL-12a* were significant, the regression models indicated that much of the variability was

Table 2. Cox regression of inflammatory risk score and *miR-21* expression with cancer-specific mortality on combined Hong Kong test cohort and Maryland validation cohort

Variable (comparison/referent)	All cases, regardless of TNM stage			
	Univariate analysis*		Multivariate analysis [†]	
	HR (95% CI)	P	HR (95% CI)	P
IRS (high/low)	2.4 (1.4-4.2)	0.001	2.2 (1.3-3.8)	0.005
<i>miR-21</i> expression (high/low)	3.0 (1.7-5.1)	<0.0005	3.0 (1.7-5.2)	<0.0005
Tumor stage (III-IV/I-II)	4.7 (2.5-8.8)	<0.0005	4.0 (2.1-7.5)	<0.0005
Age in y (50/<50)	1.1 (0.6-2.1)	0.82		
Gender (male/female)	1.9 (1.0-3.5)	0.06		
Tumor location (proximal/distal)	0.8-(0.4-1.7)	0.58		
Variable (comparison/referent)	Stage II cases, adjusted for cohort membership			
	Univariate analysis*		Multivariate analysis [†]	
	HR (95% CI)	P	HR (95% CI)	P
IRS (high/low)	5.4 (1.7-17.2)	0.005	7.5 (2.2-25.6)	0.001
<i>miR-21</i> expression (high/low)	4.8 (1.4-16.1)	0.01	6.5 (1.9-21.9)	0.002
Age (50/<50)	2.9 (0.4-24.2)	0.31		
Gender (male/female)	1.4 (0.4-4.7)	0.57		
Tumor location (proximal/distal)	0.4 (0.1-1.7)	0.20		

*Univariate analysis is adjusted for cohort membership only.

[†]Multivariate analysis is adjusted for cohort membership, IRS, *miR-21* expression, and (where appropriate) TNM stage. Multivariate analysis used stepwise addition and removal of clinical covariates found to be associated with survival in univariate models ($P < 0.10$) and final models include only those covariates that were significantly associated with survival (Wald statistic, $P < 0.05$). *miR-21* measurements were available for 115 of 117 patients, including all 42 stage II patients, and only those patients are included in multivariate analyses.

explained by these models. This indicates that other mechanisms for gene regulation contribute to the expression of these genes. For example, the *miR-21* promoter contains putative binding sites for the transcription factors AP-1, Ets/PU.1, SRF, TP53, C/EBP α , and STAT3 (40), and *miR-21* expression can also be influenced by epidermal growth factor receptor activity (41). Therefore, it is likely that the expression of *miR-21* is influenced by many of these and other factors in the context of cancer.

The expression of inflammatory genes is altered in colon adenocarcinoma. Although this study does not address the causal relationship between changes in inflammatory gene expression and carcinogenesis, these changes are consistent with published work showing a potential causal relationship between altered expression of inflammatory genes and carcinogenesis. For example, *IL-8* showed the highest fold-increase in both colon tumors and adenomas in our study. Previous studies have shown *IL-8* to be a proinflammatory chemokine that is expressed at elevated levels in tumors (42). *IL-8* expression has been shown to enhance cell proliferation, cell survival, and angiogenesis through induction of the multiple signaling pathways. Conversely, *IL-2* showed the largest reduction in tumors and adenomas. *IL-2* expression had been found to inhibit tumor growth *in vivo*, and high dose *IL-2* therapy has shown some promise to reduce tumor burden in patients (43) exhibiting a causal role between aberrant expression of *IL-2* and cancer.

Expression of inflammatory genes in tumors and the surrounding noncancerous tissues is associated with prognosis in colon adenocarcinoma. This cooperation of tumor and noncancerous expression of inflammatory genes was observed in our previous investigation of lung adenocarcinoma (22). Higher expression of *IL-10* in noncancerous tissues was associated with worse survival in that study and the current study. *IL-10* is an anti-inflammatory cytokine that can suppress cell-mediated immunity (44). Therefore, elevated *IL-10* in noncancerous tissue may create an inflammatory environment primed for metastasis and disease progression.

High levels of *IL-23a* and *IL-12a* in the noncancerous tissue were associated with improved survival. Both are members of the *IL-12* family of proinflammatory cytokines (45). *IL-12* activity is important for host resistance to tumors (46), therefore high levels of *IL-12* in the tumor macroenvironment may lead to resistance of tumor progression and metastasis through induction of IFN- γ and activation of natural killer cells and cytotoxic T cells. In contrast, elevated levels of *IL-23a* and *IL-17a* in cancerous tissues were associated with worse survival and may promote a microenvironment that suppresses any host antitumor response. *IL-23a* can stimulate Th17 cells to increase the production of *IL-17a*, and overexpression of *IL-17a* in cervical cancer (47), non-small cell lung cancer (48), or fibrosarcoma (49) cell lines increases tumor formation and/or tumor growth in xenograft mouse models. These cytokines are associated with a Th17 response. Therefore, a Th17 response in tumors may create a favorable condition for tumor progression.

There are limitations to the current study. First, patients with mucinous or adenosquamous histologies were excluded from this study; therefore IRS may not be applicable to these patients. Additionally, the IRS was built using relatively few cases ($n = 53$). Developing molecular signatures of these genes on larger cohorts may strengthen the accuracy and precision of this biomarker, although the validation of this biomarker in two independent cohorts shows its potential clinical utility. It will be

important to begin exploring the relationship between IRS and other clinical covariates, such as microsatellite instability status, p53 mutational status, or *K-ras* mutational status, to determine if a combination of these markers can provide more clinically useful information. It will also be useful to do immunohistochemistry on patient samples to determine localization patterns of these inflammatory genes to gain insights into the cell types responsible for this gene signature. This will provide mechanistic insights as to how the combination of these genes may contribute to the worse prognosis in high-IRS patients.

We found the association between IRS and survival to be strongest in TNM stage II patients. The reason for this association in stage II patients and not stage III/IV patients is unclear. One possibility is that high IRS is associated with a favorable inflammatory environment for metastasis. This is consistent in that we see more advanced-stage patients are more likely to be classified as high IRS. In this context, it may be a useful marker only in stage II patients, for which metastasis is not yet detectable by current clinical methods. In stage III or IV patients, metastasis is evident and therefore an IRS score to predict metastasis is not meaningful.

Cancer immunotherapy is a promising field of research for colon cancer (50). As in any therapy, successful stratification of patients into groups that are more or less likely to respond will increase the chances of developing successful immunotherapies. IRS is based on the expression of inflammatory genes, and the expression of these genes is likely to be correlated with the current state of the immune system. It is possible that IRS may be associated with a patient's response to immunotherapy. Although future investigation of this is needed, there is a potential that IRS, or a similar inflammatory gene biomarker, may be able to identify patients more or less likely to respond to immunotherapy.

There is a need for better ways of diagnosing early-stage colon cancer patients with undetectable micrometastases. Therefore, we propose that a subset of stage II patients would benefit from therapeutic intervention as their disease will likely progress; but for others, therapeutic intervention unnecessarily harms quality of life and continued screening would be sufficient. We found IRS was significantly associated with prognosis in stage II patients. Previously, we identified *miR-21* as a prognostic biomarker for stage II patients (28). The combination of IRS and *miR-21* expression was a better predictor of prognosis than either alone. Therefore, IRS and *miR-21*, alone or in combination, have potential to help diagnose stage II patients and assist in choosing treatment options. Prospective studies to evaluate this potential are warranted.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

We thank Drs. Howard Young and Giorgio Trinchieri for critical reading of the manuscript and thoughtful suggestions, and Drs. Xin Wang and Stefan Ambs (National Cancer Institute) for invaluable discussion. We are grateful to Raymond Jones, Audrey Salabes, and John Cottrell of the University of Maryland Medical Center and the Surgery and Pathology Departments at UMD Hospital, Baltimore Veterans Administration Medical Center, Sinai Hospital, Union Memorial Hospital, St. Agnes Hospital, Northwest Hospital Center, and Mercy Medical Center for contributions to this study.

References

1. Aaltonen LA, Hamilton SR, World Health Organization, International Agency for Research on Cancer. Pathology and genetics of tumours of the digestive system. Lyon Oxford: IARC Press; Oxford University Press (distributor); 2000.
2. Jemal A, Siegel R, Ward E, et al. Cancer statistics, 2008. *CA Cancer J Clin* 2008;58:71–96.
3. Chua YJ, Zalcborg JR. Progress and challenges in the adjuvant treatment of stage II and III colon cancers. *Expert Rev Anticancer Ther* 2008;8:595–604.
4. Wolpin BM, Mayer RJ. Systemic treatment of colorectal cancer. *Gastroenterology* 2008;134:1296–310.
5. Hussain SP, Harris CC. Inflammation and cancer: an ancient link with novel potentials. *Int J Cancer* 2007;121:2373–80.
6. DeNardo DG, Johansson M, Coussens LM. Immune cells as mediators of solid tumor metastasis. *Cancer Metastasis Rev* 2008;27:11–8.
7. Ekblom A, Helmick C, Zack M, Adami HO. Increased risk of large-bowel cancer in Crohn's disease with colonic involvement. *Lancet* 1990;336:357–9.
8. Ekblom A, Helmick C, Zack M, Adami HO. Ulcerative colitis and colorectal cancer. A population-based study. *N Engl J Med* 1990;323:1228–33.
9. Bernstein CN, Blanchard JF, Kliwler E, Wajda A. Cancer risk in patients with inflammatory bowel disease: a population-based study. *Cancer* 2001;91:854–62.
10. Cha YI, DuBois RN. NSAIDs and cancer prevention: targets downstream of COX-2. *Annu Rev Med* 2007;58:239–52.
11. Lin WW, Karin M. A cytokine-mediated link between innate immunity, inflammation, and cancer. *J Clin Invest* 2007;117:1175–83.
12. Mantovani A, Romero P, Palucka AK, Marincola FM. Tumour immunity: effector response to tumour and role of the microenvironment. *Lancet* 2008;371:771–83.
13. Jass JR, Love SB, Northover JM. A new prognostic classification of rectal cancer. *Lancet* 1987;1:1303–6.
14. Galon J, Costes A, Sanchez-Cabo F, et al. Type, density, and location of immune cells within human colorectal tumors predict clinical outcome. *Science* 2006;313:1960–4.
15. Pages F, Berger A, Camus M, et al. Effector memory T cells, early metastasis, and survival in colorectal cancer. *N Engl J Med* 2005;353:2654–66.
16. Landi S, Bottari F, Gemignani F, et al. Interleukin-4 and interleukin-4 receptor polymorphisms and colorectal cancer risk. *Eur J Cancer* 2007;43:762–8.
17. Lurje G, Zhang W, Schultheis AM, et al. Polymorphisms in VEGF and IL-8 predict tumor recurrence in stage III colon cancer. *Ann Oncol* 2008.
18. Wilkening S, Tavelin B, Canzian F, et al. Interleukin promoter polymorphisms and prognosis in colorectal cancer. *Carcinogenesis* 2008;29:1202–6.
19. Pages F, Berger A, Henglein B, et al. Modulation of interleukin-18 expression in human colon carcinoma: consequences for tumor immune surveillance. *Int J Cancer* 1999;84:326–30.
20. Berghella AM, Contasta I, Pellegrini P, Del Beato T, Adorno D. Are immunological mechanisms involved in colon cancer and are they possible markers for biotherapy improvement? *Cancer Biother Radiopharm* 2006;21:468–87.
21. Budhu A, Forgues M, Ye QH, et al. Prediction of venous metastases, recurrence, and prognosis in hepatocellular carcinoma based on a unique immune response signature of the liver microenvironment. *Cancer Cell* 2006;10:99–111.
22. Seike M, Yanaiharu N, Bowman ED, et al. Use of a cytokine gene expression signature in lung adenocarcinoma and the surrounding tissue as a prognostic classifier. *J Natl Cancer Inst* 2007;99:1257–69.
23. Calin GA, Ferracin M, Cimmino A, et al. A MicroRNA signature associated with prognosis and progression in chronic lymphocytic leukemia. *N Engl J Med* 2005;353:1793–801.
24. Johnson SM, Grosshans H, Shingara J, et al. RAS is regulated by the let-7 microRNA family. *Cell* 2005;120:635–47.
25. Volinia S, Calin GA, Liu CG, et al. A microRNA expression signature of human solid tumors defines cancer gene targets. *Proc Natl Acad Sci U S A* 2006;103:2257–61.
26. Yanaiharu N, Caplen N, Bowman E, et al. Unique microRNA molecular profiles in lung cancer diagnosis and prognosis. *Cancer Cell* 2006;9:189–98.
27. He L, Thomson JM, Hemann MT, et al. A microRNA polycistron as a potential human oncogene. *Nature* 2005;435:828–33.
28. Schetter AJ, Leung SY, Sohn JJ, et al. MicroRNA expression profiles associated with prognosis and therapeutic outcome in colon adenocarcinoma. *JAMA* 2008;299:425–36.
29. Moschos SA, Williams AE, Perry MM, Birrell MA, Belvisi MG, Lindsay MA. Expression profiling *in vivo* demonstrates rapid changes in lung microRNA levels following lipopolysaccharide-induced inflammation but not in the anti-inflammatory action of glucocorticoids. *BMC Genomics* 2007;8:240.
30. Wu H, Neilson JR, Kumar P, et al. miRNA profiling of naive, effector and memory CD8 T cells. *PLoS ONE* 2007;2:e1020.
31. Loffler D, Brocke-Heidrich K, Pfeifer G, et al. Interleukin-6 dependent survival of multiple myeloma cells involves the Stat3-mediated induction of microRNA-21 through a highly conserved enhancer. *Blood* 2007;110:1330–3.
32. Lossos IS, Czerwinski DK, Alizadeh AA, et al. Prediction of survival in diffuse large-B-cell lymphoma based on the expression of six genes. *N Engl J Med* 2004;350:1828–37.
33. Holm S. A simple sequentially rejective multiple test procedure. *Scand Journal of Statist* 1979;6:65–70.
34. Hata K, Andoh A, Shimada M, et al. IL-17 stimulates inflammatory responses via NF- κ B and MAP kinase pathways in human colonic myofibroblasts. *Am J Physiol Gastrointest Liver Physiol* 2002;282:G1035–44.
35. Lo CH, Lee SC, Wu PY, et al. Antitumor and antimetastatic activity of IL-23. *J Immunol* 2003;171:600–7.
36. Hofseth LJ, Saito S, Hussain SP, et al. Nitric oxide-induced cellular stress and p53 activation in chronic inflammation. *Proc Natl Acad Sci U S A* 2003;100:143–8.
37. Le Gouvello S, Bastuji-Garin S, Aloulou N, et al. High prevalence of Foxp3 and IL17 in MMR-proficient colorectal carcinomas. *Gut* 2008;57:772–9.
38. Friedmann RC, Farh KK, Burge CB, Bartel DP. Most mammalian mRNAs are conserved targets of microRNAs. *Genome Res* 2009;19:92–105.
39. Betel D, Wilson M, Gabow A, Marks DS, Sander C. The microRNA.org resource: targets and expression. *Nucleic Acids Res* 2008;36:D149–53.
40. Fujita S, Ito T, Mizutani T, et al. miR-21 Gene expression triggered by AP-1 is sustained through a double-negative feedback mechanism. *J Mol Biol* 2008;378:492–504.
41. Seike M, Goto A, Okano T, et al. MiR-21 is an EGFR-regulated anti-apoptotic factor in lung cancer from never-smokers. *Proc Natl Acad Sci U S A* 2009;106:12085–90.
42. Waugh DJ, Wilson C. The interleukin-8 pathway in cancer. *Clin Cancer Res* 2008;14:6735–41.
43. McDermott DF. The application of high-dose interleukin-2 for metastatic renal cell carcinoma. *Med Oncol* 2009;26 Suppl 1:13–7.
44. O'Garra A, Barrat FJ, Castro AG, Vicari A, Hawrylowicz C. Strategies for use of IL-10 or its antagonists in human disease. *Immunol Rev* 2008;223:114–31.
45. Oppmann B, Lesley R, Blom B, et al. Novel p19 protein engages IL-12p40 to form a cytokine, IL-23, with biological activities similar as well as distinct from IL-12. *Immunity* 2000;13:715–25.
46. Trinchieri G. Interleukin-12 and the regulation of innate resistance and adaptive immunity. *Nat Rev Immunol* 2003;3:133–46.
47. Tartour E, Fossiez F, Joyeux I, et al. Interleukin 17, a T-cell-derived cytokine, promotes tumorigenicity of human cervical tumors in nude mice. *Cancer Res* 1999;59:3698–704.
48. Numasaki M, Watanabe M, Suzuki T, et al. IL-17 enhances the net angiogenic activity and *in vivo* growth of human non-small cell lung cancer in SCID mice through promoting CXCR-2-dependent angiogenesis. *J Immunol* 2005;175:6177–89.
49. Numasaki M, Fukushi J, Ono M, et al. Interleukin-17 promotes angiogenesis and tumor growth. *Blood* 2003;101:2620–7.
50. Weiner LM. Cancer immunotherapy—the endgame begins. *N Engl J Med* 2008;358:2664–5.