

## The Novel Akt Inhibitor API-1 Induces c-FLIP Degradation and Synergizes with TRAIL to Augment Apoptosis Independent of Akt Inhibition

Bo Li<sup>1,2</sup>, Hui Ren<sup>1,3</sup>, Ping Yue<sup>1</sup>, Mingwei Chen<sup>3</sup>, Fadlo R. Khuri<sup>1</sup>, and Shi-Yong Sun<sup>1</sup>

### Abstract

API-1 (pyrido[2,3-*d*]pyrimidines) is a novel small-molecule inhibitor of Akt, which acts by binding to Akt and preventing its membrane translocation and has promising preclinical antitumor activity. In this study, we reveal a novel function of API-1 in regulation of cellular FLICE-inhibitory protein (c-FLIP) levels and TRAIL-induced apoptosis, independent of Akt inhibition. API-1 effectively induced apoptosis in tested cancer cell lines including activation of caspase-8 and caspase-9. It reduced the levels of c-FLIP without increasing the expression of death receptor 4 (DR4) or DR5. Accordingly, it synergized with TRAIL to induce apoptosis. Enforced expression of ectopic c-FLIP did not attenuate API-1-induced apoptosis but inhibited its ability to enhance TRAIL-induced apoptosis. These data indicate that downregulation of c-FLIP mediates enhancement of TRAIL-induced apoptosis by API-1 but is not sufficient for API-1-induced apoptosis. API-1-induced reduction of c-FLIP could be blocked by the proteasome inhibitor MG132. Moreover, API-1 increased c-FLIP ubiquitination and decreased c-FLIP stability. These data together suggest that API-1 downregulates c-FLIP by facilitating its ubiquitination and proteasome-mediated degradation. Because other Akt inhibitors including API-2 and MK2206 had minimal effects on reducing c-FLIP and enhancement of TRAIL-induced apoptosis, it is likely that API-1 reduces c-FLIP and enhances TRAIL-induced apoptosis independent of its Akt-inhibitory activity. *Cancer Prev Res*; 5(4); 612–20. ©2012 AACR.

### Introduction

API-1 (pyrido[2,3-*d*]pyrimidines) is a recently identified small-molecule inhibitor of Akt, which acts through binding to Akt and blocking its membrane translocation (1). A previous study has shown that API-1 possesses promising anticancer activity, evidenced by its ability to suppress cell growth, induce apoptosis, and inhibit the growth of cancer xenografts, particularly those with activated Akt, in nude mice (1).

TRAIL (also called APO-2L) is a member of the TNF family and is currently being tested in phase I oncology trials due to its unique ability to trigger apoptosis in various types of cancer cells with limited toxicity toward normal cells (2, 3). However, many primary tumors are inherently resistant to TRAIL-mediated apoptosis and require additional sensitization (4, 5).

TRAIL initiates apoptosis by binding to cell surface death receptor 4 (DR4) or 5 (DR5); this induces oligomerization of the death receptors and formation of the death-inducing signaling complex (DISC), involving recruitment of the adaptor molecule FADD and subsequent caspase-8. DISC assembly promotes the autocleavage and activation of caspase-8, leading to further activation of the effector caspases (e.g., caspase-3) that eventually drive apoptotic death (6). Cellular FLICE-inhibitory protein (c-FLIP) is a truncated form of caspase-8 that lacks enzymatic activity. It can also be recruited to DISC but suppresses apoptosis by blocking the activation of caspase-8 through competing with caspase-8 for binding to FADD (7). It has been well documented that elevated c-FLIP expression protects cells from death receptor-mediated apoptosis, whereas downregulation of c-FLIP by chemicals or siRNA sensitizes cells to death receptor-mediated apoptosis (7, 8). Therefore, c-FLIP acts as a key inhibitor of TRAIL/death receptor-induced apoptosis. c-FLIP has multiple splice variants, however, only 2 of them have been well characterized at the protein levels: the 26 kDa short form (c-FLIP<sub>s</sub>) containing 2 death effector

**Authors' Affiliations:** <sup>1</sup>Department of Hematology and Medical Oncology, Winship Cancer Institute, Emory University School of Medicine, Atlanta, Georgia; <sup>2</sup>Department of Pathology, Xiangya Medical School, Central South University, Changsha, Hunan; and <sup>3</sup>Department of Respiration, First Affiliated Hospital of Medical College of Xi'an Jiaotong University, Xi'an, Shaanxi, PR China

**Note:** Supplementary data for this article are available at Cancer Prevention Research Online (<http://cancerprevres.aacrjournals.org/>).

B. Li and H. Ren share the first authorship.

F.R. Khuri and S-Y. Sun are Georgia Cancer Coalition Distinguished Cancer Scholars.

**Corresponding Author:** Shi-Yong Sun, Emory University School of Medicine, Winship Cancer Institute, 1365-C Clifton Road NE, C-3088, Atlanta, GA 30322. Phone: 404-778-2170; Fax: 404-778-5520; E-mail: [ssun@emory.edu](mailto:ssun@emory.edu)

doi: 10.1158/1940-6207.CAPR-11-0548

©2012 American Association for Cancer Research.

domains and the 55 kDa long form (c-FLIP<sub>L</sub>) containing an inactive caspase-like domain in addition to the 2 death effector domains (7, 9). The levels of c-FLIP, including both FLIP<sub>L</sub> and FLIP<sub>S</sub> are regulated by ubiquitin/proteasome-mediated degradation (10–12).

Although cancer cells possess intrinsic resistance to TRAIL, many anticancer agents can sensitize cancer cells to TRAIL-induced apoptosis through various mechanisms such as induction of DR5 and/or DR4 expression and/or downregulation of c-FLIP levels (13, 14). Akt has been suggested to positively regulate c-FLIP expression because activation or suppression of Akt accordingly increased or decreased the levels of c-FLIP (15). Recently, Akt1 was shown to directly interact with FLIP<sub>L</sub> and to phosphorylate it at S273, leading to stabilization of FLIP<sub>L</sub> (16). Thus, the current study primarily focused on determining whether API-1 negatively regulates c-FLIP levels and sensitizes cancer cells to TRAIL-induced apoptosis. Moreover, we have revealed the mechanisms by which API-1 reduces c-FLIP levels and enhances TRAIL-induced apoptosis.

## Materials and Methods

### Reagents

API-1 (NSC177233) was obtained from the National Cancer Institute (Bethesda, MD). API-2 (17) was provided by Dr. J.Q. Cheng (H. Lee Moffitt Cancer Center and Research Institute, Tampa, FL). MK2206 was purchased from Active Biochem. The soluble recombinant human TRAIL was purchased from PeproTech, Inc. The proteasome inhibitor MG132 and the protein synthesis inhibitor cycloheximide were purchased from Sigma Chemical Corporation. Monoclonal anti-FLIP antibody (NF6) was obtained from Alexis Biochemicals. Mouse monoclonal anti-caspase-8 and rabbit polyclonal anti-caspase-9, anti-PARP, and anti-Akt antibodies were purchased from Cell Signaling Technology, Inc. Mouse monoclonal anti-caspase-3 antibody was purchased from Imgenex. Rabbit polyclonal anti-DR5 antibody was obtained from ProSci Inc. Mouse monoclonal anti-DR4 antibody (B-N28) was purchased from Diaclone. Rabbit monoclonal anti-p-Akt (S473) antibody was purchased from Epitomics, Inc. Both polyclonal and monoclonal anti-actin antibodies were purchased from Sigma Chemical Corporation.

### Cell lines and cell culture

Human non-small cell lung carcinoma (NSCLC) cell lines (H157, Calu-1, and H1299) and head and neck squamous cell carcinoma (HNSCC) cell lines (22A, Tr146, and SqCC/Y1) were described in our previous work (18). H157 cells were recently authenticated by Genetica DNA Laboratories, Inc. by analyzing short tandem repeat DNA profile. The other cell lines have not been authenticated. The H157-LacZ-5, H157-FLIP<sub>L</sub>-21, and H157-FLIP<sub>S</sub>-1 stable transfectants were established as described previously (19, 20). The 22A cells (pool) stably expressing LacZ, FLIP<sub>L</sub>, and FLIP<sub>S</sub> were described previously (21). These cell lines were cultured in PMRI-1640 or Dulbecco's Modified Eagle's

Medium (DMEM)/Ham's F-12 medium containing 5% FBS at 37°C in a humidified atmosphere of 5% CO<sub>2</sub> and 95% air.

### Cell survival and apoptosis assays

Cells were seeded in 96-well cell culture plates and treated the next day with the given agents. The viable cell number was determined by sulforhodamine B (SRB) assay as described previously (22). Combination index (CI) for drug interaction (e.g., synergy) was calculated with the CompuSyn software (ComboSyn, Inc.). Apoptosis was evaluated with the Annexin V-PE Apoptosis Detection Kit purchased from BD Biosciences. We also detected caspase and PARP cleavage by Western blot analysis as described later, as additional indicators of apoptosis.

### Western blot analysis

The procedures for preparation of whole-cell protein lysates and for Western blotting were the same as described before (23, 24). The quantification of Western blotting results was done with NIH ImageJ software.

### Immunoprecipitation for detection of ubiquitinated c-FLIP


H157-FLIP<sub>L</sub>-21 cells were transfected with hemagglutinin (HA)-ubiquitin plasmid with Lipofectamine 2000 transfection reagent (Invitrogen) based on the manufacturer's instructions. After 24 hours, the cells were treated with API-1 or API-1 plus MG132 for 3 hours. Cells were collected and lysed for immunoprecipitation of Flag-FLIP<sub>L</sub> using Flag M2 monoclonal antibody (Sigma) as previously described (19, 25), followed by detection of ubiquitinated FLIP<sub>L</sub> with Western blot analysis using anti-HA antibody (Abgent).

## Results

### API-1 effectively inhibits the growth and induces apoptosis of human NSCLC and HNSCC cells

We first evaluated the single-agent activity of API-1 on the growth of a panel of NSCLC and HNSCC cell lines. A 3-day exposure to API-1 effectively inhibited the growth of 5 (H1299, H157, SqCC/Y1, 22A, and Tr146) of 6 tested cancer cell lines (Fig. 1A). The effective concentrations that decreased cell numbers by 50% (IC<sub>50</sub>s) ranged between 2 and 5 μmol/L for these sensitive cell lines (Fig. 1A). Calu-1 was relatively insensitive to API-1 with an IC<sub>50</sub> value greater than 10 μmol/L.

We then determined whether API-1 induces apoptosis in these cell lines. Treatment of the representative H1299, Calu-1, and SqCC/Y1 cell lines with different concentrations of API-1 for 24 hours dose dependently increased Annexin V-positive (or apoptotic) cells in H1299 and SqCC/Y1 cells (>40% at 10 μmol/L) but did so only minimally in Calu-1 cells (<15% at 10 μmol/L; Fig. 1B). In agreement, we detected dose-dependent increase in cleavage of caspase-8, caspase-9, caspase-3, and PARP in H1299 and SqCC/Y1 cells, but this was not apparent in Calu-1 cells (Fig. 1C). These results clearly indicate that API-1 effectively induces apoptosis in API-1-sensitive NSCLC and HNSCC cell lines.




**Figure 1.** API-1 inhibits the (A) growth of and (B) induces apoptosis including caspase (Casp) activation (C) in human cancer cells. A, the indicated cancer cell lines were seeded in 96-well cell culture plates and treated the next day with API-1 at 0.625, 1.25, 2.5, 5, and 10  $\mu\text{mol/L}$ . After 3 days, cell numbers were estimated by the SRB assay. Cell survival was expressed as the percentage of control (DMSO treated) cells. Data are the means of 4 replicate determinations. Bars,  $\pm$ SDs. B and C, the indicated cell lines were treated with the given concentrations of API-1 for 24 hours and then harvested for measurement of apoptosis using Annexin V staining (A) and for detection of caspase activation with Western blotting (C). Data are the means of duplicate experiments. Bars,  $\pm$ SDs. CF, cleaved fragment.

**API-1 reduces c-FLIP levels without induction of DR4 and DR5 expression**

Because API-1 effectively activates caspase-8, we then asked whether API-1 modulates the levels of key proteins (e.g., c-FLIP, DR4, and DR5) involved in the death receptor-mediated apoptotic pathway. As presented in Fig. 2A, API-1 at 5  $\mu\text{mol/L}$  reduced the levels of c-FLIP in H157, H1299,


SqCC/Y1, and Tr146 cells but not in Calu-1 cells. c-FLIP levels in 22A cells were too low to be detected. API-1 did not increase the expression of either DR5 or DR4 in any of the cell lines. Rather, API-1 reduced the levels of DR4 in some cell lines (H1299, SqCC/Y1, and Tr146). Moreover, we conducted detailed dose-course and time-course studies of the effects of API-1 on the levels of c-FLIP, DR4, and DR5 in H157



**Figure 2.** API-1 downregulates c-FLIP levels without increasing DR4 and DR5 expression. A, the given cell lines were treated with 5  $\mu\text{mol/L}$  API-1 for 24 hours. B, H157 cells were treated with different concentrations of API-1 as indicated for 12 hours. C, H157 cells were treated with 5  $\mu\text{mol/L}$  API-1 for the indicated times. After the aforementioned treatments, the cells were subjected to preparation of whole-cell protein lysates and subsequent Western blot analysis for the given proteins. LE, longer exposure.

Downloaded from <http://aacrjournals.org/cancerpreventionresearch/article-pdf/5/4/612/22251689/612.pdf> by guest on 06 October 2024

**Figure 3.** API-1 enhances TRAIL-induced apoptosis as evaluated by cell survival (A), caspase activation (B), and Annexin V staining (C). A, the indicated cell lines were seeded in 96-well cell culture plates and treated the next day with the given concentrations of API-1 alone, TRAIL alone, or their respective combinations. After 24 hours, cell numbers were estimated by the SRB assay. Data are the means of 4 replicate determinations. Bars,  $\pm$ SDs. B and C, the indicated cell lines were treated with 20 ng/mL TRAIL alone, 5  $\mu$ mol/L API-1 alone, or their combination (A + T). After 24 hours, the cells were harvested for preparation of whole-cell protein lysates and subsequent Western blot analysis (B) or for measurement of apoptosis using Annexin V staining (C). CF, cleaved fragment. Columns, means of duplicate determinations; bars,  $\pm$ SDs.



cells. We found that API-1 could reduce c-FLIP levels even at 1.25  $\mu$ mol/L (Fig. 2B). The apparent reduction of c-FLIP in cells occurred after 4-hour exposure to API-1 (Fig. 2C). Under these conditions, we did not see that API-1 increased the expression of either DR4 or DR5 (Fig. 2B and C). Thus, API-1 downregulates c-FLIP levels without induction of DR4 and DR5 expression in NSCLC and HNSCC cells.

In addition, we compared the effects of API-1 on Akt phosphorylation among the 6 NSCLC and HNSCC cell lines. As presented in Fig. 2A, API-1 inhibited the phosphorylation of Akt in all of the cell lines, albeit with varied potencies. Thus, it is clear that API-1 inhibits Akt phosphorylation.

#### API-1 synergizes with TRAIL to induce apoptosis


Given that API-1 reduces c-FLIP levels, we determined whether API-1 enhances TRAIL-induced apoptosis. As presented in Fig. 3A, the combination of API-1 at 5 or 2.5  $\mu$ mol/L and TRAIL (25–100 ng/mL) was much more effective than either agent alone in decreasing the survival of the tested NSCLC and HNSCC cell lines, except for Calu-1 cells. The CIs for these combinations were less than 1 (Supplementary Fig. S1), indicating that the combination of API-1 and TRAIL synergistically decreased the survival of these cancer cells. In agreement, the combination of API-1 (e.g., 5  $\mu$ mol/L) and TRAIL (e.g., 20 ng/mL) was also much more potent than either agent alone in increasing cleavage

of caspase-8, caspase-9, caspase-3, and PARP in Western blot analysis (Fig. 3B) and in increasing the proportion of Annexin V–positive cells as detected by the Annexin V assay (Fig. 3C) in 2 representative cell lines, H1229 and 22A. For example, we detected approximately 5% and 12% of apoptotic cells in cells treated with TRAIL and API-1, respectively, but more than 50% of apoptotic cells in H1299 cells exposed to the combination of API-1 and TRAIL (Fig. 3C), which is clearly greater than the sum of apoptosis induced by both single agents, further indicating that the combination of API-1 and TRAIL exerts more than additive (i.e., synergistic) apoptosis-inducing activity. Taken together, it is clear that the API-1 synergizes with TRAIL to induce apoptosis in NSCLC and HNSCC cells.

#### Enforced expression of ectopic c-FLIP attenuates the ability of API-1 to augment TRAIL-induced apoptosis

To show whether c-FLIP downregulation contributes to enhancement of TRAIL-induced apoptosis by API-1, we compared the effect of API-1 plus TRAIL on cell survival and apoptosis induction among 22A stable transfectants that express LacZ (control), FLIP<sub>L</sub>, and FLIP<sub>S</sub>. As shown earlier, the combination of API-1 and TRAIL was more effective than either agent alone in decreasing the survival of 22A-LacZ cells but not of 22A-FLIP<sub>L</sub> or 22A-FLIP<sub>S</sub> cells (Fig. 4A). In agreement, the combination of API-1 and TRAIL was much more effective in increasing Annexin





**Figure 4.** Enforced expression of ectopic c-FLIP attenuates the ability of API-1 to decrease cell survival (A) and induce apoptosis (B) including activation of caspases (C). A, the indicated cell lines were seeded in 96-well cell culture plates and treated the next day with DMSO, different concentrations of API-1 alone, 20 ng/mL TRAIL alone, or API-1 plus TRAIL. After 24 hours, cell numbers were estimated by the SRB assay. Cell survival was expressed as the percentage of control (DMSO treated) cells. Data are the means of 4 replicate determinations. Bars,  $\pm$ SDs. B and C, the indicated cell lines were treated with DMSO, 5  $\mu$ mol/L API-1 alone, 20 ng/mL TRAIL alone, or API-1 plus TRAIL. After 24 hours, the cells were harvested for detection of apoptosis with Annexin V staining (B) and for preparation of whole-cell protein lysates and subsequent Western blot analysis to detect the given proteins (C). Columns, means of duplicate experiments; bars,  $\pm$ SDs. CF, cleaved fragment.


V-positive (apoptotic) cells and inducing the cleavage of caspase-8, caspase-9, caspase-3, and PARP in 22A-LacZ cells than in 22A-FLIP<sub>S</sub> and 22A-FLIP<sub>L</sub> cells (Fig. 4B and C). Similar results were also generated from H157 cells that stably express LacZ or FLIP<sub>L</sub> (Supplementary Fig. S2). Enforced expression of FLIP<sub>L</sub> attenuated the ability of API-1 to augment the effect of TRAIL on decreasing cell survival (Supplementary Fig. S2B), on increasing apoptotic populations (Supplementary Fig. S2C), and on inducing cleavage of caspase-8, caspase-9, caspase-3, and PARP (Supplementary Fig. S2D). These data taken together indicate that overexpression of c-FLIP protects cells from apoptosis induced by the API-1 and TRAIL combination, implying that c-FLIP downregulation contributes to enhancement of TRAIL-induced apoptosis by API-1.

We also determined whether overexpression of c-FLIP confers resistance to API-1 alone and found that enforced expression of FLIP<sub>L</sub> or FLIP<sub>S</sub> did not affect the ability of API-1 to decrease cell survival in both H157 and 22A cells (data not shown). This suggests that c-FLIP downregulation may not be sufficient for API-1 to trigger apoptosis.

#### API-1 reduces c-FLIP levels through facilitating ubiquitin/proteasome-mediated degradation

To elucidate the mechanism by which API-1 reduces c-FLIP levels, we first tested whether proteasomal degradation is involved in this process, as c-FLIP is known

to be regulated by an ubiquitin/proteasome-dependent mechanism. Thus, we treated H157 cells with API-1 in the absence and presence of the proteasome inhibitor MG132 and then detected c-FLIP with Western blot analysis. In the absence of MG132, API-1 decreased c-FLIP levels; however, the presence of MG132 increased basal levels of c-FLIP, particularly FLIP<sub>S</sub>, and prevented c-FLIP from reduction by API-1 (Fig. 5A). These data suggest that API-1 induces c-FLIP reduction through a proteasome-dependent mechanism. We next determined whether API-1 increases c-FLIP degradation by measuring its stability. To this end, cycloheximide was added to H157 cells 5 hours after dimethyl sulfoxide (DMSO) or API-1 treatment. The cells were then harvested at the indicated times after cycloheximide for analysis of the c-FLIP degradation rate. These data shown in Fig. 5B and C revealed that the half-lives of FLIP<sub>S</sub> and FLIP<sub>L</sub> in DMSO-treated samples were about 34 and 120 minutes, respectively; on the contrary, in API-1-treated samples, their half-lives were reduced to approximately 24 and 69 minutes, respectively. Therefore, it is apparent that API-1 reduces c-FLIP protein stability. Furthermore, we determined whether API-1 increases c-FLIP ubiquitination. As presented in Fig. 5D, the highest level of ubiquitinated FLIP<sub>L</sub> was detected in H157-FLIP<sub>L</sub>-21 cells (which stably express ectopic flag-FLIP<sub>L</sub>) treated with API-1 plus MG132 compared with API-1 alone or MG132 alone, indicating that API-1 increases c-FLIP ubiquitination. Collectively, we conclude



**Figure 5.** API-1 promotes ubiquitin/proteasome-mediated c-FLIP degradation (A and D) and decreases c-FLIP stability (B and C). A, H157 cells were pretreated with 20  $\mu\text{mol/L}$  MG132 for 30 minutes and then cotreated with 5  $\mu\text{mol/L}$  API-1 for another 4 hours. The cells were then harvested for preparation of whole-cell protein lysates and subsequent Western blot analysis. B and C, H157 cells were treated with DMSO or 5  $\mu\text{mol/L}$  API-1 for 5 hours. The cells were then washed with PBS 3 times and refed with fresh medium containing 10  $\mu\text{g/mL}$  cycloheximide (CHX). At the indicated times, the cells were harvested for preparation of whole-cell protein lysates and subsequent Western blot analysis (B). Protein levels were quantified with the NIH ImageJ software and were normalized to actin. The results were plotted as the relative c-FLIP levels compared with those at time 0 of cycloheximide treatment (C). D, H157-FLIP<sub>L</sub>-21 cells, which stably express ectopic flag-FLIP<sub>L</sub>, were transfected with HA-ubiquitin plasmid with FuGENE 6 transfection reagent for 24 hours. The cells were then pretreated with 20  $\mu\text{mol/L}$  MG132 for 30 minutes and then cotreated with 5  $\mu\text{mol/L}$  API-1 for 4 hours. Whole-cell protein lysates were then prepared for immunoprecipitation (IP) using anti-Flag antibody followed by Western blot (WB) analysis using anti-HA antibody for detection of ubiquitinated FLIP<sub>L</sub> (Ub-FLIP<sub>L</sub>) and anti-Flag antibody for detection of ectopic FLIP<sub>L</sub>. LE, longer exposure.

that API-1 facilitates ubiquitin/proteasome-mediated c-FLIP degradation, leading to downregulation of c-FLIP.


#### Other Akt inhibitors do not downregulate c-FLIP and enhance TRAIL-induced apoptosis

Because API-1 is an Akt inhibitor, we wanted to know whether the effects of API-1 on reducing c-FLIP levels and enhancing TRAIL-induced apoptosis are consequences of Akt inhibition. To this end, we tested whether 2 other allosteric Akt inhibitors, MK2206 (26) and API-2 (17), can also reduce c-FLIP levels and enhance TRAIL-mediated cell killing. Both MK2206 and API-2 at concentrations ranging from 1 to 5  $\mu\text{mol/L}$  (MK2206) or 2.5 to 10  $\mu\text{mol/L}$  (API-2) effectively inhibited Akt phosphorylation in H1299 and H157 cells but did not reduce c-FLIP levels in these cell lines (Fig. 6A and B). MK2206 did not increase the expression of DR5 or DR4 either (Fig. 6A). API-2 did not increase DR5 expression but elevated DR4 levels (Fig. 6B). When combined with TRAIL, both MK2206 and API-2 exhibited only minimal increase in cell killing (i.e., decreasing cell survival) in comparison with cell-killing effects by either agent alone (Fig. 6C and D). Thus, it is clear that other Akt inhibitors do not

function in the same manner as API-1 in downregulating c-FLIP and in enhancing TRAIL-inducing cell killing.

#### Discussion

In this study, we have shown that API-1 effectively inhibits the growth of most NSCLC and HNSCC cell lines tested, with  $\text{IC}_{50}$  values ranging from 1 to 5  $\mu\text{mol/L}$ . Moreover, API-1 effectively induces apoptosis in some NSCLC and HNSCC cell lines (Fig. 1). Thus, API-1 possesses promising single-agent activity against NSCLC and HNSCC cells. When combined with TRAIL, synergistic induction of apoptosis, including decreased cell survival, induction of caspase cleavage, and increased Annexin V-positive cells, occurred in most of the tested cell lines (Fig. 3). To the best of knowledge, this is the first report of the synergistic induction of apoptosis by the combination of API-1 and TRAIL in cancer cells. Given that TRAIL is being tested as a cancer therapeutic agent in clinical trials (6, 27), the further study of the potential application of the API-1 and TRAIL combination in cancer therapy (e.g., NSCLC and HNSCC) is warranted. Recently, targeting the Akt protein kinase or the TRAIL-mediated apoptotic pathway has been emerged as attractive strategies



**Figure 6.** MK2206 and API-2 do not reduce c-FLIP levels (A and B) and only minimally enhance TRAIL-mediated cell killing (C and D). A and B, the indicated cell lines were treated with different concentrations of MK2206 or API-2 as indicated for 24 hours and then subjected to preparation of whole-cell protein lysates and subsequent Western blot analysis for the given proteins. C and D, the indicated cell lines were seeded in 96-well cell culture plates and treated the next day with the given concentrations of MK2206 or API-2 alone, TRAIL alone, or TRAIL plus MK2206 or API-2. After 24 hours, cell numbers were estimated by the SRB assay. Data are the means of 4 replicate determinations. Bars,  $\pm$ SDs.

for cancer chemoprevention (13, 28–30). Indeed, a phase 0 chemoprevention trial on an orally active Akt inhibitor has been successfully conducted recently (31). Thus, the potential of the API-1 alone or in combination with TRAIL in cancer chemoprevention needs investigation as well.

We noted that, among the tested cancer cell lines, Calu-1 was the only cell line that exhibited resistance to API-1 alone or the combination of API-1 and TRAIL (Figs. 1 and 3). Thus understanding of the mechanisms by which API-1 induces apoptosis, including modulation of TRAIL-induced apoptosis, will be very helpful for guiding effective application of API-1 in future treatment of cancer in the clinic.

It is well known that cells can die of apoptosis primarily through the extrinsic death receptor-induced pathway and/or the intrinsic mitochondria-mediated pathway. Cross-talk between these 2 pathways is mediated by the truncated proapoptotic protein Bid (32). The activation of caspase-8 is the key step in the death receptor-mediated apoptosis, whereas caspase-9 activation is the key even in the mitochondria-mediated apoptotic pathway. Activated

caspase-8 can also induce caspase-9 activation through Bid-mediated activation of the mitochondria-mediated apoptotic pathway (32). In this study, we found that API-1 activated both caspase-8 and caspase-9 (Fig. 1C), suggesting that API-1 either activates the death receptor-mediated apoptotic pathway or both the death receptor- and mitochondria-mediated apoptotic pathways, leading to induction of apoptosis.

DR4, DR5, and c-FLIP are key components in the regulation of TRAIL-induced apoptosis (6, 33). Modulation of the levels of these proteins in general (e.g., upregulation of DR4 and/or DR5 and/or downregulation of c-FLIP) results in sensitization of cancer cells to TRAIL-induced apoptosis (13, 14). We found that API-1 reduced c-FLIP levels without increasing DR4 or DR5 expression in the sensitive cancer cell lines (Fig. 2). Interestingly, Calu-1 cells, which are relatively resistant to API-1 or API-1 plus TRAIL, expressed the highest basal levels of c-FLIP, which was not reduced by API-1 (Fig. 2A). These results suggest that c-FLIP downregulation may play a critical role in mediating apoptosis induced by API-1 or by the combination of API-1 and

TRAIL. Enforced expression of ectopic FLIP<sub>L</sub> or FLIP<sub>S</sub> did not confer resistance to API-1 alone, but indeed attenuated or abolished the effect of API-1 on enhancing TRAIL-induced apoptosis in both 22A and H157 cells (Fig. 4 and Supplementary Fig. S2). Therefore, c-FLIP downregulation may not be sufficient for API-1 to initiate apoptosis, suggesting that other mechanisms are needed for API-1–induced apoptosis. However, it is clear that c-FLIP downregulation apparently plays a critical role in mediating synergistic induction of apoptosis by API-1 and TRAIL.

It is known that enhancement of TRAIL-induced apoptosis can be achieved through other mechanisms (e.g., inhibition of Bcl-2 family members) beyond downregulation of c-FLIP (34). Here, we claim a critical role of c-FLIP downregulation in mediating enhancement of TRAIL-induced apoptosis by API-1 but does not exclude other potential mechanisms. We noted that c-FLIP protein was not detected in 22A cells (Fig. 2) and API-1 clearly enhanced TRAIL-induced apoptosis in this cell line (Fig. 3). It is possible that the downregulation of c-FLIP by API-1 in this cell line was not detected because of the sensitivity limitation of the assay. Of course, whether other mechanisms play a more important role than downregulation of c-FLIP in mediating enhancement of TRAIL-induced apoptosis by API-1 in this cell line cannot be ruled out and needs further investigation.

It is known that c-FLIP, including FLIP<sub>L</sub> and FLIP<sub>S</sub>, are rapidly turnover proteins subjected to regulation through ubiquitin/proteasome-mediated protein degradation (10–12). Some small molecules negatively regulate c-FLIP levels through this mechanism (19, 35, 36). In this study, we found that API-1 failed to decrease c-FLIP levels in the presence of a proteasome inhibitor, increased c-FLIP ubiquitination, and reduced the stability of c-FLIP protein (Fig. 5). Thus, we conclude that API-1 reduces c-FLIP levels by facilitating its degradation through the ubiquitin/proteasome-dependent pathway. In the current study, we cannot rule out additional mechanisms accounting for c-FLIP downregulation induced by API-1 such as transcriptional regulation even though they are unlikely to be the primary mechanisms.

It has been suggested that Akt positively regulates c-FLIP expression (15). Recently, Akt1 was shown to directly interact with FLIP<sub>L</sub> and to phosphorylate it at S273, leading to stabilization of FLIP<sub>L</sub> (16). Given that API-1 is an Akt inhibitor, it is reasonable to speculate that API-1 may downregulate c-FLIP due to its Akt-inhibitory activity. To explore this, we tested the effects of 2 additional Akt

inhibitors, MK2206 and API-2, on modulation of c-FLIP levels and TRAIL-induced apoptosis. Unfortunately, both MK2206 and API-2 failed to reduce c-FLIP levels or to detectably enhance TRAIL-induced cell killing although they effectively reduced p-Akt levels (Fig. 6), suggesting that inhibition of Akt does not necessarily result in c-FLIP downregulation and enhancement of TRAIL-induced apoptosis. Accordingly, we suggest that the effects of API-1 on downregulation of c-FLIP and enhancement of TRAIL-induced apoptosis are unlikely secondary to Akt inhibition. Furthermore, we noted that API-1 downregulation of c-FLIP is not associated with its activity against Akt. In Calu-1 cells, API-1 did not reduce c-FLIP levels but inhibited Akt phosphorylation (Fig. 2A). These data further support the notion that API-1 downregulates c-FLIP independent of Akt inhibition.

In summary, the current study has revealed a novel function of API-1 that induces c-FLIP degradation and synergizes with TRAIL to induce apoptosis of cancer cells. Moreover, our results warrant further evaluation of the potential of API-1 and TRAIL combination against cancer in the clinic.

#### Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

#### Authors' Contributions

**Conception and design:** F.R. Khuri, S.-Y. Sun

**Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.):** B. Li, H. Ren

**Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis):** B. Li, H. Ren, F.R. Khuri, S.-Y. Sun

**Writing, review, and/or revision of the manuscript:** F.R. Khuri, S.-Y. Sun

**Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases):** P. Yue

**Study supervision:** F.R. Khuri, S.-Y. Sun

#### Acknowledgments

The authors thank Dr. J.Q. Cheng for generously providing some reagents and Dr. A. Hammond for editing the manuscript.

#### Grant Support

The study was supported by Georgia Cancer Coalition Distinguished Cancer Scholar award (to S.-Y. Sun) and National Cancer Institute NIH SPORE P50 grant CA128613 (project 2 to S.-Y. Sun and F.R. Khuri).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received November 30, 2011; revised January 16, 2012; accepted February 8, 2012; published OnlineFirst February 16, 2012.

#### References

- Kim D, Sun M, He L, Zhou QH, Chen J, Sun XM, et al. A small molecule inhibits Akt through direct binding to Akt and preventing Akt membrane translocation. *J Biol Chem* 2010;285:8383–94.
- Kelley SK, Ashkenazi A. Targeting death receptors in cancer with Apo2L/TRAIL. *Curr Opin Pharmacol* 2004;4:333–9.
- Wajant H, Gerspach J, Pfizenmaier K. Tumor therapeutics by design: targeting and activation of death receptors. *Cytokine Growth Factor Rev* 2005;16:55–76.
- Newsom-Davis T, Prieske S, Walczak H. Is TRAIL the holy grail of cancer therapy? *Apoptosis* 2009;14:607–23.
- Van Geelen CM, de Vries EG, de Jong S. Lessons from TRAIL-resistance mechanisms in colorectal cancer cells: paving the road to patient-tailored therapy. *Drug Resist Updat* 2004;7:345–58.
- Ashkenazi A, Holland P, Eckhardt SG. Ligand-based targeting of apoptosis in cancer: the potential of recombinant human apoptosis



- ligand 2/tumor necrosis factor-related apoptosis-inducing ligand (rhApo2L/TRAIL). *J Clin Oncol* 2008;26:3621–30.
7. Wajant H. Targeting the FLICE inhibitory protein (FLIP) in cancer therapy. *Mol Interv* 2003;3:124–7.
  8. Bagnoli M, Canevari S, Mezzanzanica D. Cellular FLICE-inhibitory protein (c-FLIP) signalling: a key regulator of receptor-mediated apoptosis in physiologic context and in cancer. *Int J Biochem Cell Biol* 2010;42:210–3.
  9. Kataoka T. The caspase-8 modulator c-FLIP. *Crit Rev Immunol* 2005;25:31–58.
  10. Kim Y, Suh N, Sporn M, Reed JC. An inducible pathway for degradation of FLIP protein sensitizes tumor cells to TRAIL-induced apoptosis. *J Biol Chem* 2002;277:22320–9.
  11. Poukkula M, Kaunisto A, Hietakangas V, Denessiouk K, Katajamaki T, Johnson MS, et al. Rapid turnover of c-FLIPshort is determined by its unique C-terminal tail. *J Biol Chem* 2005;280:27345–55.
  12. Chang L, Kamata H, Solinas G, Luo JL, Maeda S, Venuprasad K, et al. The E3 ubiquitin ligase itch couples JNK activation to TNFalpha-induced cell death by inducing c-FLIP(L) turnover. *Cell* 2006;124:601–13.
  13. Sun SY. Chemopreventive agent-induced modulation of death receptors. *Apoptosis* 2005;10:1203–10.
  14. Elrod HA, Sun SY. Modulation of death receptors by cancer therapeutic agents. *Cancer Biol Ther* 2007;7:163–73.
  15. Panka DJ, Mano T, Suhara T, Walsh K, Mier JW. Phosphatidylinositol 3-kinase/Akt activity regulates c-FLIP expression in tumor cells. *J Biol Chem* 2001;276:6893–6.
  16. Shi B, Tran T, Sobkoviak R, Pope RM. Activation-induced degradation of FLIP(L) is mediated via the phosphatidylinositol 3-kinase/Akt signaling pathway in macrophages. *J Biol Chem* 2009;284:14513–23.
  17. Yang L, Dan HC, Sun M, Liu Q, Sun XM, Feldman RI, et al. Akt/protein kinase B signaling inhibitor-2, a selective small molecule inhibitor of Akt signaling with antitumor activity in cancer cells overexpressing Akt. *Cancer Res* 2004;64:4394–9.
  18. Sun SY, Yue P, Mao L, Dawson MI, Shroot B, Lamph WW, et al. Identification of receptor-selective retinoids that are potent inhibitors of the growth of human head and neck squamous cell carcinoma cells. *Clin Cancer Res* 2000;6:1563–73.
  19. Chen S, Liu X, Yue P, Schonthal AH, Khuri FR, Sun SY. CHOP-dependent DR5 induction and ubiquitin/proteasome-mediated c-FLIP downregulation contribute to enhancement of TRAIL-induced apoptosis by dimethyl-celecoxib in human non-small cell lung cancer cells. *Mol Pharmacol* 2007;72:1269–79.
  20. Raja SM, Chen S, Yue P, Acker TM, Lefkove B, Arbiser JL, et al. The natural product honokiol preferentially inhibits cellular FLICE-inhibitory protein and augments death receptor-induced apoptosis. *Mol Cancer Ther* 2008;7:2212–23.
  21. Zhao L, Yue P, Lonial S, Khuri FR, Sun SY. The NEDD8-activating enzyme inhibitor, MLN4924, cooperates with TRAIL to augment apoptosis through facilitating c-FLIP degradation in head and neck cancer cells. *Mol Cancer Ther* 2011;10:2415–25.
  22. Sun SY, Yue P, Dawson MI, Shroot B, Michel S, Lamph WW, et al. Differential effects of synthetic nuclear retinoid receptor-selective retinoids on the growth of human non-small cell lung carcinoma cells. *Cancer Res* 1997;57:4931–9.
  23. Sun SY, Yue P, Wu GS, El-Deiry WS, Shroot B, Hong WK, et al. Mechanisms of apoptosis induced by the synthetic retinoid CD437 in human non-small cell lung carcinoma cells. *Oncogene* 1999;18:2357–65.
  24. Liu X, Yue P, Zhou Z, Khuri FR, Sun SY. Death receptor regulation and celecoxib-induced apoptosis in human lung cancer cells. *J Natl Cancer Inst* 2004;96:1769–80.
  25. Liu X, Yue P, Schonthal AH, Khuri FR, Sun SY. Cellular FLICE-inhibitory protein down-regulation contributes to celecoxib-induced apoptosis in human lung cancer cells. *Cancer Res* 2006;66:11115–9.
  26. Hirai H, Sootome H, Nakatsuru Y, Miyama K, Taguchi S, Tsujioka K, et al. MK-2206, an allosteric Akt inhibitor, enhances antitumor efficacy by standard chemotherapeutic agents or molecular targeted drugs *in vitro* and *in vivo*. *Mol Cancer Ther* 2010;9:1956–67.
  27. Bellail AC, Qi L, Mulligan P, Chhabra V, Hao C. TRAIL agonists on clinical trials for cancer therapy: the promises and the challenges. *Rev Recent Clin Trials* 2009;4:34–41.
  28. Crowell JA, Steele VE, Fay JR. Targeting the AKT protein kinase for cancer chemoprevention. *Mol Cancer Ther* 2007;6:2139–48.
  29. Huang S, Ren X, Wang L, Zhang L, Wu X. Lung-cancer chemoprevention by induction of synthetic lethality in mutant KRAS premalignant cells *in vitro* and *in vivo*. *Cancer Prev Res* 2011;4:666–73.
  30. Zhang L, Ren X, Alt E, Bai X, Huang S, Xu Z, et al. Chemoprevention of colorectal cancer by targeting APC-deficient cells for apoptosis. *Nature* 2010;464:1058–61.
  31. Reid JM, Walden CA, Qin R, Ziegler KL, Haslam JL, Rajewski RA, et al. Phase 0 clinical chemoprevention trial of the Akt inhibitor SR13668. *Cancer Prev Res* 2011;4:347–53.
  32. Hengartner MO. The biochemistry of apoptosis. *Nature* 2000;407:770–6.
  33. Falschlehner C, Ganten TM, Koschny R, Schaefer U, Walczak H. TRAIL and other TRAIL receptor agonists as novel cancer therapeutics. *Adv Exp Med Biol* 2009;647:195–206.
  34. Zhang L, Fang B. Mechanisms of resistance to TRAIL-induced apoptosis in cancer. *Cancer Gene Ther* 2005;12:228–37.
  35. Fan S, Li Y, Yue P, Khuri FR, Sun SY. The eIF4E/eIF4G interaction inhibitor 4EGI-1 augments TRAIL-mediated apoptosis through c-FLIP Down-regulation and DR5 induction independent of inhibition of cap-dependent protein translation. *Neoplasia* 2010;12:346–56.
  36. Lin Y, Liu X, Yue P, Benbrook DM, Berlin KD, Khuri FR, et al. Involvement of c-FLIP and survivin down-regulation in flexible heteroarotinoind-induced apoptosis and enhancement of TRAIL-initiated apoptosis in lung cancer cells. *Mol Cancer Ther* 2008;7:3556–65.