

6-Thioguanine Selectively Kills BRCA2-Defective Tumors and Overcomes PARP Inhibitor Resistance

Natalia Issaeva¹, Huw D. Thomas², Tatjana Djurenovic¹, Janneke E. Jaspers³, Ivaylo Stoimenov¹, Suzanne Kyle², Nicholas Pedley⁴, Ponnari Gottipati⁴, Rafal Zur^{2,5}, Kate Sleeth⁴, Vicky Chatzakos¹, Evan A. Mulligan², Cecilia Lundin⁴, Evgenia Gubanova¹, Ariena Kersbergen³, Adrian L. Harris⁶, Ricky A. Sharma⁴, Sven Rottenberg³, Nicola J. Curtin², and Thomas Helleday^{1,4,7}

Abstract

Familial breast and ovarian cancers are often defective in homologous recombination (HR) due to mutations in the *BRCA1* or *BRCA2* genes. Cisplatin chemotherapy or poly(ADP-ribose) polymerase (PARP) inhibitors were tested for these tumors in clinical trials. In a screen for novel drugs that selectively kill BRCA2-defective cells, we identified 6-thioguanine (6TG), which induces DNA double-strand breaks (DSB) that are repaired by HR. Furthermore, we show that 6TG is as efficient as a PARP inhibitor in selectively killing BRCA2-defective tumors in a xenograft model. Spontaneous BRCA1-defective mammary tumors gain resistance to PARP inhibitors through increased P-glycoprotein expression. Here, we show that 6TG efficiently kills such BRCA1-defective PARP inhibitor-resistant tumors. We also show that 6TG could kill cells and tumors that have gained resistance to PARP inhibitors or cisplatin through genetic reversion of the *BRCA2* gene. Although HR is reactivated in PARP inhibitor-resistant BRCA2-defective cells, it is not fully restored for the repair of 6TG-induced lesions. This is likely to be due to several recombinogenic lesions being formed after 6TG. We show that BRCA2 is also required for survival from mismatch repair-independent lesions formed by 6TG, which do not include DSBs. This suggests that HR is involved in the repair of 6TG-induced DSBs as well as mismatch repair-independent 6TG-induced DNA lesion. Altogether, our data show that 6TG efficiently kills BRCA2-defective tumors and suggest that 6TG may be effective in the treatment of advanced tumors that have developed resistance to PARP inhibitors or platinum-based chemotherapy. *Cancer Res*; 70(15); 6268–76. ©2010 AACR.

Introduction

Breast cancer is the most common cancer in women in the Western world today, and in the United Kingdom, breast cancer incidence rates have increased by more than 50% over the last 25 years. Familial mutations in the breast cancer susceptibility genes *BRCA1* or *BRCA2* are associated with an increased risk of several cancers, particularly

breast, ovarian, and prostate cancer (1). The proteins encoded by these genes play important roles in homologous recombination (HR) repair (2–4) and it is likely that their tumor suppressor function is explained by their role in reducing mutation rates (5). This hypothesis is also supported by the observation that proteins with related functions have also been linked with a predisposition to developing breast cancer, i.e., *CHEK2* (6), *ATM* (7), *PALB2* (*FANCN*; refs. 8–10), and *BRIP1* (*BACH1*; ref. 11).

HR-defective cells are characterized by hypersensitivity to cross-linking agents, which is thought to be related to the role of HR in bypassing inter-strand cross-links during DNA replication (12). HR-defective cells are also hypersensitive to poly(ADP-ribose) polymerase (PARP) inhibitors (13, 14). This involves PARP1 having a role in DNA single-strand break repair (15), which results in the suppression of HR (16). PARP inhibitors might increase the amount of single-strand breaks, which collapse into DNA double-strand breaks (DSB) at replication forks, requiring HR for repair (17). In the absence of HR, these DSBs are not repaired, resulting in HR-defective cells such as *BRCA1*- and *BRCA2*-mutated cancers, which are therefore hypersensitive to PARP inhibitors (13). In the clinic, PARP inhibitors efficiently killed *BRCA1*- and *BRCA2*-defective tumors in a phase I/II clinical trial (18). However, acquired resistance to PARP inhibitor is a problem and may involve either

Authors' Affiliations: ¹Department of Genetics, Microbiology, and Toxicology, Stockholm University, Stockholm, Sweden; ²Northern Institute for Cancer Research, Newcastle University, Medical School, Newcastle upon Tyne, United Kingdom; ³Division of Molecular Biology, The Netherlands Cancer Institute, Amsterdam, the Netherlands; ⁴Cancer Research UK-Medical Research Council, Gray Institute for Radiation Oncology and Biology; ⁵Department of Molecular Cell Genetics, Collegium Medicum, N. Copernicus University, Bydgoszcz, Poland; ⁶Department of Medical Oncology, University of Oxford, Oxford, United Kingdom; and ⁷Science for Life Laboratory, Stockholm University, Solna, Sweden

Note: Supplementary data for this article are available at Cancer Research Online (<http://cancerres.aacrjournals.org/>).

Current address for N. Issaeva: Department of Radiation Oncology, University of Michigan Medical School, Ann Arbor, MI 48109.

Corresponding Author: Thomas Helleday, University of Oxford, ORCRB, Roosevelt Drive, Oxford OX3 7DQ, United Kingdom. Phone: 44-1865-617-324; Fax: 44-1865-617-334; E-mail: thomas.helleday@rob.ox.ac.uk.

doi: 10.1158/0008-5472.CAN-09-3416

©2010 American Association for Cancer Research.

additional mutations in the *BRCA1* or *BRCA2* genes that result in restoration of the COOH-terminal part of the proteins (19–21) or upregulation of *Abcb1a/b* genes encoding P-glycoprotein efflux pumps (22). Although the extent of resistance that can be acquired by these mechanisms is unclear in humans, it suggests that the discovery of agents that may overcome such resistance mechanisms merit further investigation.

Materials and Methods

Chemicals

All chemicals were obtained from Sigma unless stated otherwise. AG014699 was provided by Pfizer GRD, and KU0058948 and olaparib were from KuDOS Pharmaceuticals. AG014699 and KU0058948 compounds were dissolved at 10 mmol/L in 100% dry DMSO and stored at -20°C , it was diluted in culture medium to give the final desired drug concentration in 1% DMSO with control cultures exposed to 1% DMSO alone. The National Cancer Institute (NCI) diversity and mechanistic set were obtained from NCI (Bethesda, MA) and stocks were maintained in 96-well plates in DMSO [stock concentration 1 mmol/L (mechanistic set) and 10 mmol/L (diversity set)].

Cell culture and isolation of PARP inhibitor-resistant cells. HCT116 and HCT116+Chr3 were obtained from Dr. Bert Vogelstein, U2OS cell line was obtained from American Type Culture Collection, capan-1, and resistant capan-1 clones from Dr. Toshiyasu Taniguchi, AA8, irs1SF, and CXR3 cells were from Dr. Larry Thompson, whereas V-C8 and V-C8B2 cells were previously isolated (4). All cells were grown in DMEM with 10% fetal bovine serum and penicillin/streptomycin at 37°C under an atmosphere containing 5% CO_2 .

shRNA depletion of BRCA2

Depletion of BRCA2 expression in U2OS or HCT116 cells was obtained from the stably integrated regulatable expression of BRCA2 short hairpin (shRNA) using the BLOCK-iT Inducible H1 RNAi Entry Vector Kit from Invitrogen, according to the manufacturers protocol (Invitrogen, Sweden). The target sequence introduced was AAC AAC AAU UAC GAA CCA AAC UU (23).

Western blot

Proteins from cell lysates were separated and detected using Western blotting as previously described (24). The primary antibody was an anti-rabbit BRCA2 antibody (Santa Cruz Biotechnology) diluted 1:500 in blocking solution.

Screening procedure

U2OS BRCA2 shRNA cells were grown in the presence or absence of 2 $\mu\text{g}/\text{mL}$ of doxycyclin for 2 days, plated in 96-well plates (2,000 cells/well) in the presence or absence of doxycyclin, and the next day, treated with test compounds (at a concentration 10 $\mu\text{mol}/\text{L}$) from the NCI library. After 72 hours, WST-1 cell proliferation reagent was used to deter-

mine the cells viability as described earlier (25). Compounds from the library that selectively suppressed the growth of BRCA2-defective cells, but had only modest effects on BRCA2-proficient cells were selected.

Colony formation assay

Cells were plated into six-well plates at a density of 200 cells/well. The next day, the cells were treated with selected compounds at a range of concentrations, for different time periods (from 3 hours up to 6 days). After 1 week [for V-C8 and V-C8+B2 cells and PARP inhibitor-resistant (PIR) clones] or 2 weeks (for U2OS cells), when colonies could be observed, the colonies were fixed and stained with methylene blue in methanol (4 g/L). Colonies consisting of more than 30 cells were subsequently counted. In addition, exponentially growing V-C8, V-C8+B2, and PIR clones 1C and 2B were exposed to varying concentrations of AG014699 for 24 hours or 6-thioguanine (6TG) for 48 hours prior to seeding in 90 mm dishes in drug-free medium for colony formation. Colonies were fixed and stained with crystal violet 10 to 14 days later and counted on an automatic colony counter (Oxford Optronix).

Pulsed-field gel electrophoresis

Cells were treated with 6TG, collected by trypsinization, resuspended in 1% InCert-agarose (in 37°C PBS) to a final concentration of 1.5 million cells/100 μL and agarose plugs were separated by pulsed-field gel electrophoresis as previously described (26).

Immunofluorescence

Cells were grown on coverslips, treated, fixed, immunostained, and analyzed as previously described (24). The primary antibodies used were mouse monoclonal anti- γH2AX at a dilution of 1:1,000 and rabbit polyclonal anti-Rad51 (H-92; Santa Cruz Biotechnology) at a dilution of 1:1,000.

Propidium iodide staining and fluorescence-activated cell sorting analysis

Cells (1×10^6) were treated (or not treated) with compounds (such as B9 or 6TG), collected by trypsinization, and fixed in ice-cold 70% ethanol overnight at -20°C . The cells were then rehydrated in PBS and stained with 50 $\mu\text{g}/\text{mL}$ of propidium iodide and 100 $\mu\text{g}/\text{mL}$ of RNase A in PBS for 30 minutes at room temperature. Samples were further analyzed on a BD Biosciences FACScan. The data was analyzed with WinMDI software version 2.8.

In vivo experiments

Exponentially growing VC8 or VC8-B2 cells (1×10^7) were injected i.m. into the thigh of each CD1 nude (Charles River) mouse in 50 μL of PBS and handled and analyzed as previously described (13). Tumor growth was assessed using the ratio of the diameter of the right (tumor bearing) to the left (normal) thighs. When thigh ratio reached between 1.3 and 1.5, mice were divided into groups for the following treatments: AG014699 10 mg/kg (made up on day of use at 1 mg/mL in water) or 6TG 1.5 mg/kg (made up on

day of use at 0.15 mg/mL in PBS) or saline (control) administered daily \times 10 i.p.

Generation of mammary tumors

Brcal ^{$\Delta 5-13/\Delta 5-13$} ; *p53* ^{$\Delta 2-10/\Delta 2-10$} mammary tumors were generated in *K14cre;Brcal* ^{$F5-13/F5-13$} ; *p53* ^{$F2-10/F2-10$} mice and genotyped as described (27). Orthotopic transplantations of tumor fragments into syngeneic animals and caliper measurements of mammary tumors have been previously reported (22).

Results

BRCA2-defective cells are hypersensitive to 6TG

Individuals with inherited mutations in either *BRCA1* or *BRCA2* alleles have a high risk of developing breast cancer (1, 28). Here, we wanted to identify novel compounds to selectively kill BRCA2-defective cells. We developed a shRNA system to deplete the BRCA2 protein upon removal of doxycycline in U2OS sarcoma cells, and we assayed the mechanistic and diversity set compound libraries from the NCI for compounds that selectively killed BRCA2-depleted cells. Mercury-(2-amino-1, 9-dihydro-6H-purine-6-thionato-N7,S6)hexyl (B9) was identified in the screen as the most efficient compound to selectively kill BRCA2-depleted U2OS cells (Supplementary Fig. S1). We also found that this compound selectively killed BRCA2-defective V-C8 cells as compared with V-C8+B2, the isogenic cell line expressing BRCA2^{WT} protein (V-C8+B2; ref. 4; Supplementary Fig. S1). The reason for the selective killing of BRCA2-defective cells is likely to be explained by the role of BRCA2 in HR because cells defective in the RAD51 paralogs

were similarly sensitive to the B9 compound (Supplementary Fig. S1).

Structural analysis of the B9 compound revealed a strong resemblance to 6TG, which is a well-established chemotherapy used to treat hematologic malignancies in children and adults (29). We therefore decided to test the sensitivity of BRCA2-deficient cells to 6TG and found that survival was significantly lower than that of BRCA2-expressing cells (Fig. 1A). Furthermore, we found that the BRCA2 protein is required to prevent apoptosis induced by 6TG, measured by sub-G₁ population (Fig. 1B) and the terminal deoxynucleotidyl transferase-mediated dUTP nick end labeling assay (Fig. 1C).

6TG selectively kills BRCA2-defective tumors

To test the hypothesis that 6TG is useful for selective treatment of BRCA2-deficient tumors, we treated mice bearing xenografts derived from BRCA2-deficient V-C8 and wild-type BRCA2-complemented V-C8+B2 cells. Consistent with the hypothesis, we found that neither the PARP inhibitor, AG014699, nor 6TG had any effect on the outgrowth of the BRCA2-proficient tumors (Fig. 1D); in contrast, all mice with BRCA2-defective tumors responded to both 6TG and the PARP inhibitor equally, with significant growth delay and three of five complete tumor regressions in both groups. These results suggest that 6TG is as effective as the PARP inhibitor, AG014699, in selectively killing BRCA2-defective tumors. However, it should be noted that 6TG at 1.5 mg/kg caused a greater loss of body weight than AG014699 (Supplementary Table S1) and was not tolerated at a higher dose of 3 mg/kg (data not shown).





Figure 1. HR-defective cells are hypersensitive to chemical 6TG. A, 6TG selectively kills BRCA2-defective VC-8 cells in a colony formation assay. 6TG induces apoptosis in HR-defective VC-8 cells as measured by fluorescence-activated cell sorting analysis of the sub-G₁ population (B) and terminal deoxynucleotidyl transferase-mediated dUTP nick end labeling (TUNEL) staining (C). Columns, averages from at least three experiments; bars, SD. D, tumor outgrowth in xenograft mice (five per group) following injection of V-C8 and BRCA2 complemented V-C8+B2 upon i.p. treatment with 6TG and PARP inhibitor.

Figure 2. HR is needed to repair the damage induced by 6TG. Equal amounts of DNA damage in V-C8 and V-C8+B2 cells following 6TG treatment, as determined by γ H2AX foci (A; cells containing more than 10 foci were scored as positive) or by Western blot (B). C, RAD51 foci do not form in V-C8 cells upon 6TG-induced DNA damage. RAD51 foci do form in V-C8+B2 cells and colocalize with γ H2AX foci. D, quantification of RAD51 foci formed in V-C8 and V-C8+B2 cells after 6TG treatment. E, 6TG selectively kills XRCC3 defective *irs1SF* cells in a colony formation assay as compared with wild-type control (AA8).



6TG-induced DNA DSBs are repaired by HR

6TG is an antimetabolite of purine metabolism and is incorporated into the DNA of mammalian cells in place of guanine during DNA replication (29). The incorporated 6TG (~ 1 in 10^4 bases) is then methylated *in situ* to 6-meTG by endogenous *S*-adenosylmethionine and becomes a substrate for mismatch repair in the second replication round to mediate its toxicity (30–33). As a measure of formation of DSBs in DNA, we looked at the nuclear levels of γ H2AX in cells treated with 6TG and we found that γ H2AX foci formed with similar frequency in both BRCA2-deficient and -proficient cells (Fig. 2A and B). This similarity suggests that the hypersensitivity of BRCA2-deficient cells to 6TG is due to an inability to repair DNA damage rather than a difference in the amount of damage introduced. We also found that γ H2AX foci colocalized with RAD51 foci after 6TG treatment in BRCA2-proficient cells (Fig. 2C and D, right panels), suggesting that the RAD51 protein is recruited to DSBs to repair

the lesion by HR. In contrast, RAD51 foci did not form at 6TG-induced γ H2AX foci in BRCA2-defective V-C8 cells, suggesting that DSB repair is deficient in these cells (Fig. 2C and D, left panels). Using pulsed-field gel electrophoresis, we found that BRCA2-defective V-C8 cells were unable to repair 6TG-induced DSBs compared with BRCA2-expressing cells (Supplementary Fig. S2). To further investigate the role of HR in the 6TG-induced DNA damage, we looked at the survival of cells defective in the RAD51 paralogue, XRCC3 (*irs1SF*), and we found that these cells were considerably more sensitive to 6TG than AA8 control cells (Fig. 2E). Altogether, these findings suggest that the sensitivity of BRCA2-defective cells to 6TG was due to their inability to perform HR repair. HR is important in repairing DNA damage caused by a wide range of mono- or bifunctional alkylating anticancer agents, for example, the commonly used drugs, cisplatin and mitomycin C (34). To our knowledge, these results with 6TG represent the first time HR has been implicated in the




Figure 3. Response of the PIR *Brca1*^{Δ5-13/Δ5-13}; *p53*^{Δ2-10/Δ2-10} tumor T6-28 to 6TG. Animals carrying orthotopically transplanted tumors were treated with 1.5 mg of 6TG per kg i.p. daily on days 0 to 9 or 50 mg of olaparib per kg i.p. daily when the tumors reached a volume of 150 to 250 mm³ (100%, day 0). When tumors relapsed back to 100% (arrows), a second treatment of 1.5 mg of 6TG per kg daily for 4 consecutive days was tolerated.

repair of thiopurine antimetabolite drugs. Interestingly, it has previously been shown that HR-defective cells are more sensitive to O⁶-methyl guanine lesions than cell lines with defects in other repair pathways (35, 36), a finding which cor-

roborates the importance of HR in repairing lesions on the O⁶ position of guanine.

BRCA1-defective tumors that gained resistance to PARP inhibitors through P-glycoprotein expression remain 6TG-sensitive

Increased expression from the *Abcb1a* and *Abcb1b* genes encoding the mouse drug efflux transporter P-glycoprotein explains resistance to the PARP inhibitor olaparib in BRCA1;p53-defective mammary tumors (22). Here, we wanted to determine whether 6TG could target such PIR tumors. To test this, *Brca1*^{Δ5-13/Δ5-13}; *p53*^{Δ2-10/Δ2-10} mammary tumors derived from *K14cre;Brca1*^{F5-13/F5-13}; *p53*^{F2-10/F2-10} mice were grown out and treated with olaparib as previously described (22). Small tumor fragments of an olaparib-resistant tumor (T6-28) with 80-fold increased expression of the *Abcb1b* gene were transplanted orthotopically into syngeneic wild-type female mice. The animals were then treated with 6TG when the tumor volume reached ~200 mm³. Interestingly, we found that tumors responded to 6TG (Fig. 3). This shows that spontaneous BRCA1;p53-defective mammary tumors are sensitive to 6TG, and importantly, that PIR tumors in which resistance is caused by increased P-glycoprotein-mediated drug efflux remain




Figure 4. PARP inhibitor and cisplatin-resistant BRCA2-defective cells and tumors respond to 6TG. Clonogenic survival in BRCA2-defective V-C8, BRCA2-complemented V-C8+B2, and V-C8 PIR clones following treatment with PARP inhibitor AG014699 (A), cisplatin (B), or 6TG (C). Columns, averages from at least three experiments; bars, SD. D, tumor outgrowth in xenograft mice following injection of PIR V-C8 clone 2B upon i.p. treatment with 6TG and PARP inhibitor. 6TG retards PIR V-C8 clone 2B tumor outgrowth (statistically significant in Mann-Whitney test, *P* < 0.01). Points, averages from 10 mice in each group; bars, SE.




Figure 5. PARP inhibitor-resistant BRCA2-defective cells partially activates HR in response to 6TG. A, quantification of 6TG and PARP inhibitor induced RAD51 foci. BRCA2 complemented V-C8+B2 cells showed a significant increase in RAD51 foci compared with PIR V-C8 cells upon 6TG treatment (24 h after an 18-h treatment with 1 μ mol/L of 6TG). B, quantification of repair of 6TG-induced γ H2AX foci. HR-defective V-C8 cells fail to repair 6TG-induced γ H2AX foci, whereas V-C8+B2 cells repairs a majority of 6TG-induced γ H2AX foci. PIR V-C8 clones show an intermediate repair of 6TG-induced γ H2AX foci. Columns, averages of three independent experiments; bars, SE. Values marked with asterisks are statistically significant in *t* tests (*, $P < 0.05$; **, $P < 0.01$; and ***, $P < 0.001$).

sensitive to 6TG. After the 10-day treatment with 6TG, tumors eventually grew back. However, such tumors were still responding to a second line treatment with 6TG, indicating that the tumors did not easily obtain resistance to 6TG.

BRCA2-defective cells and tumors that gain resistance to PARP inhibitors through genetic reversion respond to 6TG treatment

The mechanisms of acquired resistance to PARP inhibitors could also evolve through genetic reversion in BRCA2-defective cancer cells (20, 21). In such cases, a mutation in the *BRCA2* gene results in which the COOH-terminal part of the protein is retained and the protein is overall functional in HR, despite missing a single-stranded DNA (ssDNA) binding domain (20, 21). To investigate this mechanism of resistance further, we used BRCA2-defective V-C8 clones selected for resistance to a PARP inhibitor.⁸ All PIR V-C8 clones harbor the same mutation, restoring the correct reading frame for *BRCA2* and at the same time introducing a mutation within a highly conserved region in exon 15. This mutation affects a highly conserved arginine that was also identified in a family with breast and ovarian cancers (37), and was described in mitomycin C-resistant V-C8 cells (38). Thus, the reverted BRCA2 still has a defective ssDNA domain in the COOH-terminal part of the protein, as described earlier (20, 21), which restored HR as indicated by the ability to form RAD51 foci in response to PARP inhibitor treatment (Supplementary Fig. S3). We tested the sensitivity of PIR-resistant clones 1C and 2B to the PARP inhibitor AG014699 and found that both clones had lost their sensitivity to PARP inhibitors (Fig. 4A).

⁸ Gottipati et al. 2010, submitted manuscript.

We then tested the sensitivity of PIR V-C8 cells for cross-resistance to cisplatin and 6TG. Parental V-C8 cells are highly sensitive to both cisplatin and 6TG, compared with the BRCA2-expressing V-C8+B2 cells (Fig. 4B and C). As expected from previous studies (20, 21), PIR V-C8 clones exhibited resistance to cisplatin (Fig. 4B). Surprisingly, we found that PIR V-C8 cells had not fully reverted to resistance to 6TG (Fig. 4C), suggesting that 6TG might still kill PIR BRCA2-defective tumors that gained resistance through genetic reversion.

We confirmed that BRCA2 revertant cells, which have acquired resistance to cisplatin, retain sensitivity to 6TG by using BRCA2-defective human pancreatic cancer cell line capan-1 and four different independent cisplatin-resistant capan-1 clones (20). Clones 6 and 12 acquired resistance through an additional mutation in the *BRCA2* gene that corrected the frameshift caused by the 6174delT mutation in capan-1 cells, whereas clones 10 and 11 did not have an additional *BRCA2* mutation, lack BRCA2 protein expression, and are likely to have reverted to cisplatin resistance through other unknown pathways (20). Interestingly, all four cisplatin-resistant clones showed equal sensitivity to 6TG as parental capan-1 cells (Supplementary Fig. S4), providing additional evidence that PIR and cisplatin-resistant BRCA2-defective cancer cells are sensitive to treatment with 6TG.

Next, we wanted to test whether 6TG could also retard the outgrowth of PIR BRCA2-defective tumors that have gained resistance through genetic reversion. To test this, we treated mice bearing xenografts derived from PIR clone 2B with a 10-day treatment of AG014699 or 6TG and found that PIR clone 2B tumors only responded to the 6TG treatment and not to the PARP inhibitor (Fig. 4D; statistically significant in Mann-Whitney test, $P < 0.01$).

Differential sensitivity to anticancer drugs in genetically reverted BRCA2-defective cells

To gain further insights into the function of the restored BRCA2 protein carrying a mutation in the ssDNA binding domain, we further investigated the sensitivity of the PIR clones to a range of cytotoxic agents (Supplementary Fig. S5A–E and F). The PIR clones exhibited similar levels of resistance to temozolomide, camptothecin, and ionizing radiation as the V-C8+B2 cells compared with the more sensitive V-C8 cells. Interestingly, the PIR clones were slightly more resistant to doxorubicin than the V-C8 cells (Supplementary Fig. S5D). Surprisingly, the V-C8 and the PIR clones were less sensitive to gemcitabine and paclitaxel than the V-C8+B2 cells (Supplementary Fig. S5E and F). These data suggest that recombination-defective BRCA1 and BRCA2 tumors would respond poorly to gemcitabine and paclitaxel treatments.




Figure 6. BRCA2 suppresses 6TG toxicity in mismatch repair-deficient cells. A, survival following continuous treatment with PARP inhibitor ANI, 6TG, and cisplatin in HCT116 and HCT116+Ch3 cells. B, γ H2AX foci formation in hMLH1-defective HCT116 and hMLH1 complemented HCT116+Ch3 cells after a 24-h treatment with 6TG and cisplatin or a 4-h treatment with PARP inhibitor ANI. C, clonogenic survival in HCT116 and BRCA2 depleted HCT116 to increasing doses of 6TG. Points, average of three independent experiments; bars, SD. Values marked with asterisks are statistically significant in *t* tests (***, $P < 0.001$).

The retained sensitivity to 6TG and resistance to gemcitabine and paclitaxel in the PIR clones is likely to be explained by the reverted *BRCA2* gene which did not revert back to wild-type, but retained a mutation in the ssDNA domain, which may impair HR induced by these agents. This ssDNA domain might be required for the BRCA2 response to 6TG and might prevent the efficient repair of gemcitabine- and paclitaxel-induced lesions.

6TG induces both mismatch repair-dependent and -independent lesions that require HR repair

Next, we wanted to understand the mechanism for PIR cells maintaining their sensitivity to 6TG. We analyzed RAD51 foci and found that PIR V-C8 cells induced RAD51 foci as efficiently as V-C8+B2 cells in response to treatment with a PARP inhibitor, but not in response to treatment with 6TG, suggesting that the BRCA2-reverted protein is not fully proficient for 6TG-induced HR (Fig. 5A). We also investigated the translocation of the RAD51 protein into the chromatin fraction, as this may be associated with the efficiency of RAD51 loading on to DNA and subsequent HR (39). We found that the RAD51 protein is more efficiently recruited to DNA after 6TG in V-C8+B2 cells than in the IC clone, and conversely, that the RAD51 protein is more efficiently recruited to DNA in the IC clone than V-C8+B2 following treatment with the PARP inhibitor, KU0058948 (Supplementary Fig. S6). Next, we investigated the repair of 6TG-induced DNA lesions by γ H2AX foci formation and find that BRCA2 is required for efficient repair (Fig. 5B). Interestingly, neither the IC or 2B clones fully repaired the 6TG-induced DNA damage, suggesting that their lack of efficient HR repair is the reason for their 6TG sensitivity.

The reason for the differential response to 6TG and PARP inhibitors might be related to the production of different recombinogenic lesions in DNA. To test this hypothesis, we treated mismatch repair-defective HCT116 colorectal cancer cells and the same cells with restored mismatch repair function (HCT116+Ch3; ref. 40) with 6TG, the PARP inhibitor 4-amino-1,8-naphthalimide (ANI) and cisplatin. We found that only the cytotoxicity of 6TG was dependent on a functional mismatch repair pathway (Fig. 6A; Supplementary Fig. S7). Furthermore, the level of recombinogenic DSBs, measured by γ H2AX foci, were dependent on mismatch repair after 6TG treatment, whereas the generation of these lesions was unaffected by PARP inhibitor and cisplatin treatment in these cell lines (Fig. 6B). Our data are in line with evidence showing that toxic DSBs induced by 6TG are mismatch repair-dependent (30), and it has previously been shown that HR induced by O⁶-guanine methylating agents depends on mismatch repair (41). This is also in line with our previous observation that 6TG-induced DSBs require HR for repair (Fig. 5B; Supplementary Fig. S2).

There is a possibility that 6TG may produce another mismatch repair-independent HR substrate, given that the PIR V-C8 clones show an intermediate HR response to 6TG (Fig. 5). To test this, we shRNA-depleted BRCA2 in HCT116 cells to test if the absence of 6TG-induced DSBs also

abolished the requirements for HR survival. Surprisingly, we find that mismatch repair–defective HCT116 still requires BRCA2 for survival to 6TG, showing that 6TG also produces a recombinogenic lesion that is independent of mismatch repair (Fig. 6C).

Discussion

Here, we report that cells and/or tumors defective in the HR genes *BRCA1*, *BRCA2*, or *XRCC3* are hypersensitive to 6TG and that, in the case of *BRCA2*, this could be reversed by the introduction of a vector expressing the BRCA2 protein. We show that 6TG induced RAD51 foci at 6TG-induced DSBs and that the DSBs were less efficiently repaired in BRCA2-defective cells, which correlates with increased toxicity in HR-defective cells. This is, to our knowledge, the first time that HR has been implicated in the repair of 6TG-induced DSBs.

Interestingly, the opposite result was previously reported: that expression of *BRCA1*^{WT} in BRCA1-mutated HCC1937 breast cancer cells increases sensitivity to 6TG (42). However, this is unrelated to any role of BRCA1 in HR (42) and the BRCA1 mutation in HCC1937 cells is unlikely to affect HR, as RAD51 foci are efficiently induced by IR in these cells (43).

Although PARP inhibitors have been shown to efficiently kill both *BRCA1*- and *BRCA2*-defective tumors, resistance to therapy may develop within 18 to 77 weeks (18). Although the exact mechanisms for PIR in patients remains unknown, this might involve the expression of P-glycoprotein efflux pumps as in mammary mice tumors (22) or through genetic reversion of either *BRCA1* or *BRCA2* (19–21). Here, we show that 6TG efficiently kills PIR BRCA1-defective mammary tumors (Fig. 3), which is likely explained by 6TG not being a substrate for P-glycoprotein (44). Furthermore, we show that genetically reverted BRCA2-defective cells and tumors respond to 6TG. Altogether, these findings suggest that 6TG might also be efficient in killing advanced and drug-resistant *BRCA1*- or *BRCA2*-defective tumors.

Here, we find that PIR V-C8 clones do not completely revert back to a functional HR phenotype in response to 6TG (Fig. 5), which likely explains their 6TG sensitivity. This sug-

gests that there may be several lesions formed following 6TG treatment that trigger HR. For instance, we recently showed that HR is involved in restart at stalled replication forks, which does not involve DSB repair (45, 46). Thus, there is a possibility that 6TG might cause replication lesions other than mismatch repair–dependent DSBs that trigger HR. In support for this notion, we find that mismatch repair–defective HCT116 cells are sensitized to 6TG by the depletion of BRCA2 in spite of already being defective in mismatch repair. This shows that both mismatch repair–dependent and –independent HR lesions are formed by 6TG. Also, this would explain the intermediate response in PIR V-C8 clones to 6TG.

In conclusion, we show that 6TG could be efficiently used to selectively kill BRCA2-defective tumors and that 6TG might be used as treatment for *BRCA1* or *BRCA2* mutant tumors which are resistant to cisplatin chemotherapy and/or PARP inhibitor therapy by various mechanisms.

Disclosure of Potential Conflicts of Interest

Two patent applications regarding usage of PARP inhibitors in BRCA1 and BRCA2 mutated tumors were filed in 2003 by Cancer Research Technology Limited and The University of Sheffield (N.J. Curtin and T. Helleday and T. Helleday alone are named inventors on these patents, respectively). The other authors disclosed no potential conflicts of interest.

Acknowledgments

We thank the NCI, Drs. Toshiyasu Taniguchi, Bert Vogelstein, Larry Thompson, Zdenek Hostomsky, and Mark O'Connor for materials and discussions.

Grant Support

The Swedish Cancer Society, the Swedish Children's Cancer Foundation, the Swedish Research Council, the Swedish Pain Relief Foundation, Cancer Research UK, the NIHR Biomedical Research Centre, Oxford, and the Medical Research Council. The authors declare that they have no competing financial interest.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received 09/15/2009; revised 05/21/2010; accepted 06/08/2010; published OnlineFirst 07/14/2010.

References

1. Wooster R, Bignell G, Lancaster J, et al. Identification of the breast cancer susceptibility gene BRCA2. *Nature* 1995;378:789–92.
2. Moynahan ME, Pierce AJ, Jasin M. BRCA2 is required for homology-directed repair of chromosomal breaks. *Mol Cell* 2001;7:263–72.
3. Xia F, Taghian DG, DeFrank JS, et al. Deficiency of human BRCA2 leads to impaired homologous recombination but maintains normal nonhomologous end joining. *Proc Natl Acad Sci U S A* 2001;98:8644–9.
4. Kraakman-van der Zwet M, Overkamp WJ, van Lange RE, et al. Brca2 (XRCC11) deficiency results in radioresistant DNA synthesis and a higher frequency of spontaneous deletions. *Mol Cell Biol* 2002;22:669–79.
5. Kraakman-van der Zwet M, Wiegant WW, Zdzienicka MZ. Brca2 (XRCC11) deficiency results in enhanced mutagenesis. *Mutagenesis* 2003;18:521–5.
6. Meijers-Heijboer H, van den Ouweland A, Klijn J, et al. Low-penetrance susceptibility to breast cancer due to CHEK2(*) 1100delC in noncarriers of BRCA1 or BRCA2 mutations. *Nat Genet* 2002;31:55–9.
7. Renwick A, Thompson D, Seal S, et al. ATM mutations that cause ataxia-telangiectasia are breast cancer susceptibility alleles. *Nat Genet* 2006;38:873–5.
8. Rahman N, Seal S, Thompson D, et al. PALB2, which encodes a BRCA2-interacting protein, is a breast cancer susceptibility gene. *Nat Genet* 2007;39:165–7.
9. Reid S, Schindler D, Hanenberg H, et al. Biallelic mutations in PALB2 cause Fanconi anemia subtype FA-N and predispose to childhood cancer. *Nat Genet* 2007;39:162–4.
10. Xia B, Dorsman JC, Ameziane N, et al. Fanconi anemia is associated with a defect in the BRCA2 partner PALB2. *Nat Genet* 2007;39:159–61.
11. Seal S, Thompson D, Renwick A, et al. Truncating mutations in the Fanconi anemia J gene BRIP1 are low-penetrance breast cancer susceptibility alleles. *Nat Genet* 2006;38:1239–41.
12. De Silva IU, McHugh PJ, Clingen PH, Hartley JA. Defining the roles of

- nucleotide excision repair and recombination in the repair of DNA interstrand cross-links in mammalian cells. *Mol Cell Biol* 2000;20:7980–90.
13. Bryant HE, Schultz N, Thomas HD, et al. Specific killing of BRCA2-deficient tumours with inhibitors of poly(ADP-ribose)polymerase. *Nature* 2005;434:913–7.
 14. Farmer H, McCabe N, Lord CJ, et al. Targeting the DNA repair defect in BRCA mutant cells as a therapeutic strategy. *Nature* 2005;434:917–21.
 15. Satoh MS, Lindahl T. Role of poly(ADP-ribose) formation in DNA repair. *Nature* 1992;356:356–8.
 16. Schultz N, Lopez E, Saleh-Gohari N, Helleday T. Poly(ADP-ribose) polymerase (PARP-1) has a controlling role in homologous recombination. *Nucleic Acids Res* 2003;31:4959–64.
 17. Arnaudeau C, Lundin C, Helleday T. DNA double-strand breaks associated with replication forks are predominantly repaired by homologous recombination involving an exchange mechanism in mammalian cells. *J Mol Biol* 2001;307:1235–45.
 18. Fong PC, Boss DS, Yap TA, et al. Inhibition of poly(ADP-ribose) polymerase in tumors from BRCA mutation carriers. *N Engl J Med* 2009;361:123–34.
 19. Swisher EM, Sakai W, Karlan BY, Wurz K, Urban N, Taniguchi T. Secondary BRCA1 mutations in BRCA1-mutated ovarian carcinomas with platinum resistance. *Cancer Res* 2008;68:2581–6.
 20. Sakai W, Swisher EM, Karlan BY, et al. Secondary mutations as a mechanism of cisplatin resistance in BRCA2-mutated cancers. *Nature* 2008;451:1116–20.
 21. Edwards SL, Brough R, Lord CJ, et al. Resistance to therapy caused by intragenic deletion in BRCA2. *Nature* 2008;451:1111–5.
 22. Rottenberg S, Jaspers JE, Kersbergen A, et al. High sensitivity of BRCA1-deficient mammary tumors to the PARP inhibitor AZD2281 alone and in combination with platinum drugs. *Proc Natl Acad Sci U S A* 2008;105:17079–84.
 23. Bruun D, Folias A, Akkari Y, Cox Y, Olson S, Moses R. siRNA depletion of BRCA1, but not BRCA2, causes increased genome instability in Fanconi anemia cells. *DNA Repair (Amst)* 2003;2:1007–13.
 24. Bryant HE, Petermann E, Schultz N, et al. PARP is activated at stalled forks to mediate Mre11-dependent replication restart and recombination. *EMBO J* 2009;28:2601–15.
 25. Issaeva N, Bozko P, Enge M, et al. Small molecule RITA binds to p53, blocks p53-HDM-2 interaction and activates p53 function in tumors. *Nat Med* 2004;10:1321–8.
 26. Lundin C, Erixon K, Arnaudeau C, et al. Different roles for nonhomologous end joining and homologous recombination following replication arrest in mammalian cells. *Mol Cell Biol* 2002;22:5869–78.
 27. Liu X, Holstege H, van der Gulden H, et al. Somatic loss of BRCA1 and p53 in mice induces mammary tumors with features of human BRCA1-mutated basal-like breast cancer. *Proc Natl Acad Sci U S A* 2007;104:12111–6.
 28. Miki Y, Swensen J, Shattuck-Eidens D, et al. A strong candidate for the breast and ovarian cancer susceptibility gene BRCA1. *Science* 1994;266:66–71.
 29. Karran P, Attard N. Thiopurines in current medical practice: molecular mechanisms and contributions to therapy-related cancer. *Nat Rev Cancer* 2008;8:24–36.
 30. Swann PF, Waters TR, Moulton DC, et al. Role of postreplicative DNA mismatch repair in the cytotoxic action of thioguanine. *Science* 1996;273:1109–11.
 31. Griffin S, Branch P, Xu YZ, Karran P. DNA mismatch binding and incision at modified guanine bases by extracts of mammalian cells: implications for tolerance to DNA methylation damage. *Biochemistry* 1994;33:4787–93.
 32. Waters TR, Swann PF. Cytotoxic mechanism of 6-thioguanine: hMutSalpha, the human mismatch binding heterodimer, binds to DNA containing S6-methylthioguanine. *Biochemistry* 1997;36:2501–6.
 33. Yan T, Berry SE, Desai AB, Kinsella TJ. DNA mismatch repair (MMR) mediates 6-thioguanine genotoxicity by introducing single-strand breaks to signal a G2-M arrest in MMR-proficient RKO cells. *Clin Cancer Res* 2003;9:2327–34.
 34. Caldecott K, Jeggo P. Cross-sensitivity of gamma-ray-sensitive hamster mutants to cross-linking agents. *Mutat Res* 1991;255:111–21.
 35. Lundin C, North M, Erixon K, et al. Methyl methanesulfonate (MMS) produces heat-labile DNA damage but no detectable *in vivo* DNA double-strand breaks. *Nucleic Acids Res* 2005;33:3799–811.
 36. Roos WP, Nikolova T, Quiros S, et al. Brca2/Xrcc2 dependent HR, but not NHEJ, is required for protection against O(6)-methylguanine triggered apoptosis, DSBs and chromosomal aberrations by a process leading to SCEs. *DNA Repair (Amst)* 2009;8:72–86.
 37. Santarosa M, Dolcetti R, Magri MD, et al. BRCA1 and BRCA2 genes: role in hereditary breast and ovarian cancer in Italy. *Int J Cancer* 1999;83:5–9.
 38. Wiegant WW, Overmeer RM, Godthelp BC, van Buul PP, Zdzienicka MZ. Chinese hamster cell mutant, V-C8, a model for analysis of Brca2 function. *Mutat Res* 2006;600:79–88.
 39. Sorensen CS, Hansen LT, Dziegielewska J, et al. The cell-cycle checkpoint kinase Chk1 is required for mammalian homologous recombination repair. *Nat Cell Biol* 2005;7:195–201.
 40. Koi M, Umar A, Chauhan DP, et al. Human chromosome 3 corrects mismatch repair deficiency and microsatellite instability and reduces N-methyl-N'-nitro-N-nitrosoguanidine tolerance in colon tumor cells with homozygous hMLH1 mutation. *Cancer Res* 1994;54:4308–12.
 41. Zhang H, Marra G, Jiricny J, Maher VM, McCormick JJ. Mismatch repair is required for O(6)-methylguanine-induced homologous recombination in human fibroblasts. *Carcinogenesis* 2000;21:1639–46.
 42. Yamane K, Schupp JE, Kinsella TJ. BRCA1 activates a G2-M cell cycle checkpoint following 6-thioguanine-induced DNA mismatch damage. *Cancer Res* 2007;67:6286–92.
 43. Yuan SS, Lee SY, Chen G, Song M, Tomlinson GE, Lee EY. BRCA2 is required for ionizing radiation-induced assembly of Rad51 complex *in vivo*. *Cancer Res* 1999;59:3547–51.
 44. Pieters R, Hongo T, Loonen AH, et al. Different types of non-P-glycoprotein mediated multiple drug resistance in children with relapsed acute lymphoblastic leukaemia. *Br J Cancer* 1992;65:691–7.
 45. Petermann E, Orta ML, Issaeva N, Schultz N, Helleday T. Hydroxyurea-stalled replication forks become progressively inactivated and require two different RAD51-mediated pathways for restart and repair. *Mol Cell* 2010;37:492–502.
 46. Helleday T. Homologous recombination in cancer development, treatment and development of drug resistance. *Carcinogenesis* 2010;31:955–60.