

# Thrombin Receptors Activate Potassium and Chloride Channels

By Richard Sullivan, Diana L. Kunze, and Michael H. Kroll

We used DAMI human megakaryocytic leukemia cells to study transmembrane ion currents activated through the G-protein-coupled thrombin receptor pathway. When the cells were stimulated by thrombin receptor-activating peptide, an increase in cytosolic  $\text{Ca}^{2+}$  ( $[\text{Ca}^{2+}]_i$ ) developed as predicted by the known effect that thrombin exerts in the platelet. We then monitored the membrane potentials of individual DAMI cells during this response and observed complex, triphasic changes that could not be accounted for by  $\text{Ca}^{2+}$  fluxes alone. These consisted of rapid hyperpolarization, followed by depolarization to values more positive than the resting potential and then by slow repolarization. For the purpose of this study, we focused on the hyperpolarizing current that

developed immediately after thrombin receptor activation. This proved to be composed of (1) a  $\text{Ca}^{2+}$ -independent, outwardly rectifying  $\text{Cl}^-$  current and (2) a strongly hyperpolarizing, inwardly rectifying,  $\text{Ba}^{2+}$ -sensitive  $\text{K}^+$  current that required an increase of  $[\text{Ca}^{2+}]_i$  for activation. By analogy with their functions in other cell systems, it is logical to conclude that these prominent  $\text{K}^+$  and  $\text{Cl}^-$  conductances may serve to regulate the complex volume changes that accompany thrombin receptor activation and/or to increase the electro-motive drive that supports  $\text{Ca}^{2+}$  influx under these conditions through hyperpolarization of the cell membrane. *This is a US government work. There are no restrictions on its use.*

**T**HROMBIN IS A POTENT physiologic activator of the release reaction in platelets.<sup>1</sup> Within 5 seconds after stimulation with thrombin, complex cytoskeletal reorganization is initiated, heralded by swelling and rounding of the platelet, and culminating in the release of stored granular contents.<sup>2</sup> A biphasic increase in cytosolic calcium ( $[\text{Ca}^{2+}]_i$ ), due to the release of internally stored  $\text{Ca}^{2+}$  as well as to  $\text{Ca}^{2+}$  influx, accompanies thrombin-mediated release of  $\alpha$  granules and platelet dense bodies<sup>1</sup> and is considered to contribute significantly to the activation process.<sup>3,4</sup>

In view of its complexity, it is not surprising that thrombin-mediated platelet activation is accompanied by multiphasic changes in membrane potential that have not been fully explained at the electrophysiologic level.<sup>5,6</sup> Experiments using potential-sensitive fluorescent dyes suggested that separate phases of hyperpolarization and depolarization may be elicited, depending on the concentration of thrombin used.<sup>5</sup> One may infer from these observations that multiple classes of ion channels may operate in the plasma membrane of the platelet during the intricate process of thrombin activation.

In support of this idea is the fact that a number of types of ion channels have now been identified in the plasma membrane of platelets by various approaches, such as the introduction of membranes into artificial lipid constructs,<sup>7-9</sup> the use of fluorescent cyanine membrane potential probes,<sup>5,6</sup> and

the patch clamp technique.<sup>10-12</sup>  $\text{Ca}^{2+}$ -dependent and -independent  $\text{Cl}^-$  channels,<sup>8,10</sup> nonselective cation channels that conduct  $\text{Ca}^{2+}$  as well as monovalent cations,<sup>12</sup> and  $\text{Ca}^{2+}$ -activated<sup>13</sup> and voltage-gated  $\text{K}^+$  channels<sup>14,15</sup> have been identified using these techniques. In addition, voltage-dependent  $\text{Ca}^{2+}$  channels and  $\text{K}^+$  channels of the delayed rectifier type have been reported in guinea pig megakaryocytes.<sup>16</sup> Nonetheless, except for the observations that anion channels are important in ligand-mediated degranulation<sup>17</sup> and in homeostasis of cytosolic pH,<sup>18</sup> the roles that these channels play in platelet function have been difficult to ascertain. In addition, owing to the small size of platelets and the consequent difficulty of impaling them with a patch clamp electrode, information regarding the operation of these channels during activation with surface-acting agonists such as thrombin is quite limited.

By using patch clamp techniques, Mahaut-Smith et al<sup>11,19</sup> successfully showed that ADP opens  $\text{Ca}^{2+}$  channels in the platelet plasma membrane. Although they could not electrophysiologically detect similar single  $\text{Ca}^{2+}$  channel currents in response to perfusion with thrombin, they were nonetheless able to measure kinetically slower influxes of  $\text{Ca}^{2+}$  and  $\text{Na}^+$  under those conditions through the use of fluorescent cytosolic probes.<sup>19</sup> However, to our knowledge, the operation of hyperpolarizing  $\text{Cl}^-$  or  $\text{K}^+$  channels during thrombin-mediated activation has not been described.

With these considerations in view, we chose to examine the ion currents activated by thrombin receptors in the human megakaryocytic cell line DAMI, which was cloned in 1988 by Greenberg et al<sup>20</sup> from a patient with acute megakaryocytic leukemia. These cells, which bear morphologic and functional resemblance to normal human megakaryocytes,<sup>20,21</sup> exhibit many features that make them ideally suited for such a study. First, the cells are large and lend themselves well to electrophysiologic experiments using patch clamp electrodes. Second, they contain G-protein-coupled thrombin receptors that are expressed on human platelets.<sup>22</sup> These receptors contain an autoactivating hexapeptide sequence ( $\text{NH}^3+$  ser-phe-leu-leu-arg-asn) within residues 42 through 47. The sequence becomes exposed when an  $\text{NH}^3+$ -terminal peptide that normally sequesters these residues is cleaved by thrombin and jettisoned from the cell surface.<sup>23</sup> This unique mechanism of activation makes the receptors ideal for func-

From the Research Service, Houston VA Medical Center, and the Departments of Medicine and of Molecular Physiology and Biophysics, Baylor College of Medicine, Houston, TX.


Submitted April 27, 1995; accepted August 30, 1995.

Supported by Grants No. HL 45880 and HL 02311 from the National Institutes of Health, by the Research Service of the Department of Veterans Affairs, and the Methodist Hospital Foundation. M.H.K. is an Established Investigator of the American Heart Association.

Address reprint requests to Richard Sullivan, MD, Medical Service 111-H, Houston VA Medical Center, 2002 Holcombe, Houston, TX 77030.

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. section 1734 solely to indicate this fact.

This is a US government work. There are no restrictions on its use. 0006-4971/96/8702-0035\$0.00/0



**Fig 1.** Effect of TRAP on  $[Ca^{2+}]_i$  in fura 2-loaded DAMI cells. Cells were suspended in  $Ca^{2+}$ -free solution and constantly stirred. At the times shown by the arrows,  $CaCl_2$  and 50  $\mu$ mol/L TRAP were added. The tracing shown is representative of six similar experiments using either 50 or 100  $\mu$ mol/L TRAP.

tional studies because they can be activated specifically by synthetic peptides that contain the relevant sequence.<sup>24</sup> For these reasons, we have considered the DAMI cell to be an excellent model for the study of the electrophysiologic events that accompany hematopoietic cellular activation through G-protein-coupled receptors.

## MATERIALS AND METHODS

### Cells

The DAMI cell line was obtained from the American Type Culture Collection (Rockville, MD) with the kind permission of Dr R.I. Handin (Brigham and Women's Hospital, Boston, MA). The cells were grown in a tissue culture incubator in plastic tissue culture flasks using RPMI 1640 medium containing 10% fetal calf serum and 1% penicillin and streptomycin.

### Agonists

The amino acid sequence of the synthetic peptide TRAP was  $NH^3+$ -ser-phe-leu-leu-arg-asn-pro-asn-asp-lys-tyr-glu-pro-phe. The

sequence of the scrambled peptide that was used as a control in these experiments was  $NH^3+$ -pro-ser-gly-phe-tyr-leu-lys-leu-asp-pro-arg-asn-phe-asn. These peptides were synthesized at the Protein Chemistry Core Facility of Baylor College of Medicine by an Applied Biosciences 430A Peptide Synthesizer using Fmoc chemistry. Human  $\alpha$ -thrombin (Enzyme Research Labs Inc, South Bend, IN) was stored at  $-20^\circ C$  in a 1:1 mix of glycerol and water at a concentration of 100 U/mL and diluted immediately before use with iced saline bath solution containing 0.1% essentially fatty acid-free bovine serum albumin (Sigma, St Louis, MO).

### Measurement of Cytosolic $Ca^{2+}$ Concentrations


$[Ca^{2+}]_i$  was measured in a Deltascan spectrofluorometer (Photon Technologies International, Princeton, NJ) having dual wavelength excitation capacity. Scrape-harvested DAMI cells were loaded with 3  $\mu$ mol/L fura 2/AM (Molecular Probes, Eugene, OR) in a shaking water bath at  $37^\circ C$  for 1 to 2 hours. Unincorporated fluorophore was removed by centrifugation at 500g for 5 minutes, and DAMI cells were resuspended at a concentration of 0.5 to  $1.0 \times 10^8$ /mL in a solution containing 6 mmol/L glucose, 130 mmol/L NaCl, 9 mmol/L  $NaHCO_3$ , 10 mmol/L  $Na^+$  citrate, 10 mmol/L Tris base, 3 mmol/L KCl, 0.9 mmol/L  $MgCl_2$ , and 2 mmol/L HEPES (pH 7.35).  $[Ca^{2+}]_i$  was measured by placing a 1.5 mL volume of the cell suspension in a stirred, warmed cuvette ( $37^\circ C$ ), exciting the sample at 340 and 380 nm, and measuring emission at 510 nm. The emission ratio (340 nm/380 nm) was used to calculate  $[Ca^{2+}]_i$ , assuming a  $K_d$  for fura 2 of 224 nmol/L, as described by Grynkiewicz et al.<sup>25</sup>

### Patch Clamp Experiments

**Solutions and reagents.** Analytical grade salts were dissolved in distilled, deionized water and sterilized by millipore filtration for storage at  $4^\circ C$ . The specific ionic composition of each solution used is included in the figure legends. The osmolarity of each solution was adjusted to 260 to 300 mOsm with glucose, and the pH was titrated to 7.3 to 7.4. Nystatin (Sigma) was dissolved in dimethyl sulfoxide (25 mg/mL) immediately before use, kept shielded from light, and diluted as needed in pipette solutions. The sodium salt of 4-acetamido-4'-isothiocyanatostilbene-2,2'-disulfonic acid (SITS; Sigma) was dissolved directly into appropriate perfusion solutions before use.

**Preparation of cells.** DAMI cells from stock cultures were replated in 33-mm plastic petri dishes for up to 6 days before use. Immediately before each experiment, culture plates were washed vigorously 3 to 4 times with bath solution, and nonadherent cells were discarded.

**Patch clamp procedure.** All data were obtained using an Axo-



**Fig 2.** Effect of TRAP on the membrane potential ( $E_m$ ) of a DAMI cell. The cell was suspended in saline bath solution (137 mmol/L NaCl, 5.4 mmol/L KCl, 1 mmol/L  $MgCl_2$ , 2 mmol/L  $CaCl_2$ , and 10 mmol/L HEPES) and allowed to seal to the tip of a pipette containing 100  $\mu$ g/mL nystatin in  $K^+$  aspartate solution with  $10^{-8}$  mol/L  $Ca^{2+}$  (145 mmol/L  $K^+$  aspartate, 5 mmol/L NaCl, 0.3 mmol/L  $CaCl_2$ , 2.2 mmol/L EGTA, 2 mmol/L  $MgCl_2$ , 5 mmol/L HEPES). After perforation of the patch was achieved, current was clamped to 0 and the electrical potential of the membrane was measured. At the times shown, the cell was briefly perfused with 50  $\mu$ mol/L TRAP. Shown are the results of one of seven similar experiments.




Fig 3. Effect of TRAP on whole cell transmembrane currents in DAMI cells. Cells were suspended in saline bath solution and then sealed to the tip of a patch electrode containing  $\text{K}^+$  aspartate solution with  $10^{-8}$  mol/L  $\text{Ca}^{2+}$ . After break-in by either suction or nystatin, voltage was clamped at  $-60$  mV and stepped in  $10$  mV increments from  $-120$  mV to  $+60$  mV. The current that flowed across the plasma membrane in response to the applied voltage steps was then recorded. In the left-sided panels, steady-state current was plotted as a function of applied voltages. After measuring the resting current ( $\circ$ ) in each experiment, the cell was perfused at time 0 with  $50$   $\mu\text{mol/L}$  (A and C) or  $100$   $\mu\text{mol/L}$  (B) TRAP suspended in saline bath solution. ( $\square$ ) Currents that developed after perfusion. In (A), the cells were bathed in saline bath solution containing  $2$  mmol/L  $\text{Ca}^{2+}$ , and currents were measured through a perforated patch using nystatin. In (B), the same bath solution was used, but currents were measured through a broken patch. In (C), a  $\text{Ca}^{2+}$ -free saline bath solution containing  $2.2$  mmol/L EGTA was used, and currents were measured through a broken patch. In the right-sided panels, the bar graphs show reversal potentials before and after TRAP. All data points represent the means  $\pm$  standard errors of the mean from combined experiments.

patch-1D patch clamp amplifier (Axon Instruments, Foster City, CA) and recorded and analyzed by computer using Pclamp software (Axon Instruments). Electrodes were pulled from 7052 glass capillary tubes (Garner, Claremont, CA) to a pipette resistance of 2 to 6 M $\Omega$ . Most experiments were performed with broken patches in the whole cell configuration using the voltage clamp mode. Cells were clamped at  $-60$  mV and switched at 1-second intervals through a series of voltage steps beginning at  $-120$  mV and proceeding at  $10$ -mV increments to  $+60$  mV. The duration of each step was 0.6 second. Experiments were performed at room temperature.

In certain experiments, cells were voltage-clamped using the perforated patch technique,<sup>26</sup> in which the pipette solution contained  $100$   $\mu\text{g/mL}$  nystatin. A tight seal was allowed to form between the pipette tip and the cell membrane, after which time perforation of the membrane by the antibiotic usually occurred within 3 to 10 minutes.

Membrane potential measurements were performed in the current-clamp mode using the perforated patch technique described above. The cells were allowed to seal to the pipette tip under voltage-clamp conditions. After perforation of the membrane by nystatin,

the current was clamped to 0. Junction potentials between perfusion and bath solutions were less than 5 mV and were not corrected.


#### Statistics

Combined data were expressed as means  $\pm$  standard errors of the mean. Corresponding means were analyzed using the Student's *t*-test.

## RESULTS

In our initial experiments, we examined the effect of TRAP on  $[\text{Ca}^{2+}]_i$  in DAMI cells loaded with the fluorescent probe fura 2. We found that, when the cells were suspended in medium containing  $2$  mmol/L  $\text{Ca}^{2+}$ , TRAP induced an increase in  $[\text{Ca}^{2+}]_i$  that was characterized by an initial peak within 30 to 45 seconds, followed by a slowly decaying sustained increase lasting several minutes (Fig 1). The initial increase in  $[\text{Ca}^{2+}]_i$  after TRAP was maximal at  $50$   $\mu\text{mol/L}$  and  $100$   $\mu\text{mol/L}$ ,  $\approx 65\%$  of maximum at  $10$   $\mu\text{mol/L}$ ,  $7E40\%$

**Fig 4.** Effect of  $\alpha$ -thrombin on whole cell currents in DAMI cells. Cells were subjected to the same protocol as described for Fig 3B except that  $\alpha$ -thrombin was used instead of TRAP and 0.1% fatty acid-free bovine serum albumin was added to the perfusion (and in some cases the bath) solutions. After a broken patch was established, the cell was briefly perfused with  $\alpha$ -thrombin (1 U/mL). The experiment shown in (A) and (B) is representative of the approximately two thirds of cells that responded to  $\alpha$ -thrombin. In (A), the resting current of the impaled DAMI cell is shown. (B) shows the hyperpolarizing current that developed  $\approx 30$  seconds after perfusion with  $\alpha$ -thrombin. By 3 minutes, the current had returned to the resting level (data not shown). In (C), whole cell currents before ( $\blacktriangle$ ) and  $\approx 30$  seconds after ( $\bullet$ ) perfusion with  $\alpha$ -thrombin are plotted as a function of applied voltage. Shown are means  $\pm$  standard errors of the mean of data obtained from 13 cells that developed hyperpolarizing current after perfusion.



at 5  $\mu\text{mol/L}$ , and  $\approx 5\%$  at 1  $\mu\text{mol/L}$ . Based on these results, 50  $\mu\text{mol/L}$  or 100  $\mu\text{mol/L}$  concentrations of TRAP were used in the remaining experiments to achieve an optimal response.

When we examined the effect of TRAP on the membrane potential of DAMI cells, we repeatedly observed the complex response shown in Fig 2. Although the resting potential varied somewhat from cell to cell, a typical response is shown in this experiment. Within seconds after the cell was briefly perfused with TRAP, the membrane hyperpolarized from resting values ( $-40 \pm 3.6$  mV [mean  $\pm$  SEM];  $n = 7$ ) to a maximum of  $-64 \pm 2.1$  mV. Thereafter, over 1 to 2 minutes, the cell slowly depolarized to potentials more positive than the resting potential ( $-31 \pm 1.9$  mV). During this phase, wide oscillations in potential were common. Thereafter, the cell typically hyperpolarized slowly to the original resting potential. As shown in Fig 2, some of the cells were capable of responding two or three times to TRAP, as assessed by changes in membrane potential.

We then proceeded to examine whole cell transmembrane ion currents in DAMI cells, focusing on the immediate hy-

perpolarization that followed cellular activation with TRAP. For these experiments, voltage across the cell membrane was clamped at  $-60$  mV through patches broken by suction or perforated with nystatin. The current that flowed across the cell membrane in response to the voltage protocol described in Materials and Methods was measured and expressed as a function of applied voltage.

In Fig 3, the change in whole cell current after stimulation of the cells with TRAP is shown under three conditions in which the increase in  $[\text{Ca}^{2+}]_i$  was manipulated differently. In Fig 3A, the cells were suspended in saline bath solution containing 2 mmol/L  $\text{Ca}^{2+}$ , and whole cell currents were measured through a patch perforated with nystatin, which permitted  $[\text{Ca}^{2+}]_i$  to increase unopposed after activation of the cell with TRAP. Under these conditions, the current that developed was strongly hyperpolarizing and exhibited prominent inward and outward components. In Fig 3B, the cells were suspended in the same saline bath solution containing 2 mmol/L  $\text{Ca}^{2+}$ , but the currents were measured through broken patches through which the interiors of the cells had been briefly dialyzed with  $\text{K}^+$  aspartate solution




Fig 5. Effect of the control peptide on whole cell currents. Cells were suspended in saline bath solution and impaled with electrodes containing  $K^+$  aspartate solution with  $10^{-8}$  mol/L  $Ca^{2+}$ . Currents were recorded before ( $\circ$ ) and after ( $\square$ ) perfusion with the scrambled peptide. Four cells from two culture dishes were studied in this manner; the means  $\pm$  standard errors of the mean obtained from these data are shown in the main graph. In one of the dishes, a separate cell was then perfused with  $100 \mu\text{mol/L}$  TRAP (inset).


containing  $Ca^{2+}$  buffered to  $10^{-8}$  mol/L. Under these conditions, both the inward and outward components of the current were quantitatively reduced, but whereas the change in reversal potential was less dramatic, the net effect was still hyperpolarization. In Fig 3C, the cells were bathed in saline solution containing no added  $Ca^{2+}$  and  $2.2$  mmol/L EGTA, and the currents were measured through broken patches containing  $10^{-8}$  mol/L  $Ca^{2+}$  as in Fig 3B. Under these conditions, the current that developed after activation of the cells with TRAP rectified outwardly, and hyperpolarization was no longer observed.

The whole cell current response to  $\alpha$ -thrombin was qualitatively similar to the response to TRAP (Fig 4). However, although it was unusual for hyperpolarizing currents not to develop in cells perfused with TRAP, only about two thirds of the cells responded to  $\alpha$ -thrombin in our experiments. Although TRAP was extremely stable at room temperature in protein-free solutions,  $\alpha$ -thrombin lost activity after approximately 30 minutes and required albumin to prevent its adherence to the tubing of the perfusion apparatus.

Two sets of control experiments were performed to assess the specificity of the effect of TRAP on whole cell currents. The first set tested the effect of the turbulence produced by the perfusion stream. In these experiments, four cells were suspended in saline bath solution, impaled with a patch pipette containing  $K^+$  aspartate solution, and then perfused with saline bath solution under temperature and flow conditions identical to those shown above in which TRAP was

used. No current was produced in any of these four cells simply by subjecting them to a jet of the bath solution, whereas a single cell in the same dish responded typically to TRAP (data not shown). In the second set of controls, the cells were perfused under identical conditions with a  $100 \mu\text{mol/L}$  solution of a peptide of the same length as TRAP that which contained amino acids scrambled into a nonsense sequence. In none of these cells did whole cell current develop after perfusion with the scrambled peptide that was different from that in the resting cell (Fig 5). As shown in the inset in Fig 5, a single cell that was in the same dish as two of the four cells that had been perfused with the scrambled peptide was then perfused with TRAP, a maneuver that resulted, as predicted, in a measurable increase in current.

The outwardly rectifying current induced by TRAP under conditions that prevented an increase in  $[Ca^{2+}]_i$  appeared to be due entirely to the opening of  $Cl^-$  channels (Fig 6). In this experiment, the cells were bathed in saline bath solution containing no added  $Ca^{2+}$  and  $2.2$  mmol/L EGTA and impaled with an electrode containing  $K^+$  aspartate solution (conditions identical to those in Fig 3C). After measuring steady state current (Fig 6A), the cell was perfused with the same bath solution containing and  $50 \mu\text{mol/L}$  TRAP (Fig 6B); under these conditions, an outwardly rectifying current immediately developed. While the current was active, the cell was quickly perfused with bath solution containing  $50 \mu\text{mol/L}$  TRAP and  $200 \mu\text{mol/L}$  SITS. As shown in Fig 6C, the current that developed in response to TRAP under these




**Fig 6.** Isolation of the Cl<sup>-</sup> current activated by TRAP. In these experiments, the cells were suspended in Ca<sup>2+</sup>-free saline bath solution containing 2.2 mmol/L EGTA to inhibit the K<sup>+</sup> current activated by TRAP. Each cell was impaled with a pipette containing K<sup>+</sup> aspartate solution with 10<sup>-8</sup> mol/L Ca<sup>2+</sup>. After measuring baseline currents (A), the cell was briefly perfused with bath solution containing 50 μmol/L TRAP (B). Before the current activated by TRAP had dissipated, the cell was constantly perfused with bath solution 50 μmol/L TRAP and 200 μmol/L SITS (C). Finally, the cell was reperfused with SITS-free bath solution containing TRAP (D). On the right are shown the means and standard errors of the means for three cells subjected to this protocol.

conditions was inhibited by SITS. Finally, the cell was reperfused with SITS-free bath solution containing TRAP (Fig 6D). The outward current was reduced by perfusing the cells with bath solutions in which Cl<sup>-</sup> was replaced by aspartate or SO<sub>4</sub><sup>2-</sup> anion (data not shown).

Under conditions that permitted [Ca<sup>2+</sup>]<sub>i</sub> to increase after TRAP activation, the strong hyperpolarization that developed appeared to be due to a K<sup>+</sup> current that rectified inwardly. In Fig 7, cells were suspended in saline bath solution containing 2 mmol/L Ca<sup>2+</sup> and impaled with pipettes filled with K<sup>+</sup> aspartate solution containing 10<sup>-8</sup> mol/L Ca<sup>2+</sup>. The cells were first perfused with TRAP in saline bath solution, and a hyperpolarizing current quickly developed. The cells were then immediately perfused with TRAP in high K<sup>+</sup> saline bath solution in which the

concentrations of K<sup>+</sup> and Na<sup>+</sup> were reversed and the concentration of K<sup>+</sup> approximated that of the cell interior. Under these conditions, the reversal potential shifted to 0, and the overall current exhibited mild inward rectification.

The strong inward rectification of the K<sup>+</sup> current activated by elevated [Ca<sup>2+</sup>]<sub>i</sub> is shown in Fig 8. In these cells, the current was elicited in the absence of TRAP by dialyzing the cell with K<sup>+</sup> aspartate solution containing 2 mmol/L Ca<sup>2+</sup>. This current was not present (in the absence of TRAP) when cells were dialyzed with our usual pipette solutions containing 10<sup>-8</sup> mol/L Ca<sup>2+</sup> and was inconsistently seen when cells were dialyzed with 10<sup>-6</sup> mol/L Ca<sup>2+</sup> (data not shown). As shown in Fig 8B and C, the current was reversibly inhibited by 1 mmol/L Ba<sup>2+</sup>.



**Fig 7.** Effect of external  $K^+$  on TRAP-induced whole cell currents. DAMI cells were suspended in saline bath solution (137 mmol/L NaCl, 5.4 mmol/L KCl, 1 mmol/L  $MgCl_2$ , 2 mmol/L  $CaCl_2$ , 10 mmol/L HEPES) and impaled with a patch clamp electrode containing  $K^+$  aspartate solution with  $10^{-8}$  mol/L  $Ca^{2+}$ . (○) Baseline currents. The cell was then perfused with saline bath solution containing 50  $\mu$ mol/L TRAP (□). While the current was fully active, the cell was perfused with 50  $\mu$ mol/L TRAP in a high  $K^+$  saline bath solution (137 mmol/L KCl, 5.4 mmol/L NaCl, 1 mmol/L  $MgCl_2$ , 2 mmol/L  $CaCl_2$ , 10 mmol/L HEPES) ( $\Delta$ ). Shown are means  $\pm$  standard errors of the mean of 6 cells.

## DISCUSSION

Our experiments showed an unexpectedly complex series of membrane potential changes after the activation of thrombin receptors. At least three phases were distinct. Immediate, pronounced hyperpolarization was followed by slow depolarization to levels more positive than the prestimulation potential; finally, the cell gradually repolarized to the resting potential. Wide oscillations were common towards the end of the depolarization phase. These findings predict that multiple classes of ion channels may operate during cellular activation via thrombin receptors. For the purposes of this study, we focused on the initial phase of hyperpolarization, which we found to be due to the development of both  $Cl^-$  and  $K^+$  currents.

The  $Cl^-$  current, which rectified outwardly and was inhibited by the anion channel blocker SITS, developed instantly on contact of TRAP with the cell. This current had little detectable effect on the resting potential and appeared to develop independently of a substantial increase in  $[Ca^{2+}]_i$ . The more strongly hyperpolarizing  $K^+$  current, which exhibited inward rectification and was blocked by 1 mmol/L  $Ba^{2+}$  in the bath, often lagged behind the  $Cl^-$  current by 5 to 10 seconds. Unlike the  $Cl^-$  current, the activation of the  $K^+$  current depended on an increase in  $[Ca^{2+}]_i$ . Together, these currents effected a strong hyperpolarization of the cell during the earliest stage of thrombin receptor activation.


The majority of  $Ca^{2+}$ -activated  $K^+$  channels described to

date exhibit either outward rectification (eg, large conductance, charybdotoxin-sensitive  $K^+$  channels)<sup>27</sup> or ohmic conductance (eg, small conductance, apamin-sensitive  $K^+$  channels).<sup>28</sup> However, many types of blood cells, including lymphocytes,<sup>29</sup> macrophages,<sup>30,31</sup> erythrocytes,<sup>32</sup> and dimethyl sulfoxide (DMSO)-induced HL60 cells,<sup>33</sup> have now been found to express  $Ca^{2+}$ -dependent, inwardly rectifying  $K^+$  channels with conductances of 25 to 50 pS. In addition, Mahaut-Smith<sup>13</sup> recently reported the discovery of 30 pS  $Ca^{2+}$ -activated  $K^+$  channels in human platelets that may belong to this group of channels. Because of their rectification properties, these channels remain open over a wide range of negative potentials. With the possible exception of the neutrophil,<sup>34</sup> the membrane potentials of most blood cells probably remain negative even during cellular activation.<sup>35-39</sup> For that reason, this class of  $K^+$  channel would seem to be ideally suited to induce continuous hyperpolarization of the membrane in response to the sustained increases in  $[Ca^{2+}]_i$  that accompany phospholipase C activation. In addition, Gallin<sup>40</sup> has postulated that these channels may account for the wide oscillations in membrane potential that she<sup>41</sup> and others<sup>42</sup> have described in macrophages and that we have observed in DAMI cells (Fig 2).

We were interested to observe that supraphysiologic concentrations of  $[Ca^{2+}]_i$  were required to activate the inwardly rectifying  $K^+$  current consistently in the absence of TRAP. Nonetheless, TRAP stimulation predictably induced the current when the bath solution contained 2 mmol/L  $Ca^{2+}$ , even though the increase in  $[Ca^{2+}]_i$  produced by TRAP was unlikely to have exceeded  $10^{-6}$  mol/L under the conditions of our experiments. We interpret this finding to indicate either that very high  $Ca^{2+}$  concentrations may be achieved in compartments within close proximity to the ion channels or that this particular  $K^+$  current may be regulated by multiple cooperative chemical messengers that converge at the channel level, as has been reported to be the case for  $K^+$  channels in endothelial cells.<sup>43</sup>

The precise physiologic role of the  $K^+$  and  $Cl^-$  channels that open in DAMI cells after thrombin receptor activation is unclear. However, considerable evidence, summarized by Grinstein and Foskett,<sup>44</sup> indicates that several types of hematopoietic cells, including human platelets,<sup>45</sup> respond to cellular swelling induced by hypoosmolar stress by activating  $K^+$  and  $Cl^-$  conductances. An electrically neutral efflux of  $K^+$  and  $Cl^-$  ions, together with osmotically obligated water molecules, subsequently permits detumescence of the cell. Because platelets are known to swell within seconds after stimulation with thrombin, it is conceivable that the concomitant opening of  $K^+$  and  $Cl^-$  channels that we observed may serve to regulate cell volume during the activation process.

Another potential role for these currents may be to support  $Ca^{2+}$  influx through hyperpolarization of the plasma membrane during cellular activation. In 1988, Penner et al<sup>46</sup> showed that, when rat peritoneal mast cells were stimulated by substance P, the resultant biphasic increase in  $[Ca^{2+}]_i$  was accompanied by an outwardly rectifying, cAMP-dependent  $Cl^-$  current. Because the increase in  $[Ca^{2+}]_i$  was enhanced at negative membrane potentials in these cells, they concluded that  $Ca^{2+}$  influx was sustained through the hyperpolarizing effects of the  $Cl^-$  current. In an analogous manner,



**Fig 8.** Inward rectification and Ba<sup>2+</sup> sensitivity of Ca<sup>2+</sup>-induced K<sup>+</sup> current. In these experiments, DAMI cells were suspended in K<sup>+</sup> aspartate bath solution (150 mmol/L K<sup>+</sup> aspartate, 2 mmol/L MgCl<sub>2</sub>, 2 mmol/L CaCl<sub>2</sub>, 10 mmol/L HEPES) and dialyzed with pipettes containing the same solution. The voltage was held at 0, and the cells were stepped through a series of brief voltage changes from -120 mV to +60 mV. Shown on the left are current traces from a single representative cell. On the right, the corresponding current-voltage relationship is shown for three separate cells (mean ± standard error of the mean). (A) shows the resting current. In (B), the cell was perfused with bath solution containing 1 mmol/L BaCl<sub>2</sub>. In (C), the cell was perfused with Ba<sup>2+</sup>-free bath solution.

Ca<sup>2+</sup> influx is well recognized to contribute to the increase in [Ca<sup>2+</sup>]<sub>i</sub> after activation of thrombin receptors.<sup>1,3,4</sup> Jenkins et al<sup>47</sup> have recently shown clearly that TRAP induces both internal Ca<sup>2+</sup> release and Ca<sup>2+</sup> influx across the plasma membranes of human osteoblast-like cells that bear the G-protein-coupled thrombin receptor. The Ca<sup>2+</sup>-activated K<sup>+</sup> current that we identified in DAMI cells strongly hyperpolarized the cell, as assessed by its effect on the reversal potential of whole cell current. Although the TRAP-induced Cl<sup>-</sup> current did not have a similar hyperpolarizing effect on the resting membrane potential, possibly owing to its strong outward rectification, it is likely that this current would serve to counteract depolarization caused by the influx of Ca<sup>2+</sup> and other cations. We postulate, therefore, that K<sup>+</sup> and Cl<sup>-</sup> currents in DAMI cells may function in a manner similar to that described by Penner et al<sup>46</sup> by increasing the electromotive drive that supports Ca<sup>2+</sup> influx through ligand-gated Ca<sup>2+</sup> channels during cellular activation.

#### ACKNOWLEDGMENT

The authors thank Elkin Romero, Marjorie Withers, and Ketia Francis for excellent technical assistance.

#### REFERENCES

1. Siess W: Molecular mechanisms of platelet activation. *Physiol Rev* 69:58, 1989
2. Stenberg PE, Bainton DF: in Phillips DR, Shuman MA (eds): *Biochemistry of Platelets*. Orlando, FL, Academic, 1986, p 257
3. Rink TJ, Sage SO: Calcium signaling in human platelets. *Annu Rev Physiol* 52:431, 1990
4. Sage SO, Sargeant P, Merritt JE, Mahaut-Smith MP, Rink TJ: Agonist-evoked Ca<sup>2+</sup> entry in human platelets. *Biochem J* 285:341, 1992
5. Horne WC, Simons ER: Probes of transmembrane potentials in platelets: Changes in cyanine dye fluorescence in response to aggregation stimuli. *Blood* 51:741, 1978
6. Greenberg-Sepersky SM, Simons ER: Cation gradient depen-

- dence of the steps in thrombin stimulation of human platelets. *J Biol Chem* 259:1502, 1984
7. Fujimoto T, Fujimura K, Kuramoto A: Electrophysiological evidence that glycoprotein IIb-IIIa complex is involved in calcium channel activation on human platelet plasma membrane. *J Biol Chem* 266:16370, 1991
  8. Manning SD, Williams AJ: Conduction and blocking properties of a predominantly anion-selective channel from human platelet surface membrane reconstituted into planar phospholipid bilayers. *J Membr Biol* 109:113, 1989
  9. Rybak ME, Renzulli LA, Bruns MJ, Cahaly DP: Platelet glycoproteins IIb and IIIa as a calcium channel in liposomes. *Blood* 72:714, 1988
  10. Mahaut-Smith MP: Chloride channels in human platelets: Evidence for activation by internal calcium. *J Membr Biol* 118:69, 1990
  11. Mahaut-Smith MP: Receptor-activated single channels in intact human platelets. *J Biol Chem* 265:10479, 1990
  12. Avdonin PV, Cheglakov IB, Tkachuk VA: Stimulation of non-selective cation channels providing  $Ca^{2+}$  influx into platelets by platelet-activating factor and other aggregation inducers. *Eur J Biochem* 198:267, 1991
  13. Mahaut-Smith MP: Calcium-activated potassium channels in human platelets. *J Physiol* 484.1:15, 1995
  14. Mahaut-Smith MP, Rink TJ, Collins SC, Sage SO: Voltage-gated potassium channels and the control of membrane potential in human platelets. *J Physiol* 428:723, 1990
  15. Maruyama Y: A patch-clamp study of mammalian platelets and their voltage-gated potassium current. *J Physiol* 391:467, 1987
  16. Kawa K: Voltage-gated calcium and potassium currents in megakaryocytes dissociated from guinea-pig bone marrow. *J Physiol* 431:187, 1990
  17. Ozaki Y, Matsumoto Y, Yatomi Y, Higashihara M, Kariya T, Shoji K: Effects of five anion channel blockers on thrombin- and ionomycin-activated platelet functions. *Biochem Pharmacol* 13: 2147, 1989
  18. Ozaki Y, Yatomi Y, Kariya T, Kume S: Anion channels contribute to the regulation of intracellular pH in human platelets. *Thromb Res* 53:221, 1989
  19. Mahaut-Smith MP, Sage SO, Rink TJ: Rapid ADP-evoked currents in human platelets recorded with the nystatin permeabilized patch technique. *J Biol Chem* 267:3060, 1992
  20. Greenberg SM, Rosenthal DS, Greeley TA, Tantravahi R, Handin RI: Characterization of a new megakaryocytic cell line: The Dami cell. *Blood* 72:1968, 1988
  21. Greenberg SM, Chandrasekhar C: Hematopoietic factor-induced synthesis of von Willebrand factor by the Dami human megakaryoblastic cell line and by normal human megakaryocytes. *Exp Hematol* 19:53, 1991
  22. Vu T-KH, Hung DT, Wheaton VI, Coughlin SR: Molecular cloning of a functional thrombin receptor reveals a novel proteolytic mechanism of receptor activation. *Cell* 64:1057, 1991
  23. Vu T-KH, Wheaton VI, Hung DT, Charo I, Coughlin SR: Domains specifying thrombin-receptor interaction. *Nature* 353:674, 1991
  24. Coughlin SR, Vu T-KH, Hung DT, Wheaton VI: Characterization of a functional thrombin receptor: Issues and opportunities. *J Clin Invest* 89:351, 1992
  25. Grynkiewicz G, Poenie M, Tsien RY: A new generation of  $Ca^{2+}$  indicators with greatly improved fluorescence properties. *J Biol Chem* 260:3440, 1985
  26. Falke LC, Gillis KD, Pressel DM, Misler S: 'Perforated patch recording' allows long-term monitoring of metabolite-induced electrical activity and voltage-dependent  $Ca^{2+}$  currents in pancreatic islet B cells. *FEBS Lett* 251:167, 1989
  27. Blatz AL, Magleby KL: Calcium-activated potassium channels. *Trends Neurosci* 10:463, 1987
  28. Romey G, Lazdunski M: The coexistence in rat muscle cells of two distinct classes of  $Ca^{2+}$ -dependent  $K^+$  channels with different pharmacological properties and different physiological function. *Biochem Biophys Res Commun* 118:669, 1984
  29. Mahaut-Smith MP, Schlichter LC:  $Ca^{2+}$ -activated  $K^+$  channels in human B lymphocytes and rat thymocytes. *J Physiol Lond* 415:69, 1989
  30. Gallin EK: Evidence for a Ca-activated inwardly rectifying K channel in human macrophages. *Am J Physiol* 257 (Cell Physiol 25):C77, 1989
  31. Hara N, Ichinose M, Sawada M, Imai K, Maeno T: Activation of single  $Ca^{2+}$ -dependent  $K^+$  channel by external ATP in mouse macrophages. *FEBS Lett* 267:281, 1990
  32. Grygorczyk R, Schwarz W: Properties of the  $Ca^{2+}$ -activated  $K^+$  conductance of human red cells as revealed by the patch-clamp technique. *Cell Calcium* 4:499, 1983
  33. Varnai P, Demaurex N, Jaconi M, Schlegel W, Lew DP, Krause KH: Highly co-operative  $Ca^{2+}$  activation of intermediate-conductance  $K^+$  channels in granulocytes from a human cell line. *J Physiol* 472:373, 1993
  34. Demaurex N, Schrenzel J, Jaconi ME, Lew DP, Krause KH: Proton channels, plasma membrane potential, and respiratory burst in human neutrophils. *Eur J Haematol* 51:309, 1993
  35. Friedhoff LT, Sonenberg M: The membrane potential of human platelets. *Blood* 61:180, 1983
  36. Wencel-Drake JD, Feinberg H: ADP induced depolarization of human platelet membrane potential. *Thromb Haemost* 53:75, 1985
  37. Fine BP, Hansen KA, Salcedo JR, Aviv A: Calcium-activated potassium channels in human platelets. *Proc Soc Exp Biol Med* 192:109, 1989
  38. Sagi-Eisenberg R, Pecht I: Membrane potential changes during IgE-mediated histamine release from rat basophilic leukemia cells. *J Membr Biol* 75:97, 1983
  39. Kanner BI, Metzger H: Crosslinking of the receptors for immunoglobulin E depolarizes the plasma membrane of rat basophilic leukemia cells. *Proc Natl Acad Sci USA* 80:5744, 1983
  40. Gallin EK: Ion channels in leukocytes. *Physiol Rev* 71:775, 1991
  41. Gallin EK, Wiederhold ML, Lipsky PE, Rosenthal AS: Spontaneous and induced membrane hyperpolarizations in macrophages. *J Cell Physiol* 86:653, 1975
  42. Persechini PM, Araujo EG, Oliveira-Castro GM: Electrophysiology of phagocytic membranes; induction of slow membrane hyperpolarizations in macrophages and macrophage polykaryons by intracellular calcium injection. *J Membr Biol* 61:81, 1981
  43. Vaca L, Schilling WP, Kunze DL: G-protein-mediated regulation of a  $Ca^{2+}$ -dependent  $K^+$  channel in cultured vascular endothelial cells. *Pflügers Arch* 422:66, 1992
  44. Grinstein S, Foskett JK: Ionic mechanisms of cell volume regulation in leukocytes. *Annu Rev Physiol* 52:399, 1990
  45. Livne A, Grinstein S, Rothstein A: Volume-regulating behavior of human platelets. *J Cell Physiol* 131:354, 1987
  46. Penner R, Matthews G, Neher E: Regulation of calcium influx by second messengers in rat mast cells. *Nature* 334:499, 1988
  47. Jenkins AL, Bootman MD, Berridge MJ, Stone SR: Differences in intracellular calcium signaling after activation of the thrombin receptor by thrombin and agonist peptide in osteoblast-like cells. *J Biol Chem* 269:17104, 1994