

## COVID-19 Research Tools

Defeat the SARS-CoV-2 Variants

InVivoGen

# The Journal of Immunology

RESEARCH ARTICLE | DECEMBER 01 2002

## The Role of FcγR Signaling in the K/B × N Serum Transfer Model of Arthritis<sup>1</sup>

FREE

Maripat Corr; ... et. al

*J Immunol* (2002) 169 (11): 6604–6609.

<https://doi.org/10.4049/jimmunol.169.11.6604>

### Related Content

Destructive Arthritis in the Absence of Both FcγRI and FcγRIII

*J Immunol* (April,2008)

Immunization with Glucose-6-Phosphate Isomerase Induces T Cell-Dependent Peripheral Polyarthritis in Genetically Unaltered Mice

*J Immunol* (April,2004)

The Cellular Source and Target of IL-21 in K/BxN Autoimmune Arthritis

*J Immunol* (September,2013)

# The Role of Fc $\gamma$ R Signaling in the K/B $\times$ N Serum Transfer Model of Arthritis<sup>1</sup>

Maripat Corr<sup>2</sup> and Brian Crain

Spontaneous arthritis in the KRN transgenic mouse (K/BxN) model is due to the autoreactivity of the transgenic TCR and subsequent induction of autoantibodies directed against glucose-6-phosphate isomerase. These autoantibodies transfer clinically apparent arthritis into most recipient mouse strains and systemic catabolism of the transferred Abs attenuates paw swelling. Although mice deficient in the common  $\gamma$ -chain of the Fc $\gamma$ R did not show clinical synovitis after receiving K/BxN sera, erosive lesions in the bone still developed. Further analysis demonstrated that Fc $\gamma$ RII<sup>-/-</sup> mice manifested accelerated arthritis whereas the Fc $\gamma$ RIII<sup>-/-</sup> mice had a more slowly progressing arthritis. Paw swelling required Fc $\gamma$ R expression by bone marrow-derived cells and mast cells substantially contributed to the acute phase of paw swelling. In the K/BxN serum transfer model of arthritis, there is a clinically apparent acute phase, which is modulated by Fc $\gamma$ RII and Fc $\gamma$ RIII, and a subacute component, which results in bone erosion, even in the absence of Fc $\gamma$ R signaling. *The Journal of Immunology*, 2002, 169: 6604–6609.

Spontaneous severe arthritis develops in KRN TCR transgenic mice that carry the IA<sup>g7</sup> MHC class II allele (K/BxN) (1, 2). This model is similar to human rheumatoid arthritis (RA)<sup>3</sup> in that both are chronic symmetric joint diseases with pannus formation, and destructive bone and cartilage erosion of predominantly the distal joints. Arthritis in this model has been shown to be dependent on MHC class II specificity and transgenic T cells to initiate disease, and B cells to secrete pathogenic Abs that maintain disease (1–3). These autoantibodies induce arthritis upon transfer to most naive syngeneic and allogeneic hosts (2).

The factors required in the development of paw swelling in this model include the presence of neutrophils, the alternative complement pathway, IL-1 signaling, and FcRs (4–8). A recent genetic screen for elements associated with this arthritis revealed high logarithm of odds scores for a region on chromosome 2 centering on the C5 locus and a broad region on chromosome 1 (4). This area on chromosome 1 contains several loci related to the family of FcRs including FcεRI and Fc $\gamma$ RII. Genetic mapping of collagen-induced arthritis in a cross with nonobese diabetic (NOD) and C57BL/10 mice also localized associated gene regions encoding C5 and Fc $\gamma$ RIIb (9). However, in the K/BxN model, the amount of serum transferred to Fc $\gamma$ RII<sup>-/-</sup> mice was reported to influence the development of joint swelling similar to that of wild-type controls (4).

We previously reported the lack of paw swelling in mice deficient for the common  $\gamma$ -chain of the FcR (Fc $\gamma$ R) following transfer of serum from arthritic mice (5). To further analyze the influence

of FcR family members in the development of this arthritis, we have injected mice lacking Fc $\gamma$ RII or Fc $\gamma$ RIII with K/BxN serum and found that the Fc $\gamma$ RIII-deficient mice had markedly diminished paw swelling. In contrast, the transfer of autoantibodies into mice lacking Fc $\gamma$ RII resulted in an accelerated onset and severity of arthritis. Using bone marrow chimeras, the presence of paw swelling was associated with the Fc $\gamma$ R expression of cells that arise in the bone marrow and not joint-associated tissue. Furthermore, mast cell-deficient mice had markedly attenuated synovitis, suggesting that these cells are also involved in the inflammatory cascade in the joints of affected mice.

## Materials and Methods

### Mice

KRN TCR transgenic mice were a kind gift from Drs. D. Mathis and C. Benoist (Harvard Medical School, Boston, MA) and the Institut de Génétique et de Biologie Moléculaire et Cellulaire (Strasbourg, France) (2) and were maintained on a C57BL/6 background (K/B). Arthritic mice were obtained by crossing K/B with NOD/Lt (N) animals (K/BxN). Progeny bearing the V $\beta$ 6 transgenic TCR were identified by cytofluorometry of PBL using anti-CD4 PE (Caltag Laboratories, Burlingame, CA) and anti-V $\beta$ 6 FITC (BD PharMingen, San Diego, CA)-labeled Abs. C57BL/6, BALB/c, WBB6F<sub>1</sub>, WBB6F<sub>1</sub>-Kit<sup>W</sup>/Kit<sup>Wv</sup>, Fc $\gamma$ RIII<sup>-/-</sup> (C57BL/6 background) (10),  $\beta_2$ -microglobulin ( $\beta_2m$ )<sup>-/-</sup> (BALB/c background) (11), CD1<sup>-/-</sup> (BALB/c background) (12), Tap-1<sup>-/-</sup> (C57BL/6 background) (13), (C57BL/6J  $\times$  129S1/SvImJ)F<sub>2</sub>, 129/Sv, and NOD/Lt mice were purchased from The Jackson Laboratory (Bar Harbor, ME). Fc $\gamma$ R<sup>-/-</sup> and Fc $\gamma$ RII<sup>-/-</sup> mice were a generous gift from Dr. J. Ravetch (Rockefeller University, New York, NY) and provided by Dr. H. Tighe (University of California, San Diego, CA) (14, 15). Tail samples of four Fc $\gamma$ RII<sup>-/-</sup> mice from our colony were analyzed for 84 genome-wide polymorphic microsatellite markers (Charles River Breeding Laboratories, Troy, NY). In these mice, 38.95–40.70% of the polymorphisms were associated with the C57BL/6 strain as compared with 129/Sv. Mice were bred and maintained under standard conditions in the University of California, San Diego Animal Facility that is accredited by the American Association for Accreditation of Laboratory Animal Care. All animal protocols receive prior approval by the institutional review board.

### Bone marrow chimeras

Adult mice were lethally irradiated with 800 rad. Bone marrow cells harvested from the femurs and tibia of donors were washed in serum-free medium and counted. The recipients were injected with 10<sup>7</sup> cells in 100  $\mu$ l of serum-free RPMI 1640 i.v. After 8 wk, the mice were checked for reconstitution by fluorocytometry. All Fc $\gamma$ R<sup>-/-</sup> that received C57BL/6 bone marrow had >99% of the peripheral F4/80<sup>+</sup> (Serotec, Oxford, U.K.)

Division of Rheumatology, Allergy, and Immunology and Sam and Rose Stein Institute for Research on Aging, University of California, San Diego, La Jolla, CA 92093

Received for publication March 8, 2002. Accepted for publication September 26, 2002.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

<sup>1</sup> This work was supported by Grants AI10682, AR44850, and AR47360 from the National Institutes of Health.

<sup>2</sup> Address correspondence and reprint requests to Dr. Maripat Corr, Division of Rheumatology, Allergy, and Immunology and Sam and Rose Stein Institute for Research on Aging, University of California, San Diego, La Jolla, CA 92093-0663. E-mail address: mcorr@ucsd.edu

<sup>3</sup> Abbreviations used in this paper: RA, rheumatoid arthritis; G6PI, glucose-6-phosphate isomerase.

mononuclear cells stain for CD16/32 (BD PharMingen), whereas the level of CD16/32 staining dropped to <40% in the C57BL/6 recipients of Fc $\gamma$ R<sup>-/-</sup> bone marrow.

### Serum transfer and arthritis scoring

Arthritic adult K/BxN mice were bled and the sera were pooled. Recipient mice were injected with 100–200  $\mu$ l i.p. as indicated in the figure legends on days 0 and 2. In the chronic inflammation study, mice were injected with 150  $\mu$ l of pooled K/BxN sera on days 0 and 2, and then were given a series of six more injections of 50 or 100  $\mu$ l to perpetuate the inflammatory stimulus when paw swelling was subsiding on days 18, 21, 32, 35, 53, and 56. These animals were sacrificed after 70 days. For each swollen paw, 1 point was given, resulting in a maximum score of 4 per mouse. Ankle thickness was measured with a caliper (Manostat, Herisau, Switzerland) in millimeters.

### Mast cell reconstitution

Bone marrow cells were harvested from the femurs and tibia of donors and cultured in RPMI 1640 with 10% FCS (Omega Scientific, Tarzana, CA), 1% penicillin/streptomycin, 0.1 mM nonessential amino acids,  $5 \times 10^5$  M 2-ME, and 3 ng/ml IL-3 (BD PharMingen) for 6 wk, serially monitoring the cultures until they were >95% pure by toluidine blue staining. The recipient W/W<sup>v</sup> mice were injected with  $10^7$  cells in 100  $\mu$ l of RPMI 1640 i.v. After 6 wk, the mice were injected with pooled K/BxN serum.

### Histology

Whole knee joints and hind paws were fixed in 10% Formalin, decalcified, trimmed, and embedded. Sections were prepared from the tissue blocks and stained with H&E or toluidine blue (Comparative Biosciences, Mountain View, CA and Biomedical Testing Service, San Diego, CA, respectively).

### ELISA

Lapine glucose-6-phosphate isomerase (G6PI) type IV (Sigma-Aldrich, St. Louis, MO) was coated on high-affinity 96-well ELISA plates (Costar, Cambridge, MA) at 10  $\mu$ g/ml in PBS. Plates were then blocked with PBS/1% BSA. Anti-G6PI IgG1 was detected with alkaline phosphatase-labeled goat anti-mouse IgG1 (Southern Biotechnology Associates, Santa Cruz, CA) followed by incubation with *p*-nitrophenyl phosphate substrate (Sigma-Fast; Boehringer Mannheim, Mannheim, Germany). Absorption was measured at 405 nm. For IgE measurement, serum samples were pre-treated with protein G-Sepharose beads to remove the effect of competing of IgG Abs for Ag and then diluted in blocking buffer. Bound murine IgE was detected by biotinylated rat anti-mouse IgE (R35-92; BD PharMingen) and streptavidin-peroxidase (Zymed, South San Francisco, CA) followed by TMB substrate (Kirkegaard & Perry Laboratories, Gaithersburg, MD). The reaction was stopped with 1 M phosphoric acid and absorbance was read at 450 nm. The pool of injected sera was used as standards, arbitrarily

set at  $10^6$  U of IgG1 and 100 U of IgE undiluted. Data were analyzed using DeltaSOFT II version 3.66 (Biometallics, Princeton, NJ).

## Results


### Sustained presence of autoantibodies is required to perpetuate arthritis

Pooled (K/B  $\times$  N) and normal BALB/c sera were transferred to BALB/c and  $\beta_2$ m<sup>-/-</sup> (BALB/c) mice. All mice that received the pooled KxB/N sera initially developed arthritis, but the synovitis in  $\beta_2$ m<sup>-/-</sup> mice quickly resolved (Fig. 1A). Similar to previous reports, the ankle thickness never returned to baseline or to that of mice that received control BALB/c serum despite the absence of visible erythema or swelling (1). Mice that were deficient in the expression of the transporter for Ag presentation (TAP) were tested in a separate experiment and did not show a similar rapid resolution of paw swelling (data not shown).


$\beta_2$ m is a soluble domain for several proteins, including MHC class I, CD1, and neonatal FcR (FcRn). We suspected that the deficiency of FcRn in the  $\beta_2$ m<sup>-/-</sup> mice was causing a rapid clearance of the autoantibody (16). Hence, BALB/c, CD1<sup>-/-</sup> (BALB/c), and  $\beta_2$ m<sup>-/-</sup> (BALB/c) mice were injected with pooled K/B  $\times$  N sera (Fig. 1B). The sera of the injected mice were checked for the presence of anti-G6PI Abs before the transfer of serum, at the peak of the arthritis, and after the arthritis had resolved in the  $\beta_2$ m<sup>-/-</sup> mice. The  $\beta_2$ m-deficient mice had significantly less measurable anti-G6PI IgG at 3 and 13 days after injection than the other two groups (Fig. 1C). However, these mice did not have less Ag-specific IgE, as this catabolism pathway is FcRn independent.

### Fc $\gamma$ RIII is predominantly associated with paw swelling in serum-transferred arthritis

We previously described our inability to transfer KRN arthritis to Fc $\gamma$ R<sup>-/-</sup> mice (5). To extend these studies, we injected Fc $\gamma$ RIII<sup>-/-</sup> and Fc $\gamma$ RII<sup>-/-</sup> mice as well as wild-type controls with pooled K/BxN sera. Fc $\gamma$ R<sup>-/-</sup> mice again did not show any paw swelling. The Fc $\gamma$ RII<sup>-/-</sup> mice developed severe accelerated arthritis, whereas the paw swelling that developed in Fc $\gamma$ RIII<sup>-/-</sup> mice was markedly attenuated with delayed kinetics (Fig. 2).



**FIGURE 1.** Transfer of arthritis resolves quickly in  $\beta_2$ m<sup>-/-</sup> mice and correlates with anti-G6PI IgG. Adult K/BxN and BALB/c mice were bled and the sera were pooled. **A**, Adult BALB/c (●, ○), and  $\beta_2$ m<sup>-/-</sup> (■, □) mice were injected on days 0 and 2 with 200  $\mu$ l of pooled K/BxN (●, ■) and normal sera (○, □) i.p. Arthritis was clinically scored and ankle thickness was measured with a caliper. The means of four mice per group  $\pm$  SEM are shown. **B**, Adult BALB/c (●), CD1<sup>-/-</sup> (▲), and  $\beta_2$ m<sup>-/-</sup> (■) mice were injected on days 0 and 2 with 200  $\mu$ l of pooled sera i.p. Arthritis was clinically scored and ankle thickness was measured with a caliper. The means of three to four mice per group  $\pm$  SEM are shown. **C**, The mice were bled before injection and then on days 3 and 13 after injection. Serum anti-G6PI Ab titers in injected BALB/c (■), CD1<sup>-/-</sup> (▨), and  $\beta_2$ m<sup>-/-</sup> (□) were determined by ELISA as described in *Materials and Methods*. Mean values  $\pm$  SEM of three to four mice per group are shown. The IgG1 on days 3 and 13 were significantly less in the  $\beta_2$ m<sup>-/-</sup> mice than in the other two groups (\*,  $p < 0.05$  by Student's *t* test). The data are representative of two independently performed experiments.



**FIGURE 2.** Transfer of arthritis by K/BxN sera is predominantly FcRIII dependent. The following groups of adult mice were injected on days 0 and 2 with 100  $\mu$ l of pooled K/BxN sera i.p.: *A* and *B*, C57BL/6 (●), Fc $\gamma$ R<sup>-/-</sup> (▲), and Fc $\gamma$ RII<sup>-/-</sup> (■); *C* and *D*, 129/Sv (cross), (B6  $\times$  129)F<sub>2</sub> (○), Fc $\gamma$ R<sup>-/-</sup> (▲), and Fc $\gamma$ RII<sup>-/-</sup> (■); *E* and *F*, C57BL/6 (●), Fc $\gamma$ R<sup>-/-</sup> (▲), and Fc $\gamma$ RIII<sup>-/-</sup> (◆). Arthritis was clinically scored and ankle thickness was measured with a caliper. The means of four mice per group  $\pm$  SEM are shown. Days when Fc $\gamma$ RII<sup>-/-</sup> and Fc $\gamma$ RIII<sup>-/-</sup> mice have significantly greater paw swelling than C57BL/6, 129/Sv, (B6  $\times$  129)F<sub>2</sub>, and Fc $\gamma$ R<sup>-/-</sup> mice are noted (\* and †, respectively;  $p < 0.05$  by Student's *t* test).

#### Bone erosion and hypercellular synovium in Fc $\gamma$ R<sup>-/-</sup> mice

In an attempt to produce chronic inflammation in the passive transfer model, Fc $\gamma$ R<sup>-/-</sup> and C57BL/6 mice were injected with 150  $\mu$ l of K/BxN serum on days 0 and 2 and were then re-injected with


smaller doses to re-establish clinical arthritis each time the paw swelling of the C57BL/6 mice had subsided. At the end of 72 days, the mice were sacrificed and the knees were examined. Shown is an example of a normal C57BL/6 mouse knee (Fig. 3, *A* and *B*) at the site where the cartilage meets the bone, which is sometimes referred to as the "bare area." The cartilage is smooth and the synovium is a single-cell layer. In contrast, the C57BL/6 K/BxN sera-treated mice showed a proliferative synovial response and an inflammatory infiltrate (Fig. 3, *C*, *D*, and *G*). Despite the absence of clinically apparent arthritis, the Fc $\gamma$ R<sup>-/-</sup> mice also had thickened hypercellular synovium, bone erosions, and cartilage damage but to a lesser extent (Fig. 3, *E*, *F*, and *H*).

#### Adoptive transfer of arthritis is dependent on FcR on bone marrow-derived cells

An interesting topic of discussion is why the arthritis is the only major disease manifestation in this model. FcRs present on synovial fibroblasts and extracellular matrix have been described previously (17–20). To evaluate whether joint inflammation is associated with bone marrow-derived elements or with other connective tissues, bone marrow chimeras were made by irradiating C57BL/6 and Fc $\gamma$ R recipients and reconstituting them with Fc $\gamma$ R and C57BL/6 donor bone marrow, respectively. After 8 wk, the animals were bled and checked for bone marrow engraftment. The chimeras and unmanipulated control mice were injected with pooled K/BxN sera. The development of arthritis required the presence of Fc $\gamma$ R on bone marrow-derived cells and not on synovial fibroblasts, which should be less radiation sensitive (Fig. 4).


#### Mast cells partially mediate KRN arthritis

The joint inflammation following the adoptive transfer of KxB/N serum is reminiscent of an Arthus reaction, an analogous disease associated with immune complexes (21). Mast cells play a dominant role in a passive cutaneous Arthus reaction (22, 23), and they are also present in the synovium of normal and K/BxN mice (Fig. 5, *A–D*). W/W<sup>v</sup> mice have a mutation affecting the *c-kit* tyrosine



**FIGURE 3.** Erosive disease in Fc $\gamma$ R<sup>-/-</sup> recipient mice. Four adult C57BL/6 and Fc $\gamma$ R<sup>-/-</sup> mice were injected with 150  $\mu$ l of pooled K/BxN sera on days 0 and 2 and were then given a series of six more injections on days 18, 21, 32, 35, 53, and 56 of 50  $\mu$ l or 100  $\mu$ l to perpetuate paw swelling. These animals were sacrificed after 70 days. Hind paws were fixed in Formalin, joints were decalcified, and tissue sections of knees were stained with H&E. Shown is an example of a normal C57BL/6 mouse knee (*A* and *B*) at the site where the cartilage meets the bone, which is sometimes referred to as the bare area. The cartilage is smooth and the synovium is a single-cell layer. In both C57BL/6 (*C* and *D*) and Fc $\gamma$ R<sup>-/-</sup> (*E* and *F*) K/BxN sera-treated mice, a hypercellular synovium and bone erosions (arrowheads) were noted. The scalloped edges of the synovium invading the bone can be seen. Damage to the cartilage (thin arrows) was seen in the C57BL/6 group (*G*) and to a lesser extent in the Fc $\gamma$ R<sup>-/-</sup> mice (*H*). In *G*, synovium is seen directly overlying the cartilage. *A*, *C*, and *E* are  $\times 100$  original magnification and *B*, *D*, and *F* are  $\times 400$  original magnification of approximately the same area. *G* and *H* are  $\times 400$  original magnification views of different areas. For orientation, the s overlies synovium, b designates bone, and c indicates cartilage.





**FIGURE 4.** FcγR on bone marrow-derived elements is critical for arthritis. Adult C57BL/6 and FcγR<sup>-/-</sup> mice were irradiated and reconstituted with FcγR<sup>-/-</sup> bone marrow (□) and C57BL/6 (○) bone marrow, respectively. They were injected along with unmanipulated C57BL/6 (●) and FcγR<sup>-/-</sup> (■) control mice with 150 μl of pooled K/BxN sera i.p. on days 0 and 2. Arthritis was clinically scored (A) and ankle thickness was measured with a caliper (B). The means of four mice per group ± SEM are shown.


kinase receptor, which is necessary for normal mast cell maturation, proliferation, and survival (24). As a result, these mice virtually lack tissue mast cells (25–27). These mast cell-deficient mice were injected with pooled KxB/N sera and observed for the development of paw swelling. Compared with control animals, these mice had markedly attenuated arthritis (Fig. 5E). Furthermore, mast cell-deficient mice that were reconstituted with bone marrow-derived mast cells developed an intermediate course of arthritis.

**Discussion**

Much interest has been engendered by the KRN model of spontaneously occurring erosive arthritis. In this model, T cells bearing a single autoreactive TCR escapes negative selection in mice bearing a specific MHC class II allele, IA<sup>g7</sup> (1–3, 28, 29). In the periphery, these T cells promote a breach in B cell tolerance and high levels of anti-G6PI are produced (3, 29). As anti-G6PI autoantibodies accumulate in the serum, a destructive and erosive arthritis similar to that seen in human rheumatoid arthritis was observed. The adoptive transfer of sera from these mice results in peripheral joint swelling in most recipient strains (2, 4). This adoptive transfer model allows the study of end-effector mechanisms in multiple strains of mice simultaneously (15, 30).

Early events that trigger paw swelling in the serum transfer model include signaling through FcRs (5, 7). In this study, FcγRIII is predominantly, but not entirely, responsible for the joint swelling seen in the adoptive transfer model. Although no swelling developed in the FcγR-deficient mice, another receptor that shares the common γ-chain, FcγRI has been reported not to influence the development of inflammation in this model (7). In contrast, FcγRII signaling suppresses joint swelling, which may be dependent on genetic background (4). A previous report mapped the genetic loci of one pair of responder/nonresponder (C57BL/6 × NOD)F<sub>2</sub> mice that conferred susceptibility to K/BxN serum-transferred arthritis. Two genomic regions were found to be major determinants with additive effects. The region on chromosome 2 was centered on the C5 locus, but the region on the distal arm of chromosome 1 was broad and contained many attractive candidate genes including *FcγRII*. In this study, FcγRII-deficient mice which were (129 × B6)F<sub>2</sub> did not have an accelerated course of arthritis. Clearly, there are multiple gene effects in the susceptibility to serum-transferred arthritis and the influence of a single gene in mice with a mixed genetic background can be muted.

The development of arthritis as the selected target in this model of immune-mediated disease is of great interest. Anti-G6PI Abs rapidly localize to the joint and an immunohistological study revealed the accumulation of extracellular G6PI along the cartilage surface (31, 32). The normal synovium is relatively hypocellular,



**FIGURE 5.** K/BxN serum-transferred arthritis is partially mast cell dependent. The hind paws of wild-type (A and B) and K/BxN (C and D) mice were fixed in Formalin, joints were decalcified, and embedded, and tissue sections were stained with toluidine blue. Mast cells notable for the dark staining granules were visualized in the synovium (original magnification, ×100 in A and C). The mast cells designated with the arrowheads are further magnified (original magnification, ×400 in B and D). E, Adult K/BxN mice were bled and the sera were pooled. Adult mast cell-deficient WBB6F<sub>1</sub>-W/W<sup>v</sup> mice (●), WBB6F<sub>1</sub> controls (■), and WBB6F<sub>1</sub>-W/W<sup>v</sup> mice reconstituted with mast cells (▲) were injected with 100 μl of pooled sera on days 0 and 2. Arthritis was clinically scored and ankle thickness was measured with a caliper. The means of three to four mice per group ± SEM are shown. \*, greater paw swelling in WBB6F<sub>1</sub> than in the W/W<sup>v</sup> mice, p < 0.05 by Student’s t test).

containing only a lining of synoviocytes and a sublining with fibroblasts, connective tissue, and blood vessels. Various cells expressing FcRs are present in joints. Reports of FcR expression on fibroblast-like synoviocytes led us to examine the role of joint-associated Fc $\gamma$ R expression in the serum transfer model (17–20). The experiments with the chimeric mice show that the development of arthritis depends on the expression of Fc $\gamma$ R by bone marrow-derived cells, which may include mesenchymal stem cells (33, 34).

Neutrophils are heavily recruited into the joint and have been described as crucial to the development of arthritis in the serum transfer model (6). Other cells that express FcRs include macrophages, B cells, and mast cells. Macrophages and monocytes are probably less critical in the acute paw swelling, but may be more prominent in the chronic phase (35). Increased numbers of mast cells are found in the synovial tissues and fluids of patients with RA and at sites of cartilage erosion (36, 37). Mast cell activation and degranulation results in the release of potent mediators, including histamine, heparin, proteinases, leukotrienes, and multi-functional cytokines. In the K/BxN passive transfer model, the arthritis was diminished in mast cell-deficient mice compared with controls.

The currently available data suggest a model whereby synovial mast cells are stimulated by passively transferred Abs and degranulate, causing vascular permeability and recruitment of inflammatory cells, predominantly neutrophils, into the joint. The acute phase requires complement and the correct balance of signaling through FcR family members (4, 5, 7). A more indolent and subacute process, not dependent on FcR signaling or paw swelling, is marked by articular erosion. The pathogenesis of this long-term complication is unclear. The mechanisms for inflammation may also be separate from those for bony erosions (38). Disruption of TNF-related activation-induced cytokine/receptor activator of NF- $\kappa$ B ligand signaling abrogates bony destruction despite active joint inflammation (38). In a RA treatment study with anti-TNF Ab, joint injury as seen on radiographs did not progress over 52 wk in the treated group, regardless of whether inflammatory measures improved or not (39).

Other reports also describe the dissociation between clinical improvement in RA patients and progression of bony erosions and cartilage loss (40, 41). Murine models of chronic and destructive polyarthritis suggest that synovial fibroblasts can be injurious in the absence of T and B lymphocytes. For instance, Rag-1-deficient DBA/1 mice immunized with collagen develop synovial hyperplasia and joint destruction, despite minimal articular inflammation (42). Furthermore, mice with a deletion of the 3' regulatory element of the TNF- $\alpha$  gene when bred onto a Rag-1<sup>-/-</sup> background still developed chronic and progressive joint destruction, implying that stromal elements and/or synovial fibroblasts play a central role in this model (43). The differences between mouse models of arthritis and human RA have to be considered. Each model, however, may provide clues to the mechanisms of the induction, inflammation, and destruction seen in the human disease.

## Acknowledgments

We thank A. Betancourt, S. Wu, P. Charos, N. Noon, and J. Uhle for their assistance. We are grateful to Drs. D. Mathis, C. Benoist, and J. Ravetch for their generous gifts of mice and to Drs. N. Zvaifler, C. Ciesielski, G. S. Firestein, and D. Carson for helpful discussion and critical reviews of articular sections and this manuscript; Drs. Diana Marquardt, Stephen Wasserman, and Ravi Malaviya for their helpful advice regarding mast cell cultures and reconstitution; and Drs. Jae Youn Cho, David Broide, and Jung Y. Cho for assistance with photomicroscopy.

## References

- Korganow, A. S., H. Ji, S. Mangialaio, V. Duchatelle, R. Pelanda, T. Martin, C. Degott, H. Kikutani, K. Rajewsky, J. L. Pasquali, et al. 1999. From systemic T cell self-reactivity to organ-specific autoimmune disease via immunoglobulins. *Immunity* 10:451.
- Kouskoff, V., A. S. Korganow, V. Duchatelle, C. Degott, C. Benoist, and D. Mathis. 1996. Organ-specific disease provoked by systemic autoimmunity. *Cell* 87:811.
- Mangialaio, S., H. Ji, A. S. Korganow, V. Kouskoff, C. Benoist, and D. Mathis. 1999. The arthritogenic T cell receptor and its ligand in a model of spontaneous arthritis. *Arthritis Rheum.* 42:2517.
- Ji, H., D. Gauguier, K. Ohmura, A. Gonzalez, V. Duchatelle, P. Danoy, H. J. Garchon, C. Degott, M. Lathrop, C. Benoist, and D. Mathis. 2001. Genetic influences on the end-stage effector phase of arthritis. *J. Exp. Med.* 194:321.
- Kyburz, D., D. A. Carson, and M. Corr. 2000. The role of CD40 ligand and tumor necrosis factor  $\alpha$  signaling in the transgenic K/BxN mouse model of rheumatoid arthritis. *Arthritis Rheum.* 43:2571.
- Wipke, B. T., and P. M. Allen. 2001. Essential role of neutrophils in the initiation and progression of a murine model of rheumatoid arthritis. *J. Immunol.* 167:1601.
- Ji, H., K. Ohmura, U. Mahmood, D. M. Lee, F. M. Hofhuis, S. A. Boackle, K. Takahashi, V. M. Holers, M. Walport, C. Gerard, et al. 2002. Arthritis critically dependent on innate immune system players. *Immunity* 16:157.
- Ji, H., A. Pettit, K. Ohmura, A. Ortiz-Lopez, V. Duchatelle, C. Degott, E. Gravalles, D. Mathis, and C. Benoist. 2002. Critical roles for interleukin 1 and tumor necrosis factor  $\alpha$  in antibody-induced arthritis. *J. Exp. Med.* 196:77.
- Johansson, A. C., M. Sundler, P. Kjellen, M. Johansson, A. Cook, A. K. Lindqvist, B. Nakken, A. I. Bolstad, R. Jonsson, M. Alarcon-Riquelme, and R. Holmdahl. 2001. Genetic control of collagen-induced arthritis in a cross with NOD and C57BL/10 mice is dependent on gene regions encoding complement factor 5 and Fc $\gamma$ RIIb and is not associated with loci controlling diabetes. *Eur. J. Immunol.* 31:1847.
- Hazenbos, W. L., J. E. Gessner, F. M. Hofhuis, H. Kuipers, D. Meyer, I. A. Heijnen, R. E. Schmidt, M. Sandor, P. J. Capel, M. Daeron, et al. 1996. Impaired IgG-dependent anaphylaxis and Arthus reaction in Fc $\gamma$ RIII (CD16) deficient mice. *Immunity* 5:181.
- Koller, B. H., P. Marrack, J. W. Kappler, and O. Smithies. 1990. Normal development of mice deficient in  $\beta_2$ M, MHC class I proteins, and CD8<sup>+</sup> T cells. *Science* 248:1227.
- Smiley, S. T., M. H. Kaplan, and M. J. Grusby. 1997. Immunoglobulin E production in the absence of interleukin-4-secreting CD1-dependent cells. *Science* 275:977.
- Van Kaer, L., P. G. Ashton-Rickardt, H. L. Ploegh, and S. Tonegawa. 1992. TAP1 mutant mice are deficient in antigen presentation, surface class I molecules, and CD4-8<sup>+</sup> T cells. *Cell* 71:1205.
- Takai, T., M. Ono, M. Hikida, H. Ohmori, and J. V. Ravetch. 1996. Augmented humoral and anaphylactic responses in Fc $\gamma$ RII-deficient mice. *Nature* 379:346.
- Takai, T., M. Li, D. Sylvestre, R. Clynes, and J. V. Ravetch. 1994. Fc $\gamma$  chain deletion results in pleiotropic effector cell defects. *Cell* 76:519.
- Israel, E. J., D. F. Wilsker, K. C. Hayes, D. Schoenfeld, and N. E. Simister. 1996. Increased clearance of IgG in mice that lack  $\beta_2$ -microglobulin: possible protective role of FcRn. *Immunology* 89:573.
- Bhatia, A., S. Blades, G. Cambridge, and J. C. Edwards. 1998. Differential distribution of Fc $\gamma$ RIIIa in normal human tissues and co-localization with DAF and fibrin-1: implications for immunological microenvironments. *Immunology* 94:56.
- Daniilova, T. A., L. M. Bartova, R. L. Panurina, and I. M. Lyampert. 1981. Studies of Fc receptors of heart valve and joint fibroblasts. *Clin. Exp. Immunol.* 46:575.
- Edwards, J. C., S. Blades, and G. Cambridge. 1997. Restricted expression of Fc $\gamma$ RIII (CD16) in synovium and dermis: implications for tissue targeting in rheumatoid arthritis (RA). *Clin. Exp. Immunol.* 108:401.
- Miyazawa, K., A. Mori, and H. Okudaira. 1998. Establishment and characterization of a novel human rheumatoid fibroblast-like synoviocyte line, MH7A, immortalized with SV40 T antigen. *J. Biochem.* 124:1153.
- Sylvestre, D. L., and J. V. Ravetch. 1994. Fc receptors initiate the Arthus reaction: redefining the inflammatory cascade. *Science* 265:1095.
- Zhang, Y., B. F. Ramos, and B. A. Jakschik. 1991. Augmentation of reverse Arthus reaction by mast cells in mice. *J. Clin. Invest.* 88:841.
- Sylvestre, D. L., and J. V. Ravetch. 1996. A dominant role for mast cell Fc receptors in the Arthus reaction. *Immunity* 5:387.
- Nocka, K., J. C. Tan, E. Chiu, T. Y. Chu, P. Ray, P. Traktman, and P. Besmer. 1990. Molecular bases of dominant negative and loss of function mutations at the murine *c-kit*/white spotting locus: W37, Wv, W41 and W. *EMBO J.* 9:1805.
- Sarvella, P. A., and L. B. Russell. 1956. Steel, a new dominant gene in the house mouse. *J. Hered.* 47:123.
- Huang, E., K. Nocka, D. R. Beier, T. Y. Chu, J. Buck, H. W. Lahm, D. Wellner, P. Leder, and P. Besmer. 1990. The hematopoietic growth factor KL is encoded by the *Sl* locus and is the ligand of the *c-kit* receptor, the gene product of the *W* locus. *Cell* 63:225.
- Hayashi, C., T. Sonoda, T. Nakano, H. Nakayama, and Y. Kitamura. 1985. Mast-cell precursors in the skin of mouse embryos and their deficiency in embryos of *Sl/Sl* genotype. *Dev. Biol.* 109:234.
- Kouskoff, V., A. S. Korganow, V. Duchatelle, C. Degott, C. Benoist, and D. Mathis. 1997. A new mouse model of rheumatoid arthritis: organ-specific disease provoked by systemic autoimmunity. *Ryumachi* 37:147.
- Matsumoto, I., A. Staub, C. Benoist, and D. Mathis. 1999. Arthritis provoked by linked T and B cell recognition of a glycolytic enzyme. *Science* 286:1732.

30. Ravetch, J. V., and S. Bolland. 2001. IgG Fc receptors. *Annu. Rev. Immunol.* 19:275.
31. Matsumoto, I., M. Maccioni, D. M. Lee, M. Maurice, B. Simmons, M. Brenner, D. Mathis, and C. Benoist. 2002. How antibodies to a ubiquitous cytoplasmic enzyme may provoke joint-specific autoimmune disease. *Nat. Immunol.* 3:360.
32. Wipke, B. T., Z. Wang, J. Kim, T. J. McCarthy, and P. M. Allen. 2002. Dynamic visualization of a joint-specific autoimmune response through positron emission tomography. *Nat. Immunol.* 3:366.
33. Corr, M., and N. J. Zvaifler. 2002. Mesenchymal precursor cells. *Ann. Rheum. Dis.* 61:3.
34. Zvaifler, N. J., L. Marinova-Mutafchieva, G. Adams, C. J. Edwards, J. Moss, J. A. Burger, and R. N. Maini. 2000. Mesenchymal precursor cells in the blood of normal individuals. *Arthritis Res.* 2:477.
35. Lee, D. M., D. Mathis, C. Benoist, and M. B. Brenner. 2001. Presence of inflammation and pannus formation in mice lacking type A synoviocytes. *Arthritis Rheum.* 44:S87.
36. Bromley, M., W. D. Fisher, and D. E. Woolley. 1984. Mast cells at sites of cartilage erosion in the rheumatoid joint. *Ann. Rheum. Dis.* 43:76.
37. Crisp, A. J. 1985. Synovial fluid mast cells. *Ann. Rheum. Dis.* 44:724.
38. Pettit, A. R., H. Ji, D. von Stechow, R. Muller, S. R. Goldring, Y. Choi, C. Benoist, and E. M. Gravallese. 2001. TRANCE/RANKL knockout mice are protected from bone erosion in a serum transfer model of arthritis. *Am. J. Pathol.* 159:1689.
39. Lipsky, P. E., D. M. van der Heijde, E. W. St Clair, D. E. Furst, F. C. Breedveld, J. R. Kalden, J. S. Smolen, M. Weisman, P. Emery, M. Feldmann, et al. 2000. Infliximab and methotrexate in the treatment of rheumatoid arthritis: anti-tumor necrosis factor trial in rheumatoid arthritis with concomitant therapy study group. *N. Engl. J. Med.* 343:1594.
40. Mulherin, D., O. Fitzgerald, and B. Bresnahan. 1996. Clinical improvement and radiological deterioration in rheumatoid arthritis: evidence that the pathogenesis of synovial inflammation and articular erosion may differ. *Br. J. Rheumatol.* 35:1263.
41. McQueen, F. M., N. Stewart, J. Crabbe, E. Robinson, S. Yeoman, P. L. Tan, and L. McLean. 1999. Magnetic resonance imaging of the wrist in early rheumatoid arthritis reveals progression of erosions despite clinical improvement. *Ann. Rheum. Dis.* 58:156.
42. Plows, D., G. Kontogeorgos, and G. Kollias. 1999. Mice lacking mature T and B lymphocytes develop arthritic lesions after immunization with type II collagen. *J. Immunol.* 162:1018.
43. Kontoyiannis, D., M. Pasparakis, T. T. Pizarro, F. Cominelli, and G. Kollias. 1999. Impaired on/off regulation of TNF biosynthesis in mice lacking TNF AU-rich elements: implications for joint and gut-associated immunopathologies. *Immunity* 10:387.