

Hierarchical Organization of Prostate Cancer Cells in Xenograft Tumors: The CD44⁺α2β1⁺ Cell Population Is Enriched in Tumor-Initiating Cells

Lubna Patrawala,¹ Tammy Calhoun-Davis,¹ Robin Schneider-Broussard,¹ and Dean G. Tang^{1,2}

¹Department of Carcinogenesis, The University of Texas M. D. Anderson Cancer Center, Science Park-Research Division, Smithville, Texas and ²Program in Environmental and Molecular Carcinogenesis, Graduate School of Biomedical Sciences, Houston, Texas

Abstract

Prostate cancer cells are heterogeneous in their tumorigenicity. For example, the side population cells isolated from LAPC9 xenografts are 100 to 1,000 times more tumorigenic than the corresponding non-side population cells. Highly purified CD44⁺ prostate cancer cells from several xenografts are also enriched in prostate cancer stem/progenitor cells. Because the CD44⁺ prostate cancer cell population is still heterogeneous, we wonder whether we could further enrich for tumorigenic prostate cancer cells in this population using other markers. Integrin α2β1 has been proposed to mark a population of normal human prostate stem cells. Therefore, we first asked whether the α2β1^{hi} cells in prostate tumors might also represent prostate cancer stem cells. Highly purified (≥98%) α2β1^{hi} cells from three human xenograft tumors, Du145, LAPC4, and LAPC9, show higher clonal and clonogenic potential than the α2β1^{lo} cells *in vitro*. However, when injected into the nonobese diabetic/severe combined immunodeficient (NOD/SCID) mouse prostate or s.c., the α2β1^{hi} prostate cancer cells are no more tumorigenic than the α2β1^{lo} cells. Immunofluorescence studies reveal that CD44 and α2β1 identify an overlapping and inclusive population of prostate cancer cells in that ~70% of α2β1^{hi} cells are CD44⁺ and 20% to 30% of CD44⁺ cells are distributed in the α2β1^{lo} cell population. Subsequently, we sorted out CD44⁺α2β1^{hi}, CD44⁺α2β1^{lo}, CD44⁻α2β1^{hi}, and CD44⁻α2β1^{lo} cells from LAPC9 tumors and carried out tumorigenicity experiments. The results revealed a hierarchy in tumorigenic potential in the order of CD44⁺α2β1^{hi} ≈ CD44⁺α2β1^{lo} > CD44⁻α2β1^{hi} ≫ CD44⁻α2β1^{lo}. These observations together suggest that prostate cancer cells are organized as a hierarchy. [Cancer Res 2007;67(14):6796–805]

Introduction

The cancer stem cell model posits that not all cells in a tumor are equal, and that tumor-initiating cells are a rare subset with a distinct phenotype (1, 2). This hierarchical model helps explain why most tumors are heterogeneous although they have a clonal origin; why it is often difficult to establish a permanent cell line from primary tumors; and why it takes tens of thousands of cancer cells

to reestablish a tumor. Indeed, cancer stem cells have been shown to be the driving force behind tumor formation in several types of cancers, including those of the blood, breast, brain, and colon (3–7). Cancer stem cells are generally thought of as self-renewing cells that are able to reinitiate a tumor for several generations in NOD/SCID mice and can give rise to a spectrum of differentiated cells (1, 2, 8, 9). Cancer stem cells, like normal stem cells, are also more likely to express antiapoptotic and drug-resistance genes, making them impervious to most anticancer therapeutics (1, 8). To completely eradicate a tumor and prevent recurrence, it is imperative that cancer stem cells be specifically targeted.

Prostate cancer is the second most common type of cancer afflicting American males, and yet, little is known about which cell types within the prostate are the targets of tumorigenic transformation (reviewed in ref. 9). The normal human prostate contains two major epithelial cell types: luminal and basal cells. The luminal cells express cytokeratins 8 and 18, androgen receptor, prostate-specific antigen, prostatic acid phosphatase, and 15-lipoxygenase 2 (9–11), whereas basal cells express cytokeratin 5, CD44 (12), Bcl-2 (13), p63 (14), telomerase (15, 16), and glutathione *S*-transferase π (17) and display ~75% of mitotic activity in the prostate (18). The human prostate epithelium has the ability to generate gland-like structures when combined with rat urogenital mesenchyme and implanted into the renal capsule (19), suggesting the presence of stem cells. Strong experimental evidence exists that putative human prostate stem cells might localize in the basal cell layer (9). Several candidate populations of prostate stem/progenitor cells have been reported, including those expressing CD44, α2β1, or CD133 (20, 21). For example, the α2β1^{hi} cells comprise ~1% to 15% of the CD44⁺ basal cell population and seem to possess higher *in vitro* colony-forming efficiency as well as an ability to generate prostate-like acini when engrafted with stromal cells into the flanks of nude mice (20). Further characterization reveals that this proliferation and developmental potential seems to be harbored preferentially by CD133-expressing cells within the CD44⁺α2β1^{hi} population (21). Our recent work also shows that primary prostate epithelial isolates contain cells that possess tremendous proliferative potential and the ability to “transdifferentiate” into other cell types (9). As the prostate joins the growing list of organs that are found to contain adult stem cells, it seems impossible to ignore the likelihood that prostate cancer development might involve these cells or their immediate progeny. In fact, it has been recently reported that stem-like cells in patient prostate tumors can be identified using the putative normal stem cell phenotype (i.e., CD44⁺α2β1^{hi}CD133⁺; ref. 22). Not only is this population rare but it also shows the highest colony-forming efficiency and the capacity to differentiate into several cell types. Unfortunately, the ability of these cells to reinitiate serially transplantable tumors, which is the gold standard to define cancer stem cells (8, 9), was not shown (22).

Note: Supplementary data for this article are available at Cancer Research Online (<http://cancerres.aacrjournals.org/>).

Requests for reprints: Dean G. Tang, Department of Carcinogenesis, Science Park-Research Division, The University of Texas M. D. Anderson Cancer Center, 1808 Park Road 1C, Smithville, TX 78957. Phone: 512-237-9575; Fax: 512-237-2475; E-mail: dtang@mdanderson.org.

©2007 American Association for Cancer Research.
doi:10.1158/0008-5472.CAN-07-0490

We have recently shown that cells in well-established prostate cancer xenografts also seem to be organized as a hierarchy with distinct subsets of cells that preferentially harbor tumorigenicity. For instance, the side population isolated from LAPC9 xenograft tumors is $\sim 1,000$ times more tumorigenic than the corresponding non-side population cells, suggesting that the side population is enriched in prostate cancer stem/progenitor cells (23). In addition, highly purified CD44⁺ cells are also enriched in tumorigenic and metastatic prostate cancer stem/progenitor cells (24). In this study, we continue to use these xenograft models in an attempt to further dissect out tumorigenic prostate cancer stem/progenitor subpopulations. The results reveal that prostate cancer cells show a hierarchy in their tumorigenic potential based on their CD44 and $\alpha 2\beta 1$ expression profiles.

Materials and Methods

Cells, reagents, and animals. LNCaP, Du145, PC3, and PPC-1 prostate cancer cell lines were obtained from American Type Culture Collection and cultured in RPMI containing 7% heat-inactivated fetal bovine serum (FBS). Xenograft human prostate tumors LAPC4 and LAPC9 were obtained from Dr. C. Sawyers (Department of Medicine, Jonsson Comprehensive Cancer Center, David Geffen School of Medicine, University of California at Los Angeles, Los Angeles, CA) and maintained in NOD/SCID mice (24). Du145 xenograft tumors were established using early-passage cells and maintained in NOD/SCID mice (23, 24). NOD/SCID mice were obtained from The Jackson Laboratory or bred in our own animal facility and maintained in standard conditions according to the institutional guidelines. Antibodies used include $\alpha 2\beta 1$ [monoclonal antibody (mAb), Chemicon], CD44 (mAb, BD PharMingen), and ABCG2 (mAb, Chemicon). The mouse phycoerythrin-conjugated anti-human $\alpha 2\beta 1$ mAb was obtained from Chemicon and the FITC-conjugated anti-CD44 antibody from BD PharMingen. The isotype control antibody and FITC-, phycoerythrin-, or AlexaFluor-conjugated secondary antibodies were from Chemicon.

Indirect immunofluorescence and flow cytometric analysis and sorting of $\alpha 2\beta 1^{+/hi}$ and $\alpha 2\beta 1^{-/lo}$ cells. Fluorescence microscopy was carried out as previously described (23, 24). For flow cytometry, cells were stained live in the staining solution containing 2% FBS and phycoerythrin-conjugated anti- $\alpha 2\beta 1$ mAb either alone or in combination with FITC-conjugated anti-CD44 antibody (15 min at 4°C). Samples were analyzed on a Coulter Epics Elite flow cytometer and $\sim 5 \times 10^6$ to 10×10^6 cells were typically sorted. Cell debris and clumps were electronically gated out. For the positive population, only the top 10% most brightly stained cells were selected. For the negative population, only the bottom 10% most dimly stained cells were selected. The purities of the sorted populations, as determined by both post-sorting flow analyses as well as restaining followed by fluorescence microscopy analyses, were generally $\geq 98\%$.

Clonal analysis and clonogenic assays. *In vitro* colony-forming and clonogenic assays were done as described before (23, 24). Briefly, tumor cells were plated at clonal density (i.e., 100–500 per well in a six-well tissue culture dish). Holoclones with >50 cells were counted at the end of 1 week. For clonogenic assays, cells were plated at 1,000 per well in six-well culture dishes coated with a thin layer of 1% solidified agar. Spheres or spheroids (i.e., colonies) that arose within 1 to 2 weeks were presented as clonogenicity (i.e., percent of the spheres/1,000 cells initially plated). Triplicate samples were run for each cell type and at least two individuals scored the clones and spheres separately in a blind fashion.

Xenograft tumor processing and *in vivo* tumorigenicity experiments. Basic procedures have previously been described (23, 24). In brief, xenograft prostate tumors (Du145, LAPC4, and LAPC9) were minced into $\sim 1\text{-mm}^3$ pieces in Iscove's modified Dulbecco's medium supplemented with 20% FBS. Tumor tissues were incubated with $1 \times$ Accumax (1,200–2,000 units/mL proteolytic activity containing collagenase and DNase; Innovative Cell Technologies, Inc.) at 10 mL/g of tissue in Dulbecco's PBS for 30 min at room temperature under rotating conditions. Single-cell suspension was obtained by filtering the supernatant through a 40- μm cell strainer and cell

suspension was then gently loaded onto a layer of Histopaque-1077 gradient and then centrifuged at $400 \times g$ for 30 min at room temperature. RBC, dead cells, and debris were removed from the bottom of the tube and live nucleated epithelial cells collected at the interface. The resultant cell mixture was depleted of lineage-positive host cells using the MACS Lineage Cell Depletion kit (Miltenyi Biotec). To that end, cells were first incubated (10 min at 4°C) in the staining solution [PBS (pH 7.2), 0.5% FBS, 0.5 $\mu\text{g/mL}$ insulin] containing biotinylated antibodies against a panel of lineage antigens (CD5, CD45R, CD11b, anti-Ly-6G, 7-4, and Ter-119). Cells were then incubated with the anti-biotin microbeads (15 min at 4°C) and the Lin⁻ cells were eluted using the mass spectrometry columns. The purified human epithelial tumor cells were used in various experiments.

For tumor experiments, cells in 40- μL solution consisting of 50% each medium and Matrigel were injected either s.c. or into the dorsal prostate of NOD/SCID mice (~ 8 weeks old). In some experiments, testosterone pellets ~ 0.2 cm in diameter (we used testosterone propionate powder purchased from Sigma to make pellets in our laboratory) were implanted dorsally under the skin of male NOD/SCID mice. Primary tumor sizes were measured with a caliper on a weekly basis. Tumorigenicity was measured mainly by tumor incidence (i.e., the number of tumors/number of injections) and latency (i.e., time from injection to detection of palpable tumors). For the double-sort experiments, all tumor-bearing animals were terminated at the same time when tumor burden became obvious for any one animal in one group. Animals were sacrificed and primary tumors dissected out, and tumor weights were determined. Animals with no sign of tumor burden were also examined on necropsy to confirm that there was no tumor development.

Results

$\alpha 2\beta 1$ expression correlates with malignancy in prostate cancer cell lines and xenografts. $\alpha 2\beta 1$, an integrin expressed in the human prostate (25) and mediating adhesion to collagen I/IV and laminin I (26), has been proposed to identify normal human prostate stem cells (21). To determine whether prostate cancer cells expressing high levels of $\alpha 2\beta 1$ (which are called $\alpha 2\beta 1^{+/hi}$ or simply $\alpha 2\beta 1^+$ in this study) in prostate tumors might also represent prostate cancer stem/progenitor cells, we examined its expression in the commonly used prostate cancer cell lines LNCaP, Du145, PC3, and PPC-1 (27), as well as in the LAPC4 and LAPC9 xenograft tumors (10), using a mAb against the $\alpha 2$ subunit. As shown in Fig. 1, LNCaP cells, which are the least aggressive (27), did not show detectable $\alpha 2\beta 1$ expression revealed by immunofluorescence staining. By contrast, PC3 and its derivative PPC-1 cells, which were the most malignant (27), showed 100% expression (Fig. 1). The three prostate cancer cell types that possess intermediate malignancy (i.e., Du145, LAPC4, and LAPC9) showed intermediate levels (i.e., ~ 1 –10%) of $\alpha 2\beta 1$ expression (Fig. 1). Importantly, $\alpha 2\beta 1$ was also expressed only in a subset of cells in these three xenograft tumors (Figs. 1B and 2A). Flow cytometry analysis of multiple xenograft tumors indicated that the LAPC9, LAPC4, and Du145 tumors expressed, on average, 2.4% ($n = 9$), 0.4% ($n = 5$), and 9.8% ($n = 4$), respectively, of the $\alpha 2\beta 1^{+/hi}$ cells (Supplementary Table S1; Fig. 2A). Note that the percentages of the $\alpha 2\beta 1^{+/hi}$ cells detected by flow cytometry were generally slightly higher than those detected by immunofluorescent staining. These results, taken together, suggest that the abundance of $\alpha 2\beta 1^+$ cells in prostate cancer cell cultures and xenografts is correlated with tumor cell malignancy.

$\alpha 2\beta 1^{+/hi}$ cells possess higher clonal and clonogenic potentials *in vitro* compared with the isogenic $\alpha 2\beta 1^{-/lo}$ cells. To determine whether the $\alpha 2\beta 1^{+/hi}$ prostate cancer cells are intrinsically different from the isogenic $\alpha 2\beta 1^{-/lo}$ (or simply $\alpha 2\beta 1^-$) prostate cancer cells, we used flow cytometry to purify





Figure 1. $\alpha 2\beta 1$ expression in prostate cancer cells positively correlates with their malignancy. *A*, $\alpha 2\beta 1$ was detected using indirect immunofluorescence in LNCaP, LAPC4, Du145, and PC3 cells. Representative images of several hundreds analyzed ($\times 200$). *DAPI*, 4',6-diamidino-2-phenylindole. *B*, quantification of cells stained for $\alpha 2\beta 1$. *Columns*, mean derived from a counting of 500 to 800 cells; *bars*, SD. LAPC4 and Du145 results were derived from either cultures (*Cells*) or xenograft tumors (*Tumors*; also see Supplementary Table S1). The LAPC9 results were also derived from xenograft tumor experiments (see Supplementary Table S1).

these two populations of cells from Du145 xenograft tumors. As in our experiments with ABCG2 (23) and CD44 (24), the purities of the $\alpha 2\beta 1^{+hi}$ and $\alpha 2\beta 1^{-lo}$ cell populations were $\geq 98\%$ and $\sim 100\%$, respectively, as revealed by post-sort flow analysis and/or immunostaining (data not shown). When compared for their cloning (i.e., the ability to establish a holoclone; ref. 9) and sphere-forming abilities, the $\alpha 2\beta 1^{+hi}$ cells showed significantly higher colony-forming efficiency (Fig. 2B) and formed bigger clones (Fig. 2C) compared with the isogenic $\alpha 2\beta 1^{-lo}$ cell population. The difference observed was not due to variations in adhesion capacity between the two subsets because even in the negative population, a roughly equal number of cells were observed to attach to the dish on initial plating. However, many of the $\alpha 2\beta 1^{-lo}$ Du145 cells formed abortive clones or no clones at all. Additionally, when plated at low densities in anchorage-independent conditions (i.e., on soft agar-coated dishes), the $\alpha 2\beta 1^{+hi}$ Du145 cells had a much higher sphere-initiating capacity than the corresponding negative population (Fig. 2D). The results in Fig. 2 are consistent with the possibility that the $\alpha 2\beta 1^{+hi}$ prostate cancer cell population might be enriched in prostate cancer stem/progenitor cells.

$\alpha 2\beta 1^{+hi}$ are no more tumorigenic than the corresponding $\alpha 2\beta 1^{-lo}$ cells. The gold standard in testing putative cancer stem cells is whether the candidate population of cells can preferentially initiate tumor development in recipient animals (8, 9). Therefore, we carried out surgical orthotopic implantation experiments by injecting varying numbers of acutely purified $\alpha 2\beta 1^{+hi}$ and

$\alpha 2\beta 1^{-lo}$ Du145 cells into the dorsal prostates of NOD/SCID mice. The dorsal prostate has been widely used as the "orthotopic" implantation site for human prostate cancer (28). As shown in Table 1, unsorted Du145 cells, when injected into the dorsal prostate, showed a cell number-dependent increase in tumorigenicity. In general, 100,000 Du145 cells had to be injected to initiate tumor development, and 0.5×10^6 to 2.0×10^6 cells were required to manifest significant tumorigenicity (Table 1). When freshly purified $\alpha 2\beta 1^{+hi}$ and $\alpha 2\beta 1^{-lo}$ Du145 cells were compared for their tumorigenicities, surprisingly, we did not observe any difference (Table 1). Due to the low percentage of $\alpha 2\beta 1^{+hi}$ cells in Du145 tumors (Fig. 1), we were not able to inject more than 10,000 $\alpha 2\beta 1^{+hi}$ cells for comparative purposes. One would expect 10,000 $\alpha 2\beta 1^{+hi}$ cells to be sufficient for tumor-initiation if these cells were truly primitive with respect to their tumorigenicity; however, this was not the case. The $\alpha 2\beta 1^{+hi}$ cells were not more tumorigenic than the same number of unsorted or $\alpha 2\beta 1^{-lo}$ cells. In fact, the $\alpha 2\beta 1^{-lo}$ population seemed to be slightly enriched in tumorigenic cells such that at 100,000 cells injected, we observed 100% tumor development compared with 25% tumor development with 100,000 unsorted cells (Table 1). The tumor latency was also shorter in the $\alpha 2\beta 1^{-lo}$ group than in the unsorted group (Table 1).

We repeated the experiment using LAPC4 and LAPC9 xenograft tumors and obtained similar results (Table 1). The unsorted LAPC4 cells purified from xenograft tumors exhibited tumor rate and latency that were similar to Du145 cells in that 100,000 cells had to be injected into the dorsal prostate to observe any tumor




Figure 2. $\alpha 2\beta 1^{+}$ prostate cancer cells possess higher clonal and clonogenic potential compared with $\alpha 2\beta 1^{-}$ cells. **A**, flow cytometry analysis of $\alpha 2\beta 1$ expression in xenograft human prostate tumors. Tumor cells purified from Du145 (a), LAPC9 (b), or LAPC4 (c) were stained live with FITC-conjugated anti- $\alpha 2\beta 1$ antibody and analyzed on a Coulter's Epics Elite flow cytometer. The percentages of $\alpha 2\beta 1^{hi/+}$ cells are indicated. **B**, Du145 cells were sorted by FACS for $\alpha 2\beta 1^{hi}$ (i.e., $\alpha 2\beta 1^{+}$) or $\alpha 2\beta 1^{lo}$ (i.e., $\alpha 2\beta 1^{-}$) cells (99% purity), and plated at clonal density (100 per well in a six-well dish) in triplicate. Seven days after plating, clones (with cell number >50) were counted and results were presented as percent cloning efficiency. Columns, mean from three independent experiments; bars, SE. *, $P = 0.038$, compared with the $\alpha 2\beta 1^{-}$ group (Student's t test). **C**, representative images of clonal analyses ($\times 10$). **D**, purified $\alpha 2\beta 1^{+}$ and $\alpha 2\beta 1^{-}$ Du145 cells were plated in triplicate at 1,000 per well in a six-well plate coated with soft agar for clonogenicity assays. Spheres were counted 2 wk after plating. Columns, mean from three independent experiments; bars, SE. *, $P < 0.001$, compared with the $\alpha 2\beta 1^{-}$ group (Student's t test).

Table 1. Tumorigenicity of $\alpha 2\beta 1^{+/hi}$ and $\alpha 2\beta 1^{-/lo}$ prostate cancer cells injected into the dorsal prostate

Cell type	No. cells injected	Tumor incidence*	Latency (d) [†]
Du145, unsorted	1,000	0/4	
	10,000	0/4	
	100,000	1/4	103
	500,000	3/5	53–59 (53)
	2,000,000	3/3	39–74 (46)
Du145- $\alpha 2\beta 1^{+/hi}$	1,000	0/4	
	10,000	0/8	
Du145- $\alpha 2\beta 1^{-/lo}$	1,000	0/4	
	10,000	0/4	
	100,000	4/4	53–93 (72)
	500,000	1/1	48
LAPC4, unsorted	100	0/4	
	1,000	0/4	
	10,000	0/4	
	100,000	1/4	103
	500,000	4/5	43–69 (46)
LAPC4- $\alpha 2\beta 1^{+/hi}$	1,000	0/4	
	10,000	0/2	
LAPC4- $\alpha 2\beta 1^{-/lo}$	1,000	0/4	
	10,000	1/4	102
	100,000	5/6	53–97 (62)
LAPC9, unsorted	100	0/3	
	1,000	0/9	
	10,000	4/8	46–75 (53)
	100,000	6/9	32–69 (44)
	1,000,000	4/4	48–69 (50)
LAPC9- $\alpha 2\beta 1^{+/hi}$	100	0/4	
	1,000	0/4	
	10,000	1/4	109
LAPC9- $\alpha 2\beta 1^{-/lo}$	100	0/4	
	1,000	0/4	
	10,000	1/4	109
	100,000	4/4	42–102 (48)

*Tumor cells were injected in Matrigel into the dorsal prostate of NOD/SCID mice. Tumor incidence refers to the number of tumors developed/number of injections.

[†] Tumor latency refers to the time (in days) from tumor cell injection to when the tumor is detected by palpation. The numbers in parentheses represent the median values.

development, and 500,000 cells were required to reliably generate tumors in a similar time frame (Table 1). LAPC9 cells, on the other hand, were considerably more tumorigenic because 10,000 cells could initiate tumors with 50% efficiency in about the same time interval required for 500,000 Du145 or LAPC4 cells to form tumors (Table 1). When tumorigenicity assays were done using the purified isogenic $\alpha 2\beta 1^{+/hi}$ and $\alpha 2\beta 1^{-/lo}$ populations in LAPC4 tumors, again, we did not observe an enrichment in tumor-initiating cells in the $\alpha 2\beta 1^{+/hi}$ subset (Table 1). Rather, the $\alpha 2\beta 1^{-/lo}$ population seemed slightly more tumorigenic because tumor formation was observed when 10,000 $\alpha 2\beta 1^{-/lo}$ cells were injected whereas no tumors formed when 10,000 $\alpha 2\beta 1^{+/hi}$ or unsorted cells were injected (Table 1). Similar experiments were carried out using sorted LAPC9 cells and, in this case, unsorted, $\alpha 2\beta 1^{+/hi}$, and $\alpha 2\beta 1^{-/lo}$ cells showed very similar tumorigenicity (Table 1).

Due to the overall low tumor development of human prostate cancer cells implanted into the dorsal prostate of NOD/SCID mice, we carried out tumor experiments by injecting cells s.c. as we have recently shown that the subcutis is very permissive to tumor regeneration relative to the mouse dorsal prostate.³ Indeed, as few as 100 LAPC9 cells injected s.c. initiated 50% tumor development (Supplementary Table S2) compared with 10,000 cells required to initiate similar levels of tumor development when injected into the dorsal prostate (Table 1). One thousand s.c. injected LAPC9 cells initiated tumor development in 100% recipient animals in ~2 months (Supplementary Table S2). Nevertheless, despite the dramatically enhanced tumor take in the s.c. implantation model, purified $\alpha 2\beta 1^{+/hi}$ and $\alpha 2\beta 1^{-/lo}$ LAPC9 cells did not show a significant difference with respect to their tumor-initiating capacities (Supplementary Table S2). If anything, the $\alpha 2\beta 1^{-/lo}$ LAPC9 cells seemed to be a bit more tumorigenic than the isogenic $\alpha 2\beta 1^{+/hi}$ cells because the latter cells, at lower cell numbers (i.e., 100 and 1,000 cells), regenerated more tumors (Supplementary Table S2).

The preceding experiments (Fig. 2; Table 1; Supplementary Table S2) reveal that the small population of $\alpha 2\beta 1^{+/hi}$ prostate cancer cells have high clonal and clonogenic capacities *in vitro* but are no more tumorigenic *in vivo* than the corresponding $\alpha 2\beta 1^{-/lo}$ prostate cancer cells. The behavior of $\alpha 2\beta 1^{+/hi}$ cells is reminiscent of that of the ABCG2⁺ cells (23) and suggests that the $\alpha 2\beta 1^{+/hi}$ prostate cancer cells, like ABCG2⁺ cells, may mark a population of fast-proliferating tumor progenitor cells. This would imply that, like the ABCG2⁻ cell population (23), the $\alpha 2\beta 1^{-/lo}$ prostate cancer cell population might contain primitive tumorigenic cells that can give rise to $\alpha 2\beta 1^{+/hi}$ cells. In support, tumors derived from 100 highly purified $\alpha 2\beta 1^{-/lo}$ Du145 cells contained a small percentage of $\alpha 2\beta 1^{+/hi}$ cells (Supplementary Table S1), suggesting that some $\alpha 2\beta 1^{-/lo}$ cells have generated $\alpha 2\beta 1^{+/hi}$ cells. We further analyzed several s.c. or surgical orthotopic implantation tumors derived from 1,000 to 100,000 $\alpha 2\beta 1^{-/lo}$ LAPC9 cells and, in every case, $\alpha 2\beta 1^{+/hi}$ cells were present at a low frequency (0.1–3.9%; Supplementary Table S1). Likewise, surgical orthotopic implantation tumors arising from 100,000 $\alpha 2\beta 1^{-/lo}$ LAPC4 and Du145 cells also contained a consistent small number of $\alpha 2\beta 1^{+/hi}$ cells (Supplementary Table S1). These results suggest that some $\alpha 2\beta 1^{-/lo}$ cells can give rise to $\alpha 2\beta 1^{+/hi}$ cells *in vivo*.

CD44, $\alpha 2\beta 1$, and ABCG2 identify overlapping and inclusive prostate cancer cell populations. We have recently shown that the CD44⁺ cell population is enriched in prostate cancer tumor stem/progenitor cells (24) whereas ABCG2 identifies fast-cycling tumor progenitor cells (23). The preceding experiments suggest that $\alpha 2\beta 1$ may also mark fast-cycling tumor progenitor cells. Next, we carried out immunostaining in an attempt to elucidate the interrelationship among the three (i.e., CD44⁺, $\alpha 2\beta 1^{+/hi}$, and ABCG2⁺) cell populations. We sorted out $\alpha 2\beta 1^{+/hi}$ and $\alpha 2\beta 1^{-/lo}$ cells from Du145 xenograft tumors, plated them onto coverslips immediately after sorting, and fixed them 3 h later to prevent them from dividing. When double-stained for CD44, we found that there was a significant overlap between the $\alpha 2\beta 1^{+/hi}$ and

³ H.W. Li, M. Jiang, T. Calhoun-Davis, L. Patrawala, G. Choy, R. Schneider-Broussard, S.W. Hayward, D.G. Tang. Crucial roles of microenvironments (transplantation sites) on reconstituting tumorigenic versus metastatic potentials of human prostate cancer (stem) cells in nonobese diabetic/severe combined immunodeficient mice. Submitted for publication.




Figure 3. CD44, $\alpha 2\beta 1$, and ABCG2 identify overlapping prostate cancer cell populations. *A* and *B*, CD44 and ABCG2 expression in $\alpha 2\beta 1^+$ and $\alpha 2\beta 1^-$ Du145 cells. Purified $\alpha 2\beta 1^+$ and $\alpha 2\beta 1^-$ Du145 cells were plated at 5,000 per 18-mm² coverslip and fixed after 4 h. *B*, cells were stained using indirect fluorescence with antibodies against CD44 (*a* and *b*) or ABCG2 (*c* and *d*). *A*, the majority of the $\alpha 2\beta 1^+$ were also positive for CD44 (*a*). ABCG2 was almost exclusively expressed in the $\alpha 2\beta 1^+$ population (*b*). *C* and *D*, $\alpha 2\beta 1$ and ABCG2 expression in CD44⁺ and CD44⁻ Du145 cells. Purified CD44⁺ and CD44⁻ Du145 cells were plated at 5,000 per 18-mm² coverslip and fixed after 4 h. *D*, cells were stained using indirect fluorescence with antibodies against $\alpha 2\beta 1$ (*a* and *b*) or ABCG2 (*c* and *f*). *C*, consistent with (*A*) and (*B*), the majority of the CD44⁺ cells were also positive for $\alpha 2\beta 1$ (*a*). The ABCG2⁺ cell population also localized mostly to the CD44⁺ population (*b*). *D*, *d*, arrowheads, ABCG2⁺ cells.

CD44⁺ populations [i.e., ~70% of the $\alpha 2\beta 1^{+/\text{hi}}$ cells were also CD44⁺ (by prediction, ~30% of the $\alpha 2\beta 1^{+/\text{hi}}$ cells were CD44⁻) and ~30% $\alpha 2\beta 1^{-/\text{lo}}$ cells were CD44⁺ (Fig. 3A,a and B,a-b)]. When double-stained for ABCG2, which normally is expressed in ~1% to 5% of the total population (23), we found that nearly all ABCG2⁺ cells lay within the $\alpha 2\beta 1^{+/\text{hi}}$ subset (Fig. 3A,b and B,c-d).

We then carried out the reciprocal experiment (i.e., we isolated CD44⁺ and CD44⁻ Du145 cells from xenograft tumors and stained them for $\alpha 2\beta 1$ and ABCG2; Fig. 3C and D). As expected, we observed that ~80% of CD44⁺ cells also expressed $\alpha 2\beta 1$ (i.e., CD44⁺ $\alpha 2\beta 1^{+}$) and that essentially all the ABCG2⁺ cells were localized in the CD44⁺ cell population. These results suggest that CD44, $\alpha 2\beta 1$, and ABCG2 mark overlapping populations of prostate cancer cells with differing tumorigenic properties.

Tumorigenic hierarchy revealed by CD44 and $\alpha 2\beta 1$ expression profiles. The above experiments suggest that prostate cancer cells can be stratified, according to their CD44 and $\alpha 2\beta 1$ expression profiles, to at least four different cell populations (i.e., CD44⁺ $\alpha 2\beta 1^{+}$, CD44⁺ $\alpha 2\beta 1^{-}$, CD44⁻ $\alpha 2\beta 1^{+}$, and CD44⁻ $\alpha 2\beta 1^{-}$). We used the LAPC9 xenograft tumor model to address whether these populations of cells have intrinsic differences with respect to their tumorigenicity. In LAPC9 tumors, these four populations of cells represent $0.28 \pm 0.2\%$, $9 \pm 4.7\%$, $1.4 \pm 0.8\%$, and $91 \pm 5.3\%$ ($n = 5$), respectively (Fig. 4A), suggesting that the bulk of the tumor

cell population is CD44⁻ $\alpha 2\beta 1^{-/\text{lo}}$ and the single-positive and double-positive cells represent the minority. Because we have previously shown that most CD44⁺ LAPC9 cells are androgen receptor negative and essentially all androgen receptor-positive cells are localized in the CD44⁻ cell population (24), the above results (Fig. 4A) suggest that the bulk of the LAPC9 tumors may be differentiated cells. Indeed, immunostaining revealed that most tumor cells were androgen receptor positive (not shown). When 1,000 LAPC9 cells of each phenotype were injected s.c. into male NOD/SCID mice supplemented with testosterone, the CD44⁺ $\alpha 2\beta 1^{+}$ and CD44⁺ $\alpha 2\beta 1^{-}$ cells showed similar tumor take (i.e., 100%) and tumor growth (Table 2; Fig. 4B). The CD44⁻ $\alpha 2\beta 1^{+}$ LAPC9 cells showed slightly lower tumor take (i.e., 90%) and initiated smaller tumors ($P = 0.022$; Table 2; Fig. 4B). The CD44⁻ $\alpha 2\beta 1^{-}$ cells showed the lowest tumor incidence (i.e., 40%) and tumors were significantly smaller than the CD44⁺ $\alpha 2\beta 1^{+}$ or CD44⁺ $\alpha 2\beta 1^{-}$ cell-initiated tumors (Table 2; Fig. 4B). When 10,000 highly purified LAPC9 cells of each phenotype were used in the same set of experiments, similar patterns in tumorigenicity were observed although tumors initiated by CD44⁺ $\alpha 2\beta 1^{-}$ and CD44⁻ $\alpha 2\beta 1^{+}$ cells were of similar sizes (Table 2; Fig. 4C). When 10,000 acutely purified CD44⁺ $\alpha 2\beta 1^{+}$ or CD44⁻ $\alpha 2\beta 1^{-}$ LAPC9 cells were injected s.c. into the male NOD/SCID mice without exogenous testosterone, the CD44⁺ $\alpha 2\beta 1^{+}$ cells generated more tumors, which were nearly 20 times bigger




Figure 4. Hierarchical organization of tumor cells in LAPC9 xenograft tumors. **A**, the percentages of tumor cells of different phenotypes in LAPC9 xenograft tumors analyzed by flow cytometry analysis. Columns, mean ($n = 5$); bars, SD. **B** and **C**, tumor experiments. One thousand (**B**) or 10,000 (**C**) acutely purified LAPC9 cells of four different phenotypes were injected s.c. in Matrigel into male NOD/SCID mice supplemented with exogenous testosterone pellets. Tumor images and incidences were indicated. See Table 2 for more details. **D**, a hypothetical model of hierarchical organization of prostate cancer cells. Prostate tumors contain the bulk differentiating and differentiated mature tumor cells that express CD57, androgen receptor (AR), and prostate-specific antigen (PSA), as well as small populations of undifferentiated tumorigenic cells that can be identified by CD44 expression. The CD44⁺ cell population encompasses both tumor progenitors that are ABCG2⁺ and $\alpha 2\beta 1^{+}$ and are highly proliferative (bottom) and slow-cycling cancer stem cells (CSC) that seem to be ABCG2⁻ and $\alpha 2\beta 1^{-}$. Cancer stem cells are hypothesized to have the ability to self-renew. See text for more details.

Table 2. Tumorigenicity of s.c implanted double-sorted LAPC9 cells

Phenotype	No. cells	Incidence (%) [*]	Termination (d) [†]	Weight (g) [‡]	<i>P</i> [§]
With dihydrotestosterone					
CD44 ⁺ $\alpha 2\beta 1$ ⁺	1,000	6/6 (100)	60	0.43 \pm 0.30 (0.02–0.72)	
CD44 ⁺ $\alpha 2\beta 11$ ⁻	1,000	10/10 (100)	60	0.45 \pm 0.57 (0.09–1.76)	0.63
CD44 ⁻ $\alpha 2\beta 11$ ⁺	1,000	9/10 (90)	60	0.19 \pm 0.17 (0.05–0.48)	0.022
CD44 ⁻ $\alpha 2\beta 11$ ⁻	1,000	2/5 (40) [¶]	60	0.03 \pm 0.05 (0.03–0.11) ^{**}	0.017
CD44 ⁺ $\alpha 2\beta 11$ ⁺	10,000	6/6 (100)	60	0.48 \pm 0.39 (0.16–1.1)	
CD44 ⁺ $\alpha 2\beta 11$ ⁻	10,000	8/8 (100)	60	0.18 \pm 0.21 (0.02–0.65)	0.09
CD44 ⁻ $\alpha 2\beta 11$ ⁺	10,000	6/8 (75)	60	0.28 \pm 0.25 (0.04–0.60)	0.15
$\alpha 2\beta 11$ ⁻ CD44 ⁻	10,000	1/6 (17) [¶]	60	0.3	0.029
No dihydrotestosterone					
$\alpha 2\beta 11$ ⁺ /CD44 ⁺	10,000	5/6 (83)	55	1.46 \pm 0.5 (0.55–1.1)	
$\alpha 2\beta 11$ ⁻ /CD44 ⁻	10,000	3/6 (50) [¶]	55	0.08 \pm 0.04 (0.03–0.11)	0.035

^{*}Tumor cells were injected in Matrigel s.c. into male NOD/SCID mice supplemented with or without testosterone pellets. Tumor incidence (indicated as percent in the parentheses) refers to the number of tumors developed/number of injections.

[†]Time (in days) when animals were terminated.

[‡]Mean \pm SD. Tumor weight range is indicated in parentheses.

[§]Unpaired Student's *t* test (all compared with double positive cells).

^{||}*P* = 0.19, compared with CD44⁺ $\alpha 2\beta 11$ ⁻.

[¶]*P* < 0.05, compared with double-positive cells.

^{**}*P* = 0.05, compared with CD44⁺ $\alpha 2\beta 11$ ⁻.

than the tumors initiated by corresponding CD44⁻ $\alpha 2\beta 1$ ⁻ cells (Table 2). Interestingly, tumors initiated by 10,000 CD44⁺ $\alpha 2\beta 1$ ⁺ cells in male NOD/SCID mice without exogenous testosterone (Table 2, *bottom*) were about thrice larger than the tumors initiated by the same number of CD44⁺ $\alpha 2\beta 1$ ⁺ cells in mice with exogenous testosterone (Table 2, *top*). Similar differences in tumorigenicity between the double-positive and double-negative LAPC9 cell populations were also observed when 10,000 CD44⁺ $\alpha 2\beta 1$ ⁺ or CD44⁻ $\alpha 2\beta 1$ ⁻ cells were implanted in female NOD/SCID mice, with average tumor weights being 1.62 and 0.029 g (*P* < 0.001), respectively.

Discussion

It has long been known that human tumors, although clonal by origin, are rather heterogeneous in their cellular composition. The cancer stem cell hypothesis helps explain this biological conundrum (1). Recent studies in breast and colon cancers as well as gliomas suggest that tumor cells *in vivo* may indeed be organized as a hierarchy with tumor-initiating cells or cancer stem cells sitting at the apex and having the ability to develop (or differentiate) into a spectrum of more mature progeny (see Introduction). Our work (9, 23, 24) and the work of Collin et al. (22) suggest that human prostate cancer cells may also be organized as a hierarchy. The present study provides concrete experimental evidence for a hierarchical organization of tumor cells in xenograft human prostate tumors.

Our previous studies have revealed that in several xenograft human tumors, the CD44⁺ cell population is enriched in tumorigenic prostate cancer stem and progenitor cells (9, 24). When highly purified cells are used in surgical orthotopic implantation experiments, most tumorigenicity and all metastatic ability are localized in the CD44⁺ population (24). Serial sphere-formation assays, label-retaining experiments, "stemness" gene expression profiling, clonal analyses, and asymmetrical segregation of CD44

indicate that the CD44⁺ prostate cancer cell population is still heterogeneous, with only ~1% of the cells in this population representing cancer stem cells and the majority representing highly proliferative tumor progenitors (9, 24). These observations suggest that the CD44⁺ prostate cancer cell population is still heterogeneous consisting of perhaps subsets of cells with differing tumor-initiating abilities (hence the name CD44⁺ prostate cancer stem/progenitor cells; ref. 9). This suggestion is consistent with studies showing that the leukemic stem cells in acute myelogenous leukemia (AML) identified by CD34⁺CD38⁻ (29) and colon cancer-initiating cells (or colon cancer stem cell) identified by CD133 (6) are heterogeneous populations of cells with true cancer stem cells representing only a minor subset (i.e., 0.1–1%).

Bearing in mind that the CD44⁺ prostate cancer cell population is heterogeneous, we seek to further dissect tumorigenic prostate cancer cell subsets in the current study. We put our focus on $\alpha 2\beta 1$ ^{+/hi} prostate cancer cells because this integrin receptor has been reported to mark a population of normal human prostate stem cells (21). Of great interest, highly purified $\alpha 2\beta 1$ ^{+/hi} prostate cancer cells *in vitro* possess higher cloning and clonogenic potentials than the corresponding $\alpha 2\beta 1$ ^{-/lo} cells. However, when put *in vivo*, either orthotopically in dorsal prostate or s.c., the $\alpha 2\beta 1$ ^{+/hi} cells exhibit very similar tumorigenicity to $\alpha 2\beta 1$ ^{-/lo} cells. The different behavior of the $\alpha 2\beta 1$ ^{+/hi} prostate cancer cells *in vitro* and *in vivo* is in sharp contrast to the CD44⁺ prostate cancer cells (ref. 24, and this study) and highlights the critical importance of carrying out tumor experiments when assaying candidate cancer stem cell populations because cells showing enhanced *in vitro* clonal capacity and clonogenic potential may not necessarily represent primitive tumor-initiating cells (and therefore should not be called cancer stem cells because tumor initiation is the "gold" standard in defining cancer stem cell; refs. 8, 9).

On the other hand, the $\alpha 2\beta 1$ ^{+/hi} prostate cancer cells behave very similarly to the ABCG2⁺ cancer cells, which represent

fast-proliferating tumor progenitors (23). Indeed, both ABCG2⁺ and $\alpha 2\beta 1^{+/\text{hi}}$ prostate cancer cells are highly proliferative and clonogenic *in vitro* but are not more tumorigenic than the corresponding marker negative cell populations (ref. 23, and this study). Double immunofluorescence staining experiments reveal that the ABCG2⁺ cells are all localized in the $\alpha 2\beta 1^-$ and CD44-expressing cell populations and that ~70% of the $\alpha 2\beta 1^+$ prostate cancer cells are also positive for CD44. Using dual fluorescence sorting, tumor cells in LAPC9 xenografts are fractionated into the bulk (i.e., >90%) CD44⁻ $\alpha 2\beta 1^-$ cells and minor subsets of single-positive or double-positive cells. Remarkably, although constituting <10% of the total tumor cell population, the CD44⁺ $\alpha 2\beta 1^+$ as well as CD44⁺ $\alpha 2\beta 1^-$ cells, on an equal cell number basis, show much higher tumor-initiating abilities than the CD44⁻ $\alpha 2\beta 1^-$ cells. These observations, together with our findings that most CD44⁺ prostate cancer cells are androgen receptor negative (24), lead us to propose a hypothetical model for the hierarchical organization of human prostate cancer cells (Fig. 4D). The bulk of the tumor cells in prostate cancer is differentiating and differentiated cells expressing androgen receptor and prostate-specific antigen. These mature cells possess low tumor-initiating activity whereas most tumorigenicity resides in the minor CD44⁺ cell population (Fig. 4D; ref. 24).

Several pieces of evidence provide support for this model. First, most tumorigenicity resides in the relatively small population of CD44⁺ cells, which range from ~1% to 20% in xenograft tumors (ref. 24, and this study). In primary patient tumors, interestingly, the percentage of CD44⁺ cells seems to correlate with the Gleason grade, with Gleason grade 6 to 9 tumors having ~3%, 9%, 18%, and 19% of CD44⁺ prostate cancer cells.⁴ Second, the CD44⁺ prostate cancer cell population is still heterogeneous, encompassing tumor progenitor cells that are ABCG2⁺ $\alpha 2\beta 1^+$ and relatively quiescent, slow-cycling cancer stem cells that are CD44⁺ABCG2⁻ $\alpha 2\beta 1^-$ (Fig. 4D; ref. 24, and this study). In support of this conjecture, all ABCG2⁺ cells and most (i.e., 70–80%) of the $\alpha 2\beta 1^+$ cells are included in the CD44⁺ cell population, and overall, the CD44⁺ $\alpha 2\beta 1^+$ and CD44⁺ $\alpha 2\beta 1^-$ LAPC9 cells have very similar tumorigenicities. In fact, the tumorigenicity of CD44⁺ (i.e., sorted using a single marker) cells is also indistinguishable from that of CD44⁺ $\alpha 2\beta 1^+$ or CD44⁺ $\alpha 2\beta 1^-$ cells (ref. 24, and this study), suggesting that fluorescence-activated cell sorting (FACS) using either CD44 alone or CD44/ $\alpha 2\beta 1$ combination is purifying practically the same prostate cancer cell population. Primary human tumors also reveal that ~75% of the $\alpha 2\beta 1^+$ cells are localized in the CD44⁺ prostate cancer cell population. Third, the $\alpha 2\beta 1^+$ and $\alpha 2\beta 1^-$ cells are not significantly different in terms of their tumorigenicity, which can be explained by the fact that ~30% of the CD44⁺ cells are localized in the $\alpha 2\beta 1^-$ cell population (Fig. 3). In fact, the $\alpha 2\beta 1^-$ population seems to be slightly enriched in tumorigenic cells. For example, 100,000 $\alpha 2\beta 1^-$ Du145 cells orthotopically implanted in the dorsal prostate can initiate tumor in four of the four injections, whereas the same number of unfractionated Du145 cells cannot initiate any tumor development (Table 1). In addition, all tumors derived from the $\alpha 2\beta 1^-$ cells contain small numbers of $\alpha 2\beta 1^+$ cells (see Supplementary Table S1). Remarkably, in tumors derived from high numbers (i.e., 100,000) of the $\alpha 2\beta 1^-$ LAPC4 or LAPC9 cells, more $\alpha 2\beta 1^+$ cells are observed than in unsorted tumors

(Supplementary Table S1). All these observations support the hypothesis that $\alpha 2\beta 1^-$ population contains more primitive cells that can “regenerate” $\alpha 2\beta 1^+$ cells. Furthermore, when injected s.c., 100 $\alpha 2\beta 1^-$ LAPC9 cells, like the unsorted cells, can initiate 50% tumor development whereas 10 times more $\alpha 2\beta 1^+$ cells are required to achieve similar tumor take. These data suggest that ~30% of the CD44⁺ prostate cancer cells that are $\alpha 2\beta 1^-$ might harbor primitive self-renewing cancer stem cells (Fig. 4D). Fourth, the CD44⁺ $\alpha 2\beta 1^-$ cells and CD44⁻ $\alpha 2\beta 1^+$ cells behave very similarly, in terms of their tumor-initiating abilities, to the $\alpha 2\beta 1^-$ and $\alpha 2\beta 1^+$ cells, respectively. In addition, we have previously shown that 1,000 CD44⁻ LAPC9 cells injected s.c. can initiate tumor development in five of the six injections (24), suggesting that there exist tumorigenic cells in the CD44⁻ cell population. In the present study, we find that 1,000 highly purified CD44⁻ $\alpha 2\beta 1^+$ cells initiate tumor development in 9 of the 10 implantations (Fig. 4B), suggesting that tumorigenic cells in CD44⁻ population might all be $\alpha 2\beta 1^+$ (i.e., having the CD44⁻ $\alpha 2\beta 1^+$ phenotype). These results emphasize the important concept that tumor progenitor cells, like the putative primitive cancer stem cells, can be tumorigenic in regular tumor assays. Presumably, exhaustive serial tumor transplantation experiments can functionally distinguish putative cancer stem cells from tumor progenitors (Fig. 4D; ref. 9). Finally, in all xenograft models (DNp53-T, Du145, LAPC4, and LAPC9) as well as primary patient samples we have studied, the percent of CD44⁺ cells is always higher than that of $\alpha 2\beta 1$ (24),⁵ supporting that CD44 marks both cancer stem cells and tumor progenitors whereas $\alpha 2\beta 1$ expression identifies a subset of tumor progenitors (Fig. 4D).

The most remarkable finding in this study is that the CD44⁺ cell population (including both CD44⁺ $\alpha 2\beta 1^+$ and CD44⁺ $\alpha 2\beta 1^-$ cells) has much higher tumorigenicity than CD44⁻ $\alpha 2\beta 1^-$ cells. Intriguingly, when the CD44⁺ $\alpha 2\beta 1^+$ cells are implanted in either male NOD/SCID mice without exogenous testosterone or in female NOD/SCID mice, tumors developed are more than thrice larger than when the double-positive cells are implanted in male NOD/SCID mice supplemented with exogenous testosterone pellets. Because the former two experimental settings are likely androgen deficient, it is tempting to speculate that under these conditions, the cancer stem cell-containing CD44⁺ $\alpha 2\beta 1^+$ cell population, mostly androgen receptor negative (24), may preferentially proliferate, leading to higher tumor growth.

An obvious question pertains to the phenotypic properties of the putative cancer stem cells in the CD44⁺ prostate cancer cell population (Fig. 4D). The CD133⁺ cells may represent good candidates because they have been reported to mark normal prostate stem cells (21) and potential prostate cancer stem cells with higher clonogenic potential (although tumorigenic potential has not been studied; ref. 22). We have also found that primary patient tumor samples contain 0.25% to 1.4% CD133⁺ cells and that the CD133⁺ prostate cancer cells purified from LAPC4 xenograft and HPCa13 patient tumors possess higher clonal and clonogenic potentials.⁴ Studies are under way to characterize the *in vivo* tumorigenicity of CD44⁺CD133⁺ prostate cancer cells and to determine whether they may represent human prostate cancer stem cells. Of particular interest, CD133 has recently been used as a marker to prospectively identify brain and colon tumor-initiating

⁴ L. Patrawala and D.G. Tang, unpublished observations.

⁵ This study; unpublished observations.

cells (5–7), suggesting that this surface molecule, whose biological functions are yet to be elucidated, may represent more or less a “universal” normal stem cell and cancer stem cell marker. Another potential candidate population of primitive prostate cancer stem cells might be in side population, which, in LAPC9 tumors, represents ~0.1% and has even higher tumorigenicity than the CD44⁺ cells (23, 24).

Regardless, our work (ref. 24, and this study) provides concrete experimental rationale for using CD44 as a marker to identify tumorigenic prostate cancer cells. This rationale is also in line with others’ studies using CD44 as the positive surface marker to identify tumor-initiating cells in breast (4), head and neck (30), and pancreatic (31) cancers. Our findings that the CD44⁺ prostate cancer cell population likely contains both cancer stem cells and tumor progenitors (ref. 24, and this study) and that the CD44⁺ $\alpha 2\beta 1$ ⁻ cells, which constitute the bulk of the tumor, are much less tumorigenic (Fig. 4D) suggest that targeting CD44⁺ prostate cancer cell population may represent a viable approach to prostate cancer therapy. This therapeutic strategy is supported

by the recent finding that CD44 is required for leukemic stem cells to engraft in the bone marrow (32, 33) and strongly encouraged by the success of using anti-CD44 antibody to target AML stem cells and nearly cure the disease in mice (33). We are currently designing experimental therapeutics to specifically target the CD44⁺ prostate cancer cells.

Acknowledgments

Received 2/5/2007; revised 3/30/2007; accepted 5/2/2007.

Grant support: NIH grant AG023374, American Cancer Society grant RSG MGO-105961, Department of Defense, Prostate Cancer Foundation, and Elsa Pardee Foundation (D.G. Tang); Cancer Center Support Grants CCSG-5 P30 CA166672 and ES07784; and a predoctoral fellowship from Department of Defense no. W81XWH-06-1-023 (L. Patrawala).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

We thank K. Claypool for assistance in FACS, the Animal Facility Core for help in tumor experiments, and members of the Tang lab for support and helpful discussion. We apologize to the colleagues whose work could not be cited due to space constraints.

References

- Reya T, Morrison SJ, Clarke MF, Weissman IL. Stem cells, cancer, and cancer stem cells. *Nature* 2001;414:105–11.
- Pardoll R, Clarke MF, Morrison SJ. Applying the principles of stem-cell biology to cancer. *Nat Rev Cancer* 2003;3:895–902.
- Bonnet D, Dick JE. Human acute myeloid leukemia is organized as a hierarchy that originates from a primitive hematopoietic cell. *Nat Med* 1997;3:730–7.
- Al-Hajj M, Wicha MS, Benito-Hernandez A, Morrison SJ, Clarke MF. Prospective identification of tumorigenic breast cancer cells. *Proc Natl Acad Sci USA* 2003;100:3983–8.
- Singh SK, Hawkins C, Clarke ID, et al. Identification of human brain tumor initiating cells. *Nature* 2004;432:396–401.
- O’Brien CA, Pollett A, Gallinger S, Dick JE. A human colon cancer cell capable of initiating tumour growth in immunodeficient mice. *Nature* 2007;445:106–10.
- Ricci-Vitiani L, Lombardi DG, Pilozzi E, et al. Identification and expansion of human colon-cancer-initiating cells. *Nature* 2007;445:111–5.
- Clarke MF, Dick JE, Dirks PB, et al. Cancer stem cells—perspectives on current status and future directions: AACR workshop on cancer stem cells. *Cancer Res* 2006;66:9339–44.
- Tang DG, Patrawala L, Calhoun T, et al. Prostate cancer stem/progenitor cells: identification, characterization, and implications. *Mol Carcinog* 2007;46:1–14.
- Reiter RE, Sawyers CL. Xenograft models and the molecular biology of human prostate cancer. In: Chung LWK, Isaacs WB, Simons JW, editors. *Prostate cancer: biology, genetics, and the new therapeutics*. Totowa (NJ): Humana Press Inc.; 2001. p. 163–74.
- Bhatia B, Tang S, Yang P, et al. Cell-autonomous induction of functional tumor suppressor 15-lipoxygenase 2 (15-LOX2) contributes to replicative senescence of human prostate progenitor cells. *Oncogene* 2005;24:3583–95.
- Liu AY, True LD, LaTray L, et al. Cell-cell interaction in prostate gene regulation and cytodifferentiation. *Proc Natl Acad Sci USA* 1997;94:10705–10.
- McDonnell TJ, Troncoso P, Brisbay SM, et al. Expression of the proto-oncogene bcl-2 in the prostate and its association with emergence of androgen-independent prostate cancer. *Cancer Res* 1992;52:6940–4.
- Signoretti S, Waltregny D, Dilks J, et al. p63 is a prostate basal cell marker and is required for prostate development. *Am J Pathol* 2000;157:1769–75.
- Myers RB, Grizzle WE. Changes in biomarker expression in the development of prostatic adenocarcinoma. *Biotech Histochem* 1997;72:86–95.
- Bui M, Reiter RE. Stem cell genes in androgen-independent prostate cancer. *Cancer Metastasis Rev* 1999;17:391–9.
- Cookson MS, Reuter VE, Linkov I, Fair WR. Glutathione *S*-transferase (GST- π) class expression by immunohistochemistry in benign and malignant prostate tissue. *J Urol* 1997;157:673–6.
- Bonkhoff H, Stein U, Remberger K. The proliferative function of basal cells in the normal and hyperplastic human prostate. *Prostate* 1994;24:42–6.
- Hayward SW, Haughney PC, Rosen MA, et al. Interactions between adult human prostate epithelium and rat urogenital sinus mesenchyme in a tissue recombination model. *Differentiation* 1998;63:131–40.
- Collins AT, Habib FK, Maitland NJ, Neal DE. Identification and isolation of human prostate epithelial stem cells based on $\alpha 2\beta 1$ -integrin expression. *J Cell Sci* 2001;114:3865–72.
- Richardson GD, Robson CN, Lang SH, Neal DE, Maitland NJ, Collins AT. CD133, a novel marker for human prostatic epithelial stem cells. *J Cell Sci* 2004;117:3539–45.
- Collins AT, Berry PA, Hyde C, Stower MJ, Maitland NJ. Prospective identification of tumorigenic prostate cancer stem cells. *Cancer Res* 2005;65:10946–51.
- Patrawala L, Calhoun T, Schneider-Broussard R, Zhou J-J, Claypool K, Tang DG. Side population (SP) is enriched in tumorigenic, stem-like cancer cells whereas ABCG2⁺ and ABCG2⁻ cancer cells are similarly tumorigenic. *Cancer Res* 2005;65:6207–19.
- Patrawala L, Calhoun T, Schneider-Broussard R, et al. Highly purified CD44⁺ prostate cancer cells from xenograft human tumors are enriched in tumorigenic and metastatic progenitor cells. *Oncogene* 2006;25:1696–708.
- Knox JD, Cress AE, Clark V, et al. Differential expression of extracellular matrix molecules and the $\alpha 6$ -integrins in the normal and neoplastic prostate. *Am J Pathol* 1994;145:167–74.
- Janes SM, Watt FM. New roles for integrins in squamous-cell carcinoma. *Nat Rev Cancer* 2006;6:175–83.
- van Bokhoven A, Varella-Garcia M, Korch C, et al. Molecular characterization of human prostate carcinoma cell lines. *Prostate* 2003;57:205–25.
- Wang SY, Revelo MP, Sudilovsky D, et al. Development and characterization of efficient xenograft models for benign and malignant human prostate. *Prostate* 2005;64:149–59.
- Hope KJ, Jin L, Dick JE. Acute myeloid leukemia originates from a hierarchy of leukemic stem cell classes that differ in self-renewal capacity. *Nat Immunol* 2004;5:738–43.
- Prince ME, Sivanandan R, Kaczorowski A, et al. Identification of a subpopulation of cells with cancer stem cell properties in head and neck squamous cell carcinoma. *Proc Natl Acad Sci USA* 2007;104:973–8.
- Li C, Heidt DG, Dalerba P, et al. Identification of pancreatic cancer stem cells. *Cancer Res* 2007;67:1030–7.
- Krause DS, Lazarides K, von Andrian UH, Van Etten RA. Requirement for CD44 in homing and engraftment of BCR-ABL-expressing leukemic stem cells. *Nat Med* 2006;12:1175–80.
- Jin L, Hope KJ, Zhai Q, Smadja-Joffe F, Dick JE. Targeting of CD44 eradicates human acute myeloid leukemic stem cells. *Nat Med* 2006;12:1167–74.