

# Occupational Therapy and Management of Multiple Chronic Conditions in the Context of Health Care Reform

Natalie E. Leland, Donald J. Fogelberg, Ashley D. Halle, Tracy M. Mroz

One in four individuals living in the United States has multiple chronic conditions (MCCs), and the already high prevalence of MCCs continues to grow. This population has high rates of health care utilization yet poor outcomes, leading to elevated concerns about fragmented, low-quality care provided within the current health care system. Several national initiatives endeavor to improve care for the population with MCCs, and occupational therapy is uniquely positioned to contribute to these efforts for more efficient, effective, client-centered management of care. By integrating findings from the literature with current policy and practice, we aim to highlight the potential role for occupational therapy in managing MCCs within the evolving health care system.

Leland, N. E., Fogelberg, D. J., Halle, A. D., & Mroz, T. M. (2017). Health Policy Perspectives—Occupational therapy and management of multiple chronic conditions in the context of health care reform. *American Journal of Occupational Therapy, 71*, 7101090010. <https://doi.org/10.5014/ajot.2017.711001>

**Natalie E. Leland, PhD, OTR/L, BCG, FAOTA**, is Assistant Professor, Mrs. T. H. Chan Division of Occupational Science and Occupational Therapy and Davis School of Gerontology, University of Southern California, Los Angeles; [nleland@usc.edu](mailto:nleland@usc.edu)

**Donald J. Fogelberg, PhD, OTR/L**, is Assistant Professor, Division of Occupational Therapy, Department of Rehabilitation Medicine, University of Washington, Seattle.

**Ashley D. Halle, OTD, OTR/L**, is Assistant Professor and Coordinator of Primary Care Residency and Services, Mrs. T. H. Chan Division of Occupational Science and Occupational Therapy, University of Southern California, Los Angeles.

**Tracy M. Mroz, PhD, OTR/L**, is Assistant Professor, Division of Occupational Therapy, Department of Rehabilitation Medicine, University of Washington, Seattle.

The prevalence of people with multiple chronic conditions (MCCs) continues to grow. Despite the disproportionately high health care spending of this population, outcomes are poor in part because of fragmented, low-quality care provided within a system that is not well designed to support efficient, effective, client-centered management of MCCs (Anderson, 2010; Gerteis et al., 2014; Parekh, Goodman, Gordon, & Koh, 2011). System redesign, such as primary care transformation and the move to true patient-centered care, is occurring, especially in relation to MCCs. Achieving efficient care, reducing costs, and improving health in this group are critical to the entire health care system because their care affects access and system costs for all. The occupational therapy role in this redesign must be nurtured in relation to MCCs to ensure that the knowledge, skills, and precepts of the profession are fully used to improve health for all patients, including people with MCCs.

Historically, the U.S. health care system has been structured around a reactive response to acute medical issues and therefore has not effectively addressed the chronic care needs of the population. Care is often siloed, with different providers addressing specific conditions or medical issues, which has

resulted in poor care for people with MCCs (Parekh et al., 2011). To address this longstanding, suboptimal care delivery system for those with MCCs and for other patients, recent health care reform initiatives have promoted prevention, targeted the improvement of population health, and incentivized the delivery of high-quality care (U.S. Department of Health and Human Services [HHS], 2010). To these ends, several national health care initiatives are focusing on improving care for people with MCCs, including enhancing primary care service delivery. Integrating findings from the literature with current policy and practice, this article highlights the potential roles for occupational therapy in addressing participation restrictions and promoting self-management of MCCs within the evolving health care system.

## Prevalence and Impact of Multiple Chronic Conditions

More than 25% of the U.S. community-living adult population has been diagnosed with MCCs, and their numbers are increasing exponentially (Anderson, 2010; Office of the Assistant Secretary for Planning and Evaluation [OASPE], 2010). Among older

adults, the prevalence is even higher, with as many as 75% of people age 65 and older having MCCs (Anderson, 2010; Lochner & Cox, 2013; Wolff, Starfield, & Anderson, 2002). Together, this population is responsible for more than 65% of U.S. health care spending, primarily as a result of progressive functional limitations and exacerbations of their chronic conditions (Anderson, 2010). People with MCCs have higher rates of emergency department visits; more outpatient visits; and longer, more expensive hospital stays (Gerteis et al., 2014; Lochner & Cox, 2013; Lochner, Goodman, Posner, & Parekh, 2013; Skinner, Coffey, Jones, Heslin, & Moy, 2016). Among Medicare beneficiaries, those with a greater number of chronic conditions have higher Medicare expenditures, emergency department visits, and 30-day hospital readmissions (Centers for Medicare and Medicaid Services [CMS], 2014a, 2014b).

Although more research is needed to better understand specific clusters of chronic conditions, people with MCCs are faced with the task of simultaneously managing a combination of chronic conditions, such as diabetes, hypertension, ischemic heart disease, heart failure, atrial fibrillation, arthritis, chronic kidney disease, chronic obstructive pulmonary disease, depression, and cancer (Vogeli et al., 2007). Although clients typically enter a health care encounter with a primary medical diagnosis, they frequently have multiple comorbid conditions that need to be taken into account to achieve a successful outcome.

Despite understanding the importance of addressing these multiple comorbid conditions to achieve successful outcomes, clinical outcomes for people with MCCs remain poor. This is due in part to aspects of the health care system that are not well structured to manage MCCs effectively (Gerteis et al., 2014). Health care is often delivered by multiple professionals with minimal coordination and communication between providers, leading to fragmentation, duplication of services, and diffusion of responsibility (Parekh et al., 2011). Episodes of care are too often focused on a single diagnosis, with limited attention paid to other conditions that may be present, diminishing the overall effectiveness and

quality of care (Anderson, 2010; OASPE, 2010). Moreover, although single-disease management programs have proliferated, limited client education on general principles of chronic disease management makes self-management especially challenging for those with MCC (Anderson, 2010; OASPE, 2010).

Not only do MCCs result in high rates of health care utilization and correspondingly high costs, they also have a negative impact on people's occupational performance and quality of life (Lochner & Shoff, 2015). For example, Barstow, Warren, Thaker, Hallman, and Batts (2015) found that clients with MCCs experience a loss of independence in self-care and of safe participation in desired social and leisure activities. These losses are further exacerbated by the psychosocial implications of a new diagnosis: People have reported feeling a loss of control over their own life, a theme reflected across multiple client populations (Brereton & Nolan, 2000; Pyatak, 2011; Wood, Connelly, & Maly, 2010). Research has demonstrated that occupational therapy providers are effective at tackling these common concerns related to MCC management by approaching these care needs from an occupational lens (Arbesman & Mosley, 2012; Clark et al., 2012). By understanding current policy directions, occupational therapy leaders can help position the profession to take a key role in these changes.

## Policy Initiatives Related to Multiple Chronic Conditions

Several policy initiatives have been implemented in response to the high level of health care utilization among people with MCCs and the persistent poor outcomes they experience. The goal of these national policy initiatives is to enhance care delivery, promote comprehensive care for this high-risk population, reduce adverse events, and improve client outcomes (Bazemore, Petterson, Peterson, & Phillips, 2015).

Efforts to enhance the health of this population are being developed and implemented from a public health approach through community-based group interventions and within the health care system (Ahn, Jiang, Smith, & Ory, 2014; Franek, 2013; Nolte & Osborne, 2013; Ory et al.,

2014). Self-management is a core target for improving health care generally but especially for management of MCCs. For instance, Stanford's peer mentored Chronic Disease Self-Management Program is known worldwide (Brady et al., 2013; Kahvecioglu, Moore, Michaelides, Ruiz, & Bertrand, 2011; Lorig, Sobel, Ritter, Laurent, & Hobbs, 2001). Occupational therapy, with its focus on habits, routines, and self-direction, fits well into any efforts to support clients in improving their own approach to health.

Within the traditional health care system, efforts are also being made to transition from a reactionary to a proactive model of care delivery. To this end, HHS (2010) has developed a strategic framework aimed at enhancing the quality of care for clients with MCCs. The framework emphasizes holistic and coordinated care for people with MCCs instead of disease-specific siloed care. Specifically, HHS has four overarching goals: targeting system redesign, which includes changes to the health care and public health systems; empowering clients to engage in their own care; and enhancing the evidence base by developing tools, training, and clinical decision supports and by funding necessary research to improve care delivery (HHS, 2010).

Primary care has been identified as one clinical area in which chronic disease management can be enhanced. CMS's Center for Medicare and Medicaid Innovation, established by the Patient Protection and Affordable Care Act of 2010 (Pub. L. 111-148), has multiple ongoing projects aimed at enhancing and expanding primary and coordinated care to enhance the quality of outcomes for clients with MCCs, such as the Comprehensive Primary Care Initiative and the recent Comprehensive Primary Care Plus model and the Patient-Centered Medical Home Initiative.

## Enhancing Care of Multiple Chronic Conditions: Occupational Therapy's Contribution

Occupational therapy has a long-standing history of treating clients with MCCs in acute and postacute settings to improve their occupational engagement. In these

settings, occupational therapy practitioners typically consider MCCs in addition to the primary medical diagnosis and integrate secondary and tertiary prevention strategies into the care plan to optimize client outcomes, reduce the risk of subsequent medical exacerbations (e.g., acute hospitalization for chronic obstructive pulmonary disease), and prevent adverse outcomes (e.g., infections, accidental falls).

Occupational therapy has an established record of addressing health management, wellness, and prevention in an effort to optimize people's quality of life (Arbesman & Mosley, 2012; Clark et al., 2012; Meyer, 1922). By drawing on the profession's experience in treating clients with MCCs in acute and postacute care settings and its broader history of health promotion and management, the occupational therapy practitioner can assess a client's current knowledge, willingness, and ability to engage in health management and maintenance and other health-promoting occupations within the client's context while taking into account habits, roles, and routines to optimize quality of life (American Occupational Therapy Association [AOTA], 2014).

Occupational therapy can contribute to national priorities aimed at enhancing care quality through the profession's holistic approach to care. Specifically, by evaluating and treating the client as an occupational being, taking into account all the factors that may affect participation, including MCCs, client factors, context and environment, and performance, the occupational therapy practitioner can help facilitate care designed for the client rather than a specific disease or setting. Moreover, the occupational therapy perspective would infuse into these systems new ideas about the interplay between MCCs and the client's habits, roles, and routines, which may affect their risk for adverse events and poor health outcomes.

Client activation and the delivery of effective holistic self-care management services are central to these national initiatives (HHS, 2010; Venkatesh, Goodrich, & Conway, 2014), both of which are at the core of occupational therapy practice. By taking a client-centered approach to the management of MCCs, occupational

therapy practitioners can collaborate with the client to develop foundational medical management knowledge and integrate healthy lifestyle approaches into the client's daily routine to promote self-management skills (Mroz, Pitonyak, Fogelberg, & Leland, 2015).

### Interdisciplinary Care Teams in Primary Care: A Place of Opportunity

In the context of current federal system change priorities, which are intended to enhance client engagement, promote care coordination, and improve client outcomes, occupational therapy is uniquely situated to be an integral part of the chronic disease management team, including in primary care. Given the profession's specialized education (e.g., mental and physical health, management, and advocacy) and skills in functional assessments, activity analysis, skill development, problem-solving barriers, environmental assessments, adaptation, compensation, and remediation, occupational therapy practitioners are well equipped to support clients in managing their MCCs (Krupa & Clark, 1995; Robinson, Fisher, & Broussard, 2016). To this end, occupational therapy is included among the professions targeted in HHS's (2015) MCC education and training framework, a product that resulted from HHS's (2010) strategic framework for MCC, which challenges professional academic programs to equip providers with the skills to optimize their scope of practice in order to improve care delivery to and patient outcomes among this population.

However, as interdisciplinary primary care teams have emerged, rehabilitation professionals have mostly been overlooked (Bazemore, Wingrove, Peterson, & Petterson, 2016; Peikes, Chen, Schore, & Brown, 2009; Wagner, 2000). Primary care teams are generally led by family physicians and most commonly include nurse practitioners, registered nurses, licensed practical nurses, and physician assistants and, to a lesser extent, pharmacists, behavioral health specialists, and social workers (Bazemore et al., 2016). Although some approaches facilitate the diversification of disciplines represented in primary care

(e.g., pharmacists and behavioral health specialists), occupational therapy practitioners need advocacy and an evidence base to support their claim to a place on the chronic disease management care team in primary care (Hildenbrand & Lamb, 2013; Peikes et al., 2016).

As members of the care team, occupational therapy practitioners can approach the client's functional and medical needs and enhance outcomes from an occupational performance perspective instead of a disease-specific approach. By demonstrating that their skills transcend the clinical context and environment, occupational therapy practitioners can be key contributors to the MCC care team; they can target self-management skills, foster client engagement, and facilitate client and caregiver training with the goal of reducing risk and optimizing participation. These opportunities include facilitating clients' self-management skills for MCCs, engaging in screening for and treatment of adverse events (e.g., readmissions, accidental falls), serving as case managers and care coordinators, and supporting client and caregiver education (Hand, Law, & McColl, 2011; Metzler, Hartmann, & Lowenthal, 2012; Richardson et al., 2014; Sanders & Van Oss, 2013; Taylor, 2004).

For example, complying with and managing medications associated with MCCs is an essential component of health management for this client population. Failure to adhere to a medication routine is associated with adverse events and hospitalizations (Malet-Larrea et al., 2016). Occupational therapy practitioners can work along with the physician and nurse to optimize medication utilization. Occupational therapy practitioners can assess and develop interventions related to the client's functional cognition, physical capacity, memory, and other issues to improve the client's ability to manage medications prescribed by the physician. The occupational therapy evaluation of medication issues can guide recommendations for environmental and personal supports to optimize medication management and adherence (AOTA, 2016). Moreover, occupational therapy practitioners can collaborate with the client to establish and implement a medication routine that

aligns with the physician-prescribed regimen and fits into the client's daily routine. Thus, successful participation in occupations can contribute to effective management of chronic conditions, helping to achieve the core goals of new primary care delivery models, the need to improve MCC care and outcomes, and other policy initiatives (Metzler et al., 2012).

## Conclusions and Next Steps for Occupational Therapy

National initiatives are emphasizing comprehensive care for the high-risk MCC population, reducing adverse events, and improving client outcomes (Bazemore et al., 2015). The occupational therapy practitioner can be a valuable member of the interdisciplinary care team by approaching care of MCCs in a comprehensive manner that transcends diagnoses and takes into account the client's habits, roles, and routines. To achieve this goal, though, the profession needs to take steps to establish a distinct role in both current systems and emerging systems as the transformation of health care continues. We need to generate evidence within the health care context that demonstrates the relationship between occupation-based MCC health management interventions and prioritized system outcomes such as reduced hospital admissions, lower frequency of physician visits, and enhanced medication compliance. In developing our evidence base, it will be imperative to look to other disciplines working in this area to integrate their standardized measures so we can subsequently compare outcomes and programs. Moving forward, if these objectives are achieved, occupational therapy will be well situated to define, deliver, and document its value as part of the interdisciplinary team promoting occupational engagement—and thus improved health—for clients with MCCs. ▲

## Acknowledgments

During the preparation of this article, Natalie E. Leland was supported by the Agency for Healthcare Research and Quality (K01 HS 022907-01A1); Donald J. Fogelberg was supported by the Eunice Kennedy Shriver National Institute of Child Health and

Human Development (K01HD076183); and Ashley D. Halle was supported by the Health Resources and Services Administration, under the Geriatrics Workforce Enhancement Program Award.

## References

- Ahn, S., Jiang, L., Smith, M. L., & Ory, M. G. (2014). Improvements in sleep problems among the chronic disease self-management program participants. *Family and Community Health, 37*, 327–335. <https://doi.org/10.1097/FCH.0000000000000045>
- American Occupational Therapy Association. (2014). Occupational therapy practice framework: Domain and process (3rd ed.). *American Journal of Occupational Therapy, 68*(Suppl. 1), S1–S48. <https://doi.org/10.5014/ajot.2014.682006>
- American Occupational Therapy Association. (2016). *Role of occupational therapy in assessing functional cognition*. Retrieved from <http://www.aota.org/Advocacy-Policy/Federal-Reg-Affairs/Resources/Role-OT-Assessing-Functional-Cognition.aspx>
- Anderson, G. F. (2010). *Chronic care: Making the case for ongoing care*. Princeton, NJ: Robert Wood Johnson Foundation. Retrieved from <http://www.rwjf.org/content/dam/farm/reports/reports/2010/rwjf54583>
- Arbesman, M., & Mosley, L. J. (2012). Systematic review of occupation- and activity-based health management and maintenance interventions for community-dwelling older adults. *American Journal of Occupational Therapy, 66*, 277–283. <https://doi.org/10.5014/ajot.2012.003327>
- Barstow, B. A., Warren, M., Thaker, S., Hallman, A., & Batts, P. (2015). Client and therapist perspectives on the influence of low vision and chronic conditions on performance and occupational therapy intervention. *American Journal of Occupational Therapy, 69*, 6903270010. <https://doi.org/10.5014/ajot.2015.014605>
- Bazemore, A., Petterson, S., Peterson, L. E., & Phillips, R. L., Jr. (2015). More comprehensive care among family physicians is associated with lower costs and fewer hospitalizations. *Annals of Family Medicine, 13*, 206–213. <https://doi.org/10.1370/afm.1787>
- Bazemore, A., Wingrove, P., Peterson, L., & Petterson, S. (2016). The diversity of providers on the family medicine team. *Journal of the American Board of Family Medicine, 29*, 8–9. <https://doi.org/10.3122/jabfm.2016.01.150229>
- Brady, T. J., Murphy, L., O'Colmain, B. J., Beauchesne, D., Daniels, B., Greenberg, M., . . . Chervin, D. (2013). A meta-analysis of health status, health behaviors, and health-care utilization outcomes of the Chronic Disease Self-Management Program. *Preventing Chronic Disease, 10*, 120112. <https://doi.org/10.5888/pcd10.120112>
- Brereton, L., & Nolan, M. (2000). "You do know he's had a stroke, don't you?" Preparation for family care-giving—The neglected dimension. *Journal of Clinical Nursing, 9*, 498–506. <https://doi.org/10.1046/j.1365-2702.2000.00396.x>
- Centers for Medicare and Medicaid Services. (2014a). *Medicare Chronic Conditions Dashboard: County level*. Retrieved from [https://www.cms.gov/Research-Statistics-Data-and-Systems/Statistics-Trends-and-Reports/Dashboard/Chronic-Conditions-County/CC\\_County\\_Dashboard.html](https://www.cms.gov/Research-Statistics-Data-and-Systems/Statistics-Trends-and-Reports/Dashboard/Chronic-Conditions-County/CC_County_Dashboard.html)
- Centers for Medicare and Medicaid Services. (2014b). *Medicare Chronic Conditions Dashboard: State level*. Retrieved from [https://www.cms.gov/Research-Statistics-Data-and-Systems/Statistics-Trends-and-Reports/Dashboard/Chronic-Conditions-State/CC\\_State\\_Dashboard.html](https://www.cms.gov/Research-Statistics-Data-and-Systems/Statistics-Trends-and-Reports/Dashboard/Chronic-Conditions-State/CC_State_Dashboard.html)
- Clark, F., Jackson, J., Carlson, M., Chou, C. P., Cherry, B. J., Jordan-Marsh, M., . . . Azen, S. P. (2012). Effectiveness of a lifestyle intervention in promoting the well-being of independently living older people: Results of the Well Elderly 2 Randomised Controlled Trial. *Journal of Epidemiology and Community Health, 66*, 782–790. <https://doi.org/10.1136/jech.2009.099754>
- Franek, J. (2013). Self-management support interventions for persons with chronic disease: An evidence-based analysis. *Ontario Health Technology Assessment Series, 13*(9), 1–60.
- Gerteis, J. M., Izrael, D., Deitz, D., LeRoy, L., Ricciardi, R., Miller, T., & Basu, J. (2014). *Multiple chronic conditions chartbook: 2010 Medical Expenditure Panel Survey data*. Rockville, MD: Agency for Healthcare Research and Quality.
- Hand, C., Law, M., & McColl, M. A. (2011). Occupational therapy interventions for chronic diseases: A scoping review. *American Journal of Occupational Therapy, 65*, 428–436. <https://doi.org/10.5014/ajot.2011.002071>
- Hildenbrand, W. C., & Lamb, A. J. (2013). Occupational therapy in prevention and wellness: Retaining relevance in a new health care world. *American Journal of Occupational Therapy, 67*, 266–271. <https://doi.org/10.5014/ajot.2013.673001>

- Kahvecioglu, D., Moore, T., Michaelides, M., Ruiz, S., & Bertrand, R. (2011). *Design and evaluation of three Administration on Aging (AoA) programs: Chronic Disease Self-Management Program evaluation design*. Rockville, MD: Agency for Healthcare Research and Quality.
- Krupa, T., & Clark, C. C. (1995). Occupational therapists as case managers: Responding to current approaches to community mental health service delivery. *Canadian Journal of Occupational Therapy, 62*, 16–22.
- Lochner, K. A., & Cox, C. S. (2013). Prevalence of multiple chronic conditions among Medicare beneficiaries, United States, 2010. *Preventing Chronic Disease, 10*, 120137. <https://doi.org/10.5888/pcd10.120137>
- Lochner, K. A., Goodman, R. A., Posner, S., & Parekh, A. (2013). Multiple chronic conditions among Medicare beneficiaries: State-level variations in prevalence, utilization, and cost, 2011. *Medicare and Medicaid Research Review, 23*(3). <https://doi.org/10.5600/mmrr.003.03.b02>
- Lochner, K. A., & Shoff, C. M. (2015). County-level variation in prevalence of multiple chronic conditions among Medicare beneficiaries, 2012. *Preventing Chronic Disease, 12*, E07. <https://doi.org/10.5888/pcd12.140442>
- Lorig, K. R., Sobel, D. S., Ritter, P. L., Laurent, D., & Hobbs, M. (2001). Effect of a self-management program on patients with chronic disease. *Effective Clinical Practice: ECP, 4*, 256–262.
- Malet-Larrea, A., Goyenechea, E., García-Cárdenas, V., Calvo, B., Arteché, J. M., Aranegui, P., . . . Benrimoj, S. I. (2016). The impact of a medication review with follow-up service on hospital admissions in aged polypharmacy patients. *British Journal of Clinical Pharmacology*. Advance online publication. <https://doi.org/10.1111/bcp.13012>
- Metzler, C. A., Hartmann, K. D., & Lowenthal, L. A. (2012). Defining primary care: Envisioning the roles of occupational therapy. *American Journal of Occupational Therapy, 66*, 266–270. <https://doi.org/10.5014/ajot.2010.663001>
- Meyer, A. (1922). The philosophy of occupation therapy. *American Journal of Physical Medicine and Rehabilitation, 1*, 1–10.
- Mroz, T. M., Pitonyak, J. S., Fogelberg, D., & Leland, N. E. (2015). Client centeredness and health reform: Key issues for occupational therapy. *American Journal of Occupational Therapy, 69*, 6905090010. <https://doi.org/10.5014/ajot.2015/695001>
- Nolte, S., & Osborne, R. H. (2013). A systematic review of outcomes of chronic disease self-management interventions. *Quality of Life Research, 22*, 1805–1816. <https://doi.org/10.1007/s11136-012-0302-8>
- Office of the Assistant Secretary for Planning and Evaluation, U.S. Department of Health and Human Services. (2010). *Individuals living in the community with chronic conditions and functional limitations: A closer look*. Falls Church, VA: Lewin Group.
- Ory, M. G., Smith, M. L., Ahn, S., Jiang, L., Lorig, K., & Whitelaw, N. (2014). National study of chronic disease self-management: Age comparison of outcome findings. *Health Education and Behavior, 41* (Suppl.), 34S–42S. <https://doi.org/10.1177/1090198114543008>
- Parekh, A. K., Goodman, R. A., Gordon, C., & Koh, H. K.; HHS Interagency Workgroup on Multiple Chronic Conditions. (2011). Managing multiple chronic conditions: A strategic framework for improving health outcomes and quality of life. *Public Health Reports, 126*, 460–471.
- Patient Protection and Affordable Care Act, Pub. L. 111–148, 42 U.S.C. §§ 18001–18121 (2010).
- Peikes, D., Chen, A., Schore, J., & Brown, R. (2009). Effects of care coordination on hospitalization, quality of care, and health care expenditures among Medicare beneficiaries: 15 randomized trials. *JAMA, 301*, 603–618. <https://doi.org/10.1001/jama.2009.126>
- Peikes, D., Taylor, E. F., Dale, S., O'Malley, S., Ghosh, A., Anglin, G., . . . Brown, R. (2016). *Evaluation of the Comprehensive Primary Care Initiative: Second annual report*. Princeton, NJ: Mathematica Policy Research. Retrieved from <https://innovation.cms.gov/Files/reports/cpci-evalrpt2.pdf>
- Pyatak, E. (2011). Participation in occupation and diabetes self-management in emerging adulthood. *American Journal of Occupational Therapy, 65*, 462–469. <https://doi.org/10.5014/ajot.2011.001453>
- Richardson, J., Loyola-Sanchez, A., Sinclair, S., Harris, J., Letts, L., MacIntyre, N. J., . . . Martin Ginis, K. (2014). Self-management interventions for chronic disease: A systematic scoping review. *Clinical Rehabilitation, 28*, 1067–1077. <https://doi.org/10.1177/0269215514532478>
- Robinson, M., Fisher, T. F., & Broussard, K. (2016). Role of occupational therapy in case management and care coordination for clients with complex conditions. *American Journal of Occupational Therapy, 70*, 7002090010. <https://doi.org/10.5014/ajot.2016.702001>
- Sanders, M. J., & Van Oss, T. (2013). Using daily routines to promote medication adherence in older adults. *American Journal of Occupational Therapy, 67*, 91–99. <https://doi.org/10.5014/ajot.2013.005033>
- Skinner, H. G., Coffey, R., Jones, J., Heslin, K. C., & Moy, E. (2016). The effects of multiple chronic conditions on hospitalization costs and utilization for ambulatory care sensitive conditions in the United States: A nationally representative cross-sectional study. *BMC Health Services Research, 16*, 77. <https://doi.org/10.1186/s12913-016-1304-y>
- Taylor, R. R. (2004). Quality of life and symptom severity for individuals with chronic fatigue syndrome: Findings from a randomized clinical trial. *American Journal of Occupational Therapy, 58*, 35–43. <https://doi.org/10.5014/ajot.58.1.35>
- U.S. Department of Health and Human Services. (2010). *Multiple chronic conditions: A strategic framework: Optimum health and quality of life for individuals with multiple chronic conditions*. Washington, DC: Author. Retrieved from <http://www.hhs.gov/ash/about-ash/multiple-chronic-conditions/addressing-multiple-chronic-conditions/index.html#>
- U.S. Department of Health and Human Services. (2015). *Multiple chronic conditions: A framework for education and training*. Washington, DC: Author. Retrieved from <http://www.hhs.gov/sites/default/files/ash/initiatives/mcc/education-and-training/framework-curriculum/framework-curriculum.pdf>
- Venkatesh, A., Goodrich, K., & Conway, P. H. (2014). Opportunities for quality measurement to improve the value of care for patients with multiple chronic conditions. *Annals of Internal Medicine, 161*(Suppl.), S76–S80. <https://doi.org/10.7326/M13-3014>
- Vogeli, C., Shields, A. E., Lee, T. A., Gibson, T. B., Marder, W. D., Weiss, K. B., & Blumenthal, D. (2007). Multiple chronic conditions: Prevalence, health consequences, and implications for quality, care management, and costs. *Journal of General Internal Medicine, 22*(Suppl. 3), 391–395. <https://doi.org/10.1007/s11606-007-0322-1>
- Wagner, E. H. (2000). The role of patient care teams in chronic disease management. *BMJ, 320*, 569–572. <https://doi.org/10.1136/bmj.320.7234.569>

Wolff, J. L., Starfield, B., & Anderson, G. (2002). Prevalence, expenditures, and complications of multiple chronic conditions in the elderly. *Archives of Internal*

*Medicine*, *162*, 2269–2276. <https://doi.org/10.1001/archinte.162.20.2269>  
Wood, J. P., Connelly, D. M., & Maly, M. R. (2010). “Getting back to real living”: A

qualitative study of the process of community reintegration after stroke. *Clinical Rehabilitation*, *24*, 1045–1056. <https://doi.org/10.1177/0269215510375901>