

Bevacizumab-Induced Vessel Normalization Hampers Tumor Uptake of Antibodies—Response

Marlous Arjaans, Sjoukje F. Oosting, Carolina P. Schröder, and Elisabeth G.E. de Vries

We would like to reply to the letter by Huang and colleagues on the impact of vascular changes induced by antiangiogenic therapy on tumor uptake of other drugs in relation to our findings of reduced antibody uptake after bevacizumab treatment (1). Their comments focus on the bevacizumab dose used and whether vessel normalization was demonstrated.

They consider the bevacizumab dose we used high and suggest that this resulted in "inadequate" rather than "normalized" tumor vessels. The concept of too high bevacizumab dosing is interesting and might have a major impact if translated to the clinic. Huang and colleagues reported on findings in a preclinical study with 10, 20, and 40 mg/kg of the mouse anti-VEGFR2 antibody DC101 where the highest dose resulted neither in vessel normalization nor improved efficacy of immunotherapy (2). It is unclear whether these findings with anti-VEGFR2 antibody can be translated directly to our results with bevacizumab, which binds VEGF-A, affecting both VEGFR1 and VEGFR2 signaling.

In our mouse model, one modest dose of 5 mg/kg bevacizumab reduced the tumor uptake of radiolabeled trastuzumab and aspecific control immunoglobulin G (IgG) by 38% and 27%, respectively, after only 2 days. On day 6, after three 5 mg/kg bevacizumab doses, this decrease was even more pronounced for radiolabeled trastuzumab, bevacizumab, and IgG (1). Moreover, these decreases are all within the described vascular normalization time window (3). We found increased pericyte coverage after bevacizumab treatment, indicating structural vessel normalization. This may not prove functional vessel normalization. However, tumor histology showed no difference in already low percentages of necrosis, tumor viability, or proliferation index between animals receiving bevacizumab or placebo. This is not in line with inadequate vasculature caused by high-dose antiangiogenic therapy (3).

Authors' Affiliations: Department of Medical Oncology, University of Groningen and University Medical Center Groningen, Groningen, the Netherlands

Corresponding Author: Elisabeth G.E. de Vries, Department of Medical Oncology, University of Groningen and University Medical Center Groningen, P.O. Box 30.001, Groningen, 9700 RB, the Netherlands. Phone: 31-50-361-2821; Fax: 31-50-361-4862; E-mail: e.g.e.de.vries@umcg.nl

doi: 10.1158/0008-5472.CAN-13-2532

©2013 American Association for Cancer Research.

References

1. Arjaans M, Oude Munnink TH, Oosting SF, Terwisscha van Scheltinga AG, Gietema JA, Garbaciak ET, et al. Bevacizumab-induced normalization of blood vessels in tumors hampers antibody uptake. *Cancer Res* 2013;73:3347–55.
2. Huang Y, Yuan J, Righi E, Kamoun WS, Ancukiewicz M, Nezivar J, et al. Vascular normalizing doses of antiangiogenic treatment reprogram the

Our findings of reduced uptake of antibodies after antiangiogenic therapy are supported by two preclinical studies. One 10 mg/kg dose of the cross-reactive anti-VEGF antibody B20-4.1 decreased tumor trastuzumab uptake by 50% after 2 days (4) and 10 mg/kg bevacizumab decreased tumor cetuximab uptake by 40% after 4 days (5). Importantly, our findings are also supported in the clinical setting, as a study in patients with renal cell carcinoma showed 47% decrease of ⁸⁹Zr-bevacizumab tumor uptake after 10 mg/kg bevacizumab (6). Furthermore, large phase III trials showed only modest effects of bevacizumab combined with trastuzumab in HER2-positive breast cancer and negative effects when combined with cetuximab or panitumumab in colorectal cancer.

Combining bevacizumab with chemotherapy, in a dosage that is high according to Huang and colleagues, showed also disappointing results, as it did not improve overall survival in breast and ovarian cancer. Nonetheless, chemotherapy with a relatively low dose of 5 mg/kg bevacizumab in metastatic colorectal and a high dose of 15 mg/kg in non-small cell lung cancer (NSCLC) every 2 weeks improved overall survival (7, 8). Jain (3) suggested that the reduced tumor uptake of radiolabeled docetaxel in patients with NSCLC may be the consequence of a too high bevacizumab dose of 15 mg/kg (9). However, also 24 hours after 7.5 mg/kg bevacizumab, radiolabeled 5-fluorouracil uptake decreased 20% in liver metastases of patients with colorectal cancer (10).

This illustrates that many aspects of the interplay between vessel normalization, antiangiogenic therapy dosing, and combination with other anticancer drugs still need to be clarified. Like Huang and colleagues, we are interested in using imaging for this purpose. Small exploratory studies could visualize effects of antiangiogenic drugs on distribution of other labeled drugs, provide serial information on whole body drug distribution, and guide rational trial design for large combinatorial studies. The ultimate goal is optimal drug delivery to individual tumors.

Disclosure of Potential Conflicts of Interest

E.G.E. de Vries has commercial research grant from Roche and is a consultant/advisory board member of Roche Genentech. No potential conflicts of interest were disclosed by the other authors.

Received September 4, 2013; accepted September 24, 2013; published OnlineFirst November 21, 2013.

- immunosuppressive tumor microenvironment and enhance immunotherapy. *Proc Natl Acad Sci U S A* 2012;109:17561–6.
3. Jain RK. Normalizing tumor microenvironment to treat cancer: bench to bedside to biomarkers. *J Clin Oncol* 2013;31:2205–18.
4. Pastuskovas CV, Mundo EE, Williams SP, Nayak TK, Ho J, Ulufatu S, et al. Effects of anti-VEGF on pharmacokinetics, biodistribution, and

- tumor penetration of trastuzumab in a preclinical breast cancer model. *Mol Cancer Ther* 2012;11:752–62.
5. Heskamp S, Boerman OC, Molkenboer-Kueneen DM, Oyen WJG, van der Graaf WTA, van Laarhoven HWM. Bevacizumab reduces tumor targeting of antiepidermal growth factor and anti-insulin-like growth factor 1 receptor antibodies. *Int J Cancer* 2013;133:307–14.
 6. Oosting SF, Brouwers AH, Van Es SC, Nagengast WB, Oude Munnink TH, Lub-de Hooge MN, et al. ^{89}Zr -bevacizumab PET imaging in metastatic renal cell carcinoma patients before and during antiangiogenic treatment. *J Clin Oncol* 30, 2012 (suppl; abstr 10581).
 7. Hurwitz H, Fehrenbacher L, Novotny W, Cartwright T, Hainsworth J, Heim W, et al. Bevacizumab plus irinotecan, fluorouracil, and leucovorin for metastatic colorectal cancer. *N Engl J Med* 2004; 350:2335–42.
 8. Sandler A, Gray R, Perry MC, Brahmer J, Schiller JH, Dowlati A, et al. Paclitaxel-carboplatin alone or with bevacizumab for non-small-cell lung cancer. *N Engl J Med* 2006;355:2542–50.
 9. van der Veldt AA, Lubberink M, Bahce I, Walraven M, de Boer MP, Greuter HN, et al. Rapid decrease in delivery of chemotherapy to tumors after anti-VEGF therapy: implications for scheduling of antiangiogenic drugs. *Cancer Cell* 2012;21:82–91.
 10. Zissen MH, Kunz P, Subbarayan M, Chin FT, Conti PS, Fisher GA, et al. ^{18}F -5-fluorouracil dynamic positron emission tomography/computed tomography shows decreased tracer activity after bevacizumab in colorectal liver metastases. *Nucl Med Commun* 2011;32:343–7.