

## Gene Expression Profiles of B-lineage Adult Acute Lymphocytic Leukemia Reveal Genetic Patterns that Identify Lineage Derivation and Distinct Mechanisms of Transformation

Sabina Chiaretti,<sup>4</sup> Xiaochun Li,<sup>2</sup> Robert Gentleman,<sup>2</sup> Antonella Vitale,<sup>4</sup> Kathy S. Wang,<sup>1</sup> Franco Mandelli,<sup>4</sup> Robin Foà,<sup>4</sup> and Jerome Ritz<sup>3</sup>

**Abstract Purpose:** To characterize gene expression signatures in acute lymphocytic leukemia (ALL) cells associated with known genotypic abnormalities in adult patients.  
**Experimental Design:** Gene expression profiles from 128 adult patients with newly diagnosed ALL were characterized using high-density oligonucleotide microarrays. All patients were enrolled in the Italian GIMEMA multicenter clinical trial 0496 and samples had >90% leukemic cells. Uniform phenotypic, cytogenetic, and molecular data were also available for all cases.  
**Results:** T-lineage ALL was characterized by a homogeneous gene expression pattern, whereas several subgroups of B-lineage ALL were evident. Within B-lineage ALL, distinct signatures were associated with ALL1/AF4 and E2A/PBX1 gene rearrangements. Expression profiles associated with ALL1/AF4 and E2A/PBX1 are similar in adults and children. BCR/ABL<sup>+</sup> gene expression pattern was more heterogeneous and was most similar to ALL without known molecular rearrangements. We also identified a set of 83 genes that were highly expressed in leukemia blasts from patients without known molecular abnormalities who subsequently relapsed following therapy. Supervised analysis of kinase genes revealed a high-level FLT3 expression in a subset of cases without molecular rearrangements. Two other kinases (PRKCB1 and DDR1) were highly expressed in cases without molecular rearrangements, as well as in BCR/ABL-positive ALL.  
**Conclusions:** Genomic signatures are associated with phenotypically and molecularly well defined subgroups of adult ALL. Genomic profiling also identifies genes associated with poor outcome in cases without molecular aberrations and specific genes that may be new therapeutic targets in adult ALL.

After more than four decades of intensive research, the cellular origins of acute lymphocytic leukemia (ALL) have been well defined, and several distinct genetic mechanisms that lead to malignant transformation of these cells have been identified (1–4). In ~25% to 30% of adult patients, ALL cells express

multiple genes associated with T-cell differentiation, and T-cell receptor genes show the presence of clonal rearrangements. In the remaining 70% to 75% of adult patients, ALL cells express markers of B-cell differentiation, and immunoglobulin genes reveal unique patterns of clonal rearrangement. These characteristics clearly define the derivation of these leukemias from distinct lymphoid precursor cells that have been committed to either T-lineage or B-lineage differentiation. Although chromosome translocations and molecular rearrangements are relatively infrequent in T-lineage ALL, these events occur commonly in B-lineage ALL and reflect distinct mechanisms of transformation. The relative frequencies of specific molecular rearrangements differ in children and adults with B-lineage ALL. In adult ALL, the BCR/ABL gene rearrangement occurs in about 25% of cases and the ALL1/AF4 gene rearrangement (also termed MLL/AF4) is present in 4% to 7% of cases (5). The BCR/ABL gene rearrangement occurs much less frequently in pediatric ALL, whereas the ALL1/AF4 or other mixed-lineage leukemia (MLL) rearrangements occur primarily in infants with this disease. The TEL/AML1 gene rearrangement, which is usually associated with a favorable outcome, is present in 25% to 30% of pediatric cases but is rare in adults (1, 6). Because these cytogenetic abnormalities reflect distinct mechanisms of transformation, these differences help to explain why children and adults with ALL have such different outcomes following conventional therapy.

**Authors' Affiliations:** Departments of <sup>1</sup>Medical Oncology and <sup>2</sup>Biostatistical Science, Dana-Farber Cancer Institute; <sup>3</sup>Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts; and <sup>4</sup>Division of Hematology, Department of Cellular Biotechnologies and Hematology, University "La Sapienza," Rome, Italy

Received 10/25/04; revised 6/9/05; accepted 7/15/05.

**Grant support:** NIH grant CA66996, Leukemia and Lymphoma Society Specialized Center for Research in Myeloid Leukemias, Associazione Italiana per la Ricerca sul Cancro, Ministero dell'Istruzione, Università e Ricerca, Fondo per gli Investimenti della Ricerca di Base project, Associazione per le Leucemie Acute dell'adulto "Cristina Bassi" e Fondazione Cassa di Risparmio di Genova e Imperia, and the Ted and Eileen Pasquarello Research Fund.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

**Note:** Supplementary data for this article are available at Clinical Cancer Research Online (<http://clincancerres.aacrjournals.org/>).

**Requests for reprints:** Jerome Ritz, Dana-Farber Cancer Institute, 44 Binney Street, Boston, MA 02115. Phone: 617-632-3465; Fax: 617-632-5167; E-mail: [jerome\\_ritz@dfci.harvard.edu](mailto:jerome_ritz@dfci.harvard.edu).

© 2005 American Association for Cancer Research.  
doi:10.1158/1078-0432.CCR-04-2165

Despite the frequent presence of chromosome translocations and gene rearrangements in ALL, no molecular rearrangements are found in ~50% of adult cases. Although the cellular origins of these leukemias can be readily determined by phenotypic studies, the mechanisms of transformation of these leukemias remain unknown. In this context, large-scale gene expression profiling provides a new approach to explore the mechanisms of transformation of malignant cells. In pediatric ALL, several reports have identified specific gene signatures that reflect known phenotypic characteristics and genetic rearrangements, such as ALL1/AF4, TEL/AML1, E2A/PBX1, and to a lesser extent BCR/ABL (7–11). This approach has also led to the identification of gene expression patterns that may predict response to therapy (8, 9, 12, 13), as well as new potential therapeutic targets (14, 15).

The current study used high-density oligonucleotide microarrays (Affymetrix U95Av2, Santa Clara, CA) to define the gene expression profile of ALL in a well-characterized population of 128 adult patients at the time of diagnosis, before initiation of intensive chemotherapy. Analysis of pretreatment ALL cells also included the uniform characterization of cell surface phenotype and molecular abnormalities. This resulted in the identification of gene expression signatures that were associated with well-defined genetic abnormalities known to play a central role in malignant transformation of these leukemia cells and are similar to those previously reported in children (9). To characterize potential pathways of transformation in ALL cells without known genetic abnormalities, we compared the gene expression profiles of these cases with that of ALL with defined gene rearrangements. This comparison showed that the expression profile of B-lineage ALL without known molecular rearrangements was more similar to cases with BCR/ABL rearrangements than ALL with other genetic rearrangements. Because the BCR/ABL rearrangement results in the formation of an activated tyrosine kinase, these observations suggest that deregulated activation of tyrosine kinase genes may also play an important role in the transformation of at least some ALL cases without other defined molecular rearrangements.

## Materials and Methods

**Patient samples.** Patients were enrolled in the Italian GIMEMA multicenter clinical trial 0496 for adult patients with ALL. Informed written consent was obtained from all patients before therapy. The present study was approved by the institutional review board of the University "La Sapienza" of Rome, Italy. Pretreatment leukemia samples from bone marrow and/or peripheral blood were centrally processed at University "La Sapienza" of Rome; mononuclear cells were isolated by density gradient centrifugation and cryopreserved with 10% DMSO. One hundred twenty-eight leukemia samples containing >90% blasts were selected for gene expression analysis from 431 patients enrolled between 1996 and 2000. Follow-up data were collected at University "La Sapienza".

**Flow cytometry, cytogenetic, and molecular studies.** Conventional immunophenotypic, cytogenetic, and molecular diagnostic studies were done on all samples (16). Flow cytometry analysis was done using monoclonal antibodies specific for the following antigens to determine lineage derivation: Tdt, HLA-DR, CD7, CD19, CD10, CD14, CD33, CD13, CD61, CD34, CD2, myeloperoxidase, and sCD3. For ALL cells assigned to either B-lineage or T-cell lineage, expression of the following panel of antigens was used to define the degree of differentiation of leukemic blasts: CD20, cyCD22, cyIgu, surface immunoglobulin, cyCD3, CD5, CD1a, CD4, and CD8.

**RNA extraction.** Cryopreserved leukemia cells were rapidly thawed and total RNA was extracted using Trizol reagent (Life Technologies Bethesda Research Laboratories, Grand Island, NY) and purified using SV Total RNA Isolation System, (Promega, Madison, WI) according to the manufacturer's protocol with minor modifications. To assess RNA quality, 2  $\mu$ L of RNA from each sample were analyzed by agarose gel electrophoresis; 260/280 ratio was >1.8 for all samples used for microarray analysis (17).

**Oligonucleotide microarray.** The detailed protocol for sample preparation and microarray processing is available from Affymetrix (<http://www.affymetrix.com>). Briefly, first-strand cDNA was synthesized from 5  $\mu$ g total RNA using a T7-(dT)<sub>24</sub> primer (Genset Corp., San Diego, CA) and reverse transcribed with the SuperScript Double-Stranded cDNA Synthesis kit (Life Technologies Bethesda Research Laboratories). After synthesis of the second strand of cDNA, the product was used an *in vitro* transcription reaction (Bioarray kit, Enzo Diagnostics, Farmingdale, NY) to generate biotinylated complementary RNA and subsequently fragmented. Fragmented complementary RNA (15  $\mu$ g/sample) was hybridized to the Affymetrix U95Av2 GeneChip, which contains ~12,600 sequences derived from the Genbank database. As suggested by the manufacturer, samples with 3'/5' glyceraldehyde-3-phosphate dehydrogenase (GAPDH) ratio of >3 were discarded. After hybridization, each microarray was washed, stained, and scanned with an argon-ion confocal laser, with excitation at 488 nm and detection at 570 nm. CEL files are available at <http://www.Bioconductor.org/Docs/Papers/2003/Chiaretti>.

**Data analysis and statistical methods.** Affymetrix gene expression data were processed with dChip (<http://www.dchip.org>), which uses an invariant set normalization method. The array with the median overall intensity was chosen as the baseline for normalization. Model-based expressions were computed for each array and probe set (18) using only *perfect match* probes. All other analyses were carried out using the R language (<http://www.r-project.org>; ref. 19), included in the Bioconductor package.

Nonspecific filtering criteria for unsupervised clustering were defined as follows: (a) gene expression level was required to be higher than 100 in >20% of the samples; (b) the ratio of SD to the mean expression across all samples was required to be between 0.6 and 10. We chose these broad criteria because we wanted to concentrate on genes that were common to many B-lineage ALLs or many T-lineage ALLs and not on small subgroups of either population.

To specifically identify genes differentially expressed in pre-B-ALL molecular groups (i.e., ALL1/AF4, BCR/ABL, and E2A/PBX1) and the remaining samples without evidence of major molecular rearrangements (labeled as NEG in the text and figures), we first computed the mean expression of the 12,625 probe sets within each group and only probes with at least one group mean above 100 were retained. Subsequently, an ANOVA analysis aimed at testing if there were any differences among the four group means was done for each probe on log-transformed expression values.

*P* adjustments for multiple comparisons were done using the step-up false discovery rate controlling method proposed by Benjamini and Hochberg (20), with false discovery rate controlled under 0.05. Hierarchical clustering and heat maps were used as described by Ihaka and Gentleman (19) and Eisen et al. (21). The distance between two genes was computed as 1 – the correlation between the standardized expression values across samples. False-color maps were used to display distance matrices between samples. The distance between two samples was computed as 1 – the correlation between the aforementioned standardized expression values of the samples. The magnitude of distance is represented in color with a continuous spectrum from blue (close) to red (far). The diagonal of the distance map represents the distance of *n* samples to themselves (i.e., 0) and is colored in the deepest shade of blue.

For prediction of biological groups, a *leave-one-out* cross-validation was used in the process of probe selection and nearest-neighbor prediction. Specifically, we left one sample out as the test set and

retained the other  $n - 1$  samples as our training set. For each training set, we developed three different gene lists; one for each comparison of the NEG group to a group with a known molecular aberration. Genes were filtered using the same criteria as for identification of probes specifically associated with molecular groups, leaving one sample out, in turn; then, the class of the left-out sample was predicted, using the set of genes obtained for its true class, using one-nearest neighbor classification.

A *t* test was used to identify genes differentially expressed between patients who are in continuous complete remission (CCR) versus those who experienced a relapse. Genes were required to have an average expression above 100 in at least one group, a fold change difference of  $>1.5$ , and an adjusted  $P = 0.05$ . To strengthen these results, a leave-one-out cross-validation was also done and only genes that were selected in 100% analyses were retained.

We also concentrated on genes annotated at the protein-tyrosine kinase node of Gene Ontology (Gene Ontology: 0004713, 299 probe sets) and used a similar procedure. Again, gene lists reflecting pairwise comparisons of the NEG group to the other three were obtained. The selection criteria included (a) an average expression value of  $>100$  in at least one group, (b) a nominal  $P < 0.05$ , and (c) a fold change difference of  $\geq 1.5$ . To identify additional tyrosine kinase genes highly expressed in the NEG group, we identified genes that were highly expressed in the following: (a) all four molecular subsets, (b) the NEG group and any two other subsets, and (c) the NEG group and any other single subset. For this analysis only, gene selection criteria required a mean expression of  $>500$  (sufficiently high values), and equal expression between groups was arbitrarily defined as a mean fold difference between 0.9 and 1.1.

**Real-time quantitative PCR.** Real-time quantitative reverse transcription-PCR analysis was done using an ABI PRISM 7700 Sequence Detection System and the SYBR green I dye (PE Biosystems, Foster City, CA) method as previously described (22). Real-time PCR conditions were as follows: one cycle at 50°C for 2 minutes, one cycle at 95°C for 10 minutes, one cycle at 95°C for 15 seconds, one cycle at 60°C for 1 minute, for a total of 40 cycles. For each sample,  $C_T$  values for GAPDH were determined for normalization purposes. For each run of samples, a correction factor was calculated by dividing by the minimum GAPDH values and applied to normalize the  $C_T$  values of the genes of interest. Primers were designed using the Primer Express 1.0 software. The following primers were used: 5' GAPDH, 5'-CCACCCATGGCAAATTC-3'; 3' GAPDH, 5'-GATGGGATTTCCATTGATGACA-3'; 5' FLT3, 5'-AAATTCACCAGCATGCCTGGTT-3'; 3' FLT3, 5'-TCCGAGTCCGGGTGTATCTG-3'; 5' DDR1, 5'-GATTTCC-CCCTAATGTGCGT-3'; 3' DDR1, 5'-TGCCATCTGCCGTAAGATC-3'; 5' PRKCB1, 5'-CAGAAGGAAGTGAGGCCAATG-3'; and 3' PRKCB1, 5'-TCTTCCGGACCTTGTTTC-3'.

## Results

**Patient characteristics.** Conventional immunophenotypic, cytogenetic, and molecular diagnostic studies were done on pretreatment leukemia samples from the 128 patients (Table 1; Supplementary Table 1). The median age of patients was 29 years (range, 15-58); 43 were females and 85 were males. Immunophenotypic analysis indicated T-cell derivation in 33 samples and B-cell lineage in 95 samples. Well-defined cytogenetic abnormalities were identified in 43 cases. Other simple or complex karyotypic abnormalities were found in 22 samples and a normal karyotype was identified in 24 samples. Thirty-nine cases did not have sufficient metaphases for cytogenetic analysis. Molecular diagnostic studies confirmed the presence of BCR/ABL gene rearrangements in all 26 patients with t(9;22) and in 11 additional samples which were not evaluable by cytogenetic analysis. The ALL1/AF4 gene

**Table 1.** Cytogenetic and molecular characteristics of pretreatment ALL samples

	Pre-B-ALL (n = 95)	T-ALL (n = 33)
Conventional cytogenetic analysis		
t(9;22)	26	0
t(4;11)	7	1
t(10;14)	0	1
t(1;19)	2	0
del(6q)	2	2
del(7q)	1	1
Simple karyotypic rearrangements	9	4
Complex karyotypic abnormalities	6	3
Normal karyotype	16	8
Not evaluable	26	13
Molecular rearrangements		
BCR/ABL	37*	0
ALL1/AF4	10	0
E2A/PBX1	5*	0
NUP-98	0	1
NEG	42*	32

Abbreviations: B-ALL, B-lineage ALL; T-ALL, T-lineage ALL.


\*p15 or p15/p16 deletions were detected in six BCR/ABL samples, one E2A/PBX1 sample, and five samples without molecular rearrangements.

rearrangement was detected in all seven pro-B ALL cases with t(4;11), in one patient who was not evaluable by conventional cytogenetic testing, and in two samples with a normal karyotype. The E2A/PBX1 rearrangement was found in five samples, including two with t(1;19), one with a normal karyotype, and two with insufficient metaphases for cytogenetic analysis. BCR/ABL, ALL1/AF4, and E2A/PBX1 gene rearrangements were only detected in leukemias with a B-lineage cell phenotype. A distinct t(4;11) rearrangement involving the *NUP98* gene was found in a single T-lineage ALL (23, 24). Thirty-six cases that did not harbor any major molecular abnormality were evaluable for long-term outcome: 12 patients were in CCR with a median follow-up of 60.5 months (range, 29-76), whereas the remaining 24 experienced a relapse within a median period of 11 months (range, 2-69).

**Gene expression profiles distinguish T-cell and B-cell lineage derivation.** Applying the filtering criteria to all 128 samples resulted in the selection of 792 genes, which showed sufficient levels of expression and variation across groups of interest. Unsupervised hierarchical clustering based on the expression of this set of genes identified two groups that coincided precisely with the phenotypic classification of T-cell and B-cell lineage determined by immunophenotypic analysis of cell surface markers (Fig. 1). As expected, T-lineage ALL samples were characterized in part by the homogeneous expression of known T-cell lineage antigens such as CD3 $\delta$ , CD3 $\zeta$ , and TCR-associated genes, and B-lineage ALL were characterized by the homogeneous expression of MHC class II, CD19, and immunoglobulin-associated genes. However, many genes not known to be associated with either T-cell or B-cell lineage were also differentially expressed in these two sets of samples. As shown in Fig. 1, the T-lineage ALL group seemed relatively

homogeneous. In contrast, the B-lineage ALL group seemed heterogeneous, and it was possible to identify smaller subsets of samples based on a variable profile of expression within this large set of differentially expressed genes.

*Gene expression profile of adult B-lineage acute lymphocytic leukemia identifies signatures of ALL1/AF4, E2A/PBX1, and BCR/ABL gene rearrangements.* To further characterize the B-lineage ALL samples, we first performed unsupervised hierarchical clustering of the 95 samples. Different filtering criteria were applied (see Data analysis and statistical methods), and regardless of the criteria used in gene filtering, cases with



**Fig. 1.** Unsupervised hierarchical clustering of 128 adult ALL samples based on differential gene expression. Filtering criteria were applied to normalized gene expression values, resulting in the selection of 475 genes for this analysis. Row, expression of a single gene; column, single ALL sample. Relative levels of gene expression are depicted with a color scale: dark blue, lowest levels of expression; red, highest levels of expression.

*E2A/PBX1* and *ALL1/AF4* gene rearrangements formed two distinct clusters that corresponded precisely to the molecular classification of these samples (data not shown). In contrast, samples with BCR/ABL rearrangements showed a more heterogeneous profile, being distributed in several clusters. Moreover, each BCR/ABL cluster also included NEG samples. Similarly, NEG cases could not be grouped into distinct clusters. These results suggested that both BCR/ABL-positive cases and NEG samples are likely to include several subgroups, and also that the gene expression profiles of these two groups were more similar to each other than to other molecularly defined groups.

Subsequently, to identify genes associated with known molecular abnormalities in the B-lineage ALL, samples were assigned to one of four groups: ALL1/AF4, E2A/PBX1, BCR/ABL, and NEG. ANOVA was applied to identify genes with significantly different mean expression values in at least one of the four B-lineage ALL groups. This resulted in the selection of 167 probes with adjusted  $P$ s  $< 0.0001$ . The top genes in each category with an adjusted  $P < 1 \times 10^{-5}$  are listed in Table 2. The complete list of probes is summarized in the Supplementary Material. As shown in Fig. 2, clustering using this set of genes led to the correct grouping of all ALL1/AF4 and E2A/PBX1 samples, whereas BCR/ABL-positive cases were again clustered together with NEG samples.

ALL1/AF4 was characterized by high levels of expression of a large set of genes ( $\sim 50$ ). As previously reported, this gene rearrangement was strongly associated with the up-regulation of several HOX family genes (9, 14, 25), as well as the *FLT3* receptor (15). Five of these 10 samples were further examined for the presence of either internal tandem duplication or D835 mutations of *FLT3* receptor and none showed any abnormalities.

The *E2A/PBX1* gene rearrangement also showed a distinct and homogeneous pattern characterized by the high level of expression of several genes ( $\sim 35$ ). The genes most strongly associated with this group were represented by *PBX1*, *FAT*, *NID2*, and *KANK* (26, 27). Interestingly, two samples without known molecular rearrangements clustered tightly with the E2A/PBX1 group. In one case, conventional cytogenetic analysis revealed a normal karyotype, but the second case did not have sufficient metaphases for cytogenetic analysis. Molecular diagnostic studies did not reveal the E2A/PBX1 rearrangement in these two samples. However, comparative genomic hybridization showed gains in the 1q region in both cases (1q23 in one case).

The BCR/ABL group was characterized by high expression of  $\sim 70$  genes. This included several tyrosine kinase genes (*ABL*, *FYN*, and *YES1*) and genes associated with cell cycle progression (*CCDN2* and *E2F2*). However, compared with the ALL1/AF4 and E2A/PBX1 groups, the gene expression profile in BCR/ABL-positive ALL was more heterogeneous. This heterogeneity was not associated with the expression of different BCR/ABL-derived fusion proteins (p190 and p210; data not shown).

*Gene expression profiles reveal similarities between BCR/ABL-positive acute lymphocytic leukemia and samples without molecular rearrangements.* Although our analysis did not identify a distinct or homogeneous gene expression pattern in NEG ALL samples, these cases seemed to have gene expression signatures that were more similar to cases with the BCR/ABL

**Table 2.** Differentially expressed genes in pre – B-ALL samples with ALL1/AF4, E2A/PBX1 and BCR/ABL molecular rearrangements

Gene	Locus link ID	Adjusted <i>P</i>	Function
ALL1/AF4			
<i>HOXA9</i>	3205	$<10^{-13}$	Transcription factor
<i>HOXA10</i>	3206	$<10^{-13}$	Transcription factor
<i>VLDLR</i>	7436	$<10^{-13}$	Receptor
<i>MEIS1</i>	4211	$<10^{-13}$	RNA polymerase II transcription factor
<i>DPYSL3</i>	1809	$<10^{-13}$	Nucleic acid metabolism
<i>PALLADIN</i>	23022	$1 \times 10^{-13}$	Cytoskeleton
<i>KIAA1157</i>	57460	$1.5 \times 10^{-12}$	Unknown
<i>PROM1</i>	8842	$2.4 \times 10^{-12}$	Membrane protein
<i>CCNA1</i>	8900	$1.3 \times 10^{-11}$	Cell cycle control
<i>LGALS1</i>	3956	$2.3 \times 10^{-10}$	Control of cell proliferation
<i>PRSS12</i>	8492	$3.2 \times 10^{-10}$	Serine protease
<i>NPTX2</i>	4885	$4.4 \times 10^{-10}$	Pentaxin
<i>FUT4</i>	2526	$1 \times 10^{-9}$	Fucosyltransferase
<i>TBC1D8</i>	11138	$4.8 \times 10^{-9}$	Positive control of cell proliferation
<i>DIAPH2</i>	1730	$8 \times 10^{-9}$	Actin binding
<i>DAD1</i>	1603	$1.2 \times 10^{-7}$	Antiapoptosis
<i>HOXA5</i>	3202	$1.5 \times 10^{-7}$	Transcription factor
<i>RHOBTB3</i>	22836	$3 \times 10^{-7}$	Unknown
<i>KIAA0125</i>	9834	$4.1 \times 10^{-7}$	Unknown
E2A/PBX1			
<i>PBX1</i>	5087	$<10^{-13}$	Transcription factor
<i>FAT</i>	2195	$<10^{-13}$	Cell adhesion
<i>KANK</i>	23189	$2.5 \times 10^{-12}$	Cell cycle control
<i>FGF9</i>	2254	$1.7 \times 10^{-11}$	Cell cycle control
<i>CRYM</i>	1428	$2.2 \times 10^{-11}$	Vision
<i>NID2</i>	22795	$2.3 \times 10^{-11}$	Cell adhesion
<i>KIAA0802</i>	23255	$8.7 \times 10^{-11}$	Unknown
<i>LAMA5</i>	3911	$1.8 \times 10^{-9}$	Differentiation
<i>AOX1</i>	316	$1.9 \times 10^{-9}$	Oxidoreductase activity
<i>SLAM</i>	6504	$3.1 \times 10^{-9}$	B-cell proliferation
<i>PRKCZ</i>	5590	$5.8 \times 10^{-8}$	Protein kinase
<i>MERTK</i>	10461	$1.3 \times 10^{-7}$	Oncogene/tyrosine kinase
<i>ALDH1A1</i>	216	$1.5 \times 10^{-7}$	Aldehyde dehydrogenase activity
<i>PKM2</i>	5315	$4.3 \times 10^{-7}$	Pyruvate kinase
Unknown	Unknown	$1.4 \times 10^{-6}$	Unknown


**Table 2.** Differentially expressed genes in pre – B-ALL samples with ALL1/AF4, E2A/PBX1 and BCR/ABL molecular rearrangements (Cont'd)

Gene	Locus link ID	Adjusted <i>P</i>	Function
<i>ACK1</i>	10188	$1.4 \times 10^{-6}$	Non – receptor tyrosine kinase
<i>NCBP2</i>	22916	$1.6 \times 10^{-6}$	RNA binding
<i>BLK</i>	640	$2.1 \times 10^{-6}$	Protein tyrosine kinase
<i>TMSNB</i>	11013	$2.4 \times 10^{-6}$	Actin binding
<i>TRIB2</i>	28951	$1.5 \times 10^{-5}$	Protein kinase
<i>CGI-49</i>	51097	$1.6 \times 10^{-5}$	Unknown
<i>LRMP</i>	4033	$1.7 \times 10^{-5}$	Lymphocyte development
<i>KIAA0889</i>	25781	$2.1 \times 10^{-5}$	Unknown
<i>E2F5</i>	1875	$2.1 \times 10^{-5}$	Transcription factor
BCR/ABL			
<i>CDW52</i>	1043	$9.4 \times 10^{-13}$	Integral to plasma membrane
<i>ABL1</i>	25	$1.3 \times 10^{-11}$	Tyrosine kinase activity
<i>YES1</i>	7525	$6.1 \times 10^{-11}$	Tyrosine kinase activity
<i>SOCS2</i>	8835	$5.7 \times 10^{-10}$	Regulation of cell growth
<i>CCND2</i>	894	$3 \times 10^{-9}$	Cell cycle progression
<i>HLA-DPA1</i>	3113	$4.8 \times 10^{-9}$	HLA class II antigen
<i>HLA-DOA</i>	3111	$3.6 \times 10^{-8}$	HLA class II antigen
<i>CD2AP</i>	23607	$4 \times 10^{-8}$	Cytoskeleton
<i>TCF8</i>	6935	$5 \times 10^{-8}$	Growth inhibition
<i>HLA-C</i>	3107	$5.9 \times 10^{-8}$	HLA class II antigen
<i>FYN</i>	2534	$1 \times 10^{-7}$	Protein kinase activity
<i>HLA-DPB1</i>	3115	$1.3 \times 10^{-7}$	HLA class II antigen
<i>GAB1</i>	2549	$1.6 \times 10^{-7}$	Cell proliferation
<i>IFITM1</i>	8519	$3.6 \times 10^{-7}$	IFN response
<i>OPTN</i>	10133	$3.6 \times 10^{-7}$	TNF inhibition
<i>ITGA6</i>	3655	$5.9 \times 10^{-7}$	Integrin complex
<i>HLA-A</i>	3105	$7.1 \times 10^{-7}$	HLA class I antigen
<i>PSAP</i>	5660	$9.2 \times 10^{-7}$	Saposin
<i>E2F2</i>	1870	$1.1 \times 10^{-6}$	Cell cycle progression
<i>CD24</i>	934	$1.8 \times 10^{-6}$	Plasma membrane

NOTE: Genes are rank ordered according to their adjusted *P*.  
Abbreviations: B-ALL, B-lineage ALL; TNF, tumor necrosis factor.

rearrangement than with other molecular rearrangements. To further characterize the differences and similarities between different B-lineage ALL subsets, we used two approaches. First, we generated a distance map based on the expression of the 167 probes selected by ANOVA (Fig. 3). A distance map compares each sample with every other; the diagonal represents the comparison of each sample with itself and establishes a

correlation of 1 (represented in blue) and hence a distance of 0. In this representation, samples with a high degree of correlation will appear blue. As shown in Fig. 3, samples with ALL1/AF4 cluster together and are quite distinct from all other groups (the large blue rectangles centered on the diagonal). Samples with E2A/PBX1 and known 1q gain also cluster together and are quite distinct. In contrast, the BCR/ABL



**Fig. 2.** Gene expression profiles of 95 pre-B-lineage ALL samples. An ANOVA filter was used to select 167 variably expressed genes. Supervised clustering revealed distinct gene expression profiles for samples with ALL1/AF4 and E2A/PBX1, and less distinct profiles for samples with BCR/ABL or no molecular abnormalities.

samples do not homogeneously cluster together and are admixed with samples that do not have known molecular rearrangements.

The knn CV prediction analysis was used to evaluate the error rate comparing each group to NEG. As summarized in Table 3, the prediction error rate was 0% for ALL1/AF4 and 4% for E2A/PBX1. One NEG sample and one E2A/PBX1 sample were misclassified. The misclassified negative sample has a known gain at 1q23. In contrast, seven NEG and five BCR/ABL cases were misclassified for BCR/ABL, and the overall classification error rate was 15%. This relatively high error rate again suggests that these two groups are more difficult to distinguish than the others at the gene expression level. Notably, the NEG cases that were misclassified did not have distinct biological characteristics. Of the seven NEG patients that were misclassified, three experienced a relapse, one died during induction chemotherapy, one following stem cell transplantation procedures, and only two are in CCR.

**Gene expression profiles identify a set of genes highly expressed in relapsed patients.** To identify genes associated with long-term outcome, we compared leukemia gene expression profiles in patients who remain in CCR with profiles from cases who subsequently relapsed after conventional therapy. Given the adverse effect of known molecular abnormalities, this analysis was done exclusively on samples that did not carry any major abnormalities. Eighty-seven probe sets, corresponding to 83 genes, were differentially expressed between patients who are still in CCR ( $n = 12$ ) versus those who experienced a relapse ( $n = 24$ ; Fig. 4). Virtually, all the identified genes were highly expressed in samples from patients who subsequently relapsed.

Genes expressed at high levels in the relapse group could be functionally subdivided in the following categories: (a) cell motility, cell interaction, and cytoskeleton organization (*SDFR1*, *DAAM1*, *IQGAP1*, *ADD3*, *TLN1*, *ITGB1*, *LGALS8*, *CAPZB*, *DNCI2*, *LSP1*, and *nexilin*); (b) membrane antigens (*CD79a*, *CD79b*, *CD24*, *CD45*, *CD62L*, and *CD85D*); (c) members of the phosphatidylinositol pathway (*PIP5K2B*, *PIK3C2B*, *PHKB*, *ITPR3*, and *PRKCB1*); (d) resistance to chemotherapy (*ASNS*, *CROP*, and *MVP*); and (e) DNA modeling and response to damage (*HIST1H2BD*, *HIST1H2AC*, *ATRX*, and *ATR*; see Supplementary Table 3 for a complete list).

**Comparative expression of kinase genes in pre-B-lineage acute lymphocytic leukemia.** Because the BCR/ABL gene rearrangement results in leukemic transformation through the formation of a constitutively activated tyrosine kinase, we specifically compared the expression of tyrosine kinase genes in the four groups of B-lineage ALL samples. ANOVA was used to compare NEG samples with every other group (ALL1/AF4, E2A/PBX1, and BCR/ABL) using exclusively genes known to have a kinase activity (as reported by Gene Ontology: 0004713, 299 probe sets, 190 transcripts). As shown in Fig. 5A, comparison of NEG and E2A/PBX1 identified nine kinases (five tyrosine kinase: *LCK*, *ACK1*, *ZAP70*, *BLK*, and *MERTK*; four serine/threonine kinases: *STK39*, *NEK4*, *KIAA0175*, and a dual protein dual specificity protein kinase, *MAP2K3*) that were more highly expressed in the E2A/PBX1 samples, but only two tyrosine kinases (*FLT3* and *RYK*) were more highly expressed in the samples without rearrangements; particularly, high levels were observed for *FLT3*. When compared with ALL1/AF4 samples (Fig. 5B), four kinases (two tyrosine kinases: *BLK* and *FLT3* and

two serine threonine kinases: *MAP3K5/ASK1* and *ULK1*) were more highly expressed in samples with this rearrangement, and eight genes (six tyrosine kinases: *FGFR1*, *FLT1*, *DDR1*, *LCK*, *FYN*, and *RYK* and two serine/threonine kinases: *PRKCB1* and *PRKY*) were more highly expressed in the NEG group. When compared with the samples with BCR/ABL rearrangements (Fig. 5C), only three tyrosine kinases were more highly expressed in the BCR/ABL group (*ABL*, *YES1*, and *FYN*), and *FLT3* was the only tyrosine kinase that was more highly expressed in the samples without BCR/ABL.

Consistent with previous reports, the *FLT3* gene was expressed at high levels in samples with ALL1/AF4 rearrangements. However, *FLT3* was also highly expressed in NEG samples when compared with BCR/ABL and E2A/PBX1 samples. In fact, the high-level expression of *FLT3* (>1,000 units) was found in 20 NEG samples (48% of cases). To confirm the results obtained by oligonucleotide microarrays, real-time quantitative PCR was done on 33 samples without molecular rearrangements and available material. *FLT3* mRNA levels measured by microarrays were highly correlated with those measured by quantitative PCR (Pearson coefficient,  $-0.8$ ).

To specifically identify kinase genes that were expressed at high levels in NEG samples, we also identified those genes within this

class (using Gene Ontology: 0004713) that were expressed at relatively high levels in any of the four subsets of B-lineage ALL (Table 4). No kinases represented on the Affymetrix U95Av2 arrays were exclusively expressed at high levels in the samples without molecular rearrangements. Four kinase genes were expressed at high levels in all four subsets; eight genes were expressed at equally high levels in three groups including NEG samples. Two genes, *PRKACA* and *AXL*, were expressed at comparably high levels in the NEG and ALL1/AF4 groups, and two other kinase genes, *DDR1* and *PRKCB1*, were highly expressed in both the NEG and BCR/ABL groups (Pearson correlation coefficient between oligonucleotide arrays and quantitative PCR:  $-0.68$  and  $-0.8$ , respectively).

## Discussion

The primary objective of our study was to establish gene expression signatures associated with known phenotypic and genetic abnormalities of adult ALL patients. Previous studies have examined gene expression in pediatric ALL (8–11, 14, 28, 29), but the genetic abnormalities that lead to malignant transformation of immature lymphoid progenitors, as well as their frequency, differ in pediatric and adult patients (4). Importantly, long-term outcome following treatment is



**Fig. 3.** Distance map based on the 167 genes selected by ANOVA filtering. This analysis revealed a strong correlation between the ALL1/AF4 and E2A/PBX1 samples and a correlation between BCR/ABL and samples without known molecular abnormalities.

**Table 3.** Prediction error rate comparing NEG with samples known to contain ALL1/AF4, E2A/PBX1, and BCR/ABL molecular rearrangements

Molecular groups	Model prediction		Error rate (%)
	ALL1/AF4	Negative	
ALL1/AF4	10	0	0
NEG*	0	42	
E2A/PBX1	Negative		4
	4	1	
NEG	1 <sup>†</sup>	41	
BCR/ABL	Negative		15
	32	5	
NEG	7	35	


\*Samples without molecular abnormalities are defined as negative.

<sup>†</sup>Comparative genomic hybridization analysis revealed a gain of 1q23.

markedly worse in adult patients. We therefore focused our attention on ALL cells obtained before therapy in a well-defined cohort of 128 adult patients enrolled in a single treatment protocol (GIMEMA 0496; refs. 30, 31). By correlating the expression of ~12,600 probes with conven-

tional phenotypic and molecular characteristics, we identified gene expression signatures associated with known phenotypic characteristics of adult ALL and also defined gene expression patterns associated with distinct mechanisms of leukemic transformation. Unsupervised hierarchical clustering of all ALL samples showed two distinct patterns of gene expression that correlated precisely with either T or B lineage. Although the gene expression profile of T-lineage ALL was relatively homogeneous, more detailed analysis of this data set previously identified distinct signatures associated with initial response to intensive induction chemotherapy, as well as with long-term maintenance of remission (12).


In contrast to T-lineage ALL, the profile of B-lineage ALL was very heterogeneous. The cellular origin seems relatively uniform in B-lineage ALL, but the molecular heterogeneity of this leukemia is well established. Molecular rearrangements of BCR/ABL, ALL1/AF4, and E2A/PBX1 were present in 55% of our population. The ALL1/AF4 rearrangement was uniquely associated with a high level of expression of several related transcription factors, which seem to be direct targets of MLL (14, 25, 32). Similar results have been reported by other investigators (9, 14, 25). In agreement with other reports (14, 15, 33, 34), our study also found that ALL1/AF4<sup>+</sup> cases were characterized by overexpression of FLT3, a tyrosine kinase whose mutations represent the most common genetic rearrangements in acute myelogenous leukemia (35).



**Fig. 4.** Identification of 83 differentially expressed genes in NEG ALL associated with long-term outcome.



**Fig. 5.** A-C, supervised analysis of tyrosine kinase genes (Gene Ontology: 0004713). Two-group comparisons between samples without molecular abnormalities and E2A/PBX1 (A), ALL1/AF4 (B), and BCR/ABL (C), using genes annotated by Gene Ontology as tyrosine kinases. Several genes are represented by multiple probe sets. Top bar: pink, NEG samples; light blue, E2A/PBX1<sup>+</sup>; green, ALL1/AF4<sup>+</sup>; red, BCR/ABL<sup>+</sup>.



Inhibitors of this tyrosine kinase have proven to be effective *in vitro* in ALL cells from pediatric as well as adult cases (15, 36, 37).

ALL cells from adult patients with the E2A/PBX1 rearrangement were also associated with a homogeneous gene expression pattern. In this group, it was also possible to identify a set of kinases, such as *PRKCZ*, *BLK*, and *MERTK*, that are consistently associated with this rearrangement in pediatric cases and may represent potential therapeutic targets (9, 10, 15, 38). Remarkably, two samples with a gain in the 1q and 1q23 region, respectively, showed a very similar profile to five samples with the E2A/PBX1 rearrangement. These results suggest that the gene expression signature associated with the E2A/PBX1 rearrangement may be primarily a reflection of increased gene dosage for PBX1. Alternatively, these cases may contain other mutations that result in the activation of PBX1 and a complex pattern of gene expression that is essentially identical to that observed in cases harboring the E2A/PBX1 rearrangement. Further characterization of the *PBX1* gene in ALL samples with gains in 1q23 will be necessary to determine the mechanism responsible for deregulation of this transcription factor in these cases. Because these cases were not detected by conventional cytogenetic analysis or molecular studies, deregulation of PBX1 may occur more commonly than previously recognized in adult B-lineage ALL.

Remarkably, for both E2A/PBX1 and ALL1/AF4 rearrangements, the gene expression signatures seem very similar in both pediatric and adult ALL. This was tested directly by combining our data set with the pediatric data set described by Yeoh et al. (9). Hierarchical clustering of 300 B-lineage samples based on the genes selected by ANOVA resulted in the tight grouping of both adult and pediatric samples with MLL rearrangements. Similarly, all samples with E2A/PBX1 clustered together, regardless of patient age (Supplementary Fig. 1). These results suggest that both genetic rearrangements lead to very similar downstream effects in both adult and pediatric cases, and are, at least for the MLL rearrangements, consistent with the poor outcome of both cohorts.

ALL cells with the BCR/ABL rearrangement were also found to have a distinct but more heterogeneous gene expression profile (10, 11). In addition to ABL, other protein kinases such as *YES1* and *FYN* are also highly expressed in these cells compared with ALL1/AF4-positive and E2A/PBX1-positive cases. The identification of additional kinases that are overexpressed in these samples is of great importance, considering that the introduction of imatinib has drastically changed the outcome of BCR/ABL<sup>+</sup> ALL (39) and that newer BCR/ABL kinase inhibitors, such as AMN107 and BMS-354825 (40, 41), which show higher efficacy *in vitro*, are presently being evaluated in phase I/II clinical trials.

**Table 4.** Kinase genes with high-level gene expression in samples without known molecular rearrangements and other subsets of pre-B-ALL

Gene symbol	Locus link ID	Chromosome location
High expression in NEG, BCR/ABL, E2A/PBX1, and ALL1/AF4		
<i>EFNA3</i>	1944	1q21-q22
<i>ARAF-1</i>	369	Xp11.4-p11.2
<i>RAF-1</i>	5894	3p25
<i>MAP2K1</i>	5604	15q22.1-q22.33
High expression in NEG, BCR/ABL, and ALL1/AF4		
<i>CLK3</i>	1198	115q24
<i>CSK</i>	1445	15q23-q25
<i>DYRK1A</i>	1859	21q22.13
<i>ROCK1</i>	6093	18q11.2
<i>OSR1</i>	9943	3p22-p21.3
<i>MAP3K11</i>	4296	11q13.1-13.3
High expression in NEG, BCR/ABL, and E2A/PBX1		
<i>PSK</i>	51677	16p12.1
High expression in NEG, ALL1/AF4, and E2A/PBX1		
<i>NTRK3</i>	4916	15q25
High expression in NEG and ALL1/AF4		
<i>PRKACA</i>	5566	19p13.1
<i>AXL</i>	558	19q13.1
High expression in NEG and BCR/ABL		
<i>DDR1</i>	780	6p21.3
<i>PRKCB1</i>	5579	16p11.2

Abbreviation: NEG, without known molecular rearrangements; B-ALL, B-lineage ALL.

When analyzed as a group, ALL cases without known molecular rearrangements did not seem to have a unique gene expression signature. Although these leukemic cells were clearly distinct from blasts with rearrangements involving specific transcription factors (ALL1/AF4 and E2A/PBX1), their pattern of gene expression was more reminiscent of BCR/ABL<sup>+</sup> cases. This was first noted in unsupervised hierarchical clustering and confirmed by direct comparison of gene expression patterns using distance maps and prediction analysis. The activation of transcription factors in both acute lymphoid (10, 11, 28, 42)

and myeloid (43–45) leukemia cells consistently results in unique gene expression signatures. The observation that NEG ALL cases, when considered as a single group, do not have a distinctive signature suggests that the transforming events do not involve a single leukemogenic transcription factor. Although the similarity of expression patterns in NEG and BCR/ABL<sup>+</sup> ALL suggests the activation of tyrosine kinases in at least some of these cases. Notably, two tyrosine kinases, *FLT3* and *DDR1* (38), were highly expressed in adult NEG cases. Increased expression of *FLT3* has also been found in a small percentage of B-lineage ALL in children without other genetic rearrangements (46, 47). Nevertheless, this seems to be a heterogeneous group and other potential mechanisms are also likely to be involved in these cases.

Finally, comparison of ALL cases without known molecular rearrangements that remained in CCR with cases that subsequently relapsed resulted in the identification of 83 genes that were differentially expressed. This analysis was done exclusively on samples without aberrations because of the well-established adverse effect of abnormalities such as ALL1/AF4, E2A/PBX1, and BCR/ABL on outcome (1–5). Notably, virtually all of these genes had increased expression in the relapse group. Similar results have previously been reported by Teuffel et al. (48), who compared gene expression in pediatric ALL stratified by different risk groups. In this study, 75% of the identified genes had high levels of expression in the high-risk group (48). Among the genes that were associated with relapse, several have previously been associated with resistance to chemotherapy. For example, *ASNS* (asparagine synthetase) is directly involved in the metabolism of L-asparaginase, a compound widely used in ALL regimens. The finding that this gene is expressed at higher levels in adult patients with a poor prognosis suggests that this gene may contribute to resistance in these patients (49). Similarly, high-level expression of a set of genes involved in the phosphatidylinositol pathway (*PIP5K2B*, *PIK3C2B*, *PHKB*, *ITPR3*, and *PRKCB1*) indicates that this pathway can also contribute to resistance to chemotherapy.

In conclusion, these results confirm the strong influence of cell lineage derivation and of specific rearrangements on gene expression in adult ALL. Furthermore, this analysis identifies a set of kinases within defined subsets of ALL that represent potential therapeutic targets for adult patients with these leukemias. Finally, our study identifies a set of genes differentially expressed in patients without molecular aberrations with a diverse outcome. Validation studies on these transcripts are ongoing.

## References

- Kersey JH. Fifty years of studies of the biology and therapy of childhood leukemia. *Blood* 1997;90:4243–51.
- Thandla S, Aplan PD. Molecular biology of acute lymphocytic leukemia. *Semin Oncol* 1997;24:45–56.
- Faderl S, Kantarjian HM, Talpaz M, Estrov Z. Clinical significance of cytogenetic abnormalities in adult acute lymphoblastic leukemia. *Blood* 1998;91:3995–4019.
- Pui CH, Relling MV, Downing JR. Acute lymphoblastic leukemia. *N Engl J Med* 2004;350:1535–48.
- Hoelzer D, Gokbuget N. New approaches to acute lymphoblastic leukemia in adults: where do we go? *Semin Oncol* 2000;27:540–59.
- McLean TW, Ringold S, Neuberg D, et al. TEL/AML-1 dimerizes and is associated with a favorable outcome in childhood acute lymphoblastic leukemia. *Blood* 1996;88:4252–8.
- Golub TR, Slonim DK, Tamayo P, et al. Molecular classification of cancer: class discovery and class prediction by gene expression monitoring. *Science* 1999;286:531–7.
- Ferrando AA, Neuberg DS, Staunton J, et al. Gene expression signatures define novel oncogenic pathways in T cell acute lymphoblastic leukemia. *Cancer Cell* 2002;1:75–87.
- Yeoh EJ, Ross ME, Shurtleff SA, et al. Classification, subtype discovery, and prediction of outcome in pediatric acute lymphoblastic leukemia by gene expression profiling. *Cancer Cell* 2002;1:133–43.
- Ross ME, Zhou X, Song G, et al. Classification of pediatric acute lymphoblastic leukemia by gene expression profiling. *Blood* 2003;102:2951–9.
- Fine BM, Stanulla M, Schrappe M, et al. Gene expression patterns associated with recurrent chromosomal translocations in acute lymphoblastic leukemia. *Blood* 2004;103:1043–9.
- Chiaretti S, Li X, Gentleman R, et al. Gene expression profile of adult T-cell acute lymphocytic leukemia identifies distinct subsets of patients with different response to therapy and survival. *Blood* 2004;103:2771–8.
- Shipp MA, Ross KN, Tamayo P, et al. Diffuse large B-cell lymphoma outcome prediction by

- gene-expression profiling and supervised machine learning. *Nat Med* 2002;8:68–74.
14. Armstrong SA, Staunton JE, Silverman LB, et al. MLL translocations specify a distinct gene expression profile that distinguishes a unique leukemia. *Nat Genet* 2002;30:41–7.
  15. Armstrong SA, Kung AL, Mabon ME, et al. Inhibition of FLT3 in MLL. Validation of a therapeutic target identified by gene expression based classification. *Cancer Cell* 2003;3:173–83.
  16. van Dongen JJ, Macintyre EA, Gabert JA, et al. Standardized RT-PCR analysis of fusion gene transcripts from chromosome aberrations in acute leukemia for detection of minimal residual disease. Report of the BIOMED-1 Concerted Action: investigation of minimal residual disease in acute leukemia. *Leukemia* 1999;13:1901–28.
  17. Kohlmann A, Schoch C, Dugas M, et al. Pattern robustness of diagnostic gene expression signatures in leukemia. *Genes Chromosomes Cancer* 2005;42:299–307.
  18. Li C, Wong WH. Model-based analysis of oligonucleotide arrays: expression index computation and outlier detection. *Proc Natl Acad Sci U S A* 2001;98:31–6.
  19. Ihaka R, Gentleman R. R: a language for data analysis and graphics. *J Computational and Graphical Statistics* 1996;5:299–314.
  20. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J Royal Statistical Society, Series B* 1995;57:289–300.
  21. Eisen MB, Spellman PT, Brown PO, Botstein D. Cluster analysis and display of genome-wide expression patterns. *Proc Natl Acad Sci U S A* 1998;95:14863–8.
  22. Husson H, Carideo EG, Neuberger D, et al. Gene expression profiling of follicular lymphoma and normal germinal center B cells using cDNA arrays. *Blood* 2002;99:282–9.
  23. Hussey DJ, Nicola M, Moore S, Peters GB, Dobrovic A. The (4;11)(q21;p15) translocation fuses the NUP98 and RAPIGDS1 genes and is recurrent in T-cell acute lymphocytic leukemia. *Blood* 1999;94:2072–9.
  24. Cimino G, Sprovieri T, Rapanotti MC, Foa R, Mecucci C, Mandelli F. Molecular evaluation of the NUP98/RAPIGDS1 gene frequency in adults with T-acute lymphoblastic leukemia. *Haematologica* 2001;86:436–7.
  25. Rozovskaia T, Feinstein E, Mor O, et al. Upregulation of Meis1 and HoxA9 in acute lymphocytic leukemias with the t(4;11) abnormality. *Oncogene* 2001;20:874–8.
  26. Dunne J, Hanby AM, Poulson R, et al. Molecular cloning and tissue expression of FAT, the human homologue of the *Drosophila* fat gene that is located on chromosome 4q34–q35 and encodes a putative adhesion molecule. *Genomics* 1995;30:207–23.
  27. Sarkar S, Roy BC, Hatano N, Aoyagi T, Gohji K, Kiyama R. A novel ankyrin repeat-containing gene (Kank) located at 9p24 is a growth suppressor of renal cell carcinoma. *J Biol Chem* 2002;277:36585–91.
  28. Moos PJ, Raetz EA, Carlson MA, et al. Identification of gene expression profiles that segregate patients with childhood leukemia. *Clin Cancer Res* 2002;8:3118–30.
  29. Tsutsumi S, Taketani T, Nishimura K, et al. Two distinct gene expression signatures in pediatric acute lymphoblastic leukemia with MLL rearrangements. *Cancer Res* 2003;63:4882–7.
  30. Mancini M, Vegna ML, Castoldi GL, et al. Partial deletions of long arm of chromosome 6: biologic and clinical implications in adult acute lymphoblastic leukemia. *Leukemia* 2002;16:2055–61.
  31. Cimino G, Elia L, Mancini M, et al. Clinico-biologic features and treatment outcome of adult pro-B-ALL patients enrolled in the GIMEMA 0496 study: absence of the ALL1/AF4 and of the BCR/ABL fusion genes correlates with a significantly better clinical outcome. *Blood* 2003;102:2014–20.
  32. Ferrando AA, Armstrong SA, Neuberger DS, et al. Gene expression signatures in MLL-rearranged T-lineage and B-precursor acute leukemias: dominance of HOX dysregulation. *Blood* 2003;102:262–8.
  33. Kottaridis PD, Gale RE, Linch DC. FLT3 mutations and leukaemia. *Br J Haematol* 2003;122:523–38.
  34. Libura M, Asnafi V, Tu A, et al. FLT3 and MLL intra-genic abnormalities in AML reflect a common category of genotoxic stress. *Blood* 2003;102:2198–204.
  35. Thiede C, Studel C, Mohr B, et al. Analysis of FLT3-activating mutations in 979 patients with acute myelogenous leukemia: association with FAB subtypes and identification of subgroups with poor prognosis. *Blood* 2002;99:4326–35.
  36. Brown P, Levis M, Shurtleff S, Campana D, Downing J, Small D. FLT3 inhibition selectively kills childhood acute lymphoblastic leukemia cells with high levels of FLT3 expression. *Blood* 2005;105:812–20.
  37. Torelli GF, Guarini A, Porzia A, et al. FLT3 inhibition in t(4;11)+ adult acute lymphoid leukaemia. *Br J Haematol* 2005;130:43–50.
  38. Kohlmann A, Schoch C, Schnittger S, et al. Pediatric acute lymphoblastic leukemia (ALL) gene expression signatures classify an independent cohort of adult ALL patients. *Leukemia* 2004;18:63–71.
  39. Druker BJ, Sawyers CL, Kantarjian H, et al. Activity of a specific inhibitor of the BCR-ABL tyrosine kinase in the blast crisis of chronic myeloid leukemia and acute lymphoblastic leukemia with the Philadelphia chromosome. *N Engl J Med* 2001;344:1038–42.
  40. Shah NP, Tran C, Lee FY, Chen P, Norris D, Sawyers CL. Overriding imatinib resistance with a novel ABL kinase inhibitor. *Science* 2004;305:399–401.
  41. Weisberg E, Manley PW, Breitenstein W, et al. Characterization of AMN107, a selective inhibitor of native and mutant Bcr-Abl. *Cancer Cell* 2005;7:129–41.
  42. Rozovskaia T, Ravid-Amir O, Tillib S, et al. Expression profiles of acute lymphoblastic and myeloblastic leukemias with ALL-1 rearrangements. *Proc Natl Acad Sci U S A* 2003;100:7853–8.
  43. Debernardi S, Lillington DM, Chaplin T, et al. Genome-wide analysis of acute myeloid leukemia with normal karyotype reveals a unique pattern of homeobox gene expression distinct from those with translocation-mediated fusion events. *Genes Chromosomes Cancer* 2003;37:149–58.
  44. Schoch C, Kohlmann A, Schnittger S, et al. Acute myeloid leukemias with reciprocal rearrangements can be distinguished by specific gene expression profiles. *Proc Natl Acad Sci U S A* 2002;99:10008–13.
  45. Kohlmann A, Schoch C, Schnittger S, et al. Molecular characterization of acute leukemias by use of microarray technology. *Genes Chromosomes Cancer* 2003;37:396–405.
  46. Armstrong SA, Mabon ME, Silverman LB, et al. FLT3 mutations in childhood acute lymphoblastic leukemia. *Blood* 2004;103:3544–6.
  47. Taketani T, Taki T, Sugita K, et al. FLT3 mutations in the activation loop of tyrosine kinase domain are frequently found in infant ALL with MLL rearrangements and pediatric ALL with hyperdiploidy. *Blood* 2004;103:1085–8.
  48. Teuffel O, Dettling M, Cario G, et al. Gene expression profiles and risk stratification in childhood acute lymphoblastic leukemia. *Haematologica* 2004;89:801–8.
  49. Stams WA, den Boer ML, Holleman A, et al. Asparagine synthetase expression is linked with L-asparaginase resistance in TEL-AML1-negative but not TEL-AML1-positive pediatric acute lymphoblastic leukemia. *Blood* 2005;105:4223–5.