

Gene Expression Differences between Colon and Rectum Tumors

Rebeca Sanz-Pamplona¹, David Cordero¹, Antonio Berenguer¹, Flavio Lejbkowitz⁶, Hedy Rennert⁶, Ramon Salazar², Sebastiano Biondo^{3,5}, Xavier Sanjuan⁴, Miguel A. Pujana¹, Laura Rozek¹², Thomas J. Giordano¹³, Ofer Ben-Izhak⁷, Hector I. Cohen⁸, Philip Trougouboff¹⁰, Jacob Bejhar¹¹, Yanina Sova⁹, Gad Rennert⁶, Stephen B. Gruber¹⁴, and Victor Moreno^{1,5}

Abstract

Purpose: Colorectal cancer studies typically include both colon and rectum tumors as a common entity, though this assumption is controversial and only minor differences have been reported at the molecular and epidemiologic level. We conducted a molecular study based on gene expression data of tumors from colon and rectum to assess the degree of similarity between these cancer sites at transcriptomic level.

Experimental Design: A pooled analysis of 460 colon tumors and 100 rectum tumors from four data sets belonging to three independent studies was conducted. Microsatellite instable tumors were excluded as these are known to have a different expression profile and have a preferential proximal colon location. Expression differences were assessed with linear models, and significant genes were identified using adjustment for multiple comparisons.

Results: Minor differences at a gene expression level were found between tumors arising in the proximal colon, distal colon, or rectum. Only several *HOX* genes were found to be associated with tumor location. More differences were found between proximal and distal colon than between distal colon and rectum.

Conclusions: Microsatellite stable colorectal cancers do not show major transcriptomic differences for tumors arising in the colon or rectum. The small but consistent differences observed are largely driven by the *HOX* genes. These results may have important implications in the design and interpretation of studies in colorectal cancer. *Clin Cancer Res*; 17(23); 7303–12. ©2011 AACR.

Introduction

Colorectal cancer (CRC) is considered a heterogeneous complex disease that comprises different tumor phenotypes (1). Attempts to classify tumors from a molecular perspective

that identify carcinogenic pathways have proposed 3 categories with some overlap as follows: chromosomal instability (CIN) tumors, microsatellite instability (MSI) tumors, and CpG island methylator phenotype (CIMP) tumors. This taxonomy plays a significant role in determining clinical, pathologic, and biological characteristics of CRC (2).

From a clinical point of view, the colon and rectal cancers are treated as distinct entities. Colon tumors are usually divided as proximal or right sided when originating proximal to the splenic flexure (cecum, ascending colon, and transverse colon) whereas distal tumors arise distal to this site (descending colon and sigmoid colon). Distal colon or left-sided tumors most often appear in the rectum sigmoid flexure, and the distinction of these from rectal tumors is not always easy. Usually, a tumor is considered rectal when arising within 15 cm from the anal sphincter (3, 4). Indeed, accumulating evidences suggest that grouping these anatomically distinct diseases could be a clinical and biological oversimplification: rectal cancers show higher rates of locoregional relapse and lung metastases, whereas colon cancers have a higher tropism for liver spread and a slightly better overall prognosis (5). Moreover, proximal location of colon cancer is a risk factor for development of metachronous CRC (6). Treatment also differs for colon and rectal tumors. Although both colon and rectal cancers benefit from adjuvant chemotherapy, radiation therapy is

Authors' Affiliations: ¹Unit of Biomarkers and Susceptibility and ²Medical Oncology Service, Catalan Institute of Oncology (ICO), IDIBELL, and CIBERESP, Departments of ³Digestive Surgery and ⁴Pathology, University Hospital Bellvitge, L'Hospitalet de Llobregat; ⁵Department of Medical Sciences, School of Medicine, University of Barcelona, Barcelona, Spain; ⁶Clalit Health Services National Cancer Control Center and Department of Community Medicine and Epidemiology, Carmel Medical Center and B. Rappaport Faculty of Medicine, Technion-Israel Institute of Technology; ⁷Rambam Medical Center; ⁸Bnai Zion Medical Center; ⁹Carmel Medical Center, Haifa; ¹⁰Western Galilee Medical Center, Nahariya; and ¹¹Haemek Medical Center, Afula, Israel; and Departments of ¹²Environmental Health Sciences, ¹³Pathology, and ¹⁴Internal Medicine, Epidemiology, Human Genetics, University of Michigan, Ann Arbor, Michigan

Note: Supplementary data for this article are available at Clinical Cancer Research Online (<http://clincancerres.aacrjournals.org/>).

R. Sanz-Pamplona and D. Cordero contributed equally to this work.

Corresponding Author: Victor Moreno, Catalan Institute of Oncology, Av. Gran Via 199, L'Hospitalet de Llobregat, Barcelona 08908, Spain. Phone: 34-93-260-71-86; Fax: 34-93-260-71-88; E-mail: v.moreno@iconcologia.net

doi: 10.1158/1078-0432.CCR-11-1570

©2011 American Association for Cancer Research.

Translational Relevance

Colorectal cancer (CRC) studies typically include both colon and rectum tumors as a common entity, though this assumption is controversial and only minor differences have been reported at the molecular and epidemiologic level. Here, we report a large sample pool study concluding that only minor differences at a gene expression level exist between microsatellite stable CRCs at different locations. These results have important implications in the design and interpretation of studies in colorectal cancer. For instance, several molecular profiles have been recently proposed to predict prognosis in patients with CRC that combine colon and rectum cases, assuming this hypothesis without the real proof. The conclusions provided by this study will help consolidate the idea that at the molecular level, the minor expression differences identified are more related to anatomic developmental differences than to tumoral mechanisms.

only indicated in locally advanced rectal tumors (7). Epidemiologic risk factors reflect somewhat more controversial distinctions between cancers of the colon and rectum: alcohol intake was significantly positively associated with higher risk in the rectum than in colon tumors (8). Other dietary risk factors differing between colon and rectum tumors have been suggested more inconsistently (9, 10).

At the molecular level, differences in expression of specific genes and proteins (cyclin A2, COX-2, and β -catenin) have been reported (reviewed in ref. 6). Moreover, colon cancers have a higher number of mutations including *KRAS* and *BRAF* mutations. The CIN pathway is far more common in rectal cancers than colon cancers, whereas MSI and CIMP cancers are more likely to be in the right colon. Some of the reported differences in gene expression probably correspond to molecular signatures of MSI, such as the correlation between *CDX2* expression and MSI (11).

Recently, several molecular profiles have been proposed to predict prognosis in patients with CRC (12–15). These studies typically combine colon and rectal cancers, but it is not known whether this combination is appropriate. Expression profiles may inform this choice. If proximal colon, distal colon, and rectal tumors share a common set of expressed transcripts, then it may be reasonable to combine data for prognostic studies, and in fact may inform choices for epidemiologic study designs. The aim of this work was to compare gene expression among CRC subsites in an attempt to identify molecular factors that correspond to differences in the clinical behavior of these tumors.

Materials and Methods

Study population

The Molecular Epidemiology of Colorectal Cancer (MECC) study is a population-based, case-control study

that included 2,138 incident CRC cases and 2,049 population controls from Northern Israel (16). A pathology review of the diagnostic slides centralized at the University of Michigan (Ann Arbor, MI) confirmed the eligibility criteria of invasive adenocarcinoma. The study was approved by the Institutional Review Boards at the University of Michigan and Carmel Medical Center in Haifa. Written and informed consent was required for inclusion.

A subset of these patients provided fresh tumor tissue samples that were analyzed for expression in 2 stages as previously described (17). Initially, a subset of 170 tumors was hybridized with the Affymetrix HG-U133A gene array (MECC-A). In a second stage, an additional sample of 232 tumors was hybridized in the HG-U133plus 2.0 gene array (MECC-P2). Of these patients, 4 from the first set and 7 from the second were excluded because they had multiple tumors in the colon and rectum, or the precise location was not provided. Expression data are available in Gene Expression Omnibus (GEO; ref. 18) repository with accession code GSE26682.

In addition of these 2 gene expression data sets (MECC-A and MECC-P2), publicly available expression data with information about subsite were searched in the GEO and ArrayExpress (19) databases. To guarantee a high quality analysis, the inclusion criteria was restricted to studies that had used Affymetrix U133 gene chips, with more than 50 samples, and a minimum number of 10 for each site. Two data sets were identified matching these criteria: GSE14333 included 290 consecutive patients with CRC [colon ($n = 250$), rectum ($n = 39$), and missing site ($n = 1$); ref. 20]. GSE13294 comprised 155 patients with CRC (122 colon, 25 rectum, and 8 missing; ref. 21). In addition, data set GSE9254 was identified, that included 19 normal mucosa samples from different colonic locations: cecum ($n = 2$), ascending ($n = 3$), transverse ($n = 3$), sigmoid ($n = 4$), and rectum ($n = 7$; ref. 22).

Quality control and normalization

Prior to data analysis, a careful quality control process following the Affymetrix recommendations was conducted (23). This procedure rejected 122 samples: 27 (16%) from MECC-A, 49 (21%) from MECC-P2, 21 (7%) from GSE14333, and 25 (16%) from GSE13294.

Data normalization were carried out with the *R* statistical software, version 2.9.0 (*R* foundation for statistical computing; <http://www.r-project.org>) and Bioconductor package (Bioconductor core group; <http://www.bioconductor.org>). Raw data from the different data sets were normalized together with the Robust Multiarray Average (RMA) method (24). To improve comparability between arrays from different studies, only the common subset of probes from the U133A array ($n = 22,283$) were selected, and data were renormalized with a quantile method.

MSI

Tumors showing MSI appear more often in right colon and are known to have a marked different expression profile

(25). In an attempt to homogenize the analysis and avoid potential biases due to this condition, MSI tumors were excluded from all data sets. For MECC cases, MSI was analyzed using 7 microsatellite markers that included the National Cancer Institute panel (26). Cases were considered MSI when more than 30% of the markers were unstable. A total of 16 cases were excluded from MECC-A and 15 from MECC-P2. A total of 61 MSI samples from data set GSE1324 were also excluded.

MSI status was not available for the public GSE14333 data set but was imputed using a molecular profiling-based approach (details in Supplementary Table S1 and Fig. S1). Out of the 268 samples, 53 (20%) were labeled as MSI and removed for further analysis. These excluded cases might not be a perfect selection of the real subset of MSI tumors, but their clinical characteristics are in agreement with the expectations: more frequent in female and older patients and with preferential location in right colon (Supplementary Table S2).

Differential expression analysis

Prior to the identification of differentially expressed probes, a filter was applied to remove those with low variability ($n = 7,509$), which mostly correspond to non-hybridized and saturated probes. The remaining 14,774 probes with SD greater than 0.3 were considered for further analysis. To test for differences in expression between sites, a linear model adjusted for gender, age, and study was fitted to each probe. To account for multiple comparisons, the Bonferroni correction was used. Also the less conservative q value method was used to control the false discovery rate (FDR).

Heterogeneity of expression profiles by tumor site across studies was evaluated for each probe by the linear models described earlier. A test for interaction between cancer site and study was conducted for each probe and, again, the q value method was used to correct the results by multiple comparisons.

Gene set enrichment analysis

The gene set enrichment analysis (GSEA) algorithm (27) was applied to identify enrichment of specific functions in

the list of genes preranked according to their P value for the test of differences in expression between subsites. The statistical significance of the enrichment score was calculated by permuting the genes 1,000 times as implemented in the GSEA software.

Classification of colon/rectum samples using differentially expressed genes

For each comparison considered, an agglomerative hierarchical clustering method was used to display the classification ability among site of the corresponding list of differentially expressed probes sets. This discriminating ability was formally tested using a linear discriminant analysis with leave-one-out crossvalidation to estimate the prediction error rate.

Results

Clinical data for the 460 colon tumors and 100 rectum tumors included in the analysis are summarized in Table 1. A principal component analysis (PCA) was done to assess global differences between each data set. The first and second components separated the samples by study, suggesting systematic differences that could not be corrected by careful homogeneous criteria and normalization (Supplementary Fig. S2). The most dissimilar data set was MECC-A, probably due to be the fact that the platform was Affymetrix H-U133 A gene chips instead of H-U133 Plus 2.0 used in the other studies. All pooled analyses were adjusted for study to account for these systematic differences.

Gene expression profiling: colon versus rectum tumors

Linear models adjusted for study, age, and gender identified only 11 of 14,774 differentially expressed probes between colon and rectum after Bonferroni correction. The less conservative q value method identified 20 probes (corresponding to 16 genes, Table 2) when a 1% FDR was used and 131 probes (111 genes) at the 5% FDR. Moreover, among these differentially expressed genes, no one had an absolute \log_2 fold change greater than one (Fig. 1A). These results suggest that the magnitude of expression differences

Table 1. Samples description

$n = 560$	Site ^a			Platform	Mean age	Gender ^b		Stage ^b			
	Right	Left	Rectum			Male	Female	I	II	III	IV
MECC-A ($n = 123$)	55 (44.7%)	57 (46.4%)	11 (8.9%)	Affy HG-U133A	72.53	68 (55.3%)	55 (44.7%)	4 (3.4%)	58 (50%)	41 (35.4%)	13 (11.2%)
MECC-P2 ($n = 161$)	58 (36.9%)	59 (37.6%)	40 (25.5%)	Affy U133 Plus 2.0	72.01	87 (54%)	74 (46%)	20 (15.4%)	55 (42.3%)	39 (30%)	16 (12.3%)
GSE14333 ($n = 215$)	79 (37.1%)	100 (46.9%)	34 (16%)	Affy U133 Plus 2.0	65.65	132 (61.4%)	83 (38.6%)	34 (15.8%)	61 (28.4%)	64 (29.8%)	56 (26%)
GSE13294 ($n = 61$)	46 (75.4%)		15 (24.6%)	Affy U133 Plus 2.0	65.43	32 (53.3%)	28 (46.7%)	0 (0%)	46 (75.4%)	7 (11.5%)	8 (13.1%)

^aSome cases were classified as "colon" with no information about specific subsite.

^bNumber may not add to total due to missing information.

Table 2. Differentially expressed genes between colon and rectum tumors

Probe	Gene	<i>q</i> value	Log ₂ fold change	Function
209844_at	<i>HOXB13</i>	3.65E-06	-0.600	Transcription factor activity
213823_at	<i>HOXA11</i>	5.91E-06	0.514	Transcription factor activity
209167_at	<i>GPM6B</i>	1.88E-05	-0.355	Cell differentiation
209170_s_at		3.15E-03	-0.366	
214651_s_at	<i>HOXA9</i>	2.32E-05	0.902	Transcription factor activity
209905_at		5.08E-05	0.673	
213147_at	<i>HOXA10</i>	2.99E-05	0.460	Transcription factor activity
213150_at		2.68E-04	0.534	
213844_at	<i>HOXA5</i>	2.40E-04	0.663	Transcription factor activity
39835_at	<i>SBF1</i>	3.20E-04	0.270	Protein amino acid dephosphorylation
218211_s_at	<i>MLPH</i>	2.52E-03	0.655	Melanosome transport
216629_at	<i>SRRM2</i>	2.78E-03	0.079	RNA splicing
205555_s_at	<i>MSX2</i>	2.89E-03	0.387	Transcription factor activity
210319_x_at		3.15E-03	0.455	
204461_x_at	<i>RAD1</i>	3.15E-03	-0.292	DNA repair
59644_at	<i>BMP2K</i>	3.65E-03	-0.291	Protein amino acid phosphorylation
215703_at	<i>CFTR</i>	5.60E-03	-0.396	Transmembrane transport
204425_at	<i>ARHGAP4</i>	7.13E-03	0.242	Apoptosis
203332_s_at	<i>INPP5D</i>	7.47E-03	0.387	Apoptosis
206854_s_at	<i>MAP3K7</i>	9.86E-03	-0.335	Signal transduction

among microsatellite stable (MSS) tumors arising in the colon and rectum is quite small.

Functionally, it was noteworthy that 5 of the top 6 genes belonged to the *HOX* family of transcription factors (Table 2). Other top differentially expressed genes displayed assorted functions such as DNA repair, transcription factor activity, intracellular transport, signal transduction, and apoptosis among others. To formally identify enriched biological processes associated with differentially expressed genes, a GSEA was done. Although no significant function was retrieved, the "*HOX* genes" set appeared with the highest gene enrichment score (Supplementary Fig. S3).

Heterogeneity across studies was explored to identify genes that might have differences in some studies but opposite direction in others that might compensate in the pooled analysis. Only 12 probes showed heterogeneity between studies at the 5% FDR and these could not be ascribed to a systematic effect of one specific study (Supplementary Fig. S4). None of these 12 heterogeneous probes corresponded to differentially expressed genes. Therefore, the 4 studies included in our analysis were considered homogeneous about their differences in expression profiles between colon and rectum.

Refining gene expression profiling: right colon versus left colon tumors and right colon versus rectum tumors

To discount the possibility that similar molecular backgrounds in left colon and rectum tumors were masking possible differences between total colon samples and rectum tumors, a more detailed analysis was conducted looking for differences between right colon, left colon, and

rectum tumors, when detailed data about cancer site were available ($n = 499$, all data sets except GSE13294).

Similar to previous results, no major differences were detected between right and left colon, reinforcing our impression that MSS colorectal tumors show very similar expression profiles regardless of their site of origin. Ten genes were found to be differentially expressed between right and left colon tumors after Bonferroni correction. The *q* value method only identified 44 probes differentially expressed corresponding to 40 genes at 1% FDR (Table 3) and 174 probes (150 genes) at 5% FDR. Interestingly, the comparison between left colon and rectum did not identify any differentially expressed gene at 1% FDR (only 3 genes were found at FDR 5%). In contrast, 54 probes (50 genes) were differentially expressed between right colon and rectum when a 1% FDR was used (Table 4) and 374 probes (324 genes) at the 5% FDR. From those, 21 probes (18 genes) passed Bonferroni correction (Fig. 1B). Functionally, those genes showed varied functions, highlighting the *HOX* family as in previous analysis.

To assess the ability of these profiles to discriminate cancer samples by location, a linear discriminant analysis model was built. Leave-one-out internal validation showed that only 37% of rectum tumors were correctly classified when using the colon versus rectum signature (Fig. 1C). Better performance was obtained using the right versus left signature, with 77% accuracy both in right and left tumors (Fig. 1D). The best classification was achieved using the right versus rectum tumors profile (with a total accuracy of 86%), indicating that the major differences exist between the most opposite locations (Fig. 1E).




Figure 1. Fold change plot and prediction ability of site-related differentially expressed genes. A, mean expression of each probe set versus its fold change between colon and rectum tumors. B, number of differentially expressed genes between each tumoral location at FDR 1%. Dendrogram illustrating the classification ability of differentially expressed genes among site in colon versus rectum (C), right versus left (D), and right versus rectum (E). Companion tables show the accuracy of each study.

Because classification of rectal tumors is controversial and misclassification could exist between rectal and sigmoid colon tumors, an analysis in which rectal and left-sided colon cancers were pooled and compared with right-sided colon cancer was also conducted. As a result, 46 probes corresponding to 35 genes were found to be differentially expressed after Bonferroni correction. The *q* value method identified 256 probes (202 genes) differentially expressed at 1% FDR (Supplementary Table S3) and 884 probes at 5% FDR. Though this comparison showed a larger number of significant probes, related to the increased sample size of the distal location group, the magnitude of the differences were very small (<10%) and probably not biologically relevant.

HOX genes

Remarkably, *HOX* appeared as the most differentially expressed genes in all transcriptomic comparisons and emerged in the intersection of the lists of differentially expressed genes. In fact, these *HOX* genes were expressed in a gradient in colorectal tumors. The *HOX* genes were more expressed in tumors from the proximal colon, and their expression decreased along more distal locations in the

gastrointestinal tract, with the exception of *HOXB13* that showed a reversed pattern (Fig. 2). Genes known to be targets of *HOX* transcription factors (28) were analyzed, but these showed no differences in expression between subsites, indicating that differences observed in *HOX* genes were not affecting a cascade of regulated genes (Supplementary Fig. S5). Also, specific GSEA analysis using *HOX*-related gene sets showed a statistically significant enrichment for genes activated by the chimeric protein NUP98-HOXA9, an aberrant *HOX* transcription factor and also an enrichment in genes with promoter regions around transcription start site containing the motif that binds with *HOX9* (Supplementary Table S4).

Interestingly, the analysis of expression for *HOX* genes in human normal colorectal mucosa in the data set GES9254 showed the same gradient along the gut than in tumor samples (Supplementary Fig. S6).

Discussion

This pool analysis of 4 data sets from 3 independent studies including a total of 560 samples suggests that there

Downloaded from http://aacrjournals.org/clincancerres/article-pdf/17/23/7307/3032006529/7303.pdf by guest on 13 January 2025

Table 3. Differentially expressed genes between right and left colon

Probe	Gene	q value	Log ₂ fold change	Function
206858_s_at	<i>HOXC6</i>	2.04E-08	0.868	Transcription factor activity
209844_at	<i>HOXB13</i>	1.18E-06	-0.521	Transcription factor activity
219109_at	<i>SPAG16</i>	6.11E-05	-0.703	Cell projection
205767_at	<i>EREG</i>	1.47E-04	-1.082	Growth factor activity
206307_s_at	<i>FOXD1</i>	2.50E-04	0.434	Transcription factor activity
209524_at		2.76E-04	-0.678	
209526_s_at	<i>HDGFRP3</i>	6.17E-04	-0.512	Growth factor activity
216693_x_at		6.31E-04	-0.496	
203988_s_at	<i>FUT8</i>	1.62E-03	0.308	N-glycan processing
205555_s_at	<i>MSX2</i>	2.01E-03	0.393	Transcription factor activity
210319_x_at		8.60E-03	0.440	
209752_at	<i>REG1A</i>	2.01E-03	1.263	Growth factor activity
217918_at	<i>DYNLRB1</i>	3.16E-03	-0.212	Microtubule-based movement
212423_at	<i>ZCCHC24</i>	3.63E-03	-0.406	Nucleic acid binding
212419_at		9.56E-03	-0.322	
219228_at	<i>ZNF331</i>	3.63E-03	-0.316	Transcription factor activity
219955_at	<i>L1TD1</i>	3.82E-03	0.878	Transposase
207457_s_at	<i>LY6G6D</i>	4.19E-03	-0.786	—
218094_s_at	<i>DBNDD2</i>	4.30E-03	-0.254	Regulation of protein kinase activity
217665_at	—	5.11E-03	-0.247	—
202925_s_at	<i>PLAGL2</i>	5.56E-03	-0.334	Transcription factor activity
208948_s_at	<i>STAU1</i>	5.56E-03	-0.171	RNA binding
217801_at	<i>ATP5E</i>	5.56E-03	-0.138	ATP synthesis
212349_at	<i>POFUT1</i>	5.98E-03	-0.252	Notch signaling pathway
204819_at	<i>FGD1</i>	6.02E-03	-0.201	Signal transduction
205815_at	<i>REG3A</i>	7.19E-03	1.011	Cell proliferation
206340_at	<i>NR1H4</i>	7.19E-03	0.177	Transcription factor activity
208979_at	<i>NCOA6</i>	7.94E-03	-0.194	Transcription regulation
2019.98_at	<i>ST6GAL1</i>	8.51E-03	-0.409	Protein amino acid glycosylation
202673_at	<i>DPM1</i>	8.51E-03	-0.239	Protein binding
217718_s_at	<i>YWHAB</i>	8.60E-03	-0.138	Signal transduction
204555_s_at	<i>PPP1R3D</i>	8.82E-03	-0.260	Protein binding
205463_s_at	<i>PDGFA</i>	8.82E-03	-0.323	Growth factor activity
205997_at	<i>ADAM28</i>	8.82E-03	0.295	Proteolysis
212234_at	<i>ASXL1</i>	8.82E-03	-0.200	Regulation of transcription
212787_at	<i>YLPM1</i>	8.82E-03	0.141	Regulation of transcription
213170_at	<i>GPX7</i>	8.82E-03	-0.287	Response to oxidative stress
214482_at	<i>ZBTB25</i>	8.82E-03	0.131	Transcription factor activity
215210_s_at	<i>DLST</i>	8.82E-03	0.238	Tricarboxylic acid cycle
218325_s_at	<i>DIDO1</i>	8.82E-03	-0.241	Apoptosis
219108_x_at	<i>DDX27</i>	8.82E-03	-0.188	RNA binding
221472_at	<i>SERINC3</i>	8.82E-03	-0.190	Protein binding
204015_s_at	<i>DUSP4</i>	9.56E-03	0.368	Signal transduction
2031.27_s_at	<i>SPTLC2</i>	9.79E-03	0.199	Lipid metabolism

are identifiable expression differences among MSS CRCs that arise in different sites within the large intestine. However, the number of statistically significant differentially expressed genes found between tumor locations was minimal, and the fold change of their expression was within random variation for most cases. With the exception of the *HOX* family, there were no identifiable functional distinc-

tions among the differentially expressed genes. Moreover, the most evident distinctions in expression profiles were those between the right colon and either the left colon or rectum. Expression profiles of MSS rectal cancers and left-sided colon cancers were virtually indistinguishable.

These results imply that anatomic differences are relevant for the clinical management of CRC, but specific molecular

Table 4. Differentially expressed genes between right colon and rectum tumors

Probe	Gene	q value	Log ₂ fold change	Function
209844_at	<i>HOXB13</i>	3.51E-09	-0.856	Transcription factor activity
205555_s_at	<i>MSX2</i>	4.30E-05	0.586	Transcription factor activity
210319_x_at		7.11E-05	0.696	
213823_at	<i>HOXA11</i>	4.30E-05	0.590	Transcription factor activity
214651_s_at	<i>HOXA9</i>	4.30E-05	1.013	Transcription factor activity
209905_at		3.98E-04	0.748	
206858_s_at	<i>HOXC6</i>	8.90E-05	1.057	Transcription factor activity
218211_s_at	<i>MLPH</i>	9.10E-05	0.856	ROS metabolism
213844_at	<i>HOXA5</i>	1.02E-04	0.806	Transcription factor activity
213150_at	<i>HOXA10</i>	1.77E-04	0.590	Transcription factor activity
213147_at		6.82E-04	0.509	
3983.5_at	<i>SBF1</i>	1.77E-04	0.343	Protein amino acid dephosphorylation
211756_at	<i>PTHLH</i>	8.02E-04	-0.167	Hormone activity
206854_s_at	<i>MAP3K7</i>	8.77E-04	-0.408	Signal transduction
219109_at	<i>SPAG16</i>	9.80E-04	-0.858	Cell projection
214598_at	<i>CLDN8</i>	9.93E-04	-0.722	Cell adhesion
209167_at	<i>GPM6B</i>	1.15E-03	-0.389	Cell differentiation
204425_at	<i>ARHGAP4</i>	1.18E-03	0.334	Apoptosis
36554_at	<i>ASMTL</i>	1.36E-03	0.263	Melatonin biosynthesis
204667_at	<i>FOXA1</i>	1.43E-03	0.481	Transcription factor activity
204042_at	<i>WASF3</i>	1.44E-03	-0.660	Actin binding
203699_s_at	<i>DIO2</i>	1.69E-03	-0.281	Hormone biosynthesis
213927_at	<i>MAP3K9</i>	1.69E-03	0.130	Signal transduction
211737_x_at	<i>PTN</i>	1.92E-03	-0.240	Growth factor activity
209465_x_at		2.34E-03	-0.367	
212840_at	<i>UBXN7</i>	2.34E-03	-0.501	Protein binding
210766_s_at	<i>CSE1L</i>	2.70E-03	-0.396	Protein transport
215703_at	<i>CFTR</i>	2.70E-03	-0.441	Respiratory gaseous exchange
216129_at	<i>ATP9A</i>	2.70E-03	-0.458	ATP biosynthesis
212234_at	<i>ASXL1</i>	3.21E-03	-0.257	Regulation of transcription
218454_at	<i>PLBD1</i>	3.57E-03	-0.375	Lipid degradation
205423_at	<i>AP1B1</i>	4.08E-03	0.204	Protein transport
206070_s_at	<i>EPHA3</i>	4.59E-03	-0.421	Receptor
203628_at	<i>IGF1R</i>	4.83E-03	-0.544	Receptor
202949_s_at	<i>FHL2</i>	4.98E-03	0.347	Transcription regulation
221738_at	<i>KIAA1219</i>	4.98E-03	-0.229	Signal transduction
202760_s_at	<i>PALM2</i>	5.30E-03	-0.503	Regulation of cell shape
219228_at	<i>ZNF331</i>	5.30E-03	-0.218	Regulation of transcription
219426_at	<i>EIF2C3</i>	6.45E-03	-0.486	RNA binding
214234_s_at	<i>CYP3A5</i>	6.64E-03	0.437	Electron carrier activity
218892_at	<i>DCHS1</i>	6.64E-03	-0.162	Cell adhesion
222015_at	<i>CSNK1E</i>	6.67E-03	0.321	Signal transduction
209195_s_at	<i>ADCY6</i>	6.76E-03	0.260	Signal transduction
215078_at	<i>SOD2</i>	7.65E-03	-0.363	Removal of superoxide radicals
203671_at	<i>TPMT</i>	7.85E-03	-0.238	Metabolism of thiopurine drugs
205767_at	<i>EREG</i>	7.85E-03	-1.211	Growth factor activity
221091_at	<i>INSL5</i>	7.85E-03	-0.406	Hormone activity
202925_s_at	<i>PLAGL2</i>	7.88E-03	-0.395	Transcription factor activity
213242_x_at	<i>KIAA0284</i>	8.06E-03	0.327	Microtubule organization
202673_at	<i>DPM1</i>	8.45E-03	-0.240	Protein binding
219955_at	<i>L1TD1</i>	8.47E-03	1.064	Transposase
201978_s_at	<i>KIAA0141</i>	8.75E-03	0.300	—
32069_at	<i>N4BP1</i>	8.75E-03	-0.220	Protein binding
211843_x_at	<i>CYP3A7</i>	9.25E-03	0.367	Electron carrier activity




Figure 2. *HOX* genes reverse gradient of expression along colorectal tumor locations. Mean expression value of *HOX* genes in right colon, left colon, and rectum tumors. Genes marked with an asterisk are represented in the microarray by more than one probe set.

profiles of MSS CRC are for the large part quite similar. It is well known that metastases from CRC develop in a stepwise process (29). Rectal cancers usually have a pattern of local recurrence, and retrospective studies show a relevant influence of the surgeon on the prognosis of these patients (30). For colon cancers, the progression pattern is more typically characterized by liver metastases, potentially explained by the fact that superior mesenteric vein drains the right colon whereas neither the left colon nor the rectal vasculature directly drains to liver (29). One might have hypothesized that molecular differences such as DNA repair, apoptosis, or angiogenesis might have distinguished rectal cancers, given the differential efficacy of radiotherapy for rectal cancers. However, our study did not reveal any such clues or signatures. The samples that were analyzed were all tumors collected prior to treatment. Although it is possible that expression profiles that predict response to radiotherapy might exist, our pretreatment data are unable to address this hypothesis. In addition, there is no known evidence of differential radiation sensitivity between colon and rectal cancers. It is only the particular topographic intrapelvic location of the rectum that renders it appropriate for radiotherapy due to the lack of small bowel interaction with the radiation field, which is the limiting factor of the radiotherapy administration in colon cancer (31, 32).

A potential concern of studies that fail to detect differences in expression patterns between tumors is the possibility of insufficient statistical power to detect clinically or biologically meaningful differences due to a small sample size. To address this issue, a pooled analysis has been conducted that included a total of 560 samples, enough to detect differences of 0.5 SD units. In practice, most of the few significant genes identified showed fold changes smaller than 0.6 or a 50% variation in expression, which is usually considered small in microarray expression analyses. Small studies also may show apparent differences that are particular to the selection of cases analyzed. The strength of meta-analyses like the one reported here is that only

consistent results remain, and these are easily identified as power is larger and heterogeneity can be explored to identify study specificities. In our analysis, heterogeneity among studies was not a concern as only 12 probes, out of almost 15,000 explored, showed significant heterogeneity and they could not be ascribed to a specific study.

MSI tumors were not included in the analysis due to their known different molecular background (21, 25, 33) and strong association with tumor location. In the case of GSE14333 data set, the researchers did not provide information about MSI status so a simple signature-based imputation was done to exclude putative MSI tumors from the analysis. This procedure had its limitations as its accuracy for MSI was only 85% (Supplementary Table S1). Thus more MSS tumors than necessary may have been excluded, and some MSI cancers from GSE14333 may have been inadvertently included by our simple imputation. This strategy of attempting to eliminate MSI CRCs was preferred to the alternative design that would have resulted in a strong biased estimation or a choice to completely exclude all 215 of the otherwise informative tumors from GSE14333. A choice to exclude these tumors would have further reduced the power to detect any possible existing differences. It is reassuring to note that tumors excluded from the analysis had clinical features related to MSI, such as a predominance of female and older patient that originate in the colon, mainly in the right side (Supplementary Table S2; ref. 34). In addition, an analysis excluding GSE14333 data set was conducted and similar results (still less significant genes) were obtained (Supplementary Table S5).

It is worth mentioning that differences between cancer sites previously reported in some studies may be related to MSI status: Komuro and colleagues found gene expression differences between right- and left-sided CRCs in genes related to MSI such as *MSH2* in right-sided tumors (35). A similar work by Birkenkamp-Demtroder and colleagues also reported differences between 25 MSS and MSI right and left tumors (36). Watanabe and colleagues describes small differences between proximal and distal MSI colorectal tumors (37). These differences are probably related to the combination of MSI and MSS tumors. *CDX2* has been reported to be more expressed in proximal structures than distal (11) but we did not find it as a right side-associated gene. However, if we include in our analysis MSI tumors and look for *CDX2* expression, it appeared as a differentially expressed gene with a *q* value less than 0.01. So, the significance of *CDX2* is probably due to MSI and not due to tumor location.

Although most of CIMP-positive tumors are MSI and therefore were not included in this analysis, there are some CIMP-positive MSS tumors that preferentially arise in the right colon (2, 38) which could explain some of the larger differences between the tumors arising in the right colon and other tumors. In an attempt to explore this possibility, a gene expression signature that differentiates MSS CIMP⁺ and MSS CIMP⁻ colorectal carcinomas was used (39) in a GSEA analysis. This revealed an association between CIMP⁺ genes and right-sided genes (Supplementary Fig. S7) and

suggests that some of the described differences could be related to CIMP phenotype.

Only *HOX* genes were found to be an enriched set associated with colon tumors. These genes (also known as homeobox genes) encode transcription factors that play essential roles in controlling cell growth and differentiation during embryonic and normal tissue development. Many homeobox genes have been reported to be deregulated in a variety of solid tumors including CRC and also to vary between normal mucosa and CRC tissue (40, 41). Interestingly, differences in *HOX* expression between carcinomas from the right colon and left colon have been reported previously (42). In normal human intestinal mucosa, *HOX-A* genes are widely expressed in undifferentiated proliferating cells at the base of the crypts (43). So, we speculate that *HOX* expression in colon tumors could be an amplification of the signal from colon cancer stem cell that drives intestinal cell differentiation. Because *HOX* expression patterns along the gut reflect pivotal roles of these genes in the regional regenerative process of the epithelial cells (44), it is possible that our results simply mirrors the *HOX* expression pattern maintained in tumors as it usually is in the normal mucosa. In fact, we observed the same gradient of expression in normal mucosa along the gut (Supplementary Fig. S6). However, despite our analysis showed no differential expression among genes targeted by *HOX*, enrichment in genes activated by NUP98-*HOXA9* was found. This is an aberrant *HOXA9* transcription factor that promotes the growth of murine hematopoietic progenitors and blocks their differentiation (45). This result might be related to a possible role of *HOX* genes in CRC right-sided tumor progression that deserves experimental exploration.

References

- Markowitz SD, Bertagnoli MM. Molecular origins of cancer: molecular basis of colorectal cancer. *N Engl J Med* 2009;361:2449–60.
- Ogino S, Goel A. Molecular classification and correlates in colorectal cancer. *J Mol Diagn* 2008;10:13–27.
- Iacopetta B. Are there two sides to colorectal cancer? *Int J Cancer* 2002;101:403–8.
- Buflin JA. Colorectal cancer: evidence for distinct genetic categories based on proximal or distal tumor location. *Ann Intern Med* 1990;113:779–88.
- Tan KK, Lopes Gde L Jr, Sim R. How uncommon are isolated lung metastases in colorectal cancer? A review from database of 754 patients over 4 years. *J Gastrointest Surg* 2009;13:642–8.
- Li FY, Lai MD. Colorectal cancer, one entity or three. *J Zhejiang Univ Sci B* 2009;10:219–29.
- Casillas S, Pelley RJ, Milsom JW. Adjuvant therapy for colorectal cancer: present and future perspectives. *Dis Colon Rectum* 1997;40:977–92.
- Hermann S, Rohrmann S, Linseisen J. Lifestyle factors, obesity and the risk of colorectal adenomas in EPIC-Heidelberg. *Cancer Causes Control* 2009;20:1397–408.
- Wei EK, Giovannucci E, Wu K, Rosner B, Fuchs CS, Willett WC, et al. Comparison of risk factors for colon and rectal cancer. *Int J Cancer* 2004;108:433–42.
- Terry P, Giovannucci E, Michels KB, Bergkvist L, Hansen H, Holmberg L, et al. Fruit, vegetables, dietary fiber, and risk of colorectal cancer. *J Natl Cancer Inst* 2001;93:525–33.
- Rozeck LS, Lipkin SM, Fearon ER, Hanash S, Giordano TJ, Greenon JK, et al. *CDX2* polymorphisms, RNA expression, and risk of colorectal cancer. *Cancer Res* 2005;65:5488–92.
- Fritzmann J, Morkel M, Besser D, Budczies J, Kosel F, Brembeck FH, et al. A colorectal cancer expression profile that includes transforming growth factor beta inhibitor *BAMBI* predicts metastatic potential. *Gastroenterology* 2009;137:165–75.
- Yamasaki M, Takemasa I, Komori T, Watanabe S, Sekimoto M, Doki Y, et al. The gene expression profile represents the molecular nature of liver metastasis in colorectal cancer. *Int J Oncol* 2007;30:129–38.
- Matsuyama T, Ishikawa T, Mogushi K, Yoshida T, Iida S, Uetake H, et al. *MUC12* mRNA expression is an independent marker of prognosis in stage II and stage III colorectal cancer. *Int J Cancer* 2010;127:2292–9.
- Salazar R, Roepman P, Capella G, Moreno V, Simon I, Dreezen C, et al. Gene expression signature to improve prognosis prediction of stage II and III colorectal cancer. *J Clin Oncol* 2011;29:17–24.
- Poynter JN, Gruber SB, Higgin PD, Almog R, Bonner JD, Rennert HS, et al. Statins and the risk of colorectal cancer. *N Engl J Med* 2005;352:2184–92.
- Vilar E, Bartnik CM, Stenzel SL, Raskin L, Ahn J, Moreno V, et al. *MRE11* deficiency increases sensitivity to poly(ADP-ribose) polymerase inhibition in microsatellite unstable colorectal cancers. *Cancer Res* 2011;71:2632–42.
- Barrett T, Edgar R. Gene expression omnibus: microarray data storage, submission, retrieval, and analysis. *Methods Enzymol* 2006;411:352–69.
- Parkinson H, Kapushesky M, Shojatalab M, Abeygunawardena N, Coulson RA, Holloway E, et al. ArrayExpress—a public database of microarray experiments and gene expression profiles. *Nucleic Acids Res* 2007;35 (Database issue):D747–50.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Grant Support

This study was supported by a grant (1R01CA81488) from the National Cancer Institute, the Catalan Institute of Oncology and the Private Foundation of the Biomedical Research Institute of Bellvitge (IDIBELL), the Instituto de Salud Carlos III (grants PI08-1635, PI08-1359, PS09-1037), CIBERESP CB06/02/2005 and the "Acción Transversal del Cancer," the Catalan Government DURSI grant 2009SGR1489, the European Commission grant FP7-COOP-Health-2007-B "HiPerDART," and the AECC (Spanish Association Against Cancer) Scientific Foundation.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received June 21, 2011; revised September 12, 2011; accepted September 13, 2011; published OnlineFirst October 5, 2011.

20. Jorissen RN, Gibbs P, Christie M, Prakash S, Lipton L, Desai J, et al. Metastasis-associated gene expression changes predict poor outcomes in patients with dukes stage B and C colorectal cancer. *Clin Cancer Res* 2009;15:7642–51.
21. Jorissen RN, Lipton L, Gibbs P, Chapman M, Desai J, Jones IT, et al. DNA copy-number alterations underlie gene expression differences between microsatellite stable and unstable colorectal cancers. *Clin Cancer Res* 2008;14:8061–9.
22. LaPointe LC, Dunne R, Brown GS, Worthley DL, Molloy PL, Wattochow D, et al. Map of differential transcript expression in the normal human large intestine. *Physiol Genomics* 2008;33:50–64.
23. Affymetrix, Inc. GeneChip expression analysis—data analysis fundamentals. Santa Clara, CA: Affymetrix, Inc. [cited 2002]. Available from: http://media.affymetrix.com/support/downloads/manuals/data_analysis_fundamentals_manual.pdf.
24. Irizarry RA, Hobbs B, Collin F, Beazer-Barclay YD, Antonellis KJ, Scherf U, et al. Exploration, normalization, and summaries of high density oligonucleotide array probe level data. *Biostatistics* 2003;4:249–64.
25. Kim H, Nam SW, Rhee H, Shan Li L, Ju Kang H, Hye Koh K, et al. Different gene expression profiles between microsatellite instability-high and microsatellite stable colorectal carcinomas. *Oncogene* 2004;23:6218–25.
26. Rozek LS, Herron CM, Greenson JK, Moreno V, Capella G, Rennert G, et al. Smoking, gender, and ethnicity predict somatic BRAF mutations in colorectal cancer. *Cancer Epidemiol Biomarkers Prev* 2010;19:838–43.
27. Subramanian A, Kuehn H, Gould J, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* 2005;102:15545–50.
28. Svngen T, Tonissen K. Hox transcription factors and their elusive mammalian gene targets. *Heredity* 2006;97:88–96.
29. Sugarbaker PH. Metastatic inefficiency: the scientific basis for resection of liver metastases from colorectal cancer. *J Surg Oncol Suppl* 1993;3:158–60.
30. Di Cataldo A, Scilletta B, Latino R, Cocuzza A, Li Destri G. The surgeon as a prognostic factor in the surgical treatment of rectal cancer. *Surg Oncol* 2007;16 Suppl 1:S53–6.
31. Aleman BM, Bartelink H, Gunderson LL. The current role of radiotherapy in colorectal cancer. *Eur J Cancer* 1995;31:1333–9.
32. Foroudi F, Tyldesley S, Barbera L, Huang J, Mackillop WJ. An evidence-based estimate of the appropriate radiotherapy utilization rate for colorectal cancer. *Int J Radiat Oncol Biol Phys* 2003;56:1295–307.
33. Dunican DS, McWilliam P, Tighe O, Parle-McDermott A, Croke DT. Gene expression differences between the microsatellite instability (MIN) and chromosomal instability (CIN) phenotypes in colorectal cancer revealed by high-density cDNA array hybridization. *Oncogene* 2002;21:3253–7.
34. Kakar S, Burgart LJ, Thibodeau SN, Rabe KG, Petersen GM, Goldberg RM, et al. Frequency of loss of hMLH1 expression in colorectal carcinoma increases with advancing age. *Cancer* 2003;97:1421–7.
35. Komuro K, Tada M, Tamoto E, Kawakami A, Matsunaga A, Teramoto K, et al. Right- and left-sided colorectal cancers display distinct expression profiles and the anatomical stratification allows a high accuracy prediction of lymph node metastasis. *J Surg Res* 2005;124:216–24.
36. Birkenkamp-Demtroder K, Olesen SH, Sorensen FB, Laurberg S, Laiho P, Aaltonen LA, et al. Differential gene expression in colon cancer of the caecum versus the sigmoid and rectosigmoid. *Gut* 2005;54:374–84.
37. Watanabe T, Kobunai T, Toda E, Yamamoto Y, Kanazawa T, Kazama Y, et al. Distal colorectal cancers with microsatellite instability (MSI) display distinct gene expression profiles that are different from proximal MSI cancers. *Cancer Res* 2006;66:9804–8.
38. Curtin K, Slattery ML, Samowitz WS. CpG island methylation in colorectal cancer: past, present and future. *Patholog Res Int* 2011;2011:902674.
39. Ferracin M, Gafà R, Miotto E, Veronese A, Pultrone C, Sabbioni S, et al. The methylator phenotype in microsatellite stable colorectal cancers is characterized by a distinct gene expression profile. *J Pathol* 2008;214:594–602.
40. Shah N, Sukumar S. The Hox genes and their roles in oncogenesis. *Nat Rev Cancer* 2010;10:361–71.
41. Samuel S, Naora H. Homeobox gene expression in cancer: insights from developmental regulation and deregulation. *Eur J Cancer* 2005;41:2428–37.
42. Kanai M, Hamada J, Takada M, Asano T, Murakawa K, Takahashi Y, et al. Aberrant expressions of HOX genes in colorectal and hepatocellular carcinomas. *Oncol Rep* 2010;23:843–51.
43. Freschi G, Taddei A, Bechi P, Faiella A, Gulisano M, Cillo C, et al. Expression of HOX homeobox genes in the adult human colonic mucosa (and colorectal cancer?). *Int J Mol Med* 2005;16:581–7.
44. Yahagi N, Kosaki R, Ito T, Mitsuhashi T, Shimada H, Tomita M, et al. Position-specific expression of Hox genes along the gastrointestinal tract. *Congenit Anom (Kyoto)* 2004;44:18–26.
45. Takeda A, Goolsby C, Yaseen NR. NUP98-HOXA9 induces long-term proliferation and blocks differentiation of primary human CD34+hematopoietic cells. *Cancer Res* 2006;66:6628–37.