

# Crizotinib in *MET*-Deregulated or *ROS1*-Rearranged Pretreated Non-Small Cell Lung Cancer (METROS): A Phase II, Prospective, Multicenter, Two-Arms Trial



Lorenza Landi<sup>1</sup>, Rita Chiari<sup>2</sup>, Marcello Tiseo<sup>3</sup>, Federica D'Inca<sup>4</sup>, Claudio Dazzi<sup>1</sup>, Antonio Chella<sup>5</sup>, Angelo Delmonte<sup>6</sup>, Laura Bonanno<sup>7</sup>, Diana Giannarelli<sup>8</sup>, Diego Luigi Cortinovis<sup>9</sup>, Filippo de Marinis<sup>10</sup>, Gloria Borra<sup>11</sup>, Alessandro Morabito<sup>12</sup>, Cesare Gridelli<sup>13</sup>, Domenico Galetta<sup>14</sup>, Fausto Barbieri<sup>15</sup>, Francesco Grossi<sup>16</sup>, Enrica Capelletto<sup>17</sup>, Gabriele Minuti<sup>1</sup>, Francesca Mazzoni<sup>18</sup>, Claudio Verusio<sup>19</sup>, Emilio Bria<sup>20,21</sup>, Greta Ali<sup>22</sup>, Rossella Bruno<sup>23</sup>, Agnese Proietti<sup>22</sup>, Gabriella Fontanini<sup>23</sup>, Lucio Crinò<sup>6</sup>, and Federico Cappuzzo<sup>1</sup>

## Abstract

**Purpose:** *MET*-deregulated NSCLC represents an urgent clinical need because of unfavorable prognosis and lack of specific therapies. Although recent studies have suggested a potential role for crizotinib in patients harboring *MET* amplification or exon 14 mutations, no conclusive data are currently available. This study aimed at investigating activity of crizotinib in patients harboring *MET* or *ROS1* alterations.

**Patients and Methods:** Patients with pretreated advanced NSCLC and evidence of *ROS1* rearrangements (cohort A) or *MET* deregulation (amplification, ratio *MET*/CEP7 >2.2 or *MET* exon 14 mutations, cohort B) were treated with crizotinib 250 mg twice daily orally. The coprimary endpoint was objective response rate in the two cohorts.

**Results:** From December 2014 to March 2017, 505 patients were screened and a total of 52 patients (26 patients

per cohort) were enrolled onto the study. At data cutoff of September 2017, in cohort A, objective response rate was 65%, and median progression-free survival and overall survival were 22.8 months [95% confidence interval (CI) 15.2–30.3] and not reached, respectively. In cohort B, objective response rate was 27%, median progression-free survival was 4.4 months (95% CI 3.0–5.8), and overall survival was 5.4 months (95% CI, 4.2–6.5). No difference in any clinical endpoint was observed between *MET*-amplified and exon 14–mutated patients. No response was observed among the 5 patients with cooccurrence of a second gene alteration. No unexpected toxicity was observed in both cohorts.

**Conclusions:** Crizotinib induces response in a fraction of *MET*-deregulated NSCLC. Additional studies and innovative therapies are urgently needed.

## Introduction

During the last 10 years several molecular events, including gene mutations, gene copy-number alterations, and gene rearrangements have been discovered in small fractions of lung adenocarcinomas, dramatically improving patient treatment (1).

This is the case of NSCLCs carrying *EGFR*-activating mutations or anaplastic lymphoma kinase (*ALK*) rearrangement, where targeted therapies have changed the natural history of the disease (2–5). Beyond *EGFR* mutations and *ALK* rearrangement, additional actionable alterations have been identified, with *ROS1*

<sup>1</sup>AUSL Romagna, Dipartimento di Oncologia ed Ematologia, Ravenna, Italy. <sup>2</sup>Ospedale Santa Maria della Misericordia, Perugia, Italy. <sup>3</sup>Dipartimento di Medicina e Chirurgia, Università degli Studi di Parma e Oncologia Medica, Azienda Ospedaliero-Universitaria di Parma, Parma, Italy. <sup>4</sup>Fondazione Ricerca Trasazionale, Roma, Italy. <sup>5</sup>Pneumologia Universitaria, Dipartimento Cardiotoracico Vascolare, Ospedale Cisanello, Pisa, Italy. <sup>6</sup>Istituto Scientifico Romagnolo per lo Studio e la Cura dei Tumori (IRST), IRCCS, Meldola, Italy. <sup>7</sup>Oncologia Medica 2, Istituto Oncologico Veneto, Padova, Italy. <sup>8</sup>IRCCS Regina Elena National Cancer Institute, Roma, Italy. <sup>9</sup>SC Oncologia Medica, SS Lung Unit Asst Ospedale San Gerardo, Monza, Italy. <sup>10</sup>Thoracic Oncology Division, Istituto Europeo di Oncologia IRCCS, Milan, Italy. <sup>11</sup>Ospedale Maggiore, Novara, Italy. <sup>12</sup>Istituto Nazionale Tumori, "Fondazione G.Pascale" IRCCS, Napoli, Italy. <sup>13</sup>Azienda Ospedaliera S.G. Moscati, Avellino, Italy. <sup>14</sup>Oncologia Medica Toracica, IRCCS Oncologico Giovanni Paolo II, Bari, Italy. <sup>15</sup>Azienda Ospedaliera-Universitaria Policlinico, Modena, Italy. <sup>16</sup>Division of Medical Oncology, Fondazione IRCCS Cà Granda Ospedale Maggiore Policlinico, Milano, Italy. <sup>17</sup>Dipartimento di Oncologia, Università di Torino, Torino, Italy. <sup>18</sup>SC Oncologia Medica, Azienda Ospedaliera Universitaria Careggi, Firenze, Italy. <sup>19</sup>ASST Valleolona, PO Saronno, Italy. <sup>20</sup>UO Oncologia, Azienda Ospedaliera Universitaria Integrata, Verona, Italy. <sup>21</sup>Fondazione Policlinico Universitario Agostino Gemelli IRCCS, Università Cattolica del Sacro Cuore, Roma, Italy. <sup>22</sup>Azienda Ospedaliero Universitaria Pisana, Pisa, Italy. <sup>23</sup>Dipartimento di Patologia Chirurgica, Medica, Molecolare e dell'Area Critica, Università di Pisa, Pisa, Italy.

**Note:** Supplementary data for this article are available at Clinical Cancer Research Online (<http://clincancerres.aacrjournals.org/>).

**Corresponding Author:** Federico Cappuzzo, Azienda AUSL della Romagna, Ravenna, Viale Randi 5, Ravenna 48121, Italy. Phone: 0039-0544-285206; Fax: 0039-0544-285217; E-mail: federico.cappuzzo@auslromagna.it

Clin Cancer Res 2019;25:7312–9

doi: 10.1158/1078-0432.CCR-19-0994

©2019 American Association for Cancer Research.

### Translational Relevance

Treatment of patients with *MET*-deregulated non-small cell lung cancer (NSCLC) represents an urgent need because of lack of effective targeted therapies and unfavorable prognosis. The METROS trial is a prospective study evaluating the efficacy of crizotinib in two cohorts of patients: individuals with *MET* exon 14 mutations or amplification or individuals with *ROS1* rearrangements. In the *MET*-deregulated cohort, although response rate was 27%, a remarkable result for a pretreated NSCLC population, median PFS and most importantly, median OS were disappointing with all patients rapidly progressing and dying. Interestingly, no difference for any clinical endpoint was observed between *MET*-amplified and exon 14-mutated patients. Overall, our data highlight the urgent need for more effective strategies in patients with *MET* mutations or amplification.

rearrangement and *MET* amplification or *MET* exon 14 mutations being the most appealing (5–8).

Crizotinib has been the first ALK inhibitor entered in clinical development within the PROFILE program and then approved worldwide for first-line treatment of *ALK*-positive NSCLC (3). Moreover, crizotinib has high specificity for *ROS1* and *MET* kinase domains and it potently inhibits tumor growth and invasion in cell models of *MET*- or *ROS1*-addicted NSCLC (9). In the phase I PROFILE 1001, enrolling 50 patients with *ROS1*-positive NSCLC, objective response rate (ORR) was 72% and median progression-free survival (PFS) exceeded 19 months (10). These results mirror those observed in *ALK*-positive NSCLC and are comparable with what was reported in subsequent retrospective and prospective trials (11–15). At present, crizotinib has a well-established role in *ROS1*-positive NSCLC and is available worldwide. Conversely, its role in *MET*-addicted NSCLC is not demonstrated. Initial observations in solid tumors with *MET* overexpression or *MET* amplification showed potential efficacy only in *de novo MET*-amplified tumors (16–18). In NSCLC, crizotinib produced an ORR of approximately 40%, with evidence of activity only against tumors with intermediate or high levels of *MET* amplification, defined as a ratio *MET*/centromere 7 (*MET*/CEP7) of  $> 2.2$ – $< 5$  or  $\geq 5$ , respectively (19). Interestingly, preliminary results of the phase II AcSé trial, evaluating crizotinib in *MET*-amplified NSCLC, showed an ORR of only 32% (20). Nevertheless, the criteria adopted for defining *MET* amplification were different than in the PROFILE 1001 study, providing a possible explanation for the lower drug efficacy. Moreover, AcSé data suggested that levels of *MET* amplification could be relevant for defining patients with the highest sensitivity to the drug. In addition to *MET* amplification, a recent study showed that anti-*MET* agents such as crizotinib or cabozantinib induced tumor shrinkage in patients harboring *MET* exon 14 mutations, a phenomenon occurring in approximately 3% of NSCLC (7, 8, 21, 22). In the phase I PROFILE 1001 study, ORR with crizotinib in *MET* exon 14 mutated patients was 44%, suggesting that the drug is effective against this alteration (23). On the basis of these premises and considering the urgent need of effective strategies for *MET*-deregulated NSCLC, we designed this study aiming at investigating crizotinib efficacy in *MET*-amplified or exon 14-mutated NSCLC. Because at the time of study design, few data were available in *ROS1*-rearranged patients and no

therapy was available in Italian clinical practice, a *ROS1*-rearranged cohort was also included.

## Patients and Methods

### Patients' selection

Eligible patients had histologically confirmed diagnosis of locally advanced or metastatic NSCLC and availability of archival tissue for biomarkers analyses. A local prescreening was allowed. For patients with *MET* exon 14 mutations, central confirmation was not required for trial inclusion, whereas central confirmation was mandatory for those patients who resulted positive for *ROS1* rearrangements or *MET* amplification at local labs. *ROS1* rearrangement and *MET* amplification were tested centrally by FISH, using the specific probes (Abbott Molecular). Briefly, criteria for FISH positivity were (i) presence of *ROS1* fusion patterns in  $\geq 15\%$  of tumor cells (24) or (ii) a *MET*/CEP7 ratio  $> 2.2$  according to Camidge criteria (19). *MET* mutational status was assessed locally using direct sequencing or other high sensitive methods. Even if central confirmation was not required for trial inclusion, at the end of the study, all *MET* exon 14-mutant cases were centrally verified using Sanger direct sequencing (Applied Biosystems—Thermo Fisher Scientific). Other inclusion criteria included: an Eastern Cooperative Oncology Group performance status (ECOG PS)  $\leq 2$ , at least one previous chemotherapy line, at least one measurable tumor lesion according to Response Evaluation Criteria in Solid Tumors (RECIST) version 1.1 (25), adequate bone marrow and organ functions. Patients with known *EGFR* or *KRAS* mutations or previously treated with *ROS1* or *MET* inhibitors or with symptomatic brain metastases were excluded. The study was done in accordance with the provisions of the Declaration of Helsinki and Good Clinical Practice guidelines. Each center received the approval of the local ethics committee, and all patients provided written informed consent before participation. The final version of the protocol including full list of study criteria is reported in the Supplementary Data.

### Treatment

Patients were treated with crizotinib 250 mg twice daily in continuous 28-day cycles until disease progression, unacceptable toxicity, withdrawal of consent, or death. Dose modifications or interruptions were considered in case of intolerable grade 2 or worse adverse events (AE). Radiologic assessment by CT scans was done at baseline and then every 8 weeks until disease progression; responses had to be confirmed by repeating assessment 4–8 weeks after initial response. All patients who discontinued study drug were followed up for subsequent treatments and survival every 12 weeks, until death or study completion. Patients were assessed for safety every 4 weeks. AEs, laboratory tests, and vital signs were graded according to the Common Terminology Criteria for Adverse Events version 4.0. The cut-off date for safety and efficacy data was September 30, 2017, which was the date of database lock.

### Outcomes

The primary endpoint was investigator-assessed overall response, defined as the percentage of patients who achieved a confirmed complete response (CR) or partial response (PR) per RECIST version 1.1 (25). Secondary endpoints included PFS based on investigator-assessed disease response OS, safety, and correlation between response and percentage of *ROS1* FISH positivity or levels of *MET* amplification [intermediate

levels, (MET/CEP7 ratio  $\geq 2.2$ – $<5$ ) vs. high levels, (MET/CEP7 ratio  $> 5$ )].

### Statistical analysis

The METROS was a phase II, two arms, noncomparative trial in which arms were determined by the presence of *ROS1* rearrangement or *MET* deregulation. The study was designed to test the hypothesis of an ORR  $\geq 50\%$  versus ORR  $\leq 10\%$  in each arm at a significance level of 5% (one sided) with a power of 98%. The study was originally designed to include only *MET*-amplified NSCLCs. However, clinical data published in 2015 suggested *MET* exon 14-mutated NSCLCs as an additional population potentially benefiting to crizotinib (8, 21). For such reason, the trial was amended to include also patients with such aberration without modification in the statistical plan. Patients and disease characteristics were analyzed using descriptive statistics and expressed as relative frequency (percentage) for discrete variables or median and interquartile range (IQR) for continuous variables. Associations among factors were evaluated with the  $\chi^2$  test while differences in distribution of quantitative variables were measured with the Mann–Whitney test. Confidence interval (95%) for ORR was calculated according to the exact method. PFS and OS were calculated from the date of starting therapy to the date of first evidence of either disease progression or death of the patient in the absence of documented disease progression (PFS), or death for any cause (OS). Patients without an event were censored at the date of last follow-up. Survival times were estimated using Kaplan–Meier analysis and expressed as medians with corresponding two-sided 95% confidence intervals (CI). Differences between curves were evaluated using the log-rank test. This trial is registered with ClinicalTrials.gov (NCT02499614).


## Results

### Patients

From December 2014 to March 2017, 505 patients were screened (Fig. 1). A total of 433 (86%) patients had tumor tissue

evaluable for *ROS1* and *MET* FISH analyses. Thirty-three individuals (7.6%) resulted *ROS1*-positive and 37 *MET*-deregulated (8.5%). Among them, 18 patients were not included due to death (3 *ROS1* and 4 *MET* patients), screening failure (4 *ROS1* and 4 *MET* patients), or unknown reason (3 *MET* patients). Twenty-six patients per cohort (cohort A, *ROS1* positive; cohort B, *MET* positive) accounted for the final population of the trial. Demographic and disease characteristics are reported in Table 1. Briefly, cohort A included mainly females, never smokers and with median age of 68 years, whereas cohort B included mainly males, current or past smokers with median age of 56 years. In both cohorts, most patients had an ECOG PS of 0–1, presented with two or more metastatic sites, and received crizotinib as second-line treatment. Fifty-four percent of patients in the *MET* cohort had progressive disease as best response to last therapy compared with less than 30% in *ROS1* cohort. A platinum-doublet regimen was the latest treatment before crizotinib in 69% and 81% of *ROS1* and *MET* deregulated patients, respectively (Supplementary Table S1). Among individuals included in the *MET* cohort, 16 patients had *MET* amplification (intermediate levels, 14 patients; high levels, 2 patients), 9 patients had exon 14 mutation (c.2962C>T, 1 case; c.3029C>T, 5 cases; c.3082G>T, 1 case; c.3082+1G>A, 1 case; c.3082+3A>T, 1 case), and one patient had cooccurrence of *MET* amplification (intermediate levels) and *MET* mutation (c.2942-19\_2961delinsC). At the end of the study, *MET* mutational status was centrally retested in all exon 14-mutant cases, with confirmation of local reports. Only one case resulted unconfirmed due to inadequate tumor sample for direct sequencing (data not shown). This patient had stable disease as best response to study drug.

Median number of crizotinib cycles and median duration of treatment were 15 (range, 0.3–34.4; IQR 5.4–24.9) and 15.2 months (95% CI, 4.7–25.2) in cohort A, and 3 (range, 0.4–28.6; IQR 2.0–5.6) and 4.0 months (95% CI, 2.0–5.5) in cohort B. At data cutoff, 9 patients in cohort A and 6 patients in cohort B were still receiving treatment.



**Figure 1.**  
Study profile.

**Table 1.** Baseline characteristics

	Cohort A ROS1 <sup>FISH+</sup>		Cohort B MET <sup>Ex14</sup> or MET <sup>FISH+</sup>		P <sup>a</sup>
	26	100%	26	100%	
Age, median (range)	68 (28–86)		56 (39–78)		0.07
M/F	10/16	38/62	17/9	65/35	0.05
ECOG PS 0/1/2	18/7/1	69/27/4	11/13/2	42/50/8	0.05
Never smoker/past smoker/current smoker	14/9/3	54/35/11	6/12/8	23/46/31	0.05
Adenocarcinoma/Other histology <sup>b</sup>	26/0	100/0	23/3	89/11	n.a.
Type of MET deregulation					
• Amplification	n.a.		16	61	
• Mutation	n.a.		9	35	
• Concurrent amplification and mutation	n.a.		1	4	
Metastatic sites, 1/2/>2	5/11/10	19/42/39	4/11/11	15/42/42	0.92
Disease sites					
Lung	22	85	22	85	0.99
Lymph node	16	61	12	46	0.40
Liver	3	11	6	23	0.46
Bone	9	35	5	19	0.35
Brain	6	23	5	19	0.73
Pleura	3	11	6	23	0.46
Adrenal gland	2	7	6	23	0.25
Prior line of therapy, 1/2/>2	20/3/3	76/12/12	21/3/2	81/11/8	0.63
Time from end of last treatment to crizotinib start (months, median)	12 (3–43)		3 (2–8)		0.02
Best response to prior therapy					
Complete response + partial response	10	38	4 <sup>c</sup>	15	0.12
Stable disease	7	27	6	23	
Progressive disease	7	27	14	54	
Unknown	2	8	2	8	

NOTE: Data are median (IQR) or *n* (%).

Abbreviation: n.a., not applicable.

<sup>a</sup>χ<sup>2</sup> test and Mann-Whitney were used for categorical items and for continuous variables, respectively. ECOG PS = Eastern Cooperative Oncology Group performance status.

<sup>b</sup>Other histologies includes two patients with NSCLC not otherwise specified (NOS) and one patient with pleomorphic carcinoma. All histology was determined by local pathologic report.

<sup>c</sup>Only partial responses.

**Efficacy**

Summary of efficacy measures is reported in Table 2. In cohort A, ORR was 65% (95% CI, 44–82), including one (4%) CR and 16 (61%) PR. Six patients (23%) obtained SD, resulting in an overall disease control rate (DCR) of 85%. With a median follow-up of 21 months (95% CI, 19.0–24.5), median PFS was 22.8 months (95% CI, 15.2–30.3), whereas median OS was not reached (Figs. 2

and 3). Median time to response (TTR) and median duration of response (DOR) were 7.9 weeks (IQR, 7.4–10.3) and 21.4 months (95% CI, 12.7–30.1). Depth of response, defined as the median percentage of reduction in target lesions from baseline, was –51.7% (IQR, –77.5% to –42.7%). In responding patients, median percentage of ROS1 FISH positivity was significantly higher than in nonresponders (50% vs. 22%, Mann-Whitney test *P* = 0.005; Supplementary Table S2).


**Table 2.** Efficacy endpoints in cohort A (ROS1 positive) and cohort B (MET deregulated)

	Cohort A-ROS1 ( <i>n</i> = 26)	Cohort B-MET ( <i>n</i> = 26)
Best overall response		
Complete response	1 (4%)	0
Partial response	16 (61%)	7 (27%)
Stable disease	6 (23%)	11 (42%)
Progressive disease	1 (4%)	6 (23%)
Not evaluable or not assessed	2 (8%)	2 (8%)
Progression-free survival (PFS)		
Number of events	14 (54%)	18 (69%)
PFS (months)	22.8 (15.2–30.3)	4.4 (3.0– 5.8)
6 months PFS rate	80.6%	30.9%
12 months PFS rate	71.9%	20.6%
Overall survival (OS)		
Number of events	10 (39%)	16 (61%)
OS (months)	NR	5.4 (4.2–6.5)
6 months OS rate	96.2%	43.9%
12 months OS rate	79.2%	26.3%

NOTE: Data are *n* (%) or median (95% CI), unless otherwise stated. NR, not reached.

In cohort B, ORR was 27% (95% CI, 11–47), including only PR. Eleven patients (42%) had SD, for an overall DCR of 69%. With a median follow-up of 21 months (95% CI, 19.0–24.5), median PFS and median OS were 4.4 months (95% CI, 3.0–5.8) and 5.4 months (95% CI, 4.2–6.5), respectively (Figs. 2 and 3). Median TTR and DOR were 7.4 weeks (IQR, 6.4–9.3) and 3.7 months (95% CI, 1.1–6.3). Depth of response was –47.9% (–56.5% to –35.6%). According to MET deregulation, responses occurred in 5 patients with MET amplification (intermediate levels only), in 1 patient with MET mutation, and in the coamplified and mutant case (Supplementary Table S3). Furthermore, we separately analyzed outcome in MET-amplified and in MET exon 14–mutated groups. In both groups, ORR, median DOR, PFS, and OS were similar to the whole MET-deregulated cohort (Supplementary Table S4).

Furthermore, to better characterize the study population, we retrospectively performed a Sequenom analysis in 48 of 52 tissue specimens collected at baseline. A second driver was found in 7 (14%) cases, including two cases with concomitant MET amplification (intermediate levels), three cases with MET exon 14



**Figure 2.** Tumor responses in *MET*-deregulated and *ROS1*-rearranged NSCLC. Maximum percentage reduction from baseline sum of lesion diameters by best investigator-assessed confirmed response in 52 patients receiving crizotinib as second-line or later treatment.


mutation, and two cases with *ROS1* rearrangement. Details are reported in Supplementary Tables S2 and S3. Among the 4 evaluable patients with *MET/KRAS* and *ROS1/KRAS* coaltered tumors, 1 achieved PR, 2 had SD, and 1 progressed. The double *MET*-amplified/*BRAF*-mutant subject progressed, whereas the *ROS1/MET*-positive patient voluntarily discontinued crizotinib after only 1 cycle without any tumor assessment.

Finally, as illustrated in Table 3, we evaluated the intracranial efficacy of the drug in the 11 patients with brain metastases at

baseline (six in cohort A and five in cohort B) and responses were observed only in *ROS1*-positive patients.

**Toxicity**

Safety profile of crizotinib was consistent with literature data and no new safety alert was reported in both cohorts (Supplementary Table S5). Treatment-related adverse events (TRAE), most of which were of grade 1 or 2, occurred in 26 (100%) patients in cohort A and in 21 (81%) patients of cohort B. In both



**Figure 3.** Kaplan-Meier curve of investigator-assessed PFS (A) and OS (B) in *ROS1*-positive (red line) and *MET*-deregulated (blue line) non-small cell lung cancer receiving crizotinib.

**Table 3.** Intracranial disease characteristics, intracranial response and pattern of failure in patients with brain metastases at baseline

Pt ID	Biomarker	Characteristics of CNS disease	Method for CNS Assessment	Prior RT	Date of RT	Date of start crizotinib	Intracranial response	Pattern of failure
MT-006-118	<i>ROS1</i>	Multiple lesions, nontarget	Brain MRI	No	NA	01 Dec 2015	CR	Extra- and intracranial
MT-001-013	<i>ROS1</i>	Single lesion, nontarget	Brain CT scan	No	NA	31 Mar 2015	SD	Intracranial only
MT-001-001	<i>ROS1</i>	Single lesion, nontarget	Brain CT scan	Yes	Jun 2012	27 Feb 2015	CR	Extra- and intracranial
MT-012-087	<i>ROS1</i>	Single lesion, nontarget	Brain CT scan	No	NA	27 Jan 2016	SD	Extracranial only
MT-001-011	<i>ROS1</i>	Single lesion, nontarget	Brain CT Scan	No	NA	20 Feb 2015	SD	Intracranial only
MT-019-238	<i>ROS1</i>	Multiple lesions, nontarget	Brain MRI	Yes	NR	25 Jul 2016	SD	Intracranial only
MT-006-079	<i>MET</i> <sup>FISH+</sup>	Single lesion, nontarget	Brain MRI	Yes	Aug 2015	10 Aug 2015	PD	Extra- and intracranial
MT-012-182	<i>MET</i> <sup>FISH+</sup>	Multiple lesions, target (1 lesion)	Brain CT scan	Yes	Apr 2016	26 Jul 2016	SD	Extracranial only
MT-004-286	<i>MET</i> <sup>Ex14</sup>	Multiple lesions, nontarget	Brain MRI	Yes	Jun 2016	14 Sep 2016	SD	NA <sup>§</sup>
MT-012-129	<i>MET</i> <sup>Ex14</sup>	Multiple lesions, target (1 lesion)	Brain CT scan	No	NA	22 Sep 2016	PD	Extra- and intracranial
MT-020-441	<i>MET</i> <sup>Ex14</sup>	Single lesion, nontarget	Brain CT scan	Yes	Jun 2017	05 Jul 2017	SD	Extracranial only

NOTE: *MET*<sup>FISH+</sup>, *MET* amplified (intermediate level only, *MET*/*CEP7* ratio 3.4 and 2.6 for MT-006-079 and MT-012-182, respectively); RT, radiotherapy; NR, not reported; NA, not applicable; § Pt ID MT-004-286 permanently discontinued crizotinib due to SAE.

cohorts, the most common TRAEs of any grade were fatigue (58% in cohort A and 31% in cohort B), peripheral edema (50% in cohort A and 31% in cohort B), nausea (46% in cohort A and 31% in cohort B), pain (30% in cohort A and 19% in cohort B), transaminases elevation (27% in both cohort A and cohort B), respiratory symptoms including dyspnea and cough (42% in cohort A and 46% in cohort B), and visual disorders (23% in cohort A and 27% in cohort B). In cohort A, TRAEs of grade 3/4 were peripheral edema, neutropenia, and respiratory symptoms each occurring in 1 patient (4%), and nausea and fatigue each occurring in 2 (8%) patients. In cohort B, TRAEs of grade 3/4 were nausea, neutropenia, anemia, and respiratory symptoms each occurring in one patient (4%), nausea and transaminases elevation occurring in 2 patients (8%). Overall, TRAEs leading to dose reduction, temporary or permanent discontinuation of the drug were reported in 8 (15%), 13 (25%), and 3 (6%) patients. Among 13 serious AEs (SAE) reported, only two were judged as related to study drug (Supplementary Table S6). Finally, we analyzed the incidence and clinical correlates of venous thromboembolism occurring prior or during crizotinib treatment in patients enrolled in the trial; the results of this analysis are the object of a separate publication (26).

## Discussion

Treatment of patients with *MET*-deregulated NSCLC represents an urgent clinical need because of lack of effective targeted therapies and unfavorable prognosis (7, 27). The METROS is a prospective study evaluating the efficacy of crizotinib in patients with *MET* exon 14 mutations or amplification. Although response rate was remarkable for a pretreated NSCLC population, median PFS and, most importantly, median OS were disappointing, with all patients rapidly progressing and dying.

In oncogene addicted NSCLCs, such as *EGFR*- or *ALK*-addicted NSCLC, targeted therapies are extending survival with medians ranging between 3–5 years (1–5). Similar outcome has been observed in *ROS1*-addicted patients and the results of our study, including also a *ROS1* cohort, confirmed that crizotinib is highly effective in such patients. The primary endpoint of ORR was met in *ROS1* cohort, where durable responses were observed in 65% of patients, median PFS exceeded 22 months and median OS was not reached, with approximately 80% of patients alive at 1 year. These data favorable compare with other trials, reinforcing the role of this agent in the treatment of *ROS1*-driven lung cancers (10–14). Interestingly, we observed a significant associ-

ation between percentage of *ROS1* FISH positivity and response to crizotinib, a phenomenon previously described only in *ALK*-positive NSCLC, deserving further investigations (28).

Different results were obtained in *MET*-deregulated patients. This cohort included both *MET*-amplified or exon 14–mutated. Although recent data suggest that these are different patient populations (19, 23), this concept did not emerge at the time of trial design and statistical hypothesis has been formulated considering *MET* deregulated as a homogeneous group. However, even with such limitation, outcome was similar in *MET*-amplified or exon 14–mutated subgroups, with limited responses and with only 1 month elapsing from time of tumor progression and patient death. These data are in agreement with other studies, such as the AcSé and the PROFILE 1001 (29, 30). Final results of the AcSé *MET* FISH–positive cohort showed an ORR of 32% and a median PFS of only 3.4 months, comparable with what was observed in our experience, even if criteria for *MET* positivity differed (29). In the last update of the PROFILE 1001, including a total of 37 *MET*-amplified patients, ORR was 27%, similar to the 31% observed in METROS. Importantly, among the 20 patients with high levels of amplification ORR was 40%, including 2 cases with CR (30). In our study, only one patient had high levels of amplification, precluding the possibility to explore the impact of the drug in presence of such characteristic. Nevertheless, high levels of *MET* amplification rarely occur in patients with NSCLC. In a previous study conducted in surgically resected NSCLC, among 435 screened patients, only 3 (0.6%) had high levels of amplification (27). METROS trial screened more than 430 advanced NSCLCs and only 0.4% displayed high levels of *MET* amplification, confirming the relative rarity of the event.

METROS study also included patients with *MET* exon 14 mutations, accounting for less than 3% of the screened population, as expected according to literature data (7, 8). In *MET* exon 14–mutated patients, the benefit produced by crizotinib in terms of ORR, PFS, and OS was limited. Although our findings seem inferior to what recently reported by Drillon and colleagues in PROFILE 1001, in which PFS exceed 7 months and OS is approximately 20 months, differences in patients selection limit comparison between the two studies (30). Exon 14 mutations include a wide range of abnormalities, such as insertion, deletion, or point mutation that generally lead the loss or attenuation of ubiquitin-mediated receptor degradation, for instance, by disrupting the splice acceptor site of intron 13 or affecting the splice donor site of intron 14 (8, 27, 32). How different mutations could affect

sensitivity to MET inhibitors, especially crizotinib, remains an unanswered question. In our cohort among the 4 patients with splicing mutations, only one responded.

METROS study also confirms the very unfavorable prognosis of *MET*-deregulated NSCLC. In 2009, our group first demonstrated that *MET* gene copy number was a negative prognostic factor in NSCLC (27). Additional studies confirmed that *MET* deregulation, including overexpression, gene copy number gain or mutation, confers an aggressive phenotype (33). In addition to an aggressive behavior, also reinforced by the scarce sensitivity to prior chemotherapy, *MET*-deregulated NSCLC demonstrated modest and transient responsiveness to crizotinib, suggesting that other factors could modulate sensitivity to *MET*-inhibition, such as cooccurrence of driver events or expression of the *MET* protein as recently reported (33–36). Particularly, in a context of *MET* amplification, Tong and colleagues demonstrated that low levels of amplification may occur in a background of *KRAS* mutation, while high levels of *MET* amplification were mutually exclusive with major oncogene drivers (33). In addition, it is not possible to exclude that other approaches or new drugs might be more effective. On this perspective, we are now conducting a phase II trial evaluating cabozantinib in both *MET*-amplified or mutated lung cancer untreated with *MET* inhibitor or refractory to crizotinib (CABinMET trial, EudraCT 2017-004157-16).

In conclusion, results of METROS trial indicate that even if crizotinib induces a tumor shrinkage in a fraction of *MET*-deregulated NSCLC, the drug minimally impacts the clinical course of the disease, at least in pretreated *MET*-mutated or *MET* with intermediate levels of amplification, whereas the efficacy of the drug in presence of high levels of amplification remains investigational. Additional studies and innovative therapies are urgently needed against this aggressive disease.

### Disclosure of Potential Conflicts of Interest

L. Landi reports receiving speakers bureau honoraria from Pfizer, AstraZeneca, and Bristol-Myers Squibb and is a consultant/advisory board member for Pfizer, AstraZeneca, Merck Sharp & Dohme, Boehringer Ingelheim, and Clovis. R. Chiari reports receiving speakers bureau honoraria from AstraZeneca, Takeda, Roche, Otsuka, Bristol-Myers Squibb, and Boehringer Ingelheim. M. Tiseo is a consultant/advisory board member for AstraZeneca, Bristol-Myers Squibb, Merck Sharp & Dohme, Boehringer Ingelheim, Takeda, Pfizer, Eli Lilly, Novartis, Roche, Otsuka, and Pierre Fabre. C. Gridelli reports receiving speakers bureau honoraria from and is a consultant/advisory board member for Merck Sharp & Dohme, Bristol-Myers Squibb, Roche, and AstraZeneca. D. Galetta reports receiving speakers bureau honoraria from Merck Sharp & Dohme, Bristol-

Myers Squibb, Roche, and Boehringer Ingelheim. F. Grossi reports receiving speakers bureau honoraria from Bristol-Myers Squibb, Merck Sharp & Dohme, Pfizer, Eli Lilly, AstraZeneca, Pierre Fabre, and Roche. E. Capelletto is a consultant/advisory board member for AstraZeneca. E. Bria reports receiving speakers bureau honoraria from AstraZeneca, Bristol-Myers Squibb, and Merck Sharp & Dohme, and is a consultant/advisory board member for Roche and Pfizer. F. Cappuzzo reports receiving speakers bureau honoraria from Roche, Pfizer, AstraZeneca, Bristol-Myers Squibb, Merck Sharp & Dohme, Takeda, and Lilly, and is a consultant/advisory board member for Roche, AstraZeneca, Takeda, Pfizer, Merck Sharp & Dohme, and Bristol-Myers Squibb. No potential conflicts of interest were disclosed by the other authors.

### Authors' Contributions

**Conception and design:** L. Landi, A. Chella, D. Galetta, G. Minuti, A. Proietti, F. Cappuzzo

**Development of methodology:** L. Landi, A. Chella, G. Minuti, G. Fontanini, F. Cappuzzo

**Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.):** L. Landi, R. Chiari, M. Tiseo, C. Dazzi, A. Chella, A. Delmonte, L. Bonanno, D.L. Cortinovia, F. de Marinis, G. Borra, A. Morabito, C. Gridelli, D. Galetta, F. Barbieri, F. Grossi, E. Capelletto, G. Minuti, F. Mazzoni, C. Verusio, E. Bria, G. Ali, R. Bruno, L. Crinò, F. Cappuzzo

**Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis):** L. Landi, R. Chiari, A. Chella, D. Giannarelli, G. Minuti, F. Cappuzzo

**Writing, review, and/or revision of the manuscript:** L. Landi, R. Chiari, M. Tiseo, F. D'Inca, C. Dazzi, A. Chella, A. Delmonte, L. Bonanno, D.L. Cortinovia, A. Morabito, C. Gridelli, F. Barbieri, F. Grossi, E. Capelletto, G. Minuti, F. Mazzoni, C. Verusio, E. Bria, L. Crinò, F. Cappuzzo

**Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases):** A. Chella, F. Grossi

**Study supervision:** L. Landi, A. Chella, A. Delmonte, D.L. Cortinovia, A. Proietti, F. Cappuzzo

### Acknowledgments

We thank the patients who participated in the METROS trial, their families and caregivers, the investigators, and site staff. Fondazione Ricerca Traslationale (FoRT) sponsored the trial, Pfizer supplied crizotinib, and Abbott Molecular supplied FISH probes for *ROS1* and *MET* analyses.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked advertisement in accordance with 18 U.S.C. section 1734 solely to indicate this fact.

Received March 26, 2019; revised June 13, 2019; accepted August 12, 2019; published first August 15, 2019.

### References

- Ferrara R, Mezquita L, Besse B. Progress in the management of advanced thoracic malignancies in 2017. *J Thorac Oncol* 2018;13:301–22.
- Soria J-C, Ohe Y, Vansteenkiste J, Reungwetwattana T, Chewaskulyong B, Lee KH, et al. Osimertinib in untreated EGFR-mutated advanced non-small-cell lung cancer. *N Engl J Med* 2018;378:113–25.
- Solomon BJ, Kim D-W, Wu Y-L, Nakagawa K, Mekhail T, Felip E, et al. Final overall survival analysis from a study comparing first-line crizotinib with chemotherapy: results from PROFILE 1014. *J Clin Oncol* 2018;36:2251–8.
- Peters S, Camidge DR, Shaw AT, Gadgeel S, Ahn JS, Kim D-W, et al. Alectinib versus crizotinib in untreated ALK-positive non-small-cell lung cancer. *N Engl J Med* 2017;377:829–38.
- Bergtholm K, Shaw AT, Ou S-HI, Katayama R, Lovly CM, McDonald NT, et al. *ROS1* rearrangements define a unique molecular class of lung cancers. *J Clin Oncol* 2012;30:863–70.
- Gainor JF, Tseng D, Yoda S, Dagogo-Jack I, Friboulet L, Lin JJ, et al. Patterns of metastatic spread and mechanisms of resistance to crizotinib in *ROS1*-positive non-small-cell lung cancer. *JCO Precis Oncol* 2017;1:1–13.
- Drilon A, Cappuzzo F, Ou S-HI, Camidge DR. Targeting *MET* in lung cancer: will expectations finally be met? *J Thorac Oncol* 2017;12:15–26.
- Frampton GM, Ali SM, Rosenzweig M, Chmielecki J, Lu X, Bauer TM, et al. Activation of *MET* via diverse exon 14 splicing alterations occurs in multiple tumor types and confers clinical sensitivity to *MET* inhibitors. *Cancer Discov* 2015;5:850–9.
- Pfizer, Inc. EU Xalkori (crizotinib) Summary of product characteristics. Available from: <https://www.ema.europa.eu>.
- Shaw AT, Ou S-HI, Bang Y-J, Camidge DR, Solomon BJ, Salgia R, et al. Crizotinib in *ROS1*-rearranged non-small-cell lung cancer. *N Engl J Med* 2014;371:1963–71.

11. Wu Y-L, Yang JC-H, Kim D-W, Lu S, Zhou J, Seto T, et al. Phase II study of crizotinib in east asian patients with ROS1-positive advanced non-small-cell lung cancer. *J Clin Oncol* 2018;36:1405–11.
12. Mazières J, Zalcman G, Crinò L, Biondani P, Barlesi F, Filleron T, et al. Crizotinib therapy for advanced lung adenocarcinoma and a ROS1 rearrangement: results from the EUROS1 cohort. *J Clin Oncol* 2015;33:992–9.
13. Moro-Sibilot D, Faivre L, Zalcman G, Pérol M, Barlesi F, Otto J, et al. Crizotinib in patients with advanced ROS1-rearranged non-small cell lung cancer (NSCLC). Preliminary results of the ACSé phase II trial. *JCO* 2015; 33:8065–8065.
14. Michels S, Gardizi M, Schmalz P, Thurat M, Pereira E, Sebastian M, et al. MA07.05 EUCROSS: a European Phase II trial of crizotinib in advanced adenocarcinoma of the lung harboring ROS1 rearrangements - preliminary results. *J Thor Oncol* 2017;12:S379–80.
15. Facchinetti F, Rossi G, Bria E, Soria J-C, Besse B, Minari R, et al. Oncogene addiction in non-small cell lung cancer: focus on ROS1 inhibition. *Cancer Treat Rev* 2017;55:83–95.
16. Ou S-HI, Kwak EL, Siwak-Tapp C, Dy J, Bergethon K, Clark JW, et al. Activity of crizotinib (PF02341066), a dual mesenchymal-epithelial transition (MET) and anaplastic lymphoma kinase (ALK) inhibitor, in a non-small cell lung cancer patient with de novo MET amplification. *J Thorac Oncol* 2011;6:942–6.
17. Lennerz JK, Kwak EL, Ackerman A, Michael M, Fox SB, Bergethon K, et al. MET amplification identifies a small and aggressive subgroup of esophagogastric adenocarcinoma with evidence of responsiveness to crizotinib. *J Clin Oncol* 2011;29:4803–10.
18. Chi AS, Batchelor TT, Kwak EL, Clark JW, Wang DL, Wilner KD, et al. Rapid radiographic and clinical improvement after treatment of a MET-amplified recurrent glioblastoma with a mesenchymal-epithelial transition inhibitor. *J Clin Oncol* 2012;30:e30–33.
19. Camidge DR, Ou S-HI, Shapiro G, Otterson GA, Villaruz LC, Villalona-Calero MA, et al. Efficacy and safety of crizotinib in patients with advanced c-MET-amplified non-small cell lung cancer (NSCLC). *J Clin Oncol* 2014; 32:8001.
20. Vassal G, Ledeley M, Tournigand C, Aparicio T, Ray-Coquard I, Taillandier L, et al. Activity of crizotinib in relapsed MET amplified malignancies: results of the french AcSé program. *JCO* 2015;33:2595–2595.
21. Paik PK, Drilon A, Fan P-D, Yu H, Rekhtman N, Ginsberg MS, et al. Response to MET inhibitors in patients with stage IV lung adenocarcinomas harboring MET mutations causing exon 14 skipping. *Cancer Discov* 2015;5:842–9.
22. Awad MM, Oxnard GR, Jackman DM, Savukoski DO, Hall D, Shivdasani P, et al. MET Exon 14 mutations in non-small-cell lung cancer are associated with advanced age and stage-dependent MET genomic amplification and c-Met overexpression. *J Clin Oncol* 2016;34:721–30.
23. Drilon AE, Camidge DR, Ou S-HI, Clark JW, Socinski MA, Weiss J, et al. Efficacy and safety of crizotinib in patients (pts) with advanced MET exon 14-altered non-small cell lung cancer (NSCLC). *JCO* 2016;34: 108–108.
24. Jurmeister P, Lenze D, Berg E, Mende S, Schäper F, Kellner U, et al. Parallel screening for ALK, MET and ROS1 alterations in non-small cell lung cancer with implications for daily routine testing. *Lung Cancer* 2015;87:122–9.
25. Eisenhauer EA, Therasse P, Bogaerts J, Schwartz LH, Sargent D, Ford R, et al. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur J Cancer* 2009;45:228–47.
26. Chiari R, Ricciuti B, Landi L, Morelli AM, et al. ROS1-rearranged non-small cell lung cancer is associated with high rate of venous thromboembolism: analysis from a phase II, prospective, multicentre, two arms trial (METROS). *Clin Lung Cancer* 2019; in press.
27. Cappuzzo F, Marchetti A, Skokan M, Rossi E, Gajopathy S, Felicioni L, et al. Increased MET gene copy number negatively affects survival of surgically resected non-small-cell lung cancer patients. *J Clin Oncol* 2009;27: 1667–74.
28. Soria J-C, Ho SN, Varella-Garcia M, Iafrate AJ, Solomon BJ, Shaw AT, et al. Correlation of extent of ALK FISH positivity and crizotinib efficacy in three prospective studies of ALK-positive patients with non-small-cell lung cancer. *Ann Oncol* 2018;29:1964–71.
29. Moro-Sibilot D, Cozic N, Pérol M, Otto J, Mazieres J, Souquet P, et al. OA12.03 activity of crizotinib in MET or ROS1 positive (+) NSCLC: results of the acsé trial. *J Thoracic Oncol* 2018;13:S348.
30. Camidge DR, Otterson GA, Clark JW, Ou S-HI, Weiss J, Ades S, et al. Crizotinib in patients (pts) with MET-amplified non-small cell lung cancer (NSCLC): Updated safety and efficacy findings from a phase 1 trial. *J Clin Oncol* 2018;36:9062.
31. Drilon A, Clark J, Weiss J, Ou S, Camidge DR, Solomon B, et al. OA12.02 updated antitumor activity of crizotinib in patients with MET exon 14-altered advanced non-small cell lung cancer. *J Thoracic Oncol* 2018;13: S348.
32. Tovar EA, Graveel CR. MET in human cancer: germline and somatic mutations. *Ann Transl Med* 2017;5:205.
33. Tong JH, Yeung SF, Chan AWH, Chung LY, Chau SL, Lung RWM, et al. MET amplification and exon 14 splice site mutation define unique molecular subgroups of non-small cell lung carcinoma with poor prognosis. *Clin Cancer Res* 2016;22:3048–56.
34. Bahcall M, Awad MM, Sholl LM, Wilson FH, Xu M, Wang S, et al. Amplification of wild-type KRAS imparts resistance to crizotinib in MET exon 14 mutant non-small cell lung cancer. *Clin Cancer Res* 2018;24: 5963–76.
35. Suzawa K, Offin M, Lu D, Kurzatkowski C, Vojnic M, Smith RS, et al. Activation of KRAS mediates resistance to targeted therapy in met exon 14-mutant non-small cell lung cancer. *Clin Cancer Res* 2019;25: 1248–60.
36. Guo R, Offin M, Brannon AR, Chow A, Delasos L, Somwar R, et al. MET inhibitor resistance in patients with MET exon 14-altered lung cancers. *J Clin Oncol* 37:15s, 2019 (suppl; abstr 9006).