

CONCISE REPORT

Diadenosine 5',5'''-*p*¹,*p*⁴-Tetraphosphate Deficiency in Blood Platelets of the Chédiak-Higashi Syndrome

By Byung K. Kim, Francis C. Chao, Randi Leavitt, Anthony S. Fauci, Kenneth M. Meyers, and Paul C. Zamecnik

Diadenosine tetraphosphate (AP₄A) is an unusual nucleotide found in a variety of cells, including platelets. It has been suggested that platelet AP₄A is stored in the dense granules and is metabolically inactive. We have studied the AP₄A content of blood platelets in two patients and three cattle with Chédiak-Higashi syndrome (CHS), a hereditary platelet defect with dense granule deficiency. Acid-soluble extractions of whole blood and platelets were neutralized. The adenosine triphosphate (ATP) level was measured by luminescence technique. To measure the AP₄A content, the neutralized extract was treated with phosphomonoesterase for removal of ATP. The AP₄A content was then measured by coupling the phosphodiesterase and luciferase reaction. The AP₄A content was 0.43 nmol/mg protein

for normal human platelets and 0.004 nmol/mg protein for CHS platelets. The ATP/AP₄A ratio was 67 for normal and 3,023 for CHS platelets. The whole blood AP₄A was reduced by 89% in CHS patients who had only a slight decrease in ATP level (26% reduction). Similarly, bovine platelets with CHS showed a marked decrease of AP₄A content and a moderate reduction of the ATP level. The platelet ATP/AP₄A ratio was 351 and 3,133 for normal and CHS cattle, respectively. Results demonstrate a marked reduction of AP₄A in CHS platelets and suggest that AP₄A may be a useful marker for the measurement of dense granule content in platelets.

© 1985 by Grune & Stratton, Inc.

DIADENOSINE 5',5'''-*p*¹,*p*⁴-tetraphosphate (AP₄A) was discovered in 1966 as a product of the back-reaction of amino acid activation.^{1,2} This compound has since been shown to be a ubiquitous component of living cells.³ Normal human platelets contain AP₄A at a concentration of 0.42 nmol/mg protein, with the ratio of adenosine triphosphate (ATP) to AP₄A at approximately 100.⁴ Some evidence suggests that platelet AP₄A may be stored in the dense granules. For example, incubation of platelets with ³H-adenosine resulted in no significant incorporation of radioactivity into AP₄A.⁴ Treatment of platelets with thrombin caused the complete release of AP₄A along with the release of the storage pool of other nucleotides.^{4,5} However, the precise subcellular localization of AP₄A in platelets has not been directly determined.

The Chédiak-Higashi syndrome (CHS) is a rare autosomal recessive congenital disorder characterized by a partial oculocutaneous albinism, an increased susceptibility to infection,^{6,7} and a bleeding tendency with the functional defect of the platelets^{8,9} showing a marked decrease of the storage pool of adenine nucleotides and serotonin.¹⁰⁻¹² This syndrome also occurs in animal species such as mice, mink, cats, and cattle.¹³⁻¹⁶ It has recently been reported that the AP₄A content in platelets is markedly decreased in animals with CHS.¹⁷ We report herein the results of the studies on the AP₄A content of platelets from two human patients and three cattle with CHS.

MATERIAL AND METHODS

Human blood was drawn from two CHS patients and four healthy volunteers and mixed with 0.1 vol of 1.5% EDTA solution. Informed consent was obtained prior to blood donation. Characterization of CHS patients and cattle has been described previously.^{9,16} Hemoglobin content of the blood was determined in a Coulter Counter (model S Plus, Coulter Electronics, Hialeah, Fla). To 0.1 mL of the whole blood was added 0.9 mL of phosphate-buffered saline (PBS) and then 1 mL of cold 7% trichloroacetic acid (TCA) solution. The samples were vortexed and kept on ice for 30 minutes with intermittent mixing. Platelet-poor plasma (PPP) was separated from the whole blood (1 mL) by centrifugation at 12,000 g for two minutes in a microfuge (Fisher model 59, Fisher Scientific Co, Pittsburgh, Pa) and mixed with an equal volume of 7% TCA. The remainder of the

whole blood was centrifuged at 220 g for 15 minutes to obtain platelet-rich plasma (PRP). Platelets were pelleted by centrifugation at 800 g for 15 minutes, washed twice with PBS containing 0.15% EDTA, resuspended in the same buffer, mixed with an equal volume of 7% TCA and incubated on ice for 30 minutes with intermittent mixing. Protein content of the washed platelet preparation was measured by the method of Lowry et al.¹⁸ Bovine blood was obtained from three CHS animals and five healthy animals. Bovine platelets were separated, washed, and then homogenized in a 5% cold TCA solution.

The TCA precipitates were centrifuged at 3,000 g for 30 minutes at 4°C. The supernatant was neutralized by extraction of TCA twice with 0.5 mol/L tri-*n*-octylamine in Freon.¹⁹ After vigorous mixing (by continuous vortexing for one minute), the reaction mixture was allowed to stand at room temperature for ten minutes to separate the aqueous (top layer) and amine-Freon phase. The aqueous phase containing nucleotides was used for determination of ATP and AP₄A content. Recovery of the trace amounts of ³H nucleotides added to whole blood and platelet preparations was greater than 99%. The measurement of ATP was performed by a luminescence technique²⁰ using the commercial reagent and procedure according to the manufacturer's instructions (Packard).

To assay AP₄A, the neutralized extract (100 μL) was first incubated with 3 units of purified phosphomonoesterase (specific activity, 1,000 U/mg protein, Calbiochem-Behring) for 30 minutes at 37°C. Between 98% to 99% of the ATP in the extract was degraded by this incubation step, following which the sample was kept in a 90°C water bath for five minutes. The AP₄A content was then determined by coupling the phosphodiesterase and luciferase reactions using a luminometer (model 6100 Picolite, Packard, Downers Grove, Ill).²⁰ An aliquot of the sample (10 μL) was mixed

From the Center for Blood Research, Boston; the National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Md; the Washington State University, Pullman; and the Worcester Foundation for Experimental Biology, Shrewsbury, Mass.

Submitted June 10, 1985; accepted June 25, 1985.

Supported by grants GM31562-02, P30-12708, HL-25547 and RR00515 from the National Institutes of Health.

Address reprint requests to Dr Byung K. Kim, The Center for Blood Research, 800 Huntington Ave, Boston, MA 02115.

© 1985 by Grune & Stratton, Inc.

0006-4971/85/6603-0041\$03.00/0

Table 1. ATP and AP₄A Contents in Whole Blood and Platelets From Subjects With Chédiak-Higashi Syndrome

Sample	Source	ATP	AP ₄ A	ATP/AP ₄ A
Human whole blood	Control (n = 4)	5.21 ± 0.29; μmol/g Hb	3.15 ± 1.12 nmol/g Hb	1,818 ± 625
	Chédiak-Higashi Syndrome (n = 2)	3.84; μmol/g Hb	0.35 nmol/g Hb	10,971
Human platelets	Control (n = 4)	28.5 ± 3.3 nmol/mg protein	0.43 ± 0.06 nmol/mg protein	67.0 ± 7.8
	Chédiak-Higashi Syndrome (n = 2)	13.3 nmol/mg protein	0.0044 nmol/mg protein	3,023
Bovine platelets	Control (n = 5)	23.6 ± 10.1 nmol/10 ⁹ platelets	0.078 ± 0.035 nmol/10 ⁹ platelets	351 ± 158
	Chédiak-Higashi Syndrome (n = 3)	16.3 ± 4.0 nmol/10 ⁹ platelets	0.006 ± 0.0036 nmol/10 ⁹ platelets	3,133 ± 1,026

with 40 μL of luciferin-luciferase reagent (Packard) and the counts were monitored every 30 seconds (integration mode). After stabilization of the luminescence background due to residual ATP, 0.01 unit of phosphodiesterase (specific activity, 52 U/mg protein, Calbiochem-Behring, San Diego, Calif) was added and the 30-second counting was continued until reaching a maximum luminescence, usually completed in two minutes. The net change in luminescence due to the addition of phosphodiesterase was used to calculate the AP₄A content based on internal standards.

RESULTS

Results of ATP and AP₄A measurements in whole blood and platelet preparations are summarized in Table 1. The mean AP₄A content in control human whole blood was 3.15 nmol/g hemoglobin (Hb) (range 1.86 to 4.39 nmol/g Hb) and 0.35 nmol/g Hb for two CHS patients (0.39 and 0.31 nmol/g Hb). There was an 89% reduction of blood AP₄A in these patients. In comparison, the ATP content in whole blood was only slightly decreased (26%) in the same patients (Table 1). The PPP from healthy volunteers contained less than 0.2% of the AP₄A and 0.01% of the ATP that were observed in the whole blood.

The average AP₄A content in human platelets was 0.43 nmol/mg protein, (range 0.353 to 0.485 nmol/mg protein) for four controls and 0.004 nmol/mg protein for two CHS patients (0.0037 and 0.005 nmol/mg protein). The ATP content in platelets was 28.5 and 13 nmol/mg protein for controls and patients, respectively. Thus, the ATP content in CHS platelets was reduced by 53% when compared to controls, whereas the decrease in AP₄A content was 99%. A similar pattern of a marked decrease in AP₄A content (92% reduction) and a modest reduction of ATP (31%) was observed in bovine platelets from three CHS cattle.

DISCUSSION

We demonstrate a marked reduction of AP₄A content in human and bovine platelets from subjects with CHS, which has the characteristics of a deficiency in the platelet dense granules.¹⁰⁻¹² The modest degree of reduction of ATP (31% to 53%) in the affected platelets, as shown herein, is consistent with a deficiency in the platelet dense granules in CHS because approximately 40% of the ATP in normal platelets is

stored in these granules.²¹ Our results of AP₄A measurement are in general agreement with the value observed by Flodgaard and Klenow in normal human platelets⁴ and with those reported by Flodgaard et al in bovine CHS platelets.¹⁷ The marked reduction (92% to 99%) of AP₄A in CHS platelets, as demonstrated in this study, together with previous results from other studies of the release and metabolism of AP₄A^{4,5} strongly support a subcellular localization of AP₄A in the platelet dense granules. It appears that AP₄A may be a useful marker for the measurement of dense granule content in blood platelets.

All the AP₄A in blood appears to be virtually cell bound since PPP contains only a trace amount of AP₄A. It should be mentioned, however, that 90% of ³²P-labeled AP₄A added to normal PPP is destroyed during a ten-minute incubation at 37 °C (P.C.Z., unpublished observations, November 1984). Whole blood AP₄A in CHS patients is reduced by 89%, whereas in normal controls platelet AP₄A accounts for up to 86% of the whole blood AP₄A content (62% and 86%, respectively). The latter results suggest that platelets are the major source of blood AP₄A. In preliminary experiments we found that human granulocytes from two normal volunteers contained 0.75 nmol/10⁹ cells of AP₄A (equivalent to 1% of the whole blood AP₄A) and 974 nmol/10⁹ cells of ATP, respectively.

The physiologic role of AP₄A in the blood platelet has not yet been defined. AP₄A has been shown to stimulate the initiation of DNA synthesis in permeabilized resting tissue culture cells²² and to bind with a subunit of DNA polymerase alpha.^{23,24} Although the platelets show a low rate of protein synthesis,^{25,26} normal human platelets contain only mitochondria and a trace amount of RNA, but no nucleus and no nuclear DNA.²⁷ Harrison et al²⁸ and more recently Luthje and Ogilvie²⁹ reported that AP₄A inhibited adenosine diphosphate (ADP)-induced platelet aggregation. We confirmed and expanded these observations.³⁰ In addition, Luthje and Ogilvie⁵ have shown that platelets also contain diadenosine 5',5''-p¹-p³ triphosphate (AP₃A), which is degraded to ADP by an enzyme in plasma and produces platelet aggregation.²⁹ It may be that AP₃A and AP₄A play some roles in the induction and dissolution of platelet aggregates in vivo.

REFERENCES

- Zamecnik PC, Stephenson ML, Janeway CM, Randerath K: Enzymatic synthesis of diadenosine tetraphosphate and diadenosine triphosphate with a purified lysyl-sRNA synthetase. *Biochem Biophys Res Commun* 24:91, 1966
- Randerath K, Janeway CM, Stephenson ML, Zamecnik PC: Isolation and characterization of dinucleoside tetra- and triphosphate formed in the presence of lysyl-sRNA synthetase. *Biochem Biophys Res Commun* 24:98, 1966

3. Zamecnik P, Stephenson ML: Nucleoside pyrophosphate compounds related to the first step in protein synthesis, in Kalckar HM, Klenow H, Munch-Petersen G, Ottesen M, Thaysen JH (eds): *The Role of Nucleotides for the Function and Conformation of Enzymes*, Proceedings of the Alfred Benzon Symposium 1, Copenhagen, 1969, p 276
4. Flodgaard H, Klenow H: Abundant amounts of diadenosine 5',5'''-p¹,p⁴-tetraphosphate are present and releasable but metabolically inactive in human platelets. *Biochem J* 208:737, 1982
5. Luthje J, Ogilvie A: The presence of diadenosine 5',5'''-p¹,p³-triphosphate (AP₃A) in human platelets. *Biochem Biophys Res Comm* 115:253, 1983
6. Blume RS, Wolff SM: The Chédiak-Higashi syndrome: Studies in four patients and a review of the literature. *Medicine (Baltimore)* 51:247, 1972
7. Klebanoff SJ, Clark RA: *The neutrophil: Function and clinical disorders*. Amsterdam, Elsevier, North Holland Biomedical, 1979, p 735
8. Bell TG, Meyers KM, Prieur DJ, Fauci AS, Wolff SM, Padgett GA: Decreased nucleotide and serotonin storage associated with defective function in Chédiak-Higashi syndrome cattle and human platelets. *Blood* 48:175, 1976
9. Costa JL, Fauci AS, Wolff SM: A platelet abnormality in the Chédiak-Higashi syndrome of man. *Blood* 48:517, 1976
10. Buchanan GR, Handin RI: Platelet function in the Chédiak-Higashi syndrome. *Blood* 47:941, 1976
11. Boxer GJ, Holmsen H, Robbin L, Bang N, Boxer LA, Bachner RL: Abnormal platelet function in Chédiak-Higashi syndrome. *Br J Haematol* 35:521, 1977
12. Rendu T, Breton-Gorius J, Lebreton M, Klebanoff C, Buriot D, Griscelli C, Levy-Toledano S, Caen JP: Evidence that abnormal platelet functions in human Chédiak-Higashi syndrome are the result of a lack of dense bodies. *Am J Pathol* 111:301, 1983
13. Lorez HP, Da Prada M: Fluorescence microscopical study of 5-hydroxy-tryptamine storage organelles in mepacrine-incubated blood platelets of beige mice (Chédiak-Higashi syndrome). *Experientia* 34:663, 1978
14. Meyers KM, Holmsen H, Seachord CL, Hopkins G, Gorham J: Characterization of platelets from normal mink and mink with the Chédiak-Higashi syndrome. *Am J Hematol* 7:137, 1979
15. Meyers KM, Seachord CL, Holmsen H, Prieur DJ: Evaluation of the platelet storage pool deficiency in the feline counterpart of the Chédiak-Higashi syndrome. *Am J Hematol* 11:241, 1981
16. Meyers KM, Holmsen H, Seachord CL, Hopkins GE, Borchard RE, Padgett GA: Storage pool deficiency in platelets from Chédiak-Higashi cattle. *Am J Physiol* 237:239, 1979
17. Flodgaard H, Zamecnik PC, Meyers K, Klenow H: AP₄A determinations as a possible tool for the diagnosis of Chédiak-Higashi disease and other platelet anomalies. *Hoppe Seylers Z Physiol Chem* 365:610, 1984 (abstr)
18. Lowry OH, Rosebrough N, Farr AL, Randall RJ: Protein measurement with Folin phenol reagent. *J Biol Chem* 193:265, 1951
19. Chen SC, Brown PR, Rosie DM: Extraction procedures for use prior to HPLC nucleotide analysis using microparticle chemically bonded packings. *J Chromatogr Sci* 15:218, 1977
20. Ogilvie A: Determination of diadenosine tetraphosphate (AP₄A) levels in subpicomole quantities by a phosphodiesterase luciferin-luciferase coupled assay: Applications as a specific assay for diadenosine tetraphosphatase. *Anal Biochem* 115:302, 1981
21. D'Souza L, Glueck HL: Measurement of nucleotide pools in platelets using high pressure liquid chromatography. *Thromb Haemost* 38:990, 1977
22. Grummt F: Diadenosine 5',5'''-p¹,p⁴-tetraphosphate triggers initiation of in vitro DNA replication in baby hamster kidney cells. *Proc Natl Acad Sci USA* 75:371, 1978
23. Grummt F, Waltl G, Jantzen HM, Hamprecht K, Hubscher U, Kuenzle CC: Diadenosine 5',5'''-p¹,p⁴-tetraphosphate, a ligand of the 57-kilodalton subunit of DNA polymerase alpha. *Proc Natl Acad Sci USA* 76:6081, 1979
24. Baril E, Bonin P, Burstein D, Mara K, Zamecnik P: Resolution of the diadenosine 5',5'''-p¹,p⁴-tetraphosphate binding subunit from a multiprotein form of HeLa cell DNA polymerase alpha. *Proc Natl Acad Sci USA* 80:4931, 1983
25. Warshaw AL, Laster L, Shulman NR: Protein synthesis by human platelets. *J Biol Chem* 242:2904, 1967
26. Booyse FM, Rafelson ME Jr: Stable messenger RNA in the synthesis of contractile protein in human platelets. *Biochim Biophys Acta* 145:188, 1967
27. Maupin B, Saint-Blancard J, Storck J: Platelet sulfure, taurine, proteins and ATP. *Rev Fr Etud Clin Biol* 7:169, 1962
28. Harrison MJ, Brossmer R, Goody RS: Inhibition of platelet aggregation and the platelet release reaction by alpha, omega diadenosine polyphosphates. *FEBS Lett* 54:57, 1975
29. Luthje J, Ogilvie A: Diadenosine triphosphate (AP₃A) mediates human platelet aggregation by liberation of ADP. *Biochem Biophys Res Comm* 118:704, 1984
30. Chao FC, Zamecnik P: Inhibition of platelet aggregation by AP₄A. *Hoppe Seylers Z Physiol Chem* 365:610, 1984 (abstr)