

# Discovery, Validation, and Application of Novel Methylated DNA Markers for Detection of Esophageal Cancer in Plasma



Yi Qin<sup>1</sup>, Chung W. Wu<sup>1</sup>, William R. Taylor<sup>1</sup>, Tarek Sawas<sup>1</sup>, Kelli N. Burger<sup>2</sup>, Douglas W. Mahoney<sup>2</sup>, Zhifu Sun<sup>2</sup>, Tracy C. Yab<sup>1</sup>, Graham P. Lidgard<sup>3</sup>, Hatim T. Allawi<sup>3</sup>, Navtej S. Buttar<sup>1</sup>, Thomas C. Smyrk<sup>1</sup>, Prasad G. Iyer<sup>1</sup>, David A. Katzka<sup>1</sup>, David A. Ahlquist<sup>1</sup>, and John B. Kisiel<sup>1</sup>

## Abstract

**Purpose:** The burden of esophageal cancer continues to rise, and noninvasive screening tools are needed. Methylated DNA markers (MDM) assayed from plasma show promise in detection of other cancers. For esophageal cancer detection, we aimed to discover and validate MDMs in tissue, and determine their feasibility when assayed from plasma.

**Experimental Design:** Whole-methylome sequencing was performed on DNA extracted from 37 tissues (28 EC; 9 normal esophagus) and 8 buffy coat samples. Top MDMs were validated by methylation specific PCR on tissue from 76 EC (41 adeno, 35 squamous cell) and 17 normal esophagus. Quantitative allele-specific real-time target and signal amplification was used to assay MDMs in plasma from 183 patients (85 EC, 98 controls). Recursive partitioning (rPART) identified MDM combinations predictive of esoph-

ageal cancer. Validation was performed *in silico* by bootstrapping.

**Results:** From discovery, 23 candidate MDMs were selected for independent tissue validation; median area under the receiver operating curve (AUC) for individual MDMs was 0.93. Among 12 MDMs advanced to plasma testing, rPART modeling selected a 5 MDM panel (*FER1L4*, *ZNF671*, *ST8SIA1*, *TBX15*, *ARHGEF4*) which achieved an AUC of 0.93 (95% CI, 0.89–0.96) on best-fit and 0.81 (95% CI, 0.75–0.88) on cross-validation. At 91% specificity, the panel detected 74% of esophageal cancer overall, and 43%, 64%, 77%, and 92% of stages I, II, III, and IV, respectively. Discrimination was not affected by age, sex, smoking, or body mass index.

**Conclusions:** Novel MDMs assayed from plasma detect esophageal cancer with moderate accuracy. Further optimization and clinical testing are warranted.

## Introduction

Esophageal cancer ranks as the 6th most common cause of cancer death worldwide, with 456,000 new cases and 400,000 deaths each year (1). Two main subtypes of esophageal cancer, esophageal adenocarcinoma (EAC) and esophageal squamous cell carcinoma (ESCC), are both increasing in incidence in different parts of the world (2).

Unfortunately, patients often present when symptomatic with advanced stage disease (3). Consequently, 5-year survival rates are less than 20% in developed countries and less than 5% in many developing countries (1, 3–5). Early-stage disease, however, is

associated with substantially improved survival of 80% to 90% at 5 years (6, 7). Therefore, presymptomatic early detection of esophageal cancer could change outcomes with this deadly disease (8).

Screening for esophageal cancer has yet to be implemented for the general population. Early detection by endoscopy may have a role in geographic regions with high prevalence, but is difficult to administer in most countries due to its invasiveness and high cost (9). Blood testing offers the appeal of noninvasiveness and ready distribution, but sufficiently accurate markers have yet to be developed.

Blood-based biomarkers for esophageal cancer have been investigated in small studies; these include circulating tumor cells, autoantibodies, microRNA, and aberrantly methylated DNA (10–14). Acquired aberrant covalent DNA methylation is an epigenetic modification, which promotes oncogenesis by altered gene expression. Hypermethylation of CpG islands in promoter regions can silence tumor suppressor genes, whereas hypomethylation of repetitive gene elements may lead to genomic instability and activation of oncogenes (15). DNA methylation events appear to be less heterogeneous than mutations; for example, studies in primary tumor tissues show that as few as 4 methylated DNA markers (MDM) achieve nearly perfect discrimination between GI neoplasms and controls (16–22). Furthermore, technological advances in assay chemistry have

<sup>1</sup>Gastroenterology and Hepatology, Mayo Clinic, Rochester, Minnesota.

<sup>2</sup>Biomedical Statistics and Informatics, Mayo Clinic, Rochester, Minnesota.

<sup>3</sup>Exact Sciences Development Company, Madison, Wisconsin.

**Note:** Supplementary data for this article are available at Clinical Cancer Research Online (<http://clincancerres.aacrjournals.org/>).

**Corresponding Author:** John B. Kisiel, Mayo Clinic, 200 First Street SW, Rochester, MN 55902. Phone: 507-284-0959; Fax: 507-284-0538; E-mail: [kisiel.john@mayo.edu](mailto:kisiel.john@mayo.edu)

Clin Cancer Res 2019;25:7396–404

doi: 10.1158/1078-0432.CCR-19-0740

©2019 American Association for Cancer Research.

### Translational Relevance

Esophageal cancer is a lethal disease for which no effective screening tool exists at a population level. An inexpensive, automatable, and noninvasive screening tool could transform the future of this disease. Ours was the first whole-methylome sequencing of both esophageal adenocarcinoma and squamous cell carcinoma. Through our discovery in tissue, differentially methylated regions highly discriminant for both types of cancer were identified. These subsequently underwent robust rounds of validation in independent tissue samples. Ultimately, plasma assays for candidate methylated DNA markers were developed; a 5-marker panel achieved an AUC of 0.81 on cross-validation, and 0.93 on best-fit for discrimination of esophageal cancer from normal controls, when assayed from plasma. In addition, marker sensitivity in plasma was directly related to cancer stage, suggesting potential for prognosis and surveillance. These data suggest feasibility of a noninvasive test for detection of esophageal cancer and open the door for further optimization and clinical investigation.

dramatically improved the analytical sensitivity for MDMs in plasma to allow detection of low copy numbers associated with early stage cancer, notably of the liver and stomach (19, 23).

Our group and others have systematically applied next-generation sequencing techniques to expand the list of candidate MDMs of gastrointestinal cancers and demonstrated feasibility for plasma testing in several phase I and II clinical studies (19, 23). Additionally, novel assay chemistry, specifically quantitative allele-specific real-time target and signal

amplification assay (QuARTS), achieves very high analytical sensitivity for circulating cell-free DNA, but has not yet been applied to esophageal cancer (20, 23).


We hypothesized that: (i) next-generation DNA sequencing would identify novel and highly discriminant MDMs of esophageal cancer (EAC and ESCC); (ii) MDMs would be confirmed in independent tissues; and (iii) MDMs would show clinical feasibility for detection of esophageal cancer when assayed from plasma-extracted DNA.

### Materials and Methods

#### Overview

The investigative design comprised four sequential case-control studies (Fig. 1). First, we performed unbiased whole methylome sequencing by reduced representation bisulfite sequencing (RRBS) on DNA extracted from frozen primary tumor and control tissues to identify novel differentially methylated regions (DMR) of the genome, followed by technical validation using quantitative methylation specific PCR (qMSP) on the same frozen samples. Technically validated DMRs were then biologically validated on independent formalin-fixed paraffin-embedded (FFPE) tissue samples. DMRs showing highest discrimination by *a priori* criteria were carried forward as candidate MDMs for plasma testing. Finally, QuARTS was used to assay DNA extracted from archival plasma samples from esophageal cancer case and control patients. For each sample, QuARTS products were normalized by plasma volume and  $\beta$ -actin (*ACTB*).

To distinguish DMRs of genomic DNA extracted from tissues (DMRs) from circulating fragments of methylated DNA used as tumor markers, we refer to the latter as MDM. All laboratory



**Figure 1.**  
Study flow diagram.

testing was performed by personnel blinded to clinical data. For the phase I clinical study, all clinical data were reviewed by a single clinician (YQ) prior to un-blinding of MDM assay results by the lead statistician (DWM).

### Patients and samples

All study procedures were conducted after approval from the Mayo Clinic Institutional Review Board. Studies were conducted in accordance with ethical principles from the Declaration of Helsinki. Tissues used in RRBS and technical validation steps were obtained from the Esophageal Adenocarcinoma and Barrett's Esophagus registry (EABE), which enrolled patients with EAC, ESCC, and healthy controls, under informed consent between December 2010 and September 2015. Esophageal tumor tissue was sampled in treatment-naïve patients prior to endoscopic therapy, surgery, radiation, or chemotherapy. Healthy control tissues were from individuals free from any prior esophageal neoplasm or Barrett's esophagus. De-identified buffy coat samples were also sequenced to control for and then exclude regions of leukocyte DNA with elevated methylation, due to their potential to cause high background in plasma (24). FFPE tissues used for biological validation were obtained from the Mayo Clinic Tissue Registry, an archive of clinical tissue specimens maintained by the Mayo Clinic Department of Anatomic Pathology. All frozen and FFPE tissues underwent research histopathology review by an expert gastrointestinal pathologist (TCS) prior to macro-dissection and DNA extraction.

For the plasma study, frozen archives of the EABE Registry provided EDTA-buffered plasma samples from patients with esophageal cancer. A second archive of patients free from cancer was enrolled from a 7-county regional population between December 2015 and July 2017. From these, we selected plasma samples from age- and sex-matched control patients.

All plasma samples were collected in EDTA buffer and processed by the Mayo Clinic Biospecimens Accession and Processing laboratory. Referent to the time of plasma collection, cases were required to be free from exposure to local/regional therapy or systemic chemotherapy, and free from any primary cancer outside of the esophagus, excluding nonmelanoma skin cancers; controls were required to be free from any cancer within 5 years of sample collection. Staging referent to the time of plasma sample collection was determined using the 8th edition of the American Joint Committee on Cancer (AJCC) classifications (25).

### Laboratory methods and assays

**Discovery and technical validation.** DNA was isolated from macro-dissected frozen tissue sections using the DNeasy 96 blood and Tissue Kit (Qiagen). DNA from buffy coat samples was purified using the QIAamp DNA Mini Kit (Qiagen). DNA yield was measured fluorometrically on a plate reader with the Quant-iT Pico-Green dsDNA assay reagents (Invitrogen). Samples were bisulfite-converted using the Zymo EZ-96 DNA Methylation Kit (Zymo Research) and amplified with SYBR Green I detection using the LightCycler 480 (Roche Diagnostics). RRBS was performed as described previously (19).

Sequencing results were processed by Illumina pipeline modules for image analysis and base calling. Secondary analysis was performed using SAAP-RRBS (26). Briefly, reads were cleaned-up using Trim-Galore (Babraham Institute) and aligned to the

GRCh37/hg19 reference genome build with whole genome bisulfite sequence mapping program (BSMAP; ref. 27). Methylation ratios were determined by calculating  $C/(C+T)$  or conversely,  $G/(G+A)$  for reads mapping to the reverse strand, for CpGs with coverage  $\geq 10\times$  and base quality score  $\geq 20$ .

For technical validation, qMSP primers for candidate DMRs were designed with Methprimer software (University of California) as described previously (26). Oligonucleotides were synthesized by IDT. Prior to use, qMSP assays were quality-control tested on bisulfite converted and unconverted methylation ( $\pm$ ) controls to verify amplification. Assay standards were dilutions of bisulfite converted and SssI treated genomic DNA, which ensured absolute quantification. Results were expressed as fractional methylation using the bisulfite-treated PCR product of *ACTB* as a marker for total DNA in each sample, and analyzed logistically.

**Biological validation in tissue.** The same qMSP assays then were run on DNA extracted from independent FFPE case and control tissues. DNA was purified using the QIAamp FFPE Tissue Kit (Qiagen), bisulfite treated with the Zymo Kit, and amplified with SYBR Green I detection using the LightCycler 480 as above.

**Plasma study.** DNA from independent patients with esophageal cancer and cancer-free controls was purified and bisulfite converted from 3 to 4 mL of plasma using a proprietary semiautomated silica bead DNA extraction method developed to maximize analyte recovery and sample throughput (20). The eluted DNA was subsequently treated with ammonium bisulfite and re-purified to remove polymerase inhibitors. The DMR sequences for 12 MDM candidates were used to construct modified QuARTS triplex assays, as described previously (21). This technology platform was selected for its high analytical sensitivity, which is at least 10-fold higher than that of conventional qMSP. Triplexes were assayed on the LightCycler 480 (Roche) and all results were normalized by plasma volume and by *ACTB* amplified from each sample.

### Statistical analysis

**Discovery.** To identify candidate DMRs from the RRBS data set, the percentage of methylation was compared between cases, tissue controls, and buffy coat controls at each mapped CpG. For this comparison, a tiled reading frame within 100 base pairs of each CpG was used to identify DMRs of interest, specifically areas where methylation in controls was  $< 5\%$ . Regions were analyzed only if the variance of methylation percentage across subgroups was  $> 0$  and the total depth of coverage per sample group was  $\geq 200$  reads (10 reads per subject on average). The sample size requirements were estimated as reported previously (19). In brief, the highest background for methylation in controls was assumed to be 5%, and a  $> 3\times$  increase in the odds ratio was considered biologically relevant. At a coverage depth of 10 reads,  $\geq 18$  samples per group were required to achieve 80% power assuming binomial variance inflation factor of 1, with a 2-sided test at a significance level of 5%. To account for loss of power in case of greater variance inflation, we further restricted sites with a minimum read depth of 20. Following variance-inflated logistic regression, candidate DMRs were ranked according to *P*-value, area under the receiver operating characteristics curve (AUC),

methylation fold-change ratio (FC), and absolute methylation difference between cases and controls ( $\Delta M$ ).

**Biological validation.** MDM candidates were first selected from those with AUCs and FC values within the top 40% of those that advanced through technical validation. In addition, MDMs that detected additional tumor samples not represented by the top 40%, or showing high discrimination for other gastrointestinal cancers, were also included (16–20, 23).

For the confirmation of each MDM in independent FFPE samples, those with a lower 95% CI bound for the estimated AUC exceeding 0.7 were deemed to have acceptable accuracy. Assuming an estimated AUC of 0.8, to achieve this bound of acceptance at a one-sided significance level of 0.05, a minimum of 35 cases per cancer subtype were required with a case: control mix of 2:1.

**Plasma study.** Marker selection for the plasma study was limited to 12 candidates which were representative of both EAC and ESCC, had the lowest amplification of leukocyte DNA, and could be assayed from the shortest (<100bp) amplicon size. Recursive partitioning (rPart) was used to develop a prediction model for cancer based on the QuARTS assay products of individual MDM and combinations of multiple MDMs. AUCs for individual MDMs and the predicted probability of cancer from the fitted rPart tree were used to assess accuracy. It was estimated that a minimum of 97 patients in the case group and 97 in the control group would provide 80% power to distinguish an estimated AUC of  $\geq 0.8$  from a null value of 0.7 with a 2-sided test of significance set at the 5% level.

*In silico* cross-validation was performed by generating 500 bootstrapped subsets (with a training: testing ratio of 2:1) of the original data. The reproducibility of the entire panel of MDMs was estimated by averaging sensitivity, specificity, and AUC across the 500 simulated datasets. The potential influence of clinical subtypes on prediction accuracy was assessed by comparing stratified AUCs. All tests were performed at the 5% level of significance.

To measure the biological relevance of MDMs brought into the plasma study, expectation maximization values of RNA sequencing data were obtained from publicly available data sets of EAC primary tumor samples (The Cancer Genome Atlas, TCGA) and normal esophagus tissues (Genotype Tissue Expression Project) using the XENA browser and the DESeq2 analysis (28, 29).

In order to evaluate the specificity of the MDMs for esophageal cancer in comparison to other common cancers, we downloaded DNA methylation data from TCGA from Genomic Data Commons (GDC) portal (<https://portal.gdc.cancer.gov/>, June 2018; refs. 30–34), and extracted data for cancers from the esophagus, colon/rectum, liver, and lung. Methylation data in TCGA was derived from tissue based studies using Infinium Human Methylation 450 K BeadChip containing 450 CpG sites. EAC and ESCC were evaluated both separately and combined. Because of the platform differences, we mapped the CpG sites from the 450 k array to the genomic coordinates of our selected MDMs, and those with at least one CpG site mapped were kept for evaluation (average methylation was obtained if more than one CpG site were mapped in a region). Tumor tissue was used as site-specific cases, and tumor-adjacent normal tissue was used as site-specific controls. Methylation

was measured as an average  $\beta$ -value, which is equivalent to methylation ratio from bisulfite sequencing. Box plots of case versus control were generated for each cancer type. One-way ANOVA along with TukeyHSD was used to determine overall and pair-wise methylation differences among different cancer types for each MDM; mean methylation difference and adjusted *P* values for multiple testing were obtained accordingly.

## Results

### Discovery and technical validation

DNA from frozen esophageal tissue was available from 28 esophageal cancer cases (18 EAC, 10 ESCC) and 17 normal controls (9 normal esophagus without synchronous Barrett's esophagus or esophageal cancer and 8 normal buffy coat samples from patients without history of cancer; Fig. 1). Sequencing yielded on average 25 million reads that mapped to the GRCh37/hg19 genome assembly with 3 million CpGs having  $\geq 10\times$  coverage. Of these, approximately one million occurred in regions designated as CpG islands. Prespecified filtering criteria for candidate DMRs included AUC  $\geq 0.90$ , FC  $\geq 50$ ,  $\Delta M \geq 0.15$ , *P*-value  $\leq 0.01$ , CpG density of 0.05 to 0.3, and a minimum range of 40 base pairs. Our software identified  $\sim 185,000$  EAC and  $\sim 99,000$  ESCC DMRs with a variance  $>0$ , DMRs were reduced to 94 (EAC) and 88 (ESCC) by these criteria.

Thirty-four EAC and 34 ESCC assays were selected for technical validation; qMSP testing on the sequenced cases and controls resulted in AUCs which ranged from 0.60 to 0.99 (median 0.88), FCs of 4 to  $>1,000$  (median 26), and  $\Delta M$  of 0.0082 to 0.54 (median 0.19). We also calculated intraclass correlation to assess the reliability of the MDM results. Coefficients ranged from 0.00016 to 0.98 (median 0.95) indicating high reproducibility for the majority of markers.

Thirteen MDMs hypermethylated in both EAC and ESCC (*ARHGEF4*, *Chr10.102*, *Chr19.402*, *DLX4*, *DMRTA2*, *GRIN2D*, *KCNA3*, *TBX15*, *TSPYL5*, *ZNF132*, *ZNF610*, *ZNF671*, *ZNF781*), 5 EAC specific MDMs (*Chr9.999*, *ELMO1*, *FGF14*, *ST8SIA1*, and *ZNF568*), and 5 MDMs that were recurrently methylated in other cancers (*BMP3*, *FER1L4*, *IKZF1*, *NDRG4*, *OPLAH*) were chosen for biological validation (16–20, 23).

### Biological validation in tissue

The independent tissue cohort included 41 patients with EAC, 35 patients with ESCC, and 17 normal controls without esophageal cancer or Barrett's esophagus (Fig. 1). Groups were age- and sex-balanced. Buffy coat samples from 24 patients free from cancer were also run as controls.

Discrimination by candidate MDMs on independent tissue validation is summarized (Table 1). AUCs for the independent EAC tissue comparison against normal esophageal mucosa were 0.44 to 0.99 (median 0.93); and 0.33 to 1.00 (median 0.87) for ESCC tissue. AUCs for EAC compared with buffy coat were 0.85 to 1.00 (median 0.97); AUCs for ESCC compared with buffy coat were 0.80 to 0.99 (median 0.91). Median FCs were 54 for EAC compared with normal esophagus, 10 for ESCC compared with normal, 811 for EAC compared with buffy, and 454 for ESCC compared with buffy.

We selected 12 MDMs (*ARHGEF4*, *ELMO1*, *ST8SIA1*, *OPLAH*, *FER1L4*, *TBX15*, *ZNF671*, *IKZF1*, *TSPYL5*, *NDRG4*, *BMP3*, and

**Table 1.** Discrimination of MDMs for esophageal cancer during biological validation in DNA extracted from tissues

MDM	AUC (EAC)	P value	AUC (ESCC)	P value
<i>ELMO1</i> <sup>a</sup>	0.99	<0.0001	0.74	0.0003
<i>FGF14</i>	0.93	<0.0001	0.45	0.0223
<i>Chr9.999</i>	0.83	<0.0001	0.51	0.0102
<i>ST8SIA1</i> <sup>a</sup>	0.98	<0.0001	0.59	0.0003
<i>ZNF568</i>	0.91	0.1588	0.75	0.8952
<i>ARHGEF4</i> <sup>a</sup>	0.79	0.0001	0.81	<0.0001
<i>DLX4</i>	0.75	0.0389	0.70	0.5563
<i>DMRTA2</i> <sup>a</sup>	1.00	<0.0001	1.00	<0.0001
<i>GRIN2D</i>	0.99	<0.0001	0.88	<0.0001
<i>KCNA3</i>	0.92	<0.0001	0.60	0.0288
<i>Chr10.102</i>	0.44	0.8974	0.33	0.7419
<i>Chr19.403</i>	0.66	0.0124	0.61	0.0154
<i>TBX15</i> <sup>a</sup>	0.93	<0.0001	0.91	<0.0001
<i>TSPYL5</i> <sup>a</sup>	0.95	<0.0001	0.90	<0.0001
<i>ZNF132</i>	0.88	<0.0001	0.75	<0.0001
<i>ZNF610</i>	0.97	<0.0001	0.87	<0.0001
<i>ZNF671</i> <sup>a</sup>	0.89	<0.0001	0.89	<0.0001
<i>ZNF781</i>	0.99	<0.0001	0.94	<0.0001
<i>OPLAH</i> <sup>a</sup>	0.94	<0.0001	0.77	0.0004
<i>IKZF1</i> <sup>a</sup>	0.92	<0.0001	0.37	0.0086
<i>FER1L4</i> <sup>a</sup>	0.92	<0.0001	0.69	0.0002
<i>NDRG4</i> <sup>a</sup>	0.96	<0.0001	0.54	0.0537
<i>BMP3</i> <sup>a</sup>	0.96	<0.0001	0.50	0.0181

<sup>a</sup>Markers advanced to further testing in plasma study.  
 AUC (EAC) Tissue: Median AUC for discrimination of esophageal adenocarcinoma from normal esophagus.  
 AUC (ESCC) Tissue: Median AUC for discrimination between esophageal squamous cell carcinoma from normal esophagus.

*DMRTA2*) for the plasma study. Gene expression analysis of publicly available RNA sequencing data in EAC and normal esophagus show significant up- or downregulation for 9 of 11 MDMs which annotate to coding regions (Supplementary Table S1). Oligonucleotides were redesigned in a modified QuARTS format to hybridize to the exact regions addressed with the qMSP primers.

**Plasma study**

Plasma samples from 85 independent and unique esophageal cancer cases and 98 normal controls without cancer or Barrett's esophagus were tested. Patient characteristics are summarized in Table 2. Clinical TNM staging algorithm was used according to AJCC 8th edition classification.

Cases included 76 EAC, 9 ESCC, and represented all stages: I (*n* = 14), II (*n* = 14), III (*n* = 31), and IV (*n* = 26). Cases and controls were well-balanced for age, sex, and BMI, but there was a significantly higher number of former smokers among case patients (*P* < 0.001).

After normalization to the *ACTB* internal control, individual MDMs had an AUC of 0.64 to 0.78. *ELMO1* and *NDRG4* were 100% specific (Fig. 2). Plots of MDM level distributions show substantial separation between cases and controls for most individual candidates (Supplementary Fig. S1).

The rPART modeling process selected a 5-marker combination (*FER1L4*, *ZNF671*, *ST8SIA1*, *TBX15*, *ARHGEF4*) normalized by *ACTB* as the most discriminant panel. At a specificity cut-off of 91% by the model, the MDM panel detected 74% of esophageal cancer overall, 74% of EAC, and 78% of ESCC (Fig. 3A). By stage, the panel was able to detect 43% stage I, 64% stage II, 77% stage III, and 92% stage IV disease (Fig. 3A). The rPART model yielded an area under the receiver operating characteristics curve of 0.93

**Table 2.** Clinical characteristics of patients in the plasma study

	Case (N = 85)	Control (N = 98)	P value
Age, median (IQR)	67 (59, 74)	68 (58, 73)	0.888
Male sex, <i>n</i> (%)	77 (90.6%)	84 (85.7%)	0.433
Caucasian race, <i>n</i> (%)	81 (95.3%)	82 (83.7%)	0.155
History of smoking, <i>n</i> (%)	54 (63.5%)	36 (36.7%)	<0.01
Current alcohol use, <i>n</i> (%)	35 (41.2%)	37 (37.8%)	0.192
BMI, median (IQR)	29.8 (25.4–33.1)	27.7 (25.2–30.8)	0.221
EAC, <i>n</i> (%)	76 (89%)		
ESCC, <i>n</i> (%)	9 (11%)		
Stage			
I, <i>n</i> (%)	14 (16%)		
II, <i>n</i> (%)	14(16%)		
III, <i>n</i> (%)	31 (37%)		
IV, <i>n</i> (%)	26 (31%)		


Abbreviations: BMI, body mass index; CI, confidence interval; IQR, interquartile range.

(95% CI, 0.89–0.96) on best-fit and 0.81 (95% CI, 0.75–0.88) on cross-validation (Fig. 3B).

Clinical covariates (age, sex, smoking, and weight) had no significant effect on the performance of our panel of biomarkers (Table 3).


**Site-specificity of MDMs for esophageal cancers**

Using microarray data from TCGA, 7 of our 12 markers were successfully mapped. Using this information, we were able to compare the methylation levels of these MDMs in tissue from EAC and ESCC, versus other common cancer types (Supplementary Fig. S2). Supplementary Table S2 shows pair-wise comparisons of methylation between esophageal cancer with each of the other cancer types. *TBX15* was the only marker from the 5-MDM panel



**Figure 2.** β-Actin corrected methylation intensity in DNA markers assayed from plasma. Each column on the x-axis represents an individual patient's sample; each row on the y-axis represents the methylation-specific PCR product by QuARTS assay for each marker on a logarithmic scale. A 91% specificity threshold was selected by the rPart model. Increasing yellow-red color spectrum reflects increasing methylation intensity above the 91st percentile value of the control patient samples.

Downloaded from http://aacrjournals.org/clincancerres/article-pdf/25/24/7396/2056799/7396.pdf by guest on 13 January 2025



**Figure 3.**

**A**, At a specificity cut-off of 91% by the model, the 5-marker panel was able to detect 43% stage I, 64% stage II, 77% stage III, and 92% stage IV disease. The marker panel detected 74% of esophageal cancer overall, 74% of esophageal adenocarcinoma, and 78% of esophageal squamous cell carcinoma. Confidence intervals are shown. **B**, Overall discrimination of esophageal cancer by MDM panel assayed from plasma: AUCs for best-fit and cross-validation analyses (and 95% confidence intervals) are shown.

that mapped to the TCGA data. *TBX15* methylation ratios were significantly higher for both EAC and ESCC compared with all other cancers in the comparison, except colorectal cancer. For *DMRTA2*, methylation ratios were significantly higher for EAC and ESCC combined than for all other cancers.

## Discussion

Current strategies for early detection and prevention of esophageal cancer rely on demographic stratification followed by

**Table 3.** Effect of clinical covariates in plasma study on AUC of the 5-marker panel, AUCs and 95% confidence intervals are shown

	No	Yes	P value
Age >65	0.95 (0.9–1)	0.91 (0.86–0.96)	0.30
Male	0.97 (0.92–1)	0.92 (0.88–0.96)	0.12
Tobacco Use	0.92 (0.87–0.97)	0.92 (0.86–0.98)	0.97
Obese	0.93 (0.87–0.98)	0.92 (0.87–0.98)	0.92

endoscopy in high-risk individuals. However, this approach is cost-prohibitive and insensitive in the general population, as over 80% of EAC is diagnosed *de novo* (35). An inexpensive, noninvasive, and standardized screening tool such as a blood test that could be applied to a wider population at risk for esophageal cancer would be transformative.

One means by which this could be achieved is through identification of highly discriminant markers, such as MDMs, for esophageal cancer that could be applied to liquid biopsies. Ours was the first comprehensive whole-methylome discovery effort on differential methylation across both ESCC and EAC. In the first phase of our study, thousands of novel DMRs for esophageal cancer were discovered through whole-methylome sequencing on DNA from frozen tissue. The top performing DMRs were then validated technically using a more clinically applicable platform, MSP. Carrying forward, the DMRs were biologically validated in independent FFPE tissue, which confirmed high discrimination. Through this rigorous filtering and independent validation process, DMRs were carefully selected as candidate MDMs with minimal risk of false discovery.

The 12 MDMs with highest discrimination in tissue were subsequently tested in a large, independent plasma study of esophageal cancer case patients and normal controls. Although the purpose of this investigation was to identify MDMs based on their discrimination for esophageal cancer and not to evaluate their function, it is noteworthy that each of the top candidate MDMs has biological associations with tumorigenesis (cell cycle, signaling, and transcriptional regulation), which lends confidence that our discovery process led to *bona fide* epigenetic biomarker candidates (Supplementary Table S1). Similar to reports in other cancers, we have observed that these MDMs alter gene expression through both up- and downregulation in esophageal cancer. Although DNA methylation canonically has been thought to silence gene expression through hypermethylation in promoter regions, more recent studies have found that methylation can occur anywhere in the genome; regulation of transcription can even be influenced by remote enhancers or repressors that only contact a gene via chromatin folding (36–39).

In plasma, a panel of 5 MDMs accurately detected esophageal cancer, demonstrating feasibility for a noninvasive MDM plasma assay for detection of esophageal cancer. Unlike earlier studies on biomarkers for esophageal cancer, updated clinical staging was determined for all cases according to AJCC 8th edition, and all stages were included in our study (25). Although our study was not designed to study the effect of MDMs on prognosis, our markers are in fact highly correlated with stage of disease, which predicts survival. MDMs may also be potentially useful as markers of residual disease or cancer recurrence following surgery for curative intent. Furthermore, MDMs are broadly informative and can be site specific (40).

Not only could an MDM panel be used to target those at risk for esophageal cancer on the basis of Barrett's esophagus or endemic ESCC, it may potentially be used for screening in the general population as part of a universal cancer screening tool. By aggregating the prevalence of multiple cancers, such a universal tool could revolutionize our approach to screening less-prevalent cancers, and prediction of tumor site can efficiently direct the evaluation of test-positive patients (40, 41).

Optimal site prediction for EAC and ESCC was not the primary goal of this study. It was anticipated from prior experiments that markers selected for esophageal cancer detection in this study would be methylated in other gastrointestinal cancers (23, 42). It is also known that there are additional MDMs that readily distinguish between foregut and hindgut neoplasms in tissue studies, plasma assays, and in stool assays (22, 43). In TCGA data examined in the present study, the MDMs we interrogated in plasma are methylated in other gastrointestinal cancers, most commonly colorectal cancer. However, *DMRTA2* showed significantly greater hypermethylation in esophageal cancers relative to all other aerodigestive cancers studied. It is therefore anticipated that a panel of MDMs can be further optimized for site-specific cancer detection, as is the focus of our ongoing research.

Although the MDM panel was less sensitive for stage I disease (43%; 95% CI, 18–71%), where detection would be the most impactful, it is worth noting that the sensitivity for other blood-based tests, and even endoscopy has not been specifically reported for stage I disease. In addition, our samples were collected and processed on generic platforms that were developed before recent advances in nucleic acid preservation technology, and therefore not optimized for recovery of cell-free tumor DNA. This could have led to DNA degradation, decreased recovery of markers, and, accordingly, an artefactual reduction in sensitivity. Newer nucleic acid preservation media are now known to prevent degradation and improve recovery of target DNA while also reducing plasma contamination from leukocyte lysis during processing, transport, and storage (44). Sample collection and processing optimization will therefore be a critical goal of future prospective studies.

Although our study suggests the feasibility of a plasma-based test for detecting esophageal cancer, there are several important limitations to consider. We were constrained by our relatively small regional population for ESCC cases. To maximize statistical power, ESCC and EAC were incorporated in the same cohort for the training of a 5-marker model in plasma. This was supported by the significant overlap of MDMs observed between the 2 cancers. All MDMs discovered in ESCC performed well in EAC (Table 1). Because our plasma cohort consisted mostly of EAC, MDMs specific for EAC were chosen for biological validation. Mapping 7 of our MDMs to methylation data available in TCGA (Supplementary Fig. S2) showed that similar to our results, all MDMs except those chosen specifically for EAC were similarly methylated in both cancers. Therefore, although we agree that EAC and ESCC are biologically distinct cancers, there is significant overlap in their methylation signatures which may reflect that MDMs are more representative of the host organ epithelium, rather than the underlying tumor biology. Similar patterns have also been observed in liver tumors arising in different underlying chronic disease states (45). These epigenetic similarities allowed us to select representative candidates for both cancer types in combination.

Our plasma cohort did not include patients with Barrett's esophagus as a disease-control group. This is because detection of MDMs in plasma appears to be dependent on invasive characteristics of tumors such as angiogenesis, tumor necrosis, and metastatic potential; thus, we do not anticipate false positive MDM results in plasma assay due to underlying nondysplastic Barrett's esophagus (46). Support of this hypothesis comes from studies of methylated *SEPTIN9* in plasma for detection of colo-

rectal neoplasms; the adenoma detection rate by this plasma marker fell at/below the false positive threshold, indicating that invasive cancer is most likely required to introduce cancer-specific methylated DNA into plasma (47). The markers we have piloted in plasma for EAC do include recently described markers of Barrett's esophagus (48).

Although the population of the present study is 80% white, we have tested the ESCC markers in larger independent cohorts of patient tissues from geographically and ethnically distinct regions (China, Iran, and India; ref. 49). DNA samples obtained from minimally invasive balloon cytology scrapings have also confirmed that MDMs for ESCC discovered in this study are represented in high grade dysplasia and ESCC in patients from central China (50).

Finally, the cross-validation AUC, while strong (0.81) was lower than that of the best fit (0.93). A larger clinical study is needed to measure MDM performance with greater precision. Prospective collection will also permit optimization and standardization of storage and processing conditions, referent to older EDTA collected archives, anticipated to improve recovery of MDM targets in early stage cancers.

In conclusion, our study demonstrates early feasibility for the noninvasive detection of esophageal cancer by assay of informative MDMs in plasma. This approach has potential value in esophageal cancer screening, prognosis, and surveillance applications. Further investigation is indicated to optimize sample collection and processing, assay configuration, and MDM panel selection, as well as to validate findings in larger prospective clinical studies.

#### Disclosure of Potential Conflicts of Interest

W.R. Taylor reports receiving commercial research grants from and holds ownership interest (including patents) in Exact Sciences. D.W. Mahoney holds ownership interest (including patents) in Exact Sciences. T.C. Yab holds ownership interest (including patents) in Exact Sciences. G.P. Lidgard is an employee/paid consultant for Exact Sciences. H.T. Allawi is an employee/paid consultant for Exact Sciences. P.G. Iyer reports receiving commercial research grants from Exact Sciences; reports receiving other commercial research support from Pentax Medical; and is an unpaid consultant/advisory board member for Medtronic and Symple Surgical. D.A. Katzka reports receiving speakers bureau honoraria from Celgene. D.A. Ahlquist reports receiving commercial research grants from, holds ownership interest (including patents) in, and is an unpaid consultant/advisory board member for Exact Sciences. J.B. Kisiel reports receiving other commercial research support from and holds ownership interest (including patents) in Exact Sciences. No potential conflicts of interest were disclosed by the other authors.

#### Authors' Contributions

**Conception and design:** Y. Qin, C.W. Wu, D.W. Mahoney, T.C. Yab, G.P. Lidgard, H.T. Allawi, N.S. Buttar, D.A. Katzka, D.A. Ahlquist, J.B. Kisiel  
**Development of methodology:** Y. Qin, C.W. Wu, W.R. Taylor, D.W. Mahoney, G.P. Lidgard, H.T. Allawi, D.A. Ahlquist, J.B. Kisiel  
**Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.):** Y. Qin, C.W. Wu, W.R. Taylor, N.S. Buttar, T.C. Smyrk, J.B. Kisiel  
**Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis):** Y. Qin, C.W. Wu, W.R. Taylor, K.N. Burger, D.W. Mahoney, Z. Sun, T.C. Yab, H.T. Allawi, D.A. Katzka, J.B. Kisiel  
**Writing, review, and/or revision of the manuscript:** Y. Qin, C.W. Wu, W.R. Taylor, T. Sawas, D.W. Mahoney, Z. Sun, T.C. Yab, H.T. Allawi, N.S. Buttar, T.C. Smyrk, P.G. Iyer, D.A. Katzka, D.A. Ahlquist, J.B. Kisiel  
**Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases):** Y. Qin, C.W. Wu, J.B. Kisiel  
**Study supervision:** Y. Qin, C.W. Wu, T.C. Yab, J.B. Kisiel  
**Other (obtained funding):** J.B. Kisiel

## Acknowledgments

This work was supported by the Maxine and Jack Zarrow Family Foundation of Tulsa Oklahoma (to JBK), R37CA214679 (to JBK), a gift from Eddie Gong and Dana Clay (to DAA), the Carol M. Gattion endowment for Digestive Diseases Research (to DAA). This publication was also made possible by UL1 TR002377 from the National Center for Advancing Translational Sciences (NCATS). RRBS sequencing costs and QuARTS assays were provided by Exact Sciences (Madison WI).

## References

- Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA Cancer J Clin* 2011;61:69–90.
- Thrift AP. The epidemic of oesophageal carcinoma: where are we now? *Cancer Epidemiol* 2016;41:88–95.
- Arnal MJD, Arenas ÁF, Arbeloa ÁL. Esophageal cancer: Risk factors, screening and endoscopic treatment in Western and Eastern countries. *World J Gastroenterol* 2015;21:7933.
- Codipilly DC, Qin Y, Dawsey SM, Kisiel J, Topazian M, Ahlquist D, et al. Screening for esophageal squamous cell carcinoma: recent advances. *Gastrointest Endosc* 2018;88:413–26.
- Hiripi E, Jansen L, Gondos A, Emrich K, Holleczeck B, Katalinic A, et al. Survival of stomach and esophagus cancer patients in Germany in the early 21st century. *Acta Oncol (Madr)* 2012;51:906–14.
- Buttar NS, Wang KK, Lutzke LS, Krishnadath KK, Anderson MA. Combined endoscopic mucosal resection and photodynamic therapy for esophageal neoplasia within Barrett's esophagus. *Gastrointest Endosc* 2001;54:682–8.
- Wang G-Q, Jiao G-G, Chang F-B, Fang W-H, Song J-X, Lu N, et al. Long-term results of operation for 420 patients with early squamous cell esophageal carcinoma discovered by screening. *Ann Thorac Surg* 2004;77:1740–4.
- Wei W-Q, Chen Z-F, He Y-T, Feng H, Hou J, Lin D-M, et al. Long-term follow-up of a community assignment, one-time endoscopic screening study of esophageal cancer in China. *J Clin Oncol* 2015;33:1951.
- Yang J, Wei W-Q, Niu J, Liu Z-C, Yang C-X, Qiao Y-L. Cost-benefit analysis of esophageal cancer endoscopic screening in high-risk areas of China. *World J Gastroenterol* 2012;18:2493.
- Xu YW, Peng YH, Chen B, Wu ZY, Wu JY, Shen JH, et al. Autoantibodies as potential biomarkers for the early detection of esophageal squamous cell carcinoma. *Am J Gastroenterol* 2014;109:36–45.
- Wang S, Du H, Li G. Significant prognostic value of circulating tumor cells in esophageal cancer patients: a meta-analysis. *Oncotarget* 2017;8:15815–26.
- Liu F, Tian T, Xia LL, Ding Y, Cormier RT, He Y. Circulating miRNAs as novel potential biomarkers for esophageal squamous cell carcinoma diagnosis: a meta-analysis update. *Dis Esophagus* 2017;30:1–9.
- Moinova HR, LaFramboise T, Lutterbaugh JD, Chandar AK, Dumot J, Faulx A, et al. Identifying DNA methylation biomarkers for non-endoscopic detection of Barrett's esophagus. *Sci Transl Med* 2018;10. pii: eaao5848.
- Chettouh H, Mowforth O, Galeano-Dalmau N, Bezawada N, Ross-Innes C, MacRae S, et al. Methylation panel is a diagnostic biomarker for Barrett's esophagus in endoscopic biopsies and non-endoscopic cytology specimens. *Gut* 2017;67:1942–49.
- Dickinson BT, Kisiel J, Ahlquist DA, Grady WM. Molecular markers for colorectal cancer screening. *Gut* 2015;64:1485–94.
- Imperiale TF, Ransohoff DF, Itzkowitz SH, Levin TR, Lavin P, Lidgard GP, et al. Multitarget stool DNA testing for colorectal-cancer screening. *N Engl J Med* 2014;370:1287–97.
- Kisiel JB, Klepp P, Allawi HT, Taylor WR, Giakoumopoulos M, Sander T, et al. Analysis of DNA methylation at specific loci in stool samples detects colorectal cancer and high-grade dysplasia in patients with inflammatory bowel disease. *Clin Gastroenterol Hepatol* 2018;17:914–21.e5.
- Kisiel JB, Raimondo M, Taylor W, Yab TC, Mahoney DW, Sun Z, et al. New DNA methylation markers for pancreatic cancer: discovery, tissue validation, and pilot testing in pancreatic juice. *Clin Cancer Res* 2015;21:4473–81.
- Kisiel JB, Dukek BA, Kanipakam RV, Ghaz HM, Yab TC, Berger CK, et al. Hepatocellular carcinoma detection by plasma methylated DNA: discovery, phase I pilot, and phase II clinical validation. *Hepatology* 2018;69:1180–92.
- Allawi HT, Giakoumopoulos M, Flietner E, Oliphant A, Volkman C, Aizenstein B, et al. Detection of lung cancer by assay of novel methylated DNA markers in plasma. *AACR*; 2017.
- Ghoz HM, Aboelsoud MM, Yab TC, Berger CK, Taylor WR, Cao X, et al. DNA methylation markers for detection of extrahepatic cholangiocarcinoma: discovery, tissue validation, and pilot testing in biliary brush samples. *Hepatology* 2015;62:295A.
- Kisiel JB, Taylor WR, Yab TC, Ghaz HM, Foote PH, Devens ME, et al. Accurate site prediction of gastrointestinal cancer by novel methylated DNA markers: Discovery & Validation. *Cancer Res* 2015;75:S-15–S-4252.
- Anderson BW, Suh Y-S, Choi B, Lee H-J, Yab TC, Taylor W, et al. Detection of gastric cancer with novel methylated DNA markers: discovery, tissue validation, and pilot testing in plasma. *Clin Cancer Res* 2018;24:5724–34.
- Li L, Choi J-Y, Lee K-M, Sung H, Park SK, Oze I, et al. DNA methylation in peripheral blood: a potential biomarker for cancer molecular epidemiology. *J Epidemiol* 2012;22:384–94.
- Rice TW, Ishwaran H, Ferguson MK, Blackstone EH, Goldstraw P. Cancer of the esophagus and esophagogastric junction: an eighth edition staging primer. *J Thorac Oncol* 2017;12:36–42.
- Kisiel JB, Yab TC, Taylor WR, Chari ST, Petersen GM, Mahoney DW, et al. Stool DNA testing for the detection of pancreatic cancer: assessment of methylation marker candidates. *Cancer* 2012;118:2623–31.
- Xi Y, Li W. BSMAP: whole genome bisulfite sequence MAPPING program. *BMC Bioinformatics* 2009;10:232.
- Goldman M, Craft B, Brooks A, Zhu J, Haussler D. The UCSC Xena Platform for cancer genomics data visualization and interpretation. *bioRxiv* 326470; doi: <https://doi.org/10.1101/326470>.
- Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 2014;15:550.
- TCGA Network. Integrated genomic characterization of oesophageal carcinoma. *Nature* 2017;541:169–75.
- TCGA Network. Comprehensive molecular characterization of human colon and rectal cancer. *Nature* 2012;487:330–37.
- TCGA Network. Comprehensive and integrative genomic characterization of hepatocellular carcinoma. *Cell* 2017;169:1327–41.e23.
- TCGA Network. Comprehensive molecular profiling of lung adenocarcinoma. *Nature* 2014;511:540–550.
- TCGA Network. Comprehensive genomic characterization of squamous cell lung cancers. *Nature* 2012;489:519–525.
- Visrodia K, Singh S, Krishnamoorthi R, Ahlquist D, Wang K, Iyer PG, et al. Systematic review with metaanalysis: prevalent vs. incident oesophageal adenocarcinoma and highgrade dysplasia in Barrett's oesophagus. *Aliment Pharmacol Ther* 2016;44:775–84.
- Lokk K, Modhukur V, Rajashekar B, Martens K, Magi R, Kolde R, et al. DNA methylome profiling of human tissues identifies global and tissue-specific methylation patterns. *Genome Biol* 2014;15:r54.
- Maunakea AK, Nagarajan RP, Bilenky M, Ballinger TJ, D'Souza C, Fouse SD, et al. Conserved role of intragenic DNA methylation in regulating alternative promoters. *Nature* 2010;466:253–7.
- Ball MP, Li JB, Gao Y, Lee JH, LeProust EM, Park IH, et al. Targeted and genome-scale strategies reveal gene-body methylation signatures in human cells. *Nat Biotechnol* 2009;27:361–8.
- Lorincz MC, Dickerson DR, Schmitt M, Groudine M. Intragenic DNA methylation alters chromatin structure and elongation efficiency in mammalian cells. *Nat Struct Mol Biol* 2004;11:1068–75.



40. Cohen JD, Li L, Wang Y, Thoburn C, Afsari B, Danilova L, et al. Detection and localization of surgically resectable cancers with a multi-analyte blood test. *Science* 2018;eaar3247.
41. Ahlquist DA. Universal cancer screening: revolutionary, rational, and realizable. *NPJ Precis Oncol* 2018;2:23.
42. Raimondo M, Yab TC, Mahoney DW, Taylor WR, Anderson KS, Ahlquist DA. 487 methylated DNA markers in pancreatic juice discriminate pancreatic cancer from chronic pancreatitis and normal controls. *Gastroenterology* 2013;144:S-90.
43. Kisiel JB, Wu CW, Taylor WR, Allawi HT, Yab TC, Foote PH, et al. 393—multi-site gastrointestinal cancer detection by stool DNA. *Gastroenterology* 2018;154:S-95.
44. Pottekat A, Allawi HT, Boragine GT, Kaiser MW, Sander T, Krueger C, et al. A comprehensive assessment of the impact of preanalytical variables on cell free DNA and circulating tumor cells in blood. *Cancer Res* 2018;78:S-13-S-4598.
45. Hlady RA, Tiedemann RL, Puszyk W, Zendejas I, Roberts LR, Choi JH, et al. Epigenetic signatures of alcohol abuse and hepatitis infection during human hepatocarcinogenesis. *Oncotarget* 2014;5:9425-43.
46. Ahlquist DA, Zou H, Domanico M, Mahoney DW, Yab TC, Taylor WR, et al. Next-generation stool DNA test accurately detects colorectal cancer and large adenomas. *Gastroenterology* 2012;142:248-56.
47. Ahlquist DA, Taylor WR, Mahoney DW, Zou H, Domanico M, Thibodeau SN, et al. The stool DNA test is more accurate than the plasma septin 9 test in detecting colorectal neoplasia. *Clin Gastroenterol Hepatol* 2012;10:272-7.e1.
48. Iyer PG, Taylor WR, Johnson ML, Lansing RL, Maixner KA, Yab TC, et al. Highly discriminant methylated DNA markers for the non-endoscopic detection of Barrett's esophagus. *Am J Gastroenterol* 2018;1.
49. Iyer PG, Buglioni A, Cao X, Foote PH, Yab TC, Taylor WR, et al. Mo1748 - Validation of novel methylated DNA markers for the detection of esophageal squamous cell carcinoma and dysplasia: multi-national tissue study. *Gastroenterology* 2018;154:S-794-S-5.
50. Qin Y, Taylor WR, Foote PH, McGlinch M, Cao X, Wu CW, et al. Methylated DNA markers detect esophageal squamous dysplasia in mucosal tissue and when sampled by nonendoscopic esophageal balloons: an exploratory study. *Gastroenterology* 2019;156:S-227-S-8.